



CHEMISTRY

BOOKS - GRB CHEMISTRY (HINGLISH)

PERIODIC TABLE

A. Development of Periodic Table, Period, Group and Block

1. According to Moseley, a straight line graph is obtained on plotting:
- A. the frequencies of characteristic line graph
 - B. the square of the frequencies of characteristic X-rays of elements against their atomic numbers
 - C. the square root of the frequencies of characteristic X-rays of elements against their atomic numbers.

D. the reciprocal of the frequencies of characteristic X-rays of elements against their atomic numbers

Answer: C



Watch Video Solution

2. The atomic volume was chosen as the basis of periodic classification of elements by:

A. Niel Bohr

B. Mendeleev

C. Lothar Meyer

D. Newland

Answer: C



Watch Video Solution

3. Element in periodix table with electronic configuration as $[Ar]^{18}3d^54s^1$ is placed in:

- A. IA,s-block
- B. VIA,s-block
- C. VIB,s-block
- D. VIB,d-block

Answer: D



Watch Video Solution

4. In a given energy level, the order of penetration effect of different orbitals is

- A. $f < d < p < s$
- B. $s = p = d = f$
- C. $s < p < d < f$

$$D. p > s > d > f$$

Answer: A



Watch Video Solution

5. Atomic number of Ag is 47. in the same group, the atomic number of elements placed above and below Ag will be:

A. 37,67

B. 29,79

C. 39,69

D. 18,28

Answer: B



Watch Video Solution

6. In the periodic table, where are non-metals located?

- A. Between groups IIA and IIIA
- B. On the lower left hand side
- C. On the upper left hand side
- D. On the upper right hand side

Answer: D



Watch Video Solution

7. Which of the following statements is wrong for the transition elements?

- A. Transition elements are placed from 3rd to 6th period
- B. Last electron enters in $(n - d)d$ subshell
- C. Exhibits variable valency
- D. General electronic configuration is $(n - 1)d^{1-10}ns^{1-2}$.

Answer: A



Watch Video Solution

8. The first element of a group in many ways differ from the heavier members of group. This is due to a)the small size b)the high electronegativity c)high ionisation potential d)the unavailability of d-orbitals

A. the small size

B. the high electronegativity and high ionisation potential

C. the unavailability of d-orbitals

D. all of the above

Answer: D



Watch Video Solution

9. The statement that is not correct for the periodic classification of elements is : a)The properties of elements are the periodic functions of their atomic numbers. b)non-metallic elements are lesser in number than metallic elements. c)the first ionisation energies of elements along a period do not vary in a regular manner with increase in atomic number. d)for transition elements the d – subshells are filled with electrons monotonically with increase in atomic number.

A. the properties of elements are the periodic functions of their atomic numbers

B. non-metallic elements are lesser in number than metallic elements

C. the third period contains 8 elements and not 18 as 4th period contains

D. for transition elements the d -subshells are filled with electrons monotonically with increase in atomic number

Answer: D



Watch Video Solution

10. Which of the following statement is not correct regarding hydrogen atom?

- A. It resembles halogens in some properties
- B. It resembles alkali metals in some properties
- C. It can be placed in 7th group of periodic table
- D. It cannot be placed in first group of periodic table

Answer: D



Watch Video Solution

11. According to modern periodic law the elements with similar chemical and physical properties repeat at regular intervals when the elements are arranged in order of :

- A. decreasing atomic number

- B. increasing atomic weight
- C. increasing atomic number
- D. decreasing atomic weight

Answer: C



Watch Video Solution

12. Which one of the following statements related to the modern periodic table is incorrect?

- A. The p-block has 6 columns, because a maximum of 6 electrons can occupy all the orbitals in a p-subshell
- B. The d-block has 8 columns, because a maximum of 8 electrons can occupy all the orbitals in a d-subshell
- C. Each block contains a number of columns equal to the number of electrons that can occupy that subshell.

D. The block indicates value of azimuthal quantum number (l) for the last subshell that received electrons in building up the electronic configuration.

Answer: B



Watch Video Solution

13. Which set does not show correct matching ?

A. $Sc^{3+} [Ne]3s^23p^6$ zero group

B. $Fe^{2+} [Ar]3d^6$ 8th group

C. $Cr [Ar]3d^54s^1$ 6th group

D. All of the above

Answer: A



Watch Video Solution

14. The atomic numbers of the metallic and non-metallic elements which are liquid at room temperature respectively are:

A. 55,87

B. 33,87

C. 35,80

D. 80,35

Answer: D



Watch Video Solution

15. Total number of elements in 6th period are:

A. 8

B. 18

C. 32

D. 28

Answer: C



Watch Video Solution

16. Atomic number of 15, 33, 51 represents the following family:

- A. carbon family
- B. nitrogen family
- C. oxygen family
- D. none of these

Answer: B



Watch Video Solution

17. The element with atomic number $Z = 118$ will be categorised as a:

- A. noble gas

B. transition metal

C. alkali metal

D. alkaline earth metal

Answer: A



Watch Video Solution

18. The places that were left empty by Mendeleev were for:

A. aluminium and silicon

B. gallium and germanium

C. arsenic and antimony

D. molybdenum and tungsten

Answer: B



Watch Video Solution

19. In which group all the elements do not have same number of valence electrons?

- A. zero
- B. second
- C. first
- D. seventh

Answer: A



Watch Video Solution

20. The elements which exhibit both vertical and horizontal similarities are:

- A. inert gas elements
- B. representative elements
- C. transition element

D. rare earth element

Answer: C



Watch Video Solution

21. Which of the following is the atomic number of a metal?

A. 35

B. 34

C. 36

D. 38

Answer: D



Watch Video Solution

22. In which element shielding effect is not possible ?

A. H

B. Be

C. B

D. N

Answer: A



Watch Video Solution

23. Which of the following is true about the element $_{33}\text{As}$?

A. It is 5th period element

B. It is p-block element

C. It belongs to 16th group

D. It is the member of VIA group

Answer: B



Watch Video Solution

24. Chalcogens are elements of:

- A. group 16
- B. p-block
- C. ns^2np^4 configuration
- D. all are correct

Answer: D



Watch Video Solution

25. Electronic configuration of M^{3+} is $[Ar]3d^{10}4s^2$, it belongs to

- A. s-block
- B. p-block
- C. d-block

D. f-block

Answer: B



Watch Video Solution

26. Of the following pairs, the one containing examples of metalloid elements in the periodic table is:

A. Na and K

B. F and Cl

C. Cu and Ag

D. As and Sb

Answer: D



Watch Video Solution

27. Which group of elements is analogous to the lanthanides?

- A. Halides
- B. Actinides
- C. Chalcogenides
- D. Borides

Answer: B



Watch Video Solution

28. Principle of modern periodic table was given by:


- A. Moseley
- B. Mendeleev
- C. Doberenier
- D. Newland

Answer: A



Watch Video Solution

29. In the Lotter Meyer graph, A,B,C,D and E elements are :



A. halogens

B. alkaline earth metals

C. alkali metals

D. transition metals

Answer: C



Watch Video Solution

30. Most of the d block elements are known as:

- A. alkali metals
- B. alkaline earth metals
- C. innertransition elements
- D. transition elements

Answer: D



Watch Video Solution

31. Which of the following represents the electronic configuration of a transition element?

- A. $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2$
- B. $1s^2, 2s^2, 2p^6, 3s^2, 3p^6 3d^2, 4s^2$
- C. $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^2$

D. $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^1$

Answer: B



Watch Video Solution

32. As applied to periodic table, which of the following sets include only magic number?

A. 2,8,8,18,19,32

B. 2,8,18,32

C. 8,8,18,18

D. 2,18,18,32

Answer: A



Watch Video Solution

33. The number of elements which should be theoretically present in 8^{th} period of the modern long form of periodic table, is

A. 32

B. 40

C. 50

D. 48

Answer: C



Watch Video Solution

34. If an orbital can have maximum 4 electrons then, how many elements can be present in 8th period?

A. 100

B. 75

C. 128

D. 64

Answer: A



Watch Video Solution

35. which of the following number for the element Unbiunnum is

A. 120

B. 121

C. 112

D. 122

Answer: B



Watch Video Solution

36. Elements A,B,C,D and E have following electronic configuration.

(A) $1s^2, 2s^2 2p^1$, (B) $1s^2, 2s^2 2p^6, 3s^2 3p^1$

(C) $1s^2, 2s^2 2p^6, 3s^2 3p^3$ (D) $1s^2, 2s^2 2p^6, 3s^2 3p^5$

(E) $1s^2, 2s^2 2p^6, 3s^2 3p^6$

Which among the belongs to same group in the periodic table. ?

A. A and C

B. A and D

C. A and E

D. A and B

Answer: D



Watch Video Solution

37. If Aufbau rule is not followed, K-19 will be placed in

A. s-block

B. p-block

C. d-block

D. f-block

Answer: C



Watch Video Solution

38. In the Lothar Meyer, which of the following option is incorrect.

A. Alkali metals occupied peak position at curve

B. Halogens occupied ascending position at curve

C. Alkaline earth elements descendings position at curve

D. Alkali metals are in the lower curve of graph

Answer: D



Watch Video Solution

39. If A, B and C are the three elements of Dobereiner's Triad and atomic weight of A and B are 7 and 15 respectively then the atomic weight of C is:

- A. 1
- B. 11
- C. 23
- D. 25

Answer: C



Watch Video Solution

40. According to the Lothar Meyer's curve which of the following statements is incorrect?

- A. The element having similar properties will occupy the same position in the curve
- B. Alkaline earth metals are at the peaks of the curve

C. Halogens are at ascending part of the curve

D. The atomic volume of the elements in a period initially decreases
and then increases

Answer: B



Watch Video Solution

41. If the atomic number of an element is 33, it will be placed in the
periodic table in the ____

A. first group

B. third group

C. fifth group

D. seventh group

Answer: C



Watch Video Solution

42. Eka-aluminium and Eka -silicon are known as :

- A. gallium and germanium
- B. aluminium and silicon
- C. iron and sulphur
- D. proton and silicon

Answer: A



Watch Video Solution

43. The electronic configuration of gadolinium (Atomic number 64) is

- A. $[Xe]4f^35d^56s^2$
- B. $[Xe]4f^75d^26s^1$
- C. $[Xe]4f^75d^16s^2$
- D. $[Xe]4f^85d^66s^2$

Answer: C



Watch Video Solution

44. The statement that is not correct for periodic classification of elements is

- A. The properties of elements are periodic function of their atomic numbers
- B. Non metallic elements are less in number than metallic elements
- C. For transition elements, the 3d-orbitals are filled with electrons after 3p-orbitals and before 4s-orbitals
- D. The first ionisation enthalpies of elements generally increase with increase in atomic number as we go along a period..

Answer: C



Watch Video Solution

45. Which of the following is not an actinoid?

- A. Curium ($Z = 96$)
- B. Californium ($Z = 98$)
- C. Uranium ($Z = 92$)
- D. Terbium ($Z = 65$)

Answer: D



Watch Video Solution

46. Which of following is not correctly matched?

- A. d-block element : electronic configuration is $ns^{0-2}(n-1)d^{1-10}$
- B. p-block element: electronic configuration is $ns^{1-2}np^{1-6}$
- C. s-block element: electronic configuration is ns^{1-2}
- D. Ce: f-block's first member

Answer: B



Watch Video Solution

47. Newland's law of octave applies to which of the following set elements ?

A. Na,K, Rb

B. F,Cl,Br

C. Be,Mg,Ca

D. B,Al,Ga

Answer: C



Watch Video Solution

48. Which are correct match?

(i) Eka silicon-Be (ii) Eka aluminium -Ga

(iii) Eka manganese-Tc (iv) Eka scandium-B

- A. (ii) and (iii)
- B. (i),(ii) and (iv)
- C. (i) and (iv)
- D. All of these

Answer: A



Watch Video Solution

49. In Lothar Meyer plot, the peaks are occupied by

- A. alkali metals
- B. highly electropositive elements
- C. elements having large atomic volume
- D. all of the above

Answer: D



[Watch Video Solution](#)

50. In Mendeleev's modified periodic table, there are_____vertical column called groups.

- A. 8
- B. 9
- C. 18
- D. 12

Answer: B



[Watch Video Solution](#)

51. The elements of group 1, 2, 13, 14, 15, 16, 17, 18 are collectively called

- A. noble gases
- B. representative or normal elements

C. transition elements

D. inner transition elements

Answer: B



Watch Video Solution

52. Which of the following is known as the bridge element of 2nd group in Mendeleev's table?

A. Zn

B. Sr

C. Mg

D. Hg

Answer: C



Watch Video Solution

53. Which pair is different from the others?

A. $Li - Mg$

B. $B - Si$

C. $Be - Al$

D. $Li - Na$

Answer: D



Watch Video Solution

54. Atomic number of 15, 33, 51 represents the following family:

A. carbon family

B. nitrogen family

C. oxygen family

D. none of these

Answer: B



Watch Video Solution

55. Three elements X,Y,Z are following Dobereiner's Triad rule. If the atomic weight of X and Y are 10 and 26 respectively, then atomic weight of Z will be:

A. 34

B. 40

C. 42

D. 18

Answer: C



Watch Video Solution

56. Which of the following numbers are called magic numbers?

A. 2,8,20,28,82,126

B. 2,8,8,18,18,32

C. 2,2,8,8,18,32

D. 2,8,18,18,32,32

Answer: B



Watch Video Solution

57. In the general electronic configuration :

$(n - 2)f^{1-14}(n - 1)d^{0-1}ns^2$, if the value of $n = 7$, then the configuration will be of :

A. lanthanides

B. actinides

C. transition elements

D. none of these

Answer: B



Watch Video Solution

58. In lanthanum ($Z = 57$), the 5th electron enters in a:

A. 6p orbital

B. 5d orbital

C. 6s orbital

D. 4f orbital

Answer: B



View Text Solution

59. Atomic number is the basis of :

(p) Lothar Meyer curve

(q) Newland octave rule

(r) Modern periodic table

(s) Dobereiner triad rule

(t) Long form of periodic table

A. (p),(q),(s)

B. (r),(t)

C. (p),(s)

D. (p),(r),(t)

Answer: B



Watch Video Solution

60. If each orbital can hold a maximum of two electrons, the number of elements in 9th period of periodic table (long form) are:

A. 48

B. 162

C. 50

D. 75

Answer: D



Watch Video Solution

61. The electronic configuration of an element is $1s^2 2s^2 2p^6, 3s^2 3p^5$. The atomic number of element present just below the above element in periodic table is:

A. 34

B. 35

C. 36

D. 30

Answer: B



Watch Video Solution

62. The number of elements known at that time when Mendeleev arranged them in the periodic table was:

A. 63

B. 60

C. 71

D. 65

Answer: A



Watch Video Solution

63. Find the odd element from the given electronic configuration.

A. $1s^2 2s^2 2p^6 3s^1$

B. $1s^2 2s^1$

C. $1s^2 2s^2 2p^6 3s^2 3p^1$

D. $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$

Answer: C



Watch Video Solution

64. Which of following is not correctly matched?

A. d-block element : electronic configuration is $ns^{0-1}(n-1)d^{1-10}$

B. p-block element : electronic configuration is $ns^{1-2}np^{1-6}$

C. s-block element : electronic configuration is ns^{1-2}

D. Ce: f-block's first member

Answer: B



Watch Video Solution

65. The period number in the long form of the periodic table is equal to

A. magnetic quantum number of any element of the period

- B. atomic number of any element of the period
- C. maximum principal quantum number of any element of the period
- D. maximum azimuthal quantum number of any element of the period

Answer: C



Watch Video Solution

66. The elements in which electrons are progressively filled in 4f-orbitals are called:

- A. actinoids
- B. transition elements
- C. lanthanides
- D. halogens

Answer: C



Watch Video Solution

67. Which of the following relation is correct for boron element? Here P = period number of boron

N = number of valence electrons is boron

A. $(P + 1) > N$

B. $(P + 1) < N$

C. $(P + 1) = N$

D. $P = N$

Answer: C



Watch Video Solution

68. Maximum value of $n + l + m$ for the last unpaired electron in an element present in 17th group and 4th period of periodic table:

A. 4

B. 5

C. 6

D. 7

Answer: C



Watch Video Solution

69. *IUPAC* name for the element with $Z = 117$ is _____ and its symbol is _____.

A. Uuo

B. Uus

C. Uuh

D. Uun

Answer: B



Watch Video Solution

70. Had three electrons been possible to be placed in an orbital, the block of potassium in periodic table would be:

A. s

B. p

C. d

D. f

Answer: B



Watch Video Solution

71. Which element has chemical properties most similar to those of P?

A. N

B. Al

C. S

D. As

Answer: D



Watch Video Solution

72. Which list gives nonmetals that are found in their elemental forms in nature?

- A. neon, phosphorus, fluorine
- B. helium, hydrogen, iodine
- C. nitrogen, oxygen, sulphur
- D. oxygen, chlorine, phosphorus

Answer: C



Watch Video Solution

73. The most abundant noble gas in the atmosphere is

A. *He*

B. *Ne*

C. *Ar*

D. *Rn*

Answer: C



Watch Video Solution

74. In which set are both elements metalloids?

A. Cr and Mo

B. Ge and As

C. Sn and Pb

D. Se and Br

Answer: B



Watch Video Solution

75. Which existing element would the chemistry of element 119 most resemble?

A. $Rn(Z = 86)$

B. $Fr(Z = 87)$

C. $Ra(Z = 88)$

D. $Ac(Z = 89)$

Answer: B



Watch Video Solution

76. Which element has the greatest electrical conductivity?

A. As

B. Ge

C. P

D. Sn

Answer: D



Watch Video Solution

77. Which element is a liquid at $25^{\circ}C$ and 1 atm?

A. Fluorine

B. Chlorine

C. Bromine

D. Iodine

Answer: C



Watch Video Solution

78. The material used in solar cells contains

A. As

B. Ge

C. P

D. Si

Answer: D



Watch Video Solution

79. Which pair of symbols identifies two elements that are metalloids?

A. Si and Ge

B. Mg and Si

C. P and As


D. Ti and V

Answer: A



Watch Video Solution

80. All of the following possess complete d shells except:



Answer: B



Watch Video Solution

81. Which element is a gas at $25^{\circ}C$ and 1 atm pressure?

- A. Chlorine
- B. Phosphorus
- C. Silicon
- D. Sulphur

Answer: A



Watch Video Solution

82. Which element has the greatest electrical conductivity?

- A. Ge
- B. Se
- C. Sn
- D. Te

Answer: C



Watch Video Solution

83. Select the option in which all the information about element ($Z = 113$) are correct:

- | | Block | group | Period | Valence shell configuration |
|----|-------|-------|--------|-----------------------------|
| A. | s | 2 | 7 | $7s^2$ |
| B. | p | 13 | 7 | $7s^2 7p^1$ |
| C. | p | 13 | 6 | $6s^2 6p^1$ |
| D. | d | 12 | 6 | $6s^2$ |

Answer: B



Watch Video Solution

84. An element z has electronic configuration:

$ns^2 np^6 (n-1)d^{10} (n-2)f^{14}$. Which value of n is not possible?

A. 6

B. 3

C. 5

D. Both (b) and (c)

Answer: D



Watch Video Solution

85. Which of the following statements is wrong?

A. No inert gas is present in 7th period

B. 3rd period contains 18 elements

C. 1st period contains two non metals

D. In p-block, metal, non-metal and metalloids are present

Answer: B



Watch Video Solution

86. Which substance is stored in contact with water to prevent it from reacting with air?

- A. Bromine
- B. Lithium
- C. Mercury
- D. Phosphorus

Answer: D



Watch Video Solution

87. Stannic chloride is a molecule in which central metal atom belongs to carbon family. What is the period and atomic number of this metallic element?

- A. 5,50
- B. 4,40

C. 6,60

D. 7,70

Answer: A



Watch Video Solution

88. Which element has an outer electron configuration of s^2p^4 ?

A. Ca

B. Cr

C. Ge

D. Se

Answer: D



Watch Video Solution

89. In the long form of periodic table all the non-metals are placed under

A. s-block

B. p-block

C. d-block

D. f-block

Answer: B



Watch Video Solution

90. Each period in the periodic table starts with a subshell of the new shell and ends with:

A. same subshell

B. different subshell of the same shell

C. the subshell of next higher shell

D. p-subshell of the same new shell

Answer: D



Watch Video Solution

91. In the long form of the periodic table the valence shell electronic configuration of $5s^25p^4$ corresponds to the element present in:

- A. group 16 and period 5
- B. group 17 and period 6
- C. group 17 and period 5
- D. group 16 and period 6

Answer: A



Watch Video Solution

92. Atomic number of the inert gas of 7^{th} period = 118. Which is correct IUPAC name of last element of halogen family in 7th period?

- A. Ununoctium
- B. Ununnilium
- C. Ununennium
- D. Ununseptium

Answer: D



Watch Video Solution

93. The valence shell of the element X contains 1 electron in 5s subshell and below that shell, 4 electrons in 4d subshell. The element belongs to which group (IUPAC) of periodic table?

- A. 4th group
- B. 5th group
- C. 6th group
- D. 7th group

Answer: B



Watch Video Solution

94. According to Moseley, a straight line graph is obtained on plotting:

Where, ν is frequency and Z is atomic number.

A. ν vs Z

B. ν^2 vs Z

C. $\sqrt{\nu}$ vs Z

D. $\frac{1}{\nu}$ vs Z

Answer: C



Watch Video Solution

95. Which pair of atomic numbers represent s-block elements?

A. 3,12

B. 6,12

C. 9,17

D. 7,15

Answer: A



Watch Video Solution

96. Suppose an orbital may accommodate three electrons, then estimate the number of elements in 2nd period:

A. 3

B. 6

C. 8

D. 12

Answer: D



[Watch Video Solution](#)

97. Maximum number of quantum numbers which can be equal for last electron of two elements of same group of periodic table (group does not contain any element having exceptional electronic configuration):

A. 2

B. 3

C. 4

D. 1

Answer: B



[Watch Video Solution](#)

98. Unniltrium belongs to which block and group of the periodic table?

A. d-block, group -10

- B. f-block, group 3
- C. p-block, group-13
- D. d-block, group-11

Answer: B



Watch Video Solution

99. The element having the lowest atomic number and a ground state electronic configuration of $(n - 1)d^6ns^2$ is placed in:

- A. fifth period
- B. fourth period
- C. sixth period
- D. third period

Answer: B



Watch Video Solution

100. Representative elements or main group elements are called:

- A. s-block element
- B. p-block element
- C. both (a) and (b)
- D. none of these

Answer: C



Watch Video Solution

101. The elements $Z = 117$ and 120 have not yet have been discovered, In which family / group would you place these elements and also give the electronic configuration in each case.

- A. group 17, $[Rn]5f^{14}, 6d^{10}, 7s^2 7p^5$
- B. group 1, $[Rn]7s^1$

C. group 17, $[Rn]7s^27p^5$

D. group 1, $[Rn]8s^1$

Answer: A



Watch Video Solution

102. Select the incorrect statement regarding LotharMeyer curve:

A. alkali metals occupy the peak position of curve

B. alkali earth metals are at the ascending part of curve

C. halogens are at the ascending part of curve

D. atomic volume of an element is not the volume of an atom of that element

Answer: B



Watch Video Solution

103. The elements after uranium are called:

- A. transuranic elements
- B. lanthanides
- C. transition elements
- D. none of the above

Answer: A



Watch Video Solution

104. Which orbitals are filled in given order in 5th period of periodic table?

- A. $5s \rightarrow 4d \rightarrow 5p$
- B. $5s \rightarrow 4d \rightarrow 5d$
- C. $4s \rightarrow 5d \rightarrow 4p$
- D. $5s \rightarrow 3d \rightarrow 4p$

Answer: A



Watch Video Solution

105. Which of the following is not f-block elements?

A. Th

B. Ce

C. Lu

D. None of these

Answer: D



Watch Video Solution

106. Consider the following 3 elements. Fe , Cu , Zn

Choose the incorrect option.

A. Only 2 are paramagnetic

B. Number of d-electron: $Fe < Cu < Zn$

C. All are transition elements

D. All are less metallic than s-block elements of same period

Answer: C



Watch Video Solution

107. Which of the following is not d-block elements?

A. Cu

B. Zn

C. Fe

D. Sn

Answer: D



Watch Video Solution

108. The electronic configuration:

$1s^2, 2s^2 2p^6, 3s^2 3p^6, 3d^{10}, 4s^2 4p^6 4d^{10}, 5s^2$ is for:

- A. f-block element
- B. d-block element
- C. p-block element
- D. s-block element

Answer: B



Watch Video Solution

B. Atomic and Ionic Radius

1. Select correct statement about radius of an atom:

- A. Values of van der Waal's radii are larger than those of covalent radii because the van der Waal's forces are much weaker than the forces operating between atoms in a covalently bonded molecule.
- B. The metallic radii are smaller than the van der Waal's radii, since the bonding forces in the metallic crystal lattice are much stronger than the van der Waal's forces.
- C. Both are correct
- D. None is correct



Answer: C



Watch Video Solution

2. Which of the following order of radii is correct?








Answer: B



Watch Video Solution

3. Which of the following is not isoelectronic series?



Answer: D



Watch Video Solution

4. Which of the following order of atomic / ionic radius is not correct?

A. $I^{-} > I > I^{+}$

B. $Mg^{2+} > Na^{+} > F^{-}$

C. $P^{5+} < P^{3+}$

D. $Li > Be > B$

Answer: B



Watch Video Solution

5. The radii of N , N^{3-} , O and O^{2-} are in the order of:

A. $O^{2-} > N^{3-} > O > N$

B. $O^{2-} > N^{3-} > N > O$

C. $N^{3-} > O^{2-} > N > O$

D. $O^{2-} > O > N^{3-} > N$

Answer: C



Watch Video Solution

6. Atomic radii of F & Ne in Angstrom are respectively given by : A) 0.72, 1.60 B) 1.60, 1.60 C) 0.72, 0.72 D) 1.60, 0.72

A. 0.72, 1.60

B. 1.60, 1.60

C. 0.72, 0.72


D. 1.60, 0.72

Answer: A



Watch Video Solution

7. Which one of the following is the smallest in size?







Answer: D



Watch Video Solution

8. Which one of the following is correct increasing order of size ?







Answer: B



Watch Video Solution

9. When the following five anions are arranged in order of decreasing ionic radius, the correct sequence is :




Answer: D



Watch Video Solution

10. Which of the following isoelectronic species has the largest radius?



D. S^{2-}

Answer: C



Watch Video Solution

11. The correct decreasing order of atomic size among the following species is :

$Ar, K^+, Cl^-, S^{2-}, Ca^{2+}$

A. $Ca^{2+} > K^+ > Ar > Cl^- > S^{2-}$

B. $K^+ > Ca^{2+} > Cl^- > Ar > S^{2-}$

C. $S^{2-} > Cl^- > Ar > K^+ > Ca^{2+}$





D. $S^{2-} > Ar > Cl^- > Ca^{2+} > K^+$

Answer: C



Watch Video Solution

12. Ionic radii of :




Answer: D



Watch Video Solution

13. The correct order of radii is:



Answer: B



Watch Video Solution

14. Arrange Ce^{3+} , La^{3+} , Pm^{3+} and Yb^{3+} in increasing order of their size -

A. $Yb^{+3} < Pm^{+3} < Ce^{+3} < La^{+3}$

B. $Ce^{+3} < Yb^{+3} < Pm^{+3} < La^{+3}$

C. $Yb^{+3} < Pm^{+3} < La^{+3} < Ce^{+3}$



D. $Pm^{+3} < La^{+3} < Ce^{+3} < Yb^{+3}$

Answer: A



Watch Video Solution

15. Which one of the following ions has the highest value of ionic radius?







Answer: A



Watch Video Solution

16. The set representing the correct order of ionic radius is



Answer: A



Watch Video Solution

17. Which of the following statement is incorrect for the isoelectronic species?

- A. They have same number of electrons
- B. They may have different number of protons
- C. Their ionic radii decrease with increase in nuclear charge
- D. None of the above

Answer: D



Watch Video Solution

18. Ionic radii are

- A. inversely proportional to effective nuclear charge
- B. inversely proportional to square of effective nuclear charge
- C. directly proportional to effective nuclear charge


D. directly proportional to square of effective nuclear charge

Answer: A



Watch Video Solution

19. The correct order of radii is:







Answer: B



Watch Video Solution

20. K^+ , Cl^- , Ca^{2+} , S^{2-} ions are isoelectronic. The decreasing order of other size is:



Answer: A



Watch Video Solution

21. Li resembles Mg due to diagonal relationship which is attributed to:

A. nearly same atomic and ionic size

B. same value of electron affinity

C. penetration of sub-shells

D. identical effective nuclear charge

Answer: A



Watch Video Solution

22. Select the correct statement.

- A. Across a transition series (from Cr to Cu), there is only a small decrease in atomic radius from one element to another due to very small increase in effective nuclear charge
- B. The rate of decrease in the size across the lanthanide series is less than that across the first transition series
- C. Both are correct statements
- D. None of the statement is correct

Answer: C



Watch Video Solution

23. Among the following species the highest radius is of:



Answer: D



Watch Video Solution

24. In which of the following compound size of cation to anion ratio is minimum?






Answer: B



Watch Video Solution

25. Choose the incorrect ionic radius order.






Answer: B



Watch Video Solution

26. Choose the correct order of atomic/ionic radii.








Answer: A



Watch Video Solution

27. Correct order of radius among the following

F^{-} , Al^{3+} , Na^{+} are



Answer: D



Watch Video Solution

28. Which of the following is correct regarding atomic radius?





Answer: A



Watch Video Solution

29. The correct order of radii is:






Answer: A



Watch Video Solution

30. Select the correct order of ionic radii.







D. None of the above

Answer: A



Watch Video Solution

31. Consider the isoelectronic species, Na^+ , Mg^{2+} , F^- and O^{2-} . The correct order of increasing length of their radii is:







Answer: B



Watch Video Solution

32. Which one of the following is correct order of the size of iodine species ?



Answer: D



33. Which statement about the radii of atoms and their ions is correct?

- A. Cations are smaller than their atoms, anions are larger.
- B. Cations and anions are both smaller than their atoms.
- C. Cations and anions are both larger than their atoms
- D. Cations are larger than their atoms, anions are smaller.

Answer: A

34. The order of magnitude of ionic radii of ions Na^+ , Mg^{2+} , Al^{3+} and Si^{4+} is

- A. $Na^+ < Mg^{2+} < Al^{3+} < Si^{4+}$
- B. $Mg^{2+} > Na^+ > Al^{3+} > Si^{4+}$



Answer: D



Watch Video Solution

35. Which ions has greatest radius in the following?



Answer: D



Watch Video Solution

36. In the isoelectronic species the ionic radii of N^{3-} , O^{2-} , F^{-} are respectively given by

A. 1.36, 1.40, 1.71

B. 1.36, 1.71, 1.40

C. 1.71, 1.40, 1.36

D. 1.70, 1.36, 1.40

Answer: C



Watch Video Solution

37. The Lanthanide contraction is responsible for the fact that

A. Zr and Y have almost the same radius

B. Zr and Nb have similar oxidation state

C. Zr and Hf have almost the same radius





D. Zr and Zn have the same oxidation state

Answer: C



Watch Video Solution

38. Mg , Mg^{2+} , Al and Al^{3+} are arranged in decreasing order of size $1 > 2 > 3 > 4$. Species which are present at 1 and 4 position respectively are:



Answer: B



Watch Video Solution

39. Which of the following order of radii is incorrect?

A. $Sc < Y < La$

B. $Ti < Zr \approx Hf$

C. $Al \approx Zn$

D. $He < Ne < Ar$

Answer: C



Watch Video Solution

40. Size of which elements are represented as non-bonded radii?

A. Halogen

B. Chalcogens

C. Noble gas

D. Pnictogens

Answer: C



Watch Video Solution

41. Which gas-phase atom has the largest radius?

A. Na

B. K

C. Mg

D. Ca

Answer: B



Watch Video Solution

42. For which pair of species are the radii most similar?

A. Li and Na

B. Na and Mg

C. Mn and Fe

D. Fe^{2+} and Fe^{3+}

Answer: C



Watch Video Solution

43. The ions below are listed in order of increasing radii except:

A. $Al^{3+} < Mg^{2+} < Na^{+}$

B. $K^{+} < Cl^{-} < Se^{2-}$

C. $Cl^{-} < Br^{-} < I^{-}$

D. Fe^{2+} and Fe^{3+}

Answer: D



Watch Video Solution

44. Which atom has the largest atomic radius?

A. S

B. Cl

C. Se

D. Br

Answer: C



Watch Video Solution

45. In which list are atoms of the elements Be , B , Mg and Al arranged from smallest to largest atomic radius?

A. $Be < B \leq Mg \leq Al$





B. $Mg < Be < Al < B$

C. $B < Be < Al < Mg$

D. $Al < Mg < B < Be$

Answer: C

46. K^+ , Cl^- , Ca^{2+} , S^{2-} ions are isoelectronic. The decreasing order of other size is:



Answer: B

47. The radius of which is closest to that of the Li^+ ions?









Answer: C



Watch Video Solution

48. For which pair of species is the difference in radii the greatest?







Answer: C



Watch Video Solution

49. In which choice are the species listed in order of increasing radius?







Answer: B



Watch Video Solution

50. In which list are the ions arranged in order of increasing size?



Answer: D



Watch Video Solution

51. In which series are the species listed in order of increasing size ?

A. N, O, F

B. Na, Mg, K

C. Cr, Cr^{2+}, Cr^{3+}

D. Cl, Cl^{-}, S^{2-}

Answer: D



Watch Video Solution

52. When the atoms: Ba, Cs, Mg, Na are arranged in order of increasing size, what is the correct order?

A. $Cs < Na < Mg < Ba$

B. $Mg < Na < Ba < Cs$

C. $Mg < Ba < Na < Cs$

D. $Ba < Mg < Na < Cs$

Answer: B



Watch Video Solution

53. Which element has the largest atomic radius?

A. Br

B. K

C. Mg

D. Na

Answer: B



Watch Video Solution

54. When the atoms, $P(Z = 15)$, $S(Z = 16)$ and $As(Z = 33)$, are arranged in order of increasing radius, what is the correct order?

A. P, S, As

B. As, S, P

C. S, P, As

D. P, As, S

Answer: C



Watch Video Solution

55. Which property decreases from left to right across the periodic table and increases from top to bottom?

A. Atomic radius

B. Electronegativity

C. Ionization energy

D. Melting point

Answer: A



Watch Video Solution

56. Consider the ions Li^+ , Na^+ , Be^{2+} and Mg^{2+} . Which two are closest to one another in size?

A. Li^+ and Na^+

B. Be^{2+} and Mg^{2+}

C. Be^{2+} and Li^+

D. Li^+ and Mg^{2+}

Answer: D



Watch Video Solution

57. Which pair of elements have chemical properties that are the most similar ?

A. Be and B

B. Al and Ga

C. Co and Cu

D. F and I

Answer: B



Watch Video Solution

58. When the atoms Li , Be , B and Na are arranged in order of increasing atomic radius, what is the correct order?

A. B , Be , Li , Na

B. Li , Be , B , Na

C. Be , Li , B , Na

D. *Be*, *B*, *Li*, *Na*

Answer: A



Watch Video Solution

59. Which property of an element is most dependent on this shielding effect?

- A. Atomic number
- B. Atomic mass
- C. Atomic radius
- D. Number of stable isotopes

Answer: C



Watch Video Solution

60. Which element has the largest atomic radius?

A. Li

B. K

C. As

D. Br

Answer: B



Watch Video Solution

61. The atomic radius of elements of which of the following series would be nearly the same

A. *Na, K, Rb, Cs*

B. *Li, Be, B, C*

C. *Fe, Co, Ni*

D. *F, Cl, Br, I*

Answer: C



Watch Video Solution

62. The radii of F , F^{-} , O and O^{-2} are in the order of

A. $O^{2-} > F^{-} > F > O$

B. $F^{-} > O^{2-} > F > O$

C. $O^{-2} > O > F^{-} > F$

D. $O^{2-} > F^{-} > O > F$

Answer: D



Watch Video Solution

63. Which of the following case the size ratio is minimum?

A. Li^{-} / Li

B. H^- / H

C. Na^- / Na

D. Can not be predict

Answer: C



Watch Video Solution

64. For the element X, student Surbhi measured its radius as $102nm$, Mr. Gupta as 113 nm and Mr. Agarwal as $100nm$ using same apparatus. Their teacher explained that measurements were correct by saying that recorded values by three students were

A. crystal, van der Waal's and covalent radii

B. covalent, crystal and van der Waal's radii

C. van der Waal's ionic and covalent

D. None is correct

Answer: A



Watch Video Solution

65. The species having smallest ionic radius is:



Answer: A



Watch Video Solution

66. In the crystals of which of the following ionic compounds would you expect maximum distance between the centres of the cations and anion?

A. CsF

B. CsI

C. LiI

D. LiF

Answer: B



Watch Video Solution

67. CONSEQUENCE OF LANTHANIDE CONTRACTION

- A. Contraction of atom of lanthanum element due to poor shielding of d-subshell electron
- B. Contraction of atom of lanthanum element due to high shielding of d-subshell electron
- C. Contraction of atom of elements after lanthanum due to poor shielding of f-subshell electron

D. Contraction of atom of elements before lanthanum due to poor shielding of f-subshell electron.

Answer: C



Watch Video Solution

68. Which of the following have total 18 electrons and it is larger than Cl^{-} ?

A. K^{+}

B. Ca^{2+}

C. Br^{-}

D. None of these

Answer: D



Watch Video Solution

69. The size of isoelectronic species $F^{\ominus}$, Ne , and Na^{+} affected by

- A. nuclear charge (Z)
- B. valence principal quantum number (n)
- C. electron-electron interaction in the outer orbitals
- D. none of the factors because their size is the same

Answer: A



Watch Video Solution

70. Which of the following order of ionic-radius is correct?


- A. $F^{-} > O^{2-} > N^{3-}$
- B. $Na^{+} > Mg^{+2} > Al^{+3}$
- C. $H^{-} > H < H^{+}$
- D. $Fe < Fe^{+2} < Fe^{+3}$

Answer: B




Watch Video Solution


71. Covalent radius of Cl is 99 pm. Select best representation for Cl_2 molecule:




A.



B.



C.



D.

Answer: C

[Watch Video Solution](#)

C. Ionisation Energy

1. The values of $IE(I)$, $IE(II)$, $IE(III)$ and $IE(IV)$ of an atom are respectively $7.5eV$, $25.6eV$, $48.6eV$ and $170.6eV$. The electronic configuration of the atom will be :

A. $1s^2, 2s^2 2p^6, 3s^1$

B. $1s^2, 2s^2 2p^6, 3s^2 3p^1$

C. $1s^2, 2s^2 2p^6, 3s^2 3p^3$

D. $1s^2, 2s^2 2p^6, 3s^2$

Answer: B

[Watch Video Solution](#)

2. IE_1 and IE_2 of Mg are 178 and $348 \text{ kcal mol}^{-1}$. The energy required for the reaction $Mg \rightarrow Mg^{2+} + 2e^-$ is

- A. +170 kcal
- B. +526 kcal
- C. -170 kcal
- D. -526 kcal

Answer: B



Watch Video Solution

3. Which represents alkali metals (ie., 1 group metals) based on $(IE)_1$ and $(IE)_2$ values?

- | | | | |
|----|---|----------|----------|
| A. | | $(IE)_1$ | $(IE)_2$ |
| | Y | 100 | 110 |
| B. | | $(IE)_1$ | $(IE)_2$ |
| | Y | 95 | 120 |
| C. | | $(IE)_1$ | $(IE)_2$ |
| | Z | 95 | 500 |

D.
$$\begin{matrix} & (IE)_1 & (IE)_2 \\ M & 200 & 250 \end{matrix}$$

Answer: C



Watch Video Solution

4. Which is most similar for the elements in a group in the periodic table?

- A. Physical state
- B. Melting point
- C. First ionization energy
- D. Ground state electronic configuration

Answer: D



Watch Video Solution

5. The first ionisation potential of Al is smaller than that of Mg because:

- A. the atomic size of $Al > Mg$
- B. the atomic size of $Al < Mg$
- C. Al has one unpaired electron in $3p$ – orbital
- D. Mg has incompletely filled $3s$ – orbital

Answer: C



Watch Video Solution

6. Which of the following gaseous atoms has the highest value of ionisation enthalpy?

- A. P
- B. Si
- C. Mg
- D. Al

Answer: A

 [Watch Video Solution](#)

7. The first ionisation potential of Na , Mg , Al and Si are in the order

A. $Na < Mg > Al < Si$

B. $Na > Mg > Al > Si$

C. $Na < Mg < Al < Si$

D. $Na > Mg > Al < Si$

Answer: A

 [Watch Video Solution](#)

8. The first ionisation potential in electron volts of nitrogen and oxygen atoms are respectively given by

A. 14.6, 13.6

B. 13.6, 14.6

C. 13.6, 13.6

D. 14.6, 14.6

Answer: A



Watch Video Solution

9. Which of the following relation is correct with respect first (I) and second (II) ionisation of sodium and magnesium?

A. $I_{Mg} = II_{Na}$

B. $I_{Na} > I_{Mg}$

C. $II_{Mg} > II_{Na}$

D. $II_{Na} > II_{Mg}$

Answer: D



Watch Video Solution

10. The ionization energy will be higher when the electron is removed fromif other factors being equal.

A. s-orbital

B. p-orbital

C. d-orbital

D. f-orbital

Answer: A



Watch Video Solution

11. Which of the following isoelectronic ion has the lowest 1st ionization energy?

A. Na^{+}

B. F^{-}

C. Mg^{2+}

D. O^{2-} and F^{-}

Answer: D



Watch Video Solution

12. A large difference between the third and fourth ionization energies indicates the presence of:

A. 4 valence electrons in an atom

B. 5 valence electrons in an atom

C. 3 valence electrons in an atom

D. 8 valence electrons in an atom

Answer: C



Watch Video Solution

13. The second ionisation enthalpies of elements are always higher than their first ionisation enthalpies because:




- A. the cation is smaller than its parent atom
- B. it is easier to remove electron from cation
- C. ionization is an endothermic process
- D. cation formed always have stable half filled or completely filled valence shell electron configuration


Answer: A



Watch Video Solution

14. Which of the following is the correct order of ionisation energy?





Answer: A



Watch Video Solution

15. The peaks in ionisation potential curves are occupied by

- A. alkali metals
- B. alkaline earth metals
- C. noble gas elements
- D. halogens

Answer: C



Watch Video Solution

16. Which metal is the most reactive?

A. Silver

B. Lead

C. Iron

D. Caesium

Answer: D



Watch Video Solution

17. The five successive energies of an element are 800, 2427, 3658, 25024 and $32824 \text{ kJ mol}^{-1}$ respectively. The number of valence electron is

A. 3

B. 5

C. 1

D. 2

Answer: A

 Watch Video Solution

18. Which has maximum first ionization potential?

A. C

B. N

C. B

D. O

Answer: B

 Watch Video Solution

19. Values of IE_1 , IE_2 and IE_3 of an element are 9.31, 18.21 and 553.83eV . What informations do these data convey?

A. The element has two electrons in the valence shell

B. The element has two p-electrons in the valence shell

C. (a) and (b) both

D. None of the two

Answer: A



Watch Video Solution

20. Which of the following is the correct order of ionisation energy?

(1) $Be^+ > Be$ (2) $Be > Be^+$

(3) $C > Be$ (4) $B > Be$

A. 2,3

B. 3,4

C. 1,3

D. None of these

Answer: C



Watch Video Solution

21. The incorrect statement Among the following is A)The first ionisation potential of Al is less than the first ionisation potential of Mg. B)The first ionisation potential of Na is less than the first ionisation potential of Mg. C)The second ionisation potential of Mg greater than the second ionisation potential of Na D)The third ionisation potential of Mg greater than the third ionisation potential of Al

- A. the IE_1 of Al is less than IE_1 of Mg
- B. the IE_2 of Mg is less than IE_2 of Mg
- C. the IE_1 of Na is less than IE_1 of Mg
- D. the IE_3 of Mg is less than IE_3 of Al

Answer: B



Watch Video Solution

22. The set representing the correct order of the first ionisation potential is

A. $K > Na > Li$

B. $Be > Mg > Ca$

C. $B > C > N$

D. $Ge > Si > C$

Answer: B



Watch Video Solution

23. The increasing order of the first ionization enthalpies of the elements B, P, S and F (lowest first) is:

A. $F < S < P < B$

B. $P < S < B < F$

C. $B < P < S < F$

$$\text{D. } B < S < P < F$$

Answer: D



Watch Video Solution

24. Which of the following is the result of f-orbital contraction effect?

- A. Similarity in the size of Al and Ga
- B. Similarity in the properties of Li and Mg
- C. High ionization energy of Tl as compared to In
- D. Decrease in atomic size of '3d' series elements

Answer: C



Watch Video Solution

25. Which of the following magnetic moment values will correspond to the highest ionisation energy for the element which is placed in 4th period and 7th group of long form of periodic table?

A. $2\sqrt{2}$

B. $\sqrt{15}$

C. $\sqrt{35}$

D. $\sqrt{24}$

Answer: A



View Text Solution

26. Which of the following s-block metals has the lowest first ionisation energy?

A. Mg

B. Be

C. Na

D. Li

Answer: C



Watch Video Solution

27. Hydrogen has high ionization energy than alkali metals, due to its:

A. large size

B. small size

C. higher nuclear charge

D. more penetration power of s-subshell electrons

Answer: C



Watch Video Solution

28. Which of the following element has maximum, first ionisation potential?

A. V

B. Ti

C. Cr

D. Mn

Answer: D



Watch Video Solution

29. An element having high first ionization energy in a given period is:

A. inert gas

B. transition element

C. representative element

D. inner transition element

Answer: A



Watch Video Solution

30. Ionisation enthalpy of Na would be same as

- A. electron affinity of Na^+
- B. electronegativity of Na^+
- C. electron affinity of He
- D. ionisation potential of Mg

Answer: A



Watch Video Solution

31. Which element having the following electronic configuration has minimum ionisation potential?

A. $1s^2, 2s^2, 2p^5$

B. $1s^2, 2s^2 2p^6, 3s^2 3p^2$

C. $1s^2, 2s^2 2p^6, 3s^1$

D. $1s^2, 2s^2 2p^6$

Answer: C



Watch Video Solution

32. Select correct statement.

A. Atomic size of the same group elements of 3d and 4d-series are almost similar

B. First ionisation energies of 5d-series elements are smaller than that of 3d and 4d-series elements

C. Both are correct

D. Both are incorrect

Answer: D



Watch Video Solution

33. Which option is correct?

A. 3rd I. E. of $C > N$

B. 2nd I.E. of $N < O$

C. 1st I.E. of $F > O$

D. All of these

Answer: D



Watch Video Solution

34. Which one of the following electronic configuration of an atom has the highest ionisation energy?

A. $1s^2, 2s^2 2p^3$

B. $1s^2, 2s^2 2p^6, 3s^1$

C. $1s^2, 2s^2 2p^6$

D. $1s^2, 2s^2 2p^5$

Answer: C



Watch Video Solution

35. Identify the incorrect order of first ionization energy.

A. $Li > Na > K$

B. $F > Cl > I$

C. $Be < B < C$





D. $O < N < F$

Answer: C



Watch Video Solution

36. Amongst the following elements whose electronic configuration are given below, the one having the highest enthalpy is



Answer: B



Watch Video Solution

37. Which of the following ion of an element E (E belongs to 2nd period) having $1s^22s^0, 2p^0$ electronic configuration in ground state? For the element E following data is given:

$$IE_1 \text{ of } E = 800 \text{ kJ/mole} \quad IE_2 \text{ of } E = 1428 \text{ kJ/mole}$$

$$IE_3 \text{ of } E = 2394 \text{ kJ/mole} \quad IE_4 \text{ of } E = 25020 \text{ kJ/mole}$$

A. E^{+}

B. E^{++}

C. E^{+++}

D. E^{++++}

Answer: C



Watch Video Solution

38. Higher values of ionisation energies of the 5d-transition elements are consistent with the:

A. relative smaller effective nuclear charge

B. relative smaller size of their atoms

C. relative smaller penetration

D. all are correct

Answer: B



[Watch Video Solution](#)

39. Metallic nature increases moving down the group because:

- A. nuclear charge decrease
- B. shielding increases
- C. (a) and (b) both
- D. none of these

Answer: B



[Watch Video Solution](#)

40. Select the correct statement.

- A. More active metals are on the left side of the periodic table
- B. Less active metals are on the left side of the periodic table

- C. Reducing power of alkali metals (g) decreases moving down the group
- D. Metallic character decreases down the group

Answer: A



Watch Video Solution

41. Which of the following sequence regarding the first ionisation potential of coinage metal is correct?

A. $Cu > Ag > Au$

B. $Cu < Ag < Au$

C. $Cu > Ag < Au$

D. $Ag > Cu > Au$

Answer: C



Watch Video Solution

42. The ionisation energy for 'Sc', 'Y' and 'La' are x , y and z kcal/mol respectively. Choose the correct relationship between them.

A. $z > x \approx y$

B. $x > y > z$

C. $z > z \approx y$


D. $z < y > z$

Answer: B




Watch Video Solution


43. Select proper graph of 13th group ionization energy vs. its elements.




A.



B.



C.



D.

Answer: C



Watch Video Solution

44. Second ionisation potential of Li , Be , B is in the order:

A. $Li > Be > B$

B. $Li > B > Be$

C. $Be > Li > B$


D. $B > Be > C$

Answer: B




Watch Video Solution


45. Select correct graph, which is plotted between ionisation energy of 2nd period elements and their atomic number.




A.



B.



C.



D.

Answer: A



Watch Video Solution

46. For metal M, first ionisation energy is equal to 38 kcal/mole. Which of the following can be a possible value of ionisation enthalpy of M^+ ion in kcal.mole at 400K. [Given: $R = 2\text{cal} / \text{mole kelvin}$]

A. $38\text{kcal} / \text{mole}$

B. $40\text{kcal} / \text{mole}$

C. $39\text{kcal} / \text{mole}$

D. $45\text{kcal} / \text{mole}$

Answer: D



View Text Solution

47. The ionisation enthalpy of lithium is much lower than that of helium despite the fact that the nuclear charge of lithium is $+3$ and that of helium is $+2$. It is due to:

- A. 2s electron in lithium is more strongly bound to the nucleus than the 1s electron in He
- B. 2s electron in lithium is equally bound to the nucleus than the 1s electron in He
- C. 2s electron in lithium is less strongly bound to the nucleus than the 1s electron in He
- D. Atomic size of He is more than Li

Answer: C



Watch Video Solution

48. The first ionisation enthalpy ($\Delta_i H^{o-}$) values of the third period elements, Na , Mg and Si are respectively 496, 737 and $786 kJ mol^{-1}$. Predict whether the first $\Delta_i H^{o-}$ value for Al will be more close to 575 or $760 kJ mol^{-1}$? Justify your answer.

A. 350

B. 860

C. 575

D. 1125

Answer: C



Watch Video Solution

49. Correct order of IE_1 are:

(P) $Li < B < Be < C$ (Q) $O < N < F$

(R) $Be < N < Ne$

A. P,Q

B. Q only


C. P,R

D. P,Q,R

Answer: D



Watch Video Solution



Then find ΔH_{eg} of X.

A. $-250kJ$

B. $-1550kJ$

C. $250kJ$

D. $+1550kJ$

Answer: A



Watch Video Solution

51. The correct order of the second ionisation potential of carbon, nitrogen, oxygen and fluorine is

A. $C < N < O < F$

B. $C < O < N < F$

C. $C < N < F < O$

D. $C < O < F < N$

Answer: C



Watch Video Solution

52. Sodium forms Na^+ ion but it does not form Na^{+2} because -

A. Very low value of $(I.E.)_1$ and $(I.E.)_2$

B. Very high value of $(I.E.)_1$ and $(I.E.)_2$

C. High value of $(I. E.)_1$ and low value of $(I. E.)_2$

D. Low value of $(I. E.)_1$ and high value of $(I. E.)_2$

Answer: D



Watch Video Solution

53. One method for calculating Z_{eff} is to use the equation:

$$Z_{eff} = \sqrt{\frac{(E)(n^2)}{1312kJ/\text{mole}}}$$

where E is the energy necessary to remove an electron from an atom and n is the principal quantum number of the electron. Use this equation to calculate Z_{eff} value for the highest energetic electron in potassium.

$$E = 418.8kJ/\text{mole}$$

A. 2.86

B. 2.26

C. 3.26

D. 2.00

Answer: B



Watch Video Solution

54. Atomic nitrogen has a higher ionization energy than atomic oxygen.

This is best explained by:

- A. by lower electron-electron repulsion in nitrogen
- B. the greater effective nuclear charge of nitrogen
- C. the fact that the electron ionized in N is from the 2s subshell, while that ionized from O is from the 2p subshell
- D. the fact that N has an odd number of electron while O has an even number.

Answer: A



Watch Video Solution

55. Which element has the smallest first-ionization energy?

A. Mg

B. Al

C. Si

D. P

Answer: B



Watch Video Solution

56. Which set is expected to show the smallest difference in IE_1 ?

A. He, Ne, Ar

B. B, N, O

C. Mg, Mg^+, Mg^{2+}

D. Fe, Co, Ni

Answer: D



[Watch Video Solution](#)

57. When the elements carbon, nitrogen and oxygen are arranged in order of increasing ionization energies, what is the correct order?

A. C, N, O

B. O, N, C

C. N, C, O

D. C, O, N

Answer: D



[Watch Video Solution](#)

58. Which has the highest ionization energy for the removal of the second electron?

A. F

B. Ne

C. Na

D. Mg

Answer: C



Watch Video Solution

59. In which list are the elements arranged in order of increasing first ionization energy?

A. *Li, Na, K*

B. *S, O, F*

C. *Na, Mg, Al*

D. *F, Ne, Na*

Answer: B



Watch Video Solution

60. Which atom has the smallest first ionization energy?

A. Na

B. K

C. Mg

D. Ca

Answer: B



Watch Video Solution

61. Which element has the largest first ionization energy?

A. Li

B. B

C. N

D. Na

Answer: C



Watch Video Solution

62. Which list gives the symbols of the element in the order of increasing first ionization energy?

A. F , Ne , Na

B. Al , Mg , Na

C. Sr , Ca , Mg





D. Cl , Br , I

Answer: C



Watch Video Solution

63. Which equation represents the first ionization of calcium?







Answer: B



Watch Video Solution

64. Which process release the most energy?



Answer: C



Watch Video Solution

65. Which element exhibits the successive ionization energies given in the table?

Ionization Energy $\text{kJ} \times \text{mol}^{-1}$

1st	738
2nd	1451
3rd	7733
4th	10540
5th	13628

A. *Na*

B. *Mg*

C. *Al*

D. *Si*

Answer: B



Watch Video Solution

66. Which element has the highest first ionization energy?

A. Al

B. Si

C. P

D. S

Answer: C



Watch Video Solution

67. The removal of an electron from which gaseous atom requires the greatest amount of energy?

A. *Na*

B. *Cl*

C. *K*

D. *Br*

Answer: B

68. Properties of the alkaline earth metals that increase from Be to Ba include which of the following?

(P) Atomic radius (Q) Ionization energy

(R) Nuclear charge

A. P and Q only

B. P and R only

C. Q and R only

D. P,Q and R

Answer: B

69. Of the atoms listed, which has the largest third ionization energy?

A. Ca

B. Mg

C. Al

D. Si

Answer: B



Watch Video Solution

70. The successive ionization energies (in kJ/mol) for an element are shown below.

E_1	E_2	E_3	E_4	E_5
577	1820	2740	11600	14800

What is the electron configuration of this element?

A. $1s^2 2s^2 2p^6 3s^1$

B. $1s^2 2s^2 2p^6 3s^2 3p^1$

C. $1s^2 2s^2 2p^6 3s^2 3p^3$

D. $1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$

Answer: B



Watch Video Solution

71. Of the elements listed, which has the highest first ionization energy?

A. *Li*

B. *Be*

C. *Na*

D. *Mg*

Answer: B



Watch Video Solution

72. Ionization energies vary from left to right across the periodic table.

Factors that contribute to this variation include which of the following?

(P) Changes in the nuclear charge

(Q) Differences in shielding by valence electrons

(R) Differences in shielding by core electrons

A. P only

B. R only

C. P and Q only

D. P,Q and R

Answer: C



Watch Video Solution

73. Which group best illustrates the transition from non-metallic to metallic behavior with increasing atomic numbers?

A. *Be, Mg, Ca, Sr*

B. *N, P, As, Sb*


C. *F, Cl, Br, I*

D. *Fe, Ru, Os, Hs*

Answer: B



Watch Video Solution



A. In (I) energy is released, (II) energy is absorbed

B. In both (I) and (II) energy is absorbed

C. In both (I) and (II) energy is released

D. In (I) energy is absorbed, (II) energy is released

Answer: C



Watch Video Solution

75. How does the reducing ability of the elements vary across the period from N to At? It:

- A. decreases steadily
- B. increases steadily
- C. decreases then increases
- D. increases then decreases

Answer: A



Watch Video Solution

76. Of the elements given, which has the lowest ionization energy?

- A. N
- B. P
- C. S
- D. Cl

Answer: C



Watch Video Solution

77. Three elements X , Y and Z have atomic numbers 19 , 37 and 55 respectively . Then the correct statements (s) is / are

- A. Their ionisation potential would increase with the increasing atomic number
- B. Y would have an ionisation potential in between those of X and Z
- C. Z would have the highest ionisation potential
- D. Y would have the highest ionisation potential

Answer: B



Watch Video Solution

78. Which element has the lowest first ionization energy?

A. B

B. C

C. Al


D. Si

Answer: C



Watch Video Solution

79. If we move from centre towards end points through three different paths (P),(Q),(R) in following figure, then which path is correct increasing order of IE_1 ?



[*N*, *S*, *C*, *O*, *P*, *Si* represent various elements of periodic table]

- A. P only
- B. Q and R
- C. R only
- D. P and R

Answer: A



Watch Video Solution

80. First, second and third I.P. values are $10eV$, $15eV$ and $150eV$. Element can be:

A. Be

B. B

C. F

D. Na

Answer: A



Watch Video Solution

81. The third ionisation energy will be maximum for:

A. nitrogen

B. boron

C. phosphorus

D. aluminium

Answer: A



Watch Video Solution

82. Find the highest ratio of IP values of given pair of elements:

A. $He:Ne$

B. $Ne:Ar$

C. $He:Xe$

D. $Kr:Xe$

Answer: C



Watch Video Solution

83. Given the correct order of initials T (true) of F (false) for following statement.

(I) Top positions of Lothar-Mayer's atomic volume curve are occupied by

Alkali metals

(II) Number of elements presents in the fifth period of the periodic table are 32.

(III) 2^{nd} I. P. of Mg is less than the 2^{nd} I.P. of Na.

(IV) A p-orbital can take maximum of six electrons.

A. TFTF

B. TFFT

C. FFTF


D. TTFF




Answer: A



Watch Video Solution

84. Which of the following transitions involves maximum amount of energy?





Answer: D



Watch Video Solution

85. The first five ionization energies of an element are 9.1, 16.2, 24.5, 35 and 205.7eV respectively then, number of valence electrons in the atom is:

A. 2

B. 3

C. 4

D. 5

Answer: C

[Watch Video Solution](#)

86. The incorrect option out of the following is:

- A. IP_2 of $Li > IP_2$ of Be
- B. EA_1 of $F > EA_1$ of Br
- C. IP_1 of $Pb < IP_1$ of Sn
- D. $N_2O_5 > CO_2$ (acidic nature)

Answer: C

[Watch Video Solution](#)

87. The third ionization energy is least for:

- A. nitrogen
- B. phosphorus
- C. aluminium

D. boron

Answer: C



Watch Video Solution

88. Ionisation energy is highest for:

A. F

B. Cl^-

C. Be^-

D. S^-

Answer: A



Watch Video Solution

89. Ionisation energies of Cl^- , Cl and Cl^+ are respectively (in kJ/mol):

A. 1251, 349, 2300

B. 2300, 349, 1251

C. 349, 1251, 2300

D. 349, 2300, 1251

Answer: C



Watch Video Solution

90. Which of the following order is correct for indicated property?

A. $I_1 > Br_2 > Cl_2 > F_2$ (Reactivity)

B. $Li(aq) > Na(aq) > K^+(aq)$ (electrical conductivity)

C. $Na > Mg > Cu$ (Electropositive character)

D. $N^{-3} > O^{-2} > Mg^{+2} > Na^+$ (Ionic size)

Answer: C



Watch Video Solution

91. Mg cannot exhibit +3 oxidation state, because its:

A. $IE_3 < IE_2$

B. $IE_1 \approx IE_2$

C. $IE_3 > IE_2$

D. $IE_3 = 0$

Answer: C



Watch Video Solution

92. Electronic configuration of valence shell of four elements, A , B , C and D are?

(A) ns^2, np^2 (D) ns^2, np^3

(B) ns^2, np^4 (D) ns^2, np^5

Arrange them in increasing order of IE_2 :

A. $A < B < C < D$

B. $A < B < D < C$

C. $B < A < D < C$

D. $B < A < C < D$

Answer: B



Watch Video Solution

93. Which of the following order is incorrect?

A. $Cd < Hg(IE_1)$

B. $Ag < Au(IE_1)$

C. $Y < La(IE_1)$

D. None of these

Answer: C



Watch Video Solution

D. Electron Gain Enthalpy (Electron Affinity)

1. The correct order of electron affinity is:

A. $Be < B < C < N$

B. $Be < N < B < C$

C. $N < Be < C < B$

D. $N < C < B < Be$

Answer: B



Watch Video Solution

2. Which of the following will have the most negative electron gain enthalpy and which the least negative? P , S , Cl , F . Explain your answer.

A. F, Cl

B. CI, F

C. CI, S

D. CI, P

Answer: D



Watch Video Solution

3. Arrange S , O and Se in ascending order of electron affinity?

A. $O > S > Se$

B. $S > Se > O$

C. $Se > O > S$

D. $S > O > Se$

Answer: B



Watch Video Solution

4. Fluorine has the highest electronegativity among the ns^2np^5 group on the Pauling scale, but the electron affinity of fluorine is less than that of chlorine because :

- A. the atomic number of fluorine is less than that of chlorine
- B. fluorine being the first member of the family behaves in an unusual manner
- C. chlorine can accommodate an electron better than fluorine by utilising its vacant 3d-orbital
- D. of small size, high electron density and an increased electron repulsion makes addition of an electron to fluorine less favourable than that in the case of chlorine in isolated stage

Answer: D



Watch Video Solution

5. Which one of the following arrangements represents the correct order of electron gain enthalpy of the given atomic species?

A. $S < O < Cl < F$

B. $O < S < F < Cl$

C. $Cl < F < S < O$



D. $F < Cl < O < S$

Answer: B



Watch Video Solution

6. For the process



A. ionisation energy of $X^-(g)$ is y

B. electron affinity of $X(g)$ is x

C. electron affinity of $X(g)$ is y

D. all are correct statements

Answer: D



Watch Video Solution

7. Considering the elements F , Cl , O and S , the correct order of their electron affinity values is:

A. $F > Cl > O > S$

B. $F > I > Cl > S$

C. $Cl > F > S > O$

D. $O > F > S > Cl$

Answer: C



Watch Video Solution

8. Which will have the maximum value of electron affinity O^x , O^y , O^z [x,y and z respectively are 0, -1 and -2]?

A. O^x

B. O^y

C. O^z

D. All have equal

Answer: A



Watch Video Solution

9. The electron gain enthalpies of halogens are as given below:

$F = -332$, $Cl = -349$, $Br = -324$, $I = -295 \text{ kJ mol}^{-1}$. The less negative value for F as compared to that of Cl is due to:

A. strong electron-electron repulsions in the compact 2-p sub shell of F

B. strong electro-electron repulsion in the bigger 3-p sub shell of Cl

C. higher electronegativity value of Cl


D. higher effective nuclear charge of F

Answer: A



Watch Video Solution

10. The formation of the oxide ion $O_{(g)}^{2-}$ requires first an exothermic and then an endothermic step as shown below.



A. O^{-} ion will tend to resist the addition of another electron of

account of same charge

B. Oxygen has high electron affinity

C. Oxygen is more electronegative

D. O^{-} ion has comparatively larger size than oxygen atom.

Answer: A



Watch Video Solution

11. For the electron affinity of halogens (with $-ve$ sign), which of the following is correct?

A. $Br > F$

B. $F > Cl$

C. $Br > I$

D. $F^- > I$

Answer: C



Watch Video Solution

12. Which of the following has minimum electron affinity?

A. O

B. S

C. *Se*

D. *Te*

Answer: A



Watch Video Solution

13. The correct order of electron affinity is:

A. $N > P > S > Cl$

B. $P > N > S > Cl$

C. $Cl > N > P > S$

D. $Cl > S > P > N$

Answer: D



Watch Video Solution

14. Which of the following statement is wrong?

- A. van der Waals' radius of iodine is more than its covalent radius.
- B. All isoelectronic ions belong to same period of the periodic table
- C. I.E (I) of N is higher than that of O while I.E (II) of O is higher than that of N
- D. The electron affinity of N is less than that of P

Answer: B



Watch Video Solution

15. Which one of the following statements is incorrect?

- A. Greater is the nuclear charge, greater is the negative electron gain enthalpy

- B. Nitrogen has almost zero electron gain enthalpy
- C. Negative electron gain enthalpy decreases from fluorine to iodine in the group
- D. Chlorine has highest negative electron gain enthalpy

Answer: C



Watch Video Solution

16. Which of the following electron affinity order is incorrect?





- A. $C < N$
- B. $Na^+ > Li^+$
- C. $O^+ < S^+$
- D. All of these

Answer: D



Watch Video Solution

17. Select the equation is endothermic in nature?



Answer: D



Watch Video Solution

18. Which one of the following statements is correct?

A. The elements like F, Cl, Br, O etc. having high values of electron affinity act as strong oxidising agent.

- B. The elements having low values of ionisation energies act as strong reducing agent
- C. The formation of $S^{2-}(g)$ is an endothermic process
- D. All of the above

Answer: D



Watch Video Solution

19. Electron affinity of first element in which of the following pairs is higher?

A. Cl, F

B. Se, S

C. O, F

D. S, Cl


Answer: A

20. The least stable in amongst the following is :



Answer: B

21. In which of the following arrangements the order is NOT according to the property indicated against it ?



C. $I < Br < F < Cl$ increasing electron gain enthalpy (with negative sign)

D. $Li < Na < K < Rb$ increasing metallic radius

Answer: B



Watch Video Solution

22. Electron gain enthalpy is positive when:

A. O^- is formed from O

B. O^{2-} is formed from O^-

C. O^+ is formed from O

D. electron affinity is always a negative value

Answer: B



Watch Video Solution

23. Following are the values of the electron gain enthalpy (in kJ mol^{-1}) of the formation of O^- and O^{2-} from O:

A. $-142, -744$

B. $-142, 744$

C. $142, 744$

D. $-142, -142$

Answer: B



Watch Video Solution

24. Electron addition would be easier in:

A. S

B. S^+

C. S^-

D. S^{2+}

Answer: D



Watch Video Solution

25. The electron affinities of N , O , S and Cl are such that:

A. $N < O < S < Cl$

B. $O < N < Cl < S$

C. $O = Cl = N = S$

D. $O < S < Cl < N$

Answer: A



Watch Video Solution

26. Consider the following statements.

Statement-1: Fluorine does not form any polyhalide as it has low F-F bond energy.

Statement-2: The chlorine has the most negative electron gain enthalpy.

Statement-3: The first ionization potentials of N and O atoms are 14.6 and 13.6eV respectively.

Which of the above statements are correct?

A. Statements 1,2 and 3

B. Statement 1 and 2

C. Statement 1 and 3



D. Statements 2 and 3



Answer: A



Watch Video Solution

27. In which of the following process, maximum amount of energy involved?








Answer: A



Watch Video Solution

28. The increasing order of electron affinity is



Answer: C



Watch Video Solution

29. Which of the following has least first electron affinity ?

A. O

B. S

C. F





D. Cl

Answer: A



Watch Video Solution

30. Which of the following processes is endoergic in nature?



Answer: A



Watch Video Solution

31. Which of the following is not the correct arrangement according to the property indicated against it ?

A. $Li < Na < K < Cs$ (metallic radius)

B. $I < Br < F < Cl$ (electron affinity)

C. $B < C < N < O$ (first ionisation energy)

D. $Al^{3+} < Mg^{2+} < Na^{+} < F^{-}$ (ionic radius)

Answer: C



Watch Video Solution

32. Electronic configuration of four elements p,q,r and s are given below

(p) $1s^2 2s^2 2p^6$ (q) $1s^2 2s^2 2p^4$

(r) $1s^2 2s^2 3s^1$ (s) $1s^2 2s^2 2p^5$

Correct order of increasing tendency of gain electron is

A. $A < C < B < D$

B. $A < B < C < D$

C. $D < B < C < A$

D. $D < A < B < C$

Answer: A



Watch Video Solution

33. Among halogens, the correct order of amount of energy released in electron gain (electron gain enthalpy) is:

A. $F > Cl > Br > I$

B. $F < Cl < Br < I$

C. $F < Cl > Br > I$

$$D. F < CI < Br < I$$

Answer: C



Watch Video Solution

34. The first ionisation potential of Na is $5.1eV$. The value of electrons gain enthalpy of Na^+ will be

A. $-2.55eV$

B. $-5.1eV$

C. $-10.2eV$

D. $2.55eV$

Answer: B



Watch Video Solution

35. The incorrect statement is :

- A. The formation of $O^{2-}(g)$ from $O(g)$ is exothermic process
- B. Phosphorous has more electron affinity than nitrogen
- C. Ne has larger radius than N
- D. $|\Delta H_{eg} \text{ of } Na^{\oplus}| = |I. E. \text{ of } Na|$

Answer: A



Watch Video Solution

36. Which of the following atom has lowest negative electron gain enthalpy?

- A. O
- B. S
- C. Se
- D. Te

Answer: A



Watch Video Solution

37. Which of the following series correctly match relationship between elements as indicated

$A \rightarrow B$

A. $F \rightarrow IEA$ decreases

B. $C \rightarrow F$ 2nd IE increases

C. $He \rightarrow Xe$ noble character decreases

D. $Sc \rightarrow Zn$ atomic radii decreases regularly

Answer: C



Watch Video Solution

38. Select the incorrect order about halogen.

A. $E. A. = CI > F > Br > I$

B. $B. E. = CI_2 > F_2 > Br_2 > I_2$

C. $I. E. = F > CI > Br > I$

D. Oxidizing strength = $F_2 > CI_2 > Br_2 > I_2$

Answer: B



Watch Video Solution

39. Most easily Fe^{2+} ions can be produced from Fe atom by absorbing energy from which of the following process. Consider equal no. of given species.

A. Formation of O^- from O-atom

B. Formation of F^- from F-atom

C. Formation of Cl^- from Cl-atom




D. Formation of S^- from S-atom

Answer: C



Watch Video Solution

40. Which of the following process is associated with best possibility of the energy release?



Answer: C



Watch Video Solution

41. The correct order of electron affinity for the different families is:

A. Halogen > carbon > nitrogen > oxygen

B. Halogen > oxygen > nitrogen > carbon

C. Halogen > nitrogen > carbon > oxygen

D. Halogen > oxygen > carbon > nitrogen

Answer: D



Watch Video Solution

42. $N_0/2$ atoms of $X(g)$ are converted into $X^+(g)$ by energy E_1 . $N_0/2$ atoms of $X(g)$ are converted into $X^-(g)$ by the energy E_2 . Hence ionisation potential and electron affinity of $X(g)$ are :

A. $\frac{2E_1}{A_0}, \frac{2(E_1 - E_2)}{A_0}$

B. $\frac{2E_1}{A_0}, \frac{2E_2}{A_0}$

C. $\frac{(E_1 - E_2)}{A_0}, \frac{2E_2}{A_0}$

D. none of these

Answer: B



Watch Video Solution

43. Which of the following statements is correct?

- A. Mg^{+2} has greater ionic radius than Li^{+}
- B. Shielding constant value for us and $3d e^{-}$ is same in scandium
- C. H^{-} is the smallest anion in periodic table
- D. F has highest negative $\Delta H_{eg}(I)$




Answer: B



Watch Video Solution

44. Which of the following process is exothermic?

- A. $F^{-} \rightarrow F$







Answer: D



Watch Video Solution

45. Which of the following processes is exothermic?



Answer: A



Watch Video Solution

46. Arrange N,O and S in order of increasing electron affinity.

A. $S < O < N$

B. $N < S < O$

C. $N < O < S$

D. $O < N < S$

Answer: C



Watch Video Solution

47. Which of the following statements is incorrect?

A. ΔH_{eg} value of noble gases are negative

B. Si is diagonally related to B

C. The element which have higher $-\Delta H_{eg}$ act as strong oxidising agent.

D. H^+ and Cs^+ are the smallest and largest cation in periodic table respectively

Answer: A



Watch Video Solution

48. Which enthalpy is not associated with following change ?

Elemental chlorine $\rightarrow$ chloride ion

A. Bond dissociation enthalpy

B. First ionisation enthalpy

C. First electron gain enthalpy


D. None of the above

Answer: B



Watch Video Solution

49. Consider the electronic configuration of 2 elements:



Choose the incorrect option:




- A. (P) has more IE than (Q)
- B. (P) has more EA than (Q)
- C. Electron addition would be easier in (Q) than (P)
- D. Removal of electron would be easier in (Q) than (P)


Answer: B



Watch Video Solution

50. Which of the following would require least energy?









Answer: B



Watch Video Solution

E. Electronegativity

1. The electronic configuration of the outemost shell of the most electronegative element is



Answer: A



Watch Video Solution

2. Following the Mulliken scale, what parameters are required to evaluate electronegativity ?

- A. Only electronegativity
- B. Only electron affinity
- C. Electron affinity and ionization energy
- D. Ionic potential and electronegativity

Answer: C



Watch Video Solution

3. The electronegativity values of the elements are useful in predicting:

- A. bond energy of a molecule
- B. polarity of a molecule
- C. nature of an oxide
- D. all of the above

Answer: D



Watch Video Solution

4. In C , N , O and F the electronegativity

- A. increase from carbon to fluorine
- B. decrease from carbon to fluorine
- C. increase up to oxygen and is minimum at fluorine
- D. is minimum at nitrogen and then increase continuously

Answer: A



Watch Video Solution

5. If x , y and z are Mulliken electronegativity, ionisation potential and electron-affinity respectively. Then the electron affinity (z) in the terms of electronegativity (x) and ionisation potential (y) will be:

A. $z = \frac{x + y}{2}$

B. $z = \frac{x - y}{2}$

C. $z = \frac{x^2 - y^2}{2}$

D. $z = 2x - y$

Answer: D



Watch Video Solution

6. The *EN* of the elements *C*, *N*, *Si* and *P* increases in the order of _____ .

A. $N < P < C < Si$

B. $B > C > Si > P$

C. $N = P > C = Si$

D. $N > C > P > Si$

Answer: D

 [Watch Video Solution](#)

7. If electro negativity of x be 3.2 and that of y be 2.2 , the percentage ionic character of xy is

A. 19.5

B. 18.5

C. 9.5

D. 29.5

Answer: A

 [Watch Video Solution](#)

8. Which of the following relation is correct-

A. $2IP - EA - EN = 0$

B. $2EN - IP - EA = 0$

C. $2EA - IP - EA = 0$

D. $EN - IP - EA = 0$

Answer: B



Watch Video Solution

9. As one moves down the group from top to bottom then which of the following will not be observed?

A. Ionisation energy increases

B. Electron affinity decreases

C. Electronegativity decreases

D. Atomic radii increase

Answer: A



Watch Video Solution

10. Which is a true statement?

- A. Larger is the value of ionisation energy easier is the formation of cation
- B. Larger is the value of electron affinity easier is the formation of anion
- C. Larger is the value of ionisation energy as well as electron affinity the smaller is the electronegativity of atom
- D. Larger is the Z_{eff} larger is the size of atom

Answer: B



Watch Video Solution

11. The correct order of electronegativity is:

- A. $F > Cl > Br > I$
- B. $F > O > N > C$

C. $S < O < Se < Te$

D. All of these

Answer: A,B



Watch Video Solution

12. The electronegativity of the following elements increases in the order

A. C, B, Si, P

B. N, Si, C, P

C. Si, P, C, N

D. P, Si, N, C

Answer: C



Watch Video Solution

13. Fluorine is more electronegativity than nitrogen. The best explanation is that:

- A. the valency electrons in F are on the average, a little farther to the nucleus than in N
- B. the charge on a F nucleus is $+9$, while that on N nucleus is $+7$
- C. the nitrogen has half filled valence shell electron configuration, ns^2np^3 where as fluorine has partially filled electron configuration ns^2np^5
- D. the electronegativity decreases from top to bottom in each of the group as the effect nuclear charge remains constant.

Answer: B



Watch Video Solution

14. Consider the following statements.

(P) $IE(I)$ of nitrogen atom is ore than $IE(I)$ of oxygen atom.

(Q) Electron affinity of oxygen is less than sulphur atom.

(R) Electronegativity of an atom has no relation with its ionization enthalpy and electron gain enthalpy.

Which of the above statements are correct?

A. P,Q

B. P,R

C. Q,S

D. P,Q and R

Answer: A



Watch Video Solution

15. Choose the incorrect statement.

A. H^+ is the smallest size cation in the periodic table.

B. H^- is the smallest size anion in the periodic table,

C. F is the most electronegative element on the Pauling scale in the periodic table.

D. [He] has highest I.E. in the periodic table.

Answer: B



Watch Video Solution

16. With respect to oxygen maximum oxidation state is shown by:

A. halogen family

B. oxygen family

C. nitrogen family

D. boron family

Answer: A



Watch Video Solution

17. Which of the following statements is incorrect?






- A. The tendency to attract bonded pair of electron in case of hybrid orbitals follow the order: $sp > sp^2 > sp^3$
- B. Alkali metals generally have negative value of electron gain enthalpy
- C. The average oxidation number of S in $Na_2S_4O_6$ is greater than in the case of $Na_2S_2O_3$.
- D. The electronegativity values for 2p-series elements is less than that for 3p-series elements on account of small size and high inter electronic repulsions.

Answer: D



Watch Video Solution

18. The group state electronic configuration of some elements, P, Q, R, S and T (these symbols represent the some of the known elements given in the periodic table) are as follows. Itbr. (P)



Match the electronic configurations of the elements with the properties given below and select the correct sequence by choosing the correct codes given.

(i) Element forms a cation which is isoelectronic with P^{3-} .

(ii) Element which in its compounds can show a maximum oxidation state of $+6$ and that is coloured too.

(iii) Element has largest atomic radius and highest first ionisation energy in the respective period.

(iv) Element which has intermediate value of electronegativity and its oxide forms salts with strong acids and bases.

A. Q R T P

B. Q S T R

C. Q R S T

D. P Q R S

Answer: B



Watch Video Solution

19. In which case bond length is shortened?

A. When multiplication occurs between atoms

B. When electronegativities are different

C. In both cases

D. In none of the cases

Answer: C



Watch Video Solution

20. Which of the following is incorrect?

A. E.N. of $F > Cl$

B. E.A. of $S > O$

C. E.A of $O > Se$

D. Atomic radius of $Li > Be$

Answer: C



Watch Video Solution

21. Calculate the % ionic character for molecule AB when E.N. different is 2.0:

A. 46 %

B. 36 %

C. 30 %

D. 50 %

Answer: A



Watch Video Solution

22. A student went to meet his friend, where he saw that his friend was doing the study of a particular chemistry book. But he could not find the theoretical value of bond length in H-F but he found that r_H and r_F are $0.37\text{\AA}$ and $0.72\text{\AA}$ respectively and electronegativity of F and H are 4.0 and 2.1 respectively. what is bond length of H-F bond?

- A. 1.09
- B. 1.784
- C. 0.92
- D. 0.46

Answer: C



Watch Video Solution

23. Which of the following options is incorrect?

A. 1st ionisation potential : $Na < Mg < Si < P$

B. Electron affinity: $N < C < O < F < Cl$

C. Hydrated radius : $Ba^{2+}(aq) < Sr^{2+}(aq) < Mg^{2+}(aq)$

D. Polarity order: $N - H < Sb - H < As - H < P - H$

Answer: D



Watch Video Solution

24. Calculate the % ionic character for metal fluoride where the electronegativity of electropositive element is 2.1:

A. 43.03 %

B. 56.8 %

C. 50 %

D. 12 %

Answer: A



Watch Video Solution

25. The most predominantly ionic compounds will obtained from the combination of elements belonging to

A. I and VII

B. II and VI

C. IV and VIII

D. III and V

Answer: A



Watch Video Solution

26. The electronegativity of H and Cl are 2.1 and 3.0 respectively. The correct statement (s) about the nature of HCl is/are:

A. HCl is 17% ionic

B. 83% ionic

C. 50% ionic

D. 100% ionic

Answer: A



Watch Video Solution

27. Which of the following set has the strongest tendency to form anions?

A. Ga, In, Fe

B. Na, Mg, Al

C. N, O, F

D. V , Cr , Mn

Answer: C



Watch Video Solution

28. Which of the following is correct option?

(p) Same number of electron in d-subshell (Mn , Cr)

(q) Same value of Z_{eff} (Li^+ , B^+)

(r) Same value of σ , shielding constant (Li^+ , Be^{2+})

(s) Same value of electronegativity (P,H)

(t) Same value of electron affinity (F,Cl)

A. (p),(r),(t)

B. (r),(s),(t)

C. (p),(r),(s)

D. All are correct statement

Answer: C

[Watch Video Solution](#)

29. The correct relation given by Pauling to calculate electronegativity of an element X is (with respect to hydrogen). Given EN of $X > EN$ of H

A. $EN(X) - EN(H) = 0.208\sqrt{BE_{H-X} - \sqrt{BE_{X-X} \times BE_{H-H}}}$

B. $EN(X) - EN(H) = 0.102\sqrt{BE_{H-X} - \sqrt{BE_{X-X} \times BE_{H-H}}}$

C. $EN(X) - EN(H) = 0.102\sqrt{BE_{H-X} - \sqrt{BE_{X-X} \times BE_{H-H}}}$

where bond energies (BE) are in kJ/mole.

D. Both (a) and (c) are correct

Answer: C

[Watch Video Solution](#)

30. If $E_1^{\delta-} - E_2^{\delta+}$ bond has given polarity, which of the following is/are correct about polarity? (Electronegativity of $E_1 = 2.4$, $Q = 3.1$, $R = 2.6$ respectively)

A. $E_1^{\delta-} - Q^{\delta+}$

B. $Q^{\delta-} - R^{\delta+}$

C. $R^{\delta+} - E_2^{\delta-}$

D. $Q^{\delta+} - E_2^{\delta-}$

Answer: B



Watch Video Solution

31. % Ionic character in the covalent bond A-B is: (Given that

$X_A = 2, X_B = 3$)

A. 12.5 %

B. 30 %




C. 19.5 %

D. data is insufficient

Answer: C

[Watch Video Solution](#)

32. Which of the following does not represent the correct order of the properties indicated?



Answer: B

[Watch Video Solution](#)

33. Which element's electronegativity is closest to that of S?

A. O

B. P

C. Cl

D. Se

Answer: D



Watch Video Solution

34. Which atom has the highest electronegativity?

A. *Na*

B. *P*

C. *Cl*

D. *Br*

Answer: C



Watch Video Solution

35. Which family of elements has solid, liquid and gaseous members at 25°C and 1 atm pressure?

- A. Alkali metals ($\text{Li} - \text{Cs}$)
- B. Phinctogens ($\text{N} - \text{Bi}$)
- C. Chalcogens ($\text{O} - \text{Te}$)
- D. Halogens ($\text{F} - \text{I}$)

Answer: D



Watch Video Solution

36. In which molecule does the chlorine have the most positive partial charge?

- A. HCl
- B. BrCl
- C. OCl_2

D. SCI_2

Answer: C



Watch Video Solution

37. Electronegativities change both down a group and across a period. In general these changes are to:





- | | | |
|----|--------------------------|--|
| A. | Down a group
increase | Across a period from left to right
increase |
| B. | Down a group
increase | Across a period from left to right
decrease |
| C. | Down a group
decrease | Across a period from left to right
increase |
| D. | Down a group
decrease | Across a period from left to right
decrease |

Answer: C



Watch Video Solution

38. Which has the largest bond dissociation energy?



Answer: A



Watch Video Solution

39. Which of these properties increase across the period from N to Cl?

(P) Atomic radius

(Q) Density

(R) Electronegativity

A. P only

B. R only

C. P and Q only

D. Q and R only

Answer: D



Watch Video Solution

40. What is the correct order when the substances O_2 , H_2O , OF_2 and H_2O_2 are arranged in order of increasing oxidation number for oxygen?

A. O_2 , H_2O , OF_2 , H_2O_2

B. H_2O , H_2O_2 , O_2 , OF_2

C. H_2O_2 , O_2 , H_2O , OF_2

D. OF_2 , O_2 , H_2O_2 , H_2O

Answer: B



Watch Video Solution

41. Which list includes elements in order of increasing metallic character?

A. Si , P , S

B. As , P , N

C. Al , Ge , Sb

D. Br , Se , As

Answer: D



Watch Video Solution

42. Which bond is expected to be the least polar?

A. $O - F$

B. $P - F$

C. $Si - N$

D. $B - Cl$

Answer: A



Watch Video Solution

43. Which of these elements has the greatest electronegativity?

A. *Br*

B. *N*

C. *O*

D. *S*

Answer: C



Watch Video Solution

44. Select the group where *EN* increases down the group

A. *Ca, Sr, Ba*

B. Zn, Cd, Hg

C. F, Cl, Br

D. Li, Na, K

Answer: B



Watch Video Solution

45. Which of the following is incorrect?

- A. An element which has high electronegativity always has higher electron gain enthalpy
- B. Electron gain enthalpy is the property of an isolated gaseous atom
- C. Electronegativity is the property of bonded atom
- D. Both electronegativity and electron gain enthalpy are generally directly related to nuclear charge and inversely related to atomic size

Answer: A



Watch Video Solution

46. The pair of elements which on combination is most likely to form an ionic compound is:

A. Na and Ca

B. K and O_2

C. O_2 and CI_2

D. AI and I_2

Answer: B



Watch Video Solution

47. Two elements A and B have values of electronegativity respectively as 4 and 1.2, then the compound A-B will be:

- A. predominantly covalent
- B. predominantly ionic
- C. % covalent character less than 50%
- D. both (b) and (c)

Answer: D



Watch Video Solution

48. Identify the only incorrect statement.

- A. Fe^{3+} has greater electron affinity than Fe^{2+}
- B. Between F and Cl, formation of Cl^- is more exothermic than formation of F^-
- C. IE_1 of Cl^- = electron affinity of Cl
- D. Electronegativity depends only on electron affinity

Answer: D



[Watch Video Solution](#)

49. According to theory of electronegativity, most suitable name for F_2O molecule will be:

- A. oxygen fluoride
- B. difluorine oxide
- C. oxygen difluoride
- D. both (b) and (c)

Answer: C



[Watch Video Solution](#)

50. Define the general formula of non-polar compound on the basis of given information, where the electronic configuration of valence shell for A and B are as follows (a) ns^2np^3 (b) ns^2np^5 :

(P) AB_3 (Q) AB_4

(R) AB_5 (S) AB_6^-

A. P,Q,R

B. Q,R,S

C. R,S,P

D. S,P,Q

Answer: B




View Text Solution

51. Using the periodic table predict the formula of compound which might be formed by the following pair of elements: silicon and bromine

A. $SiBr_2$

B. $SiBr_4$

C. $Si_2Br_8^{4-}$





Answer: B



Watch Video Solution

52. In which of the following compound Mn shows maximum radius?



Answer: C



Watch Video Solution

53. Which one is not correct statement?

A. $IE(I)$ of He is maximum among all elements.

B. $E_{\alpha}(I)$ for noble gases is zero/positive

C. Electronegativity is maximum for fluorine

D. $IE(I)$ for nitrogen is less than that of oxygen.

Answer: D



Watch Video Solution

54. Which of the following is affected by the stable electron configuration of an atom?

(P) Electronegativity (Q) Ionisation energy (Q) Electron affinity

Correct answer is:

A. only electronegativity

B. only ionisation potential

C. electron affinity and ionisation energy both

D. all of the above





Answer: C



Watch Video Solution

F. Acidic-Basic Character, Hydration etc.

1. The order of basic character of given oxides is:



Answer: A



Watch Video Solution

2. Amphoteric behaviour is shown by the oxide of:

A. Al and Ca

B. Pb and Ba

C. Cr and Mg

D. Sn and Zn

Answer: D



Watch Video Solution

3. An element X occurs in short period having configuration ns^2np^1 . The formula and nature of its oxide is:

A. XO_3 , basic

B. XO_3 , acidic

C. X_2O_3 , amphoteric

D. X_2O_3 , basic

Answer: C

4. Aqueous solution of two compounds $M_1 - O - H$ and $M_2 - O - H$ are prepared in two different beakers. If electronegativity of $M_1 = 3.4$, $M_2 = 1.2$, $O = 3.5$ and $H = 2.1$, then the nature of two solution will be respectively

- A. acidix, basic
- B. acidic, acidic
- C. basic, acidic
- D. basic, basic

Answer: A

5. which one of the following oxides is neutral?

A. CO

B. SnO_2

C. ZnO

D. SiO_2

Answer: A



Watch Video Solution

6. The correct order of acidic strength is

A. $Cl_2O_7 > SO_3 > P_4O_{10}$

B. $CO_2 > N_2O_5 > SO_3$

C. $Na_2O > MgO > Al_2O_3$





D. $K_2O > CaO > MgO$

Answer: A



Watch Video Solution

7. Identify the correct order of acidic strength of CO_2 , CuO , CaO and H_2O .





Answer: A



Watch Video Solution

8. Which one of the following is an amphoteric oxide?








Answer: A



Watch Video Solution

9. Which of the following is the most acidic in nature?







Answer: D



Watch Video Solution

10. Among Al_2O_3 , SiO_2 , P_2O_3 and SO_2 the correct order of acidic strength is:






Answer: D



Watch Video Solution

11. Which of the following oxides is amphoteric in nature?



Answer: D



Watch Video Solution

12. In which of the following arrangements, the sequence is not strictly according to the property written against it?

A. $HF < HCl < HBr < HI$: increasing acid strength

B. $NH_3 < PH_3 < AsH_3 < SbH_3$: increasing basic strength

C. $B < C < O < N$: increasing first ionization enthalpy



D. $CO_2 < SiO_2 < SnO_2 < PbO_2$: increasing oxidising power

Answer: B



View Text Solution

13. Find the incorrect acidic strength order:







Answer: B



Watch Video Solution

14. The correct order of acidic nature of the oxides of chlorine is







Answer: A



Watch Video Solution

15. The most acidic oxide is:







Answer: C



Watch Video Solution

16. The correct order of basic nature is







Answer: A



Watch Video Solution

17. Select the amphoteric oxide.



Answer: B



Watch Video Solution

18. Select the correct statement.

A. E.N. of atom X is 2.5 on Pauling scale if IE of X is $2eV$ EA of X is $3eV$

B. $|IE \text{ of } X(g)| = |EA \text{ of } X(g)^-|$

C. $F^-(aq) < Cl^-(aq) < Br^-(aq) < I^-(aq)$ (ionic mobility)

D. $Li^+(aq) < Na^+(aq) < K^+(aq) < Rb^+(aq)$ (size)

Answer: C



Watch Video Solution

19. Which of the following option is incorrect

A.

<i>List – I</i>	<i>List – II</i>
Ionisation potential	$Na < Mg < Si < P$

B.

<i>List – I</i>	<i>List – II</i>
Electron affinity	$N < C < O < F < Cl$

C.

<i>List – I</i>	<i>List – II</i>
Hydrated radius	$Ba^{2+}(aq) < Sr^{2+}(aq) < Mg^{2+}(aq)$

D.


<i>List – I</i>	<i>List – II</i>
Polarity order	$N - H < Sb - H < As - H < P - H$

Answer: D



Watch Video Solution

20. What property of the oxygen atom is represented by the equation



- A. Electronegativity
- B. First electron affinity
- C. First ionization energy
- D. Lattice energy

Answer: B



Watch Video Solution

21. Which element forms a compound with the formula H_3XO_4 ?

- A. As
- B. Cl
- C. N

D. S

Answer: A



Watch Video Solution

22. Which forms the most alkaline solution when added to water?

A. Al_2O_3

B. B_2O_3

C. CO_2

D. SiO_2

Answer: A



Watch Video Solution

23. The oxide of which element is the most ionic?

A. Al

B. B

C. C

D. Si

Answer: A



Watch Video Solution

24. The incorrect statement among the following is:

A. density increases across the period from left to right while decreases down the group

B. ionization energy depends upon the type of orbital (of same energy level) from which electron is being removed

C. generally electron affinity decreases down the group

D. moving diagonally, the charge to size ratio remains nearly same for
2 and 3rd period elements upto 14th group.

Answer: A



Watch Video Solution

25. Which of these compounds is amphoteric ?

(P) $Al(OH)_2$ (Q) $Ba(OH)_2$

(R) $Zn(OH)_2$

A. P only

B. Q only

C. P and R only

D. Q and R only

Answer: C



Watch Video Solution

26. Which oxide produces the most acidic solution when 0.1 mol is added to 1L of H_2O ?

A. BaO

B. BaO_2

C. SO_2

D. SO_3

Answer: D



Watch Video Solution

27. Element P, Q, R and S belong to the same group. The oxide of P is acidic, oxide of Q and R are amphoteric while the oxide of S is basic. Which of the following elements is the most electropositive ?

A. R

B. Q

C. P

D. S

Answer: C



Watch Video Solution

28. Increasing order of basic character of NO_2 , K_2O and ZnO is

A. $NO_2 < ZnO < K_2O$

B. $K_2O < ZnO < NO_2$

C. $NO_2 < K_2O < ZnO$

D. $K_2O < NO_2 < ZnO$

Answer: A



Watch Video Solution

29. Which of the following options is correct regarding periodic properties?

A. Ionic radii order $Cl^- > H^-$

B. Electropositivity order of elements $Li > Na > K > Rb$

C. Z_{eff} order $Li < Na < K < Rb$

D. Hydration energy order $Fe^{+3} > Fe^{+2} > Fe^{+4}$

Answer: D



Watch Video Solution

30. Condition suitable for high solubility of ionic compound in solvent should be:

A. High lattice energy and low hydration energy

B. Low lattice energy and high hydration energy

C. Lattice energy must be equal to hydration energy


D. Solubility is independent of lattice or hydration energy

Answer: B



Watch Video Solution

31. Find out the value of (P) in the following cycle:



Given

$$\Delta H_1 = -400 \text{ kJ}, \text{I.E.} = +50 \text{ kJ}, D = +50 \text{ kJ}, \Delta H_{\text{eg}} = -30 \text{ kJ}$$

lattice energy = -100 kJ

A. -400 kJ

B. -300 kJ

C. -370 kJ




D. -310kJ

Answer: C



Watch Video Solution

32. Which terms are exothermic for the formation of $\text{NaF}(s)$?



A. P only

B. Q only

C. P and R only

D. Q and R only

Answer: D



Watch Video Solution

33. In order to calculate the lattice energy of $NaCl$ using a Born-Haber cycle, which value is not needed?

- A. enthalpy of sublimation of $Na(s)$
- B. first ionization energy of $Cl(g)$
- C. bond dissociation energy of $Cl_2(g)$
- D. enthalpy of formation of $NaCl(s)$


Answer: B



Watch Video Solution

34. Which of the following represent correct of hydration energy released in aqueous state?

- A. $N^{3-} > O^{2-} > F^{-}$
- B. $F^{-} < Cl^{-} < Br^{-} < I^{-}$
- C. $Li^{+} < K^{+} < Cs^{+}$



Answer: D



Watch Video Solution

35. The following order are given along with their mentioned properties:

(P) $Mg < Na < Sr < Ba$ [order of atomic size]

(Q) $M^{+} > M^{+2} > M^{+3}$ [M = hypothetical element, order of IP_1 of respective ion]

(R) $F > Cl > Br > I$ [order of electronegativity]

(S) $Al_2O_3 < P_2O_5 < SO_3 < Cl_2O_7$ [order of basic nature]

(T) $F^{-} > Cl^{-} > Br > I^{-}$ [order of ionic mobility in aqueous solution]

Which option represents group incorrect order?

A. P,R,T

B. Q,R,T

C. Q,R,S

D. R,S,T

Answer: D



View Text Solution

36. Which of the following order is/are correct ?

A. $Li^+ < Na^+ < K^+$: Hydrated radius

B. $Na^+ < Mg^{+2} < Al^{+3}$: Ionic radius

C. $F < Cl > Br$: Electron affinity

D. $Cl - OH < Br - OH < I - OH$: Acidic character

Answer: C



Watch Video Solution

37. The following acid have arrange in the order of decreasing strength.

Identify the correct order. $\text{ClOH(I)}\text{BrOH(II)}\text{IOH(III)}$

A. $P > Q > R$

B. $Q > P > R$

C. $R > Q > P$

D. $P > R > Q$

Answer: A



Watch Video Solution

38. Which of the following is/are correct statement(s)?

A. Electron affinity of oxygen is more than that of sulphur

B. If spin quantum number was assigned three different values, the

6th period would have maximum 48 elements

- C. Acidic nature of oxides of group XIII elements increases down the group
- D. All of the above

Answer: B



Watch Video Solution

39. Which of the following is an amphoteric oxide?

- A. CO
- B. NO
- C. BeO
- D. N_2O_3

Answer: C



Watch Video Solution

40. Which is an amphoteric oxides?

A. BeO

B. SnO

C. ZnO

D. All of these

Answer: D



Watch Video Solution

41. The order of basic strength of given oxide:

A. $\text{Na}_2\text{O} > \text{MgO} > \text{Al}_2\text{O}_3 > \text{CuO}$

B. $\text{MgO} > \text{Al}_2\text{O}_3 > \text{CuO} > \text{Na}_2\text{O}$

C. $\text{Al}_2\text{O}_3 > \text{MgO} > \text{CuO} > \text{Na}_2\text{O}$

D. $\text{CuO} > \text{Na}_2\text{O} > \text{MgO} > \text{Al}_2\text{O}_3$

Answer: A



Watch Video Solution

42. Which of the following is neutral oxide?

A. SnO

B. PbO

C. PbO_2

D. N_2O

Answer: D



Watch Video Solution

43. The high oxidising power of fluorine is due to :

A. high electron affinity

B. high ionisation energy

C. both (a) and (b)

D. none of the above

Answer: A



Watch Video Solution

44. The correct statement is:

A. F has more electron affinity than $O^{\oplus}$

B. $Al^{3\oplus}$ ion has more hydration energy than $Mg^{2\oplus}$ ion

C. noble gases have positive electron affinity

D. all of the above

Answer: B



Watch Video Solution

45. A,B and C are oxides of elements X,Y and Z respectively. X,Y and Z are in the same period of the periodic table. A gives an aqueous solution which turns blue litmus red. B reacts with both strong acids and strong alkalies. C gives an aqueous solution which is strongly alkaline. Which of the following statement is/are true ?

(P) All the three elements are metals.

(Q) The electronegativities decreases from X to Y to Z.

(R) The atomic radius increases in the order $X < Y < Z$.

(S) X,Y and Z could be phosphorus, aluminium and sodium respectively.

A. P,Q,R only correct

B. P,R only correct

C. Q,S only correct

D. Q,R,S only correct

Answer: D



Watch Video Solution

1. Assertion: Electron gain enthalpy value of the 3rd period p-block elements of the modern periodic table are generally more negative than the 2nd period element of the same group.

Reason: Due to smaller atomic size of the 2nd period element, its electron density is high which eases the addition of electron.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

2. Statement-1: 'He' has lowest ionisation energy among all the elements.

Statement-2: Electron affinity values of metals are generally low while those of non-metals are high.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: D



Watch Video Solution

3. Statement-1: In a period, noble gas has largest atomic radius.

Statement-2: In case of noble gases van der Waal's radius is defined and

there is much inter electronic repulsions.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-3
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-3
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

4. Statement-1: The 5th period of periodic table contains 18 elements are 32.

Statement-1: $n = 5, l = 0, 1, 2, 3$. The order in which the energy of available orbitals $4d, 5s$ and $5p$ increases is $5s < 4d < 5p$ and the total

number of orbitals available are 9 and thus 18 electrons can be accommodated.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-4
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-4
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

5. Assertion: the $4f^-$ and $5f^-$ inner transition series of elements are placed separately at the bottom of the modern periodic table.

Reason : (i) Position of f-block elements prevents the undue expansion of the modern periodic table, ie, maintains its structure.

(ii) Position of f-block elements preserves the principle of classification by keeping elements with similar properties in a single column.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-5
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-5
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

6. Statement-1: Manganese (atomic number 25) a less favourable electron affinity than its neighbours on either side.

Statement-2: The magnitude of an element's electron affinity only depends on the element's valence shell electronic configuration.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-6
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-6
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

7. Assertion: In general, for an element, $IE_1 < IE_2 < IE_3 \dots$

Reason: After the removal of each successive electron remaining electrons are held more tightly by the nucleus so removal of next electron becomes difficult.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-7

- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-7
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

8. Statement-1: Second electron gain enthalpy of halogens is always positive.

Statement-2: Fluorine has most negative electron gain enthalpy.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-8
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-8
- C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

9. Statement-1: Third ionisation energy of phosphorous is larger than sulphur.

Statement-2: There is a larger amount of stability associated with filled s- and p-sub-shells (a noble gas electron configuration) which corresponds to having eight electrons in the valence shell of an atom or ion.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-9

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-9

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: D



Watch Video Solution

10. Statement-1: First ionisation energy of $O^+(g)$ is greater than that of $N(g)$ whereas vice versa is true for the first ionisation energy of $N^+(g)$ and $N(g)$ respectively.

Statement-2: Electronic configuration of $N(g)$ is more stable than that of $N^+(g)$.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: D



[Watch Video Solution](#)

11. Statement-1: Down the group, electronegativity decreases from B to Tl.

Statement-2: Electronegativity of an element depends on its electron affinity and ionisation energy.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: B



[Watch Video Solution](#)

12. Statement-1: $Sc(Z = 21)$ is placed as d-block element.

Statement-2: Last filling electron goes into 3d-subshell.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-12

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-12

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

13. Statement-1: Na^+ and Al^{3+} are isoelectronic but the magnitude of ionic radius of Al^{3+} is less than that of Na^+ .

Statement-2: The magnitude of effective nuclear charge of the outer most shell electrons in Al^{3+} is greater than that of Na^{+} .

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

14. Statement-1: The atomic radii of the elements of the oxygen family are smaller than the atomic radii of the corresponding elements of the nitrogen family.

Statement-2: The members of the oxygen family are more electronegative

and thus have lower values of nuclear charge than those of the nitrogen family.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

15. Statement-1: Metallic character of first group metals increases down the group with the decreasing ionisation energy.

Statement-2: The loss and gain of electron(s) can be used in explaining the reducing and oxidising behaviour of the element respectively.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

16. Statement-1: Fluorine has a less negative electron gain enthalpy than chlorine.

Statement-2: There is relative greater effectivebess of 2p-electrons in the small F atom to repel the additional electron entering the atom than to 3p-electrons in the large Cl atom.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-16
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-16
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

17. Statement-1: Fluorine forms only one oxoacid, HOF,
Statement-2: Fluorine has small size and high electronegativity.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-17

- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-17
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

18. Statement-1: Li and Mg shows diagonal relationship.

Statement-2: Li and Mg are diagonal to each other in the periodic table.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-18
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-18
- C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

19. Statement-1: The decreasing order of acidic character of CO_2 , N_2O_5 , SiO_2 and SO_3 is $SO_3 > N_2O_5 > CO_2 > SiO_2$.

Statement-2: Oxides of the metals are generally basic and oxides of the non-metals are acidic.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-1

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

20. Statement-1: Oxidising property of F_2 is greater than Cl_2 .

Statement-2: $(\Delta H)_{eg}$ (electron gain enthalpy) of F is greater than Cl.

Screen reader support enabled.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

21. Statement-1: Generally, ionisation enthalpy increases from left to right in a period.

Statement-2: When successive electrons are added to the orbitals in the same principal quantum level, the shielding effect of inner core of electrons does not increase very much to compensate for the increased attraction of the electron to the nucleus.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

22. Statement-1: Boron has a smaller first ionisation enthalpy than beryllium.

Statement-2: The penetration of a 2s electron to the nucleus is more than the 2p electron hence 2p electron is more shielded by the inner core of electrons than the 2s electrons.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

23. Assertion: Electron gain enthalpy always becomes less negative as we go down a group in Modern periodic table.

Reason: The size of the atom increase on going down the group in Modern periodic table and the added electron would be farther from the nucleus.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-23

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-23

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: D



Watch Video Solution

24. Assertion: The first ionisation energy of Be is greater than that of B .

Reason: 2p-orbital is lower in energy than 2s-orbital.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-24

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-24

C. Statement-1 is True, Statement-2 is False


D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

25. Statement-1: The acidic strength order of hydro acids is



Statement-2: The E.N. of halogens are $F > Cl > Br > I$.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

26. Statement-1: The decreasing order of acidic character of CO_2 , N_2O_5 , SiO_2 and SO_3 is $SO_3 > N_2O_5 > CO_2 > SiO_2$.

Statement-2: As electronegativity difference (E-O) decreases, acidic character of the oxide increases.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2

- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A



Watch Video Solution

27. Statement-1: The ionisation potential of Sn is greater than Pb.

Statement-2: Usually, ionisation energy decreases down the group.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2
- B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2
- C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: D



Watch Video Solution

28. Statement-1: $(I. E.)_n$ of an atom is always greater than $(I. E.)_{n-1}$ ($n \in I^+$)

Statement-2: ne/Z ratio decreases on successive elimination of electrons.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: A



[Watch Video Solution](#)

29. Statement-1: Nitrogen atom has higher ionization energy than fluorine atom.

Statement-2: Nitrogen atom has extra stable electronic configuration due to half filled p-subshell.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-2

B. Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: D



[Watch Video Solution](#)

1. Which of the following statements is/are true about the modern form of periodic table?





- A. Third group of periodic table accommodates maximum number of elements.
- B. Due to presence of half filled and fully filled sub shells in electronic configuration electronegativity of atom increases
- C. The element of 13 th group and 7th period will have atomic number 113.
- D. Diagonal relationship in 2nd and 3rd period element is found due to similar charge to size ratio

Answer: A::C::D



Watch Video Solution

2. Which of the following orders is correct for the size ?






Answer: A::B::D



Watch Video Solution

3. In which of the following arrangements, the order is not correct according to the property indicated against it.



Answer: B::D



Watch Video Solution

4. Which of the following statement(s) is/are correct?

- A. Ionization energy may be negative for some elements
- B. Second electron gain enthalpy always remains positive for all the elements
- C. Negative value of electron gain enthalpy of fluorine is minimum in its group
- D. Ionization energy of Ga is slightly more than Al

Answer: B::D



Watch Video Solution

5. Ionisation energy of atoms A and B are 350 and 250kcalmol^{-1} respectively. The electron affinities of these atoms are 70kcalmol^{-1} and 90kcalmol^{-1} respectively then:

- A. Electron cloud is more attracted by A
- B. Electron cloud is more attracted by B
- C. Electronegativity of A is more than B
- D. Electronegativity of A is less than B

Answer: A::C



Watch Video Solution

6. Which of the following statements is/are true for the long form of the periodic table?

- A. it reflects the sequence of filling the electrons in the order of sub energy level, s,p,d and f

- B. it helps to predict the stable valency states of the elements
- C. it reflects trends in physical and chemical properties of elements
- D. it helps to predict the relative ionicity of the bond between any two elements

Answer: A::B::C



Watch Video Solution

7. Which of the following statements is/are true about the modern form of periodic table?

- A. Properties of the elements are the periodic functions of their atomic number
- B. There are 7 periods
- C. There are 8 groups
- D. It has separate positions for the isotopes

Answer: A::B



Watch Video Solution

8. The factors that influence the ionisation energies are:

- A. the size of the atom
- B. the charge on the nucleus
- C. how effectively the inner electron shell screen the nuclear charge
- D. stability of electronic configuration

Answer: A::B::C::D



Watch Video Solution

9. Which of the following statement(s) is/are correct?

- A. The second ionization energy of sulphur is greater than that of chlorine
- B. The third ionization energy of phosphorus is greater than that of aluminium
- C. The first ionization energy of aluminium is nearly the same as that of gallium
- D. The second ionization energy of boron is greater than that of carbon.

Answer: A::B::C::D



Watch Video Solution

10. Ionization energy of an element is:

- A. equal in magnitude but opposite in sign to the electron gain enthalpy of the cation of the element

- B. same as electrin affinity of the element
- C. energy required to remove one valence electron from an isolated gaseous atom in its ground state
- D. equal in magnitude but opposite in sign to the electron gain enthalpy of the anion of the element

Answer: A::C



Watch Video Solution

11. The properties which are common to both groups 1 and 17 elements in the periodic table are:

- A. metallic character increases down the groups
- B. reactivity decreases from top to bottom in groups
- C. atomic radii increases as the atomic number increases
- D. electronegativity decreases on moving down a group

Answer: A::C::D



Watch Video Solution

12. Select the incorrect statement(s).

- A. $IE(I)$ of nitrogen atom is less than $IE(I)$ of oxygen atom.
- B. Electron affinity of oxygen is less than selenium atom.
- C. Electronegativity on Pauling scale is 2.8 times greater than electronegativity on Muliken scale.
- D. The Cr^{6+} is smaller than Cr^{3+}

Answer: A::C



Watch Video Solution

13. Which of the following order(s) is/are correct?

A. $O^{2-} < F^{-} < Na^{+} < Mg^{2+}$ increasing $Z_{\text{effective}}$

B. $Mg^{2+} < Na^{+} < O^{2-} < F^{-}$ increasing size

C. $O^{2-} < F^{-} < Na^{+} < Mg^{2+}$ increasing ionisation energy

D. $O^{2-} < F^{-} < Na^{+} < Mg^{2+}$ increasing electron affinity

Answer: A::C::D



Watch Video Solution

14. Which of the following is/are true order (s)?

A. $B^{+} < B < B^{-}$ Size

B. $I < Br < Cl < F$ Electron affinity

C. $O^{2-} < O^{-} < O^{+}$ $Z_{\text{effective}}$

D. $Na < Al < Mg < Si$ Ionisation potential

Answer: A::C::D



Watch Video Solution

15. Poor shielding of nuclear charge by d or f-orbital elements is responsible for which of the following facts?


- A. Atomic radius of Nb(4-d series) is comparable to that of Ta(5-d series)
- B. The 1st ionisation energy of copper is less than that of zinc
- C. The value of electron gain enthalpy is more negative for sulphur than for oxygen.
- D. The 1st ionisation energy for gold is greater than that of silver.



Answer: A::D



Watch Video Solution

16. Select equations having endothermic step:








Answer: A::B::C::D



Watch Video Solution

17. Which is not correct order for the properties specified ?



Answer: A::C



Watch Video Solution

18. Select the correct statement(s).

- A. The normal oxide formed by the element on extreme left is the most basic
- B. The tendency of group IA (i.e., 1st group) to form oxygen rich compounds increases from top to bottom
- C. Oxides of metals are called as basic anhydrides
- D. In general metallic oxides (O^{2-}) peroxides (O_2^{2-}) and super oxides (O_2^-) are ionic

Answer: A::B::C::D



Watch Video Solution

19. Degenerate orbital have:

A. n same

B. l same

C. m different

D. s different

Answer: A::B::C



Watch Video Solution

20. Which of the following pairs were incorrectly arranged in Mendeleev's Periodic table?

A. $Ar - K$

B. $Te - I$

C. $Th - Pa$

D. $Co - Ni$

Answer: A::B::C::D

21. Which option is correct for atomic radius?

A. $Ne > F$

B. $Ar > Cl$

C. $F > Ne$

D. $Cl > Ar$

Answer: A::B

22. Choose the correct order of ionic size:

A. $Na^+ > Mg^{2+} > Al^{3+}$

B. $Na^+ < Mg^{+2} < Al^{3+}$

C. $N^{3-} > O^{2-} > F^-$



Answer: A::C



Watch Video Solution

23. The ionic radii depends upon the following factors:

A. charge of cation

B. charge of anion

C. shell number of valence shell electron(s) of the ion

D. number of atoms for multiatomic ion

Answer: A::B::C::D



Watch Video Solution

24. Which of the following is correct order of EA.

A. $N < C < O < F$

B. $F > Cl > Br > I$

C. $Cl > F > Br > I$

D. $C < N < O < F$

Answer: A::C



Watch Video Solution

25. Which of the following is/are correct?

A. For $A(g) + e^- \rightarrow A^-(g) \Delta H$ may be negative

B. For $A^-(g) + e^- \rightarrow A^{2-}(g) \Delta H$ may be negative

C. For $A^-(g) + e^- \rightarrow A^{2-}(g) \Delta H$ must be positive

D. For $A^{+3}(g) + e^- \rightarrow A^{+2}(g) \Delta H$ must be negative

Answer: A::C::D



Watch Video Solution

26. According to Slater's rule, which of the following has correct order of Z_{eff} ?

A. $Na > K$

B. $Li > Na$

C. $Na < Mg$

D. $Li < Be$

Answer: C::D



Watch Video Solution

27. Which of the following is/are correctly matched?

A. $E > O$ (Electronegativity)

B. $Mg > Al$ (1st ionisation energy)

C. $HCl > HI$ (Thermal stability)

D. Diamond > Graphite (Electrical conductance)

Answer: A::B::C



Watch Video Solution

28. Which of the following statements is/are incorrect ?

A. Ionisation energy of A^- is greater than A when A is a halogen atom.

B. Ionisation energy of A^+ is greater than that of A^{2+} when A is the number of alkali metals

C. Successive ionisation energy is always increasing for 1st and 2nd period element

D. Electron affinity value of A^+ is numerically identical with the ionisation potential of A^- [for any atom]

Answer: A::B::D



[Watch Video Solution](#)

29. Mendeleev left the space for elements in periodic table for the elements:

A. *Ga*

B. *Sc*

C. *Ge*

D. *Tc*

Answer: A::B::C::D



[Watch Video Solution](#)

30. Which of the following properties are the properties of metal?

A. They are sonorous

B. They are in general poor conductor of heat and electricity

C. They are malleable and ductile

D. They are hard

Answer: A::C::D



Watch Video Solution

31. For which of the following species the contribution of s electron to the shielding constant is 0.3 ?

A. H^+

B. He

C. Li^+

D. H^-

Answer: B::C::D



Watch Video Solution

32. Which of the following represents the correct order of the properties indicated ?

A. $F > O > N > C$ (Electronegativity)

B. $Sc^{3+} > Ti^{4+} > K^{+}$ (Size)

C. $Ni^{2+} < Co^{2+} < Fe^{2+} < Mn^{2+}$ (Unpaired electron)

D. $HNO_3 > HNO_2$ (Acidic strength order)

Answer: A::C::D



Watch Video Solution

33. Select the element (s) which belongs to d-block but not transition elements:

A. Zn

B. Cd

C. Hg

D. P^+

Answer: A::B::C



Watch Video Solution

34. The species having 4 or more unpaired electrons are:

A. Fe^{2+}

B. Co^{3+}

C. Mn^{2+}

D. Ni^{2+}

Answer: A::B::C



Watch Video Solution

35. Which option is/are correct?

- A. Atomic radius decreases with increase in Z_{eff}
- B. The atomic number of 50 elements is present in 5th period
- C. Alkaline earth elements are IA group elements
- D. Alkali metals are IIA group elements

Answer: A::B



Watch Video Solution

36. The elements $Z = 116$ and 119 have been discovered. In which family/group, would you place these elements and also give the electronic configuration in each case?

- A. $Z = 116$, oxygen family, configuration = $[Rn]5f^{14}6d^{10}7s^27p^4$
- B. $Z = 119$, alkali metal, configuration = $[Uuo]8s^1$
- C. $Z = 116$, halogen family, configuration = $[Rn]5f^{14}6d^{10}7s6(2)7p^4$
- D. $Z = 119$, alkali metal , configuration = $[Uuo]8s^2$

Answer: A::B



Watch Video Solution

37. Those elements impart colour to the flame on heating in it, the atoms of which require low energy for the ionisation (i.e., absorb energy in the visible region of spectrum). The elements of which of the following groups will impart colour to the flame?

A. 2

B. 13

C. 1

D. 17

Answer: A::C



Watch Video Solution

38. Which of the following sequences contain atomic numbers of only representative elements?

A. 3,33,53,87

B. 2,10,22,36

C. 7,17,25,37,48

D. 9,35,51,88

Answer: A::D



Watch Video Solution

39. Which of the following elements will gain one electron more readily in comparison to other elements of their group?

A. $S(g)$

B. $Na(g)$

C. $O(g)$

D. $CI(g)$

Answer: A::D



Watch Video Solution

40. Which of the following statements are correct?





- A. Helium has the highest first ionisation enthalpy in the periodic table
- B. Chlorine has less negative electron gain enthalpy than fluorine
- C. Mercury and bromine are liquids at room temperature
- D. In any period, atomic radius of alkali metal is the highest

Answer: A::C::D



Watch Video Solution

41. Which of the following sets contain only isoelectronic ions?







Answer: B::C



Watch Video Solution

42. In which of the following options order of arrangement does not agree with the variation of property indicated against it?



Answer: B::C



Watch Video Solution

43. Which of the following have no unit?

- A. Electrongativity
- B. Electron gain enthalpy
- C. Ionisation enthalpy
- D. Metallic character

Answer: A::D



Watch Video Solution

44. Ionic radii vary in

- A. inverse proportional to the effective nuclear charge

- B. inverse proportional in the square of effective nuclear charge
- C. direct proportional to the screening effect
- D. direct proportional to the square of screening effect

Answer: A::C



Watch Video Solution

45. An element belongs to 3rd period and group-13 of the periodic table.

Which of the following properties will be shown by the element?

- A. Good, conductor of electricity
- B. Liquid, metallic
- C. Solid, metallic
- D. Solid, non metallic

Answer: A::C



Watch Video Solution

46. Which of the following statements are correct?

- A. F is the most electronegative and Cs is the most electropositive element
- B. The EN of halogens decreases from F to I
- C. The E.A of Cl is higher than that of F through their EN values are in the reverse order
- D. The E.A of noble gases is low

Answer: A::B::C::D



Watch Video Solution

47. Which of the following statements (s)is/are true?

- A. Ionisation energy $\propto \frac{1}{\text{Screening effect}}$

- B. The first ionisation energies of Be and Mg are more than ionisation energies of B and Al respectively
- C. Atomic and ionic radii of niobium and tantalum are almost same
- D. Metallic and covalent radii of potassium are $2.3\text{\AA}$ and $2.03\text{\AA}$ respectively.

Answer: A::B::C::D



Watch Video Solution

48. Which of the following statement(s) is/are true for IA elements?





- A. Na is regarded as a typical/representative element
- B. Basic character of their oxides increases down the group
- C. Among alkali metal ions, degree of hydration is highest in Li
- D. In general, electron affinity values decreases from top to bottom in the group

Answer: A::B::C::D



Watch Video Solution

49. The process(es) requiring the absorption of energy is/are:



Answer: B::D



Watch Video Solution

50. Predict block of element whose outer electron set of quantum number n, l, m is 6,0,0 respectively:

A. s-block

B. p-block

C. d-block

D. f-block

Answer: A::C::D



Watch Video Solution

51. Which of the following are neutral oxide?

A. N_2O

B. NO

C. CO

D. H_2O

Answer: A::B::C::D



Watch Video Solution

52. Which of the following statement(s) is/are correct?

- A. Th' is transuranic element
- B. He' has highest first ionization energy among all elements
- C. Li' is highest metal
- D. F' has positive second electron gain enthalpy

Answer: B::C::D



Watch Video Solution

53. In which of the following pairs, the first atom is larger than the second ?

- A. *Br, Cl*
- B. *Na, Mg*
- C. *Sr, Ca*

D. N, P

Answer: A::B::C



Watch Video Solution

54. Among the given elements, select those, which are not at the peak of Lothar Meyer's curves?

A. La

B. K

C. Cl

D. Xe

Answer: A::C::D



Watch Video Solution

55. An element has electronic configuration as $1s^2, 2s^2 2p^6, 3s^2 3p^6, 4s^2, 3d^7$.

Correct statement regarding element is:

- A. Element must be Co
- B. Element belongs to group number '9' according to the long form of periodic table
- C. Maximum number of electrons in element having $m = +1$ are 6
- D. Element has magnetic moment $\sqrt{15}BM$

Answer: A::B::C::D




Watch Video Solution

56. The correct order of radii is:

A. $N < Be < B$

B. $F^- < O^{2-} < N^{3-}$



Answer: B::C::D



Watch Video Solution

57. Choose the correct statement.

A. Be and Al are not in same group

B. All the transition metals correspond to d-block

C. Be and Al are having lot of similarities in their properties

D. The atomic radius gradually decreases from Sc to Zn.

Answer: A::B::C



Watch Video Solution

58. Which of the following pair of elements are not of same group of periodic table?

A. *Li*, *Na*

B. *Be*, *B*

C. *N*, *As*

D. *O*, *At*

Answer: B::D



Watch Video Solution

59. Which of the following statements are incorrect?

A. The elements after thorium are called transuranium elements

B. $P < Si < Be < Mg < Na$, the order of increasing metallic character

C. The f-block elements are called transition elements





D. Zn , Cd and Hg are called transition elements

Answer: A::C::D



Watch Video Solution

60. Which of the following processes involve absorption of energy?



Answer: B::C::D



Watch Video Solution

61. In which of the following arrangements the order which is correct according to the property indicated against it.

A. $Al^{+3} < Mg^{+2} < Na^{+} < F^{-}$: Increasing ionic size

B. $B < C < N < O$: Increasing first ionisation enthalpy

C. $I < Br < F < Cl$: Increasing electron gain enthalpy (magnitude only)

D. $Li < Na < K < Rb$: Increasing metallic radius

Answer: A::C::D



Watch Video Solution

62. which of the following order of ionisation energy is correct?

A. $Li > Be(III. E.)$

B. $N > O(III. E.)$

C. $Be < B(III. E.)$

D. $F < O(III. E.)$

Answer: A::C::D



Watch Video Solution

63. Which of the following statements is/are correct?

A. Elements Ce to Lu are called lanthanides

B. $P < Si < Be < Mg < Na$ in the increasing order of metallic character

C. Mg has the largest size and Al^{3+} has smallest size among Mg, Mg^{+2}, Al, Al^{+3}





D. Among P, S, Cl, F , phosphorous has least negative electron gain enthalpy.

Answer: A::B::C::D



Watch Video Solution

64. Which of the following are endothermic process?



Answer: A::B::C::D







Watch Video Solution

Comprehension 1

1. The periodicity is related to the electronic configuration. That is, all chemical and physical properties are a manifestation of the electronic configuration of the elements. The atomic and ionic radii generally decrease in a period from left to right. As a consequence, the ionization

enthalpies generally increase and electron gain enthalpies become more negative across a period. In other words, the ionization enthalpy of the extreme left element in a period is the least and the electron gain enthalpy of the element on the extreme right is the highest negative. This results into high chemical reactivity at the two extremes and the lowest in the centre. Similarly, down the group, the increase in ionization enthalpies and a regular decrease (with exception in some third period elements) in electron gain enthalpies in the case of main group elements. The loss and gain of electrons can be co-related with the reducing and oxidising behaviour, and also with metallic and non-metallic character respectively of the elements.

The correct order of the metallic character is:



Answer: A



Comprehension

1. The periodicity is related to the electronic configuration. That is, all chemical and physical properties are a manifestation of the electronic configuration of the elements. The atomic and ionic radii generally decrease in a period from left to right. As a consequence, the ionization enthalpies generally increase and electron gain enthalpies become more negative across a period. In other words, the ionization enthalpy of the extreme left element in a period is the least and the electron gain enthalpy of the element on the extreme right is the highest negative. This results into high chemical reactivity at the two extremes and the lowest in the centre. Similarly, down the group, the increase in ionization enthalpies and a regular decrease (with exception in some third period elements) in electron gain enthalpies in the case of main group elements. The loss and gain of electrons can be co-related with the reducing and oxidising behaviour, and also with metallic and non-metallic character

respectively of the elements.

The correct order of the non-metallic character is:

A. $B > C > Si > N > F$

B. $Si > C > B > N > F$

C. $F > N > C > B > Si$

D. $F > N > C > Si > B$

Answer: C




Watch Video Solution

2. The periodicity is related to the electronic configuration. That is, all chemical and physical properties are a manifestation of the electronic configuration of the elements. The atomic and ionic radii generally decrease in a period from left to right. As a consequence, the ionization enthalpies generally increase and electron gain enthalpies become more negative across a period. In other words, the ionization enthalpy of the extreme left element in a period is the least and the electron gain

enthalpy of the element on the extreme right is the highest negative. This results into high chemical reactivity at the two extremes and the lowest in the centre. Similarly, down the group, the increase in ionization enthalpies and a regular decrease (with exception in some third period elements) in electron gain enthalpies in the case of main group elements. The loss and gain of electrons can be co-related with the reducing and oxidising behaviour, and also with metallic and non-metallic character respectively of the elements.

Which of the following statements is incorrect?

- A. Oxide of aluminium (Al_2O_3), and arsenic (As_2O_3) are amphoteric
- B. Oxide of chlorine (Cl_2O_7) is less acidic than oxide of nitrogen (N_2O_5)
- C. Oxide of carbon (CO_2) is more acidic than oxide of silica (SiO_2).
- D. The correct increasing order of basic character of various oxides is



Answer: B



Watch Video Solution

3. The amount of energy required to remove, an electron from the last orbit of an isolated (free) atom in gaseous state is known as ionisation energy or first ionisation of the element. Similarly the energy required for the removal of the electron from the unipositive ion (M^+ produced above) is referred to as second ionization energy and thus the third, fourth etc. The ionisation energy depends on various factors like nuclear charge, size of atom, type configurations, screening effect and penetration power of the electrons.

The domination factor responsible for the decreasing ionisation energies of the elements of moving down the group is:

- A. atomic radius
- B. type of electron to be removed
- C. the valence sheel electron configuration
- D. all of the above

Answer: A



4. The amount of energy required to remove, an electron from the last orbit of an isolated (free) atom in gaseous state is known as ionisation energy or first ionisation of the element. Similarly the energy required for the removal of the electron from the unipositive ion (M^+ produced above) is referred to as second ionization energy and thus the third, fourth etc. The ionisation energy depends on various factors like nuclear charge, size of atom, type configurations, screening effect and penetration power of the electrons.

Which of the following order is not correct?

- A. $IE(I)$ of $Be > IE(I)$ of B but $IE(II)$ of $Be < IE(II)$ of B
- B. $IE(I)$ of $Be < IE(I)$ of B but $IE(II)$ of $Be < IE(II)$ of B
- C. $IE(II)$ of O $>$ $IE(II)$ of N
- D. $IE(I)$ of Mg $>$ $IE(I)$ of Al

Answer: B



5. Numerous forms of the periodic table have been revised from time to time. A modern version, which is most convenient and widely used is the long or extended form of the periodic table. The aufbau principle (electrons are filled in the progressive order of their increasing energy i.e., by $n + l$ rule) and the electronic configuration of atom provides a theoretical foundation for the periodic classification. The horizontal rows are called periods. There are altogether seven periods. The first period consists of 2 elements. The subsequent periods consist of 2, 8, 8, 18 and 32 elements respectively. The seventh period has a maximum of 32 elements. Elements having similar outer electronic configurations in their atoms are grouped in vertical columns. These are referred to as groups or families. According to the recommendation of IUPAC, the groups are numbered 1 to 18 replacing the older notation of groups I, II, ... VIIA, VIII, IB, ... VII B. Each successive period in the periodic table is associated with the filling up of next higher principal energy level following aufbau sequence. The number of elements in each period is twice the number of atomic orbitals available in the energy level that is being filled. All the elements are

classified into four blocks, i.e., s-block, p-block, d-block and f-block depending on the type of atomic orbitals that are being filled with the last electron of the element.

Elements a,b,c,d and e have the following electronic configurations.

(a) $1s^2, 2s^2 2p^1$

(b) $1s^2, 2s^2 2p^1 (6), 3s^2 3p^1$

(c) $1s^2, 2s^2 2p^6, 3s^2 3p^1$

(d) $1s^2, 2s^2 2p^6, 3s^2 3p^5$

(e) $1s^2, 2s^2 2p^6, 3s^2 3p^5$

Which among these will belong to the same group in periodic table?

A. a and c

B. a and b

C. a and b

D. d and e

Answer: B



Watch Video Solution

6. Numerous forms of the periodic table have been revised from time to time. A modern version, which is most convenient and widely used is the long or extended form of the periodic table. The aufbau principle (electrons are filled in the progressive order of their increasing energy i.e., by $n + l$ rule) and the electronic configuration of atom provides a theoretical foundation for the periodic classification. The horizontal rows are called periods. There are altogether seven periods. The first period consists of 2 elements. The subsequent periods consist of 2, 8, 8, 18 and 32 elements respectively. The seventh period is incomplete and has a maximum of 32 elements. Elements having similar outer electronic configurations in their atoms are grouped in vertical columns. These are referred to as groups or families. According to the recommendation of IUPAC, the groups are numbered 1 to 18 replacing the older notation of groups I, II, III, IV, V, VI, VII, VIII, IX, X, XI, XII, XIII, XIV, XV, XVI, XVII, XVIII, and XIX. Each successive period in the periodic table is associated with the filling up of next higher principal energy level following aufbau sequence. The number of elements in each period is twice the number of atomic orbitals available in the energy level that is being filled. All the elements are classified into four blocks, i.e., s-block, p-block, d-block and f-block depending on the type of atomic orbitals that are being filled with the

last electron of the element.

If aufbau rule is not followed, Ca-20 will be placed in.....block.

A. s

B. p

C. d

D. f

Answer: C



Watch Video Solution

7. Numerous forms of the periodic table have been revised from time to time. A modern version, which is most convenient and widely used is the long or extended form of the periodic table. The aufbau principle (electrons are filled in the progressive order of their increasing energy i.e., by $n + l$ rule) and the electronic configuration of atom provides a theoretical foundation for the periodic classification. The horizontal rows are called periods. There are altogether seven periods. The first period consists of 2

elements. The subsequent periods consist of 8, 8, 18 and 32 elements respectively. The seventh maximum of 32 elements. Elements having similar outer electronic configurations in their atoms are grouped in vertical columns. These are referred to as groups or families. According to the recommendation of IUPAC, the groups are numbered 1 to 18 replacing the older notation of groups O, IA, IIA VIIA, VIII, IB, VII B. Each successive period in the periodic table is associated with the filling up of next higher principal energy level following aufbau sequence. The number of elements in each period is twice the number of atomic orbitals available in the energy level that is being filled. All the elements are classified into four blocks, i.e., s-block, p-block, d-block and f-block depending on the type of atomic orbitals that are being filled with the last electron of the element.

What is the position of the element in the periodic table satisfying the electronic configuration $(n-1)d^1ns^2$ for $n = 4$?

- A. 3rd period and 3rd group
- B. 4th period and 4th group
- C. 3rd period and 2nd group

Answer: D



Watch Video Solution

8. Numerous forms of the periodic table have been revised from time to time. A modern version, which is most convenient and widely used is the long or extended form of the periodic table. The aufbau principle (electrons are filled in the progressive order of their increasing energy i.e., by $n + l$ rule) and the electronic configuration of atom provides a theoretical foundation for the periodic classification. The horizontal rows are called periods. There are altogether seven periods. The first period consists of 2 elements. The subsequent periods consist of 8, 8, 18 and 32 elements respectively. The seventh maximum of 32 elements. Elements having similar outer electronic configurations in their atoms are grouped in vertical columns. These are referred to as groups or families. According to the recommendation of IUPAC, the groups are numbered 1 to 18 replacing the older notation of groups 0, IA, IIA VIIA, VIII, IB, VII B.

Each successive period in the periodic table is associated with the filling up of next higher principal energy level following aufbau sequence. The number of elements in each period is twice the number of atomic orbitals available in the energy level that is being filled. All the elements are classified into four blocks, i.e., s-block, p-block, d-block and f-block depending on the type of atomic orbitals that are being filled with the last electron of the element.

In Mendeleev's periodic table, silver belongs to IB group. The group to which silver belongs in long form of periodic table is (Atomic number =47):

- A. first
- B. tenth
- C. sixteenth
- D. eleventh

Answer: D



Watch Video Solution

9. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ = sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Group	All Higher Group	Same Group	Group $n - 1$	Group $\leq n - 2$
$1s$	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in n th shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

Z^* for a 1st electron in Fe atom is:

- A. 2.85
- B. 25.70
- C. 25.65
- D. 3.75

Answer: B



Watch Video Solution

10. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ =sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Grpoup	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
1s	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in nth shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

The magnitude of first ionisation energy for Na (according to formula given) is equal to:

A. energy of its 3s electron

B. energy of its 1s electron

C. energy of its 2s electron

D. energy of its 2p electron

Answer: A



Watch Video Solution

11. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ = sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Group	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
$1s$	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in n th shell of

an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

The size of isoelectronic species F^- , Na^+ and Mg^{2+} is effected by:

- A. nuclear charge
- B. valence principal quantum number (n)
- C. electron-electron interaction in the outer orbitals
- D. none of the above

Answer: A



Watch Video Solution

12. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the

total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ =sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Grpoup	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
1s	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in nth shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

The correct order of radii is:

A. $Na < Li < K$

B. $O > S > Se$

C. $Cl < F < Li$

D. $Fe^{3+} < Fe^{2+} < Fe$

Answer: D



Watch Video Solution

13. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ = sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Group	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
$1s$	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in n th shell of

an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

Atomic radii of the noble gases are larger than the precedent elements of the same periods because:

- A. atomic radius of noble gas is expressed as van der Waals' radius
- B. valence shell electrons are completely filled so there is interelectronic repulsions
- C. both (a) and (b)
- D. none of the above

Answer: C



Watch Video Solution

14. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of

shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ = sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Group	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
1s	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in nth shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

Which of the following statements is correct?

A. Metallic radius refer to metals only is greater than covalent radius.

B. Metallic radius refer to metals only and is smaller than covalent radius

C. Generally covalent radius refer to non-metals as well as metals in bonded state (covalent bond).

D. Atomic radii of noble gases are expressed as van der Waal's radii which are smaller than metallic radii.

Answer: A



Watch Video Solution

15. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ =sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :




Electron Grpoup	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
1s	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in nth shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

Which one among the following sets of ions represents the collection of isoelectronic species?



D. (a) and (b) both


Answer: D



Watch Video Solution



16. The amount of energy required to remove the most loosely bound electron from an isolated gaseous atom is called as first ionization energy (IE_1). Similarly the amount of energies required to knock out second, third etc. electrons from the isolated and $IE_3 > IE_2 > IE_1$.

(i) Nuclear charge (ii) Atomic size (iii) penetration effect of the electrons (iv) shielding effect of the inner electrons and (b) electronic configurations (exactly half filled and completely filled configurations are extra stable) are the important factors which affect the ionisation energies. Similarly, the amount of energy released when a neutral isolated gaseous atom accepts an extra electron to form gaseous anion is called electron affinity.



A positive electron affinity indicates that the ion X^- has a lower more negative energy than the neutral atom X. The second electron affinity for the addition of a second electron to an initially neutral atom is negative

because the electron repulsion outweighs the nuclear attraction, e.g.,



The electron affinity of an element depends upon (i) atomic size (ii) nuclear charge and (iii) electronic configuration. In general, in a group, ionisation energy and electron affinity decrease as the atomic size increases. The members of third period have some higher (e.g., S and Cl) electron affinity values than the members of second period (e.g., O and F) because second period elements have very small atomic size. Hence, there is tendency of electron-electron repulsion, which results in less evolution of energy in the formation of corresponding anion.

The first ionisation energy of *Na*, *Mg*, *Al* and *Si* are in the order of:

A. $Na < Mg > Al < Si$

B. $Na > Mg > Al > Si$

C. $Na < Mg < Al < Si$


D. $Na > Mg > Al > Si$

Answer: A





17. The amount of energy required to remove the most loosely bound electron from an isolated gaseous atom is called as first ionization energy (IE_1). Similarly the amount of energies required to knock out second, third etc. electrons from the isolated and $IE_3 > IE_2 > IE_1$.

(i) Nuclear charge (ii) Atomic size (iii) penetration effect of the electrons (iv) shielding effect of the inner electrons and (b) electronic configurations (exactly half filled and completely filled configurations are extra stable) are the important factors which affect the ionisation energies. Similarly, the amount of energy released when a neutral isolated gaseous atom accepts an extra electron to form gaseous anion is called electron affinity.



A positive electron affinity indicates that the ion X^- has a lower more negative energy than the neutral atom X. The second electron affinity for the addition of a second electron to an initially neutral atom is negative because the electron repulsion outweighs the nuclear attraction, e.g.,





The electron affinity of an element depends upon (i) atomic size (ii) nuclear charge and (iii) electronic configuration. In general, in a group, ionisation energy and electron affinity decrease as the atomic size increases. The members of third period have some higher (e.g., S and Cl) electron affinity values than the members of second period (e.g., O and F) because second period elements have very small atomic size. Hence, there is tendency of electron-electron repulsion, which results in less evolution of energy in the formation of corresponding anion.

Which one of the following statements is incorrect in relation to ionisation enthalpy?

- A. Ionization enthalpy increase for each successive electron.
- B. The greatest increase in ionization enthalpy is experienced on removal of electron from core of noble gas configuration
- C. End of valence electrons is marked by a big jump in ionization enthalpy

D. Removal of electron from orbitals bearing lower n value is easier than from orbital having higher n value.

Answer: D



Watch Video Solution

18. The minimum amount of energy which is required to remove an outermost electron from any isolated neutral gaseous atom is known as first ionisation energy. These are the following factors which effect ionisation energy.

(i) Ionisation energy $\propto \frac{1}{\text{Principal quantum number}}$

(ii) Ionisation energy $\propto Z_{eff}$

(iii) If orbitals are fully filled or half filled so stability will be more and ionisation energy will be high.

(iv) If penultimate electron will effectively shield the nucleus, ionisation energy will be less and vice versa.

Which of the following has maximum 1st ionisation energy?

A. He

B. Mg

C. N

D. Na

Answer: A



Watch Video Solution

19. The ionisation energy (IE_1) of an atom is defined as the energy needed to remove an electron from gaseous atom in its ground state. The 2nd ionisation energy (IE_2) is the additional energy needed to remove the 2nd electron and so on. The successive ionisation energy of any species is increasing always.

Which represents the correct order of first ionisation potential of third period elements?

A. $Na > Mg > Al > Si$

B. $Na < Mg < Al < Si$

C. $Na < Si < Al < Mg$

D. $Na < Al < Mg < Si$

Answer: D




Watch Video Solution

20. The ionisation energy (IE_1) of an atom is defined as the energy needed to remove an electron from gaseous atom in its ground state. The 2nd ionisation energy (IE_2) is the additional energy needed to remove the 2nd electron and so on. The successive ionisation energy of any species is increasing always.

In the graph below elements of period given with their ionisation energy.

Select the one which represents an alkali metal:



A. X

B. Y

C. Z

D. L

Answer: A



Watch Video Solution

21. An atom is assumed to be spherical in shape and thus, the size of atom is generally given in terms of radius of the sphere and is called atomic radius. It is usually defined as the distance between the centre of the nucleus and outermost shell where electron are present. The exact measure of atomic radius is not easy due to following reasons:




- (i) The atom does not have well defined boundary. the probability of finding the electron is never zero even at large distance from the nucleus.
- (ii) It is not possible to get an isolated atom. the electron density around an atom is affected by the presence of neighbouring atoms, i.e., the size of the atom changes in going from one set of environment to another.
- (iii) the size of an atom is very small, of the order of about $1.2 \text{ \AA}$, i.e., $1.2 \times 10^{-10} m$.

An estimate of the size of the atom can, however, be made by knowing the distance between the atoms in the combined state. the distance between the atoms, i.e., bond length are generally measured by the application of techniques such as X-ray diffraction, electron diffraction, infrared spectroscopy, nuclear magnetic resonance spectroscopy, etc. However, bond lengths change with different type of bonding. Three

types of radius are commonly used, i.e.,

(a) Covalent radius (b) crystals radius (c) Vander waal's radius

Choose incorrect option regarding atomic radius



Answer: C



Watch Video Solution

22. An atom is assumed to be spherical in shape and thus, the size of atom is generally given in terms of radius of the sphere and is called atomic radius. It is usually defined as the distance between the centre of the nucleus and outermost shell where electron are present. The exact measure of atomic radius is not easy due to following reasons:

(i) The atom does not have well defined boundary. the probability of

finding the electron is never zero even at large distance from the nucleus.

(ii) It is not possible to get an isolated atom. the electron density around an atom is affected by the presence of neighbouring atoms, i.e., the size of the atom changes in going from one set of environment to another.

(iii) the size of an atom is very small, of the order of about $1.2 \text{ \AA}$, i.e., $1.2 \times 10^{-10} m$.

An estimate of the size of the atom can, however, be made by knowing the distance between the atoms in the combined state. the distance between the atoms, i.e., bond length are generally measured by the application of techniques such as X-ray diffraction, electron diffraction, infrared spectroscopy, nuclear magnetic resonance spectroscopy, etc. However, bond lengths change with different type of bonding. Three types of radius are commonly used, i.e.,

(a) Covalent radius (b) crystals radius (c) Vander waal's radius

Atomic radius of atoms in a period decreases with the increases in Z_{eff}

which of the following is incorrect order of atomic radius ?

A. $Li > Be > B > C$

B. $Na < Mg < Al < Si$

C. $Si > P > S > Cl$

D. $K > Ca$

Answer: B



Watch Video Solution

23. An atom is assumed to be spherical in shape and thus, the size of atom is generally given in terms of radius of the sphere and is called atomic radius. It is usually defined as the distance between the centre of the nucleus and outermost shell where electron are present. The exact measure of atomic radius is not easy due to following reasons:

- (i) The atom does not have well defined boundary. the probability of finding the electron is never zero even at large distance from the nucleus.
- (ii) It is not possible to get an isolated atom. the electron density around an atom is affected by the presence of neighbouring atoms, i.e., the size of the atom changes in going from one set of environment to another.
- (iii) the size of an atom is very small, of the order of about $1.2 \text{ \AA}$, i.e., $1.2 \times 10^{-10} m$.

An estimate of the size of the atom can, however, be made by knowing the distance between the atoms in the combined state. The distance between the atoms, i.e., bond length are generally measured by the application of techniques such as X-ray diffraction, electron diffraction, infrared spectroscopy, nuclear magnetic resonance spectroscopy, etc. However, bond lengths change with different type of bonding. Three types of radius are commonly used, i.e.,

(a) Covalent radius (b) crystals radius (c) Vander waal's radius

Choose incorrect option regarding atomic size

A. $Zr = Hf$

B. $Fe = Co = Ni$

C. $Y = La$

D. All of these

Answer: C



Watch Video Solution

24. An atom is assumed to be spherical in shape and thus, the size of atom is generally in term of radius of the sphere and is called atomic radius. It is usually defined as the distance between the centre of the nucleus and outermost shell where electron or electrons are present. The exact measure of atomic radius is not easy due to following reasons :

(a) The atom does not have well defined boundary. The propability of finding the electron is never zero even at large distance from the nucleus.

(b) It is not distances from to get an isolated atom. The electron density around an atom is affected by the presence of neighbouring atoms, i.e., the size of the atom changes in going from one set of environment to another.

(c) The size of an atom is very small, of the order of about $1.2\text{\AA}$, i.e., $1.2 \times 10^{-10}m$.



An estimate of the size of the atom can, however, be made by knowing the distance between the atoms in the combined state. The distance between the atoms, i.e., bond lengths are generally measured by the application of techniques such as X-ray diffraction, electron diffraction, infrared spectroscopy, nuclear magnetic resonance spectroscopy, etc.

However, bond length change with different type of bonding. Three types of radius are commonly used, i.e.,

(a) Covalent radius (b) Crystal radius

(c) van der Waal's radius

Which of the following set of ions have the same value of screening constant for the valence electron, calculated from Slater's rule?



Answer: B



Watch Video Solution

25. Some of the properties which depend on electronic configuration elements are - atomic radii, ionisation potential, and electronegativity etc.

In period, the ionisation energy is lowest for the:

A. noble gases

B. halogens

C. alkaline earth metals

D. alkali metals

Answer: D



Watch Video Solution

26. Some of the properties which depend on electronic configuration elements are - atomic radii, ionisation potential, and electronegativity etc. The values of electronegativity of atoms A and B are 1.20 and 4 respectively. The percentage ionic character of A-B bond is:

A. 50 %

B. 72.2 %

C. 55.3 %

D. 43.0 %


Answer: B






Watch Video Solution

27. In the modern period table, elements are arranged in order of increasing atomic number which is related to the electric configuration. Depending upon the type of orbitals receiving the last electron, the elements in the periodic table have been divided into four blocks viz s,p,d and f. The modern periodic table consists of 7 periods and 18 groups. Each period begins with the filling of a new energy shell. In accordance with the Aufbau principle, the seven periods (1 to 7) have 2, 8, 8, 18, 18, 32 and 32 elements respectively. The seventh period is still incomplete. To avoid the periodic table being too long, the two series of f-block elements, called lanthanoids and actinoids are placed at the bottom of the main body of the periodic table.

The last element of p-block in 6th period is represented by the outermost electronic configuration.





Answer: C



Watch Video Solution

28. In the modern period table, elements are arranged in order of increasing atomic number which is related to the electric configuration. Depending upon the type of orbitals receiving the last electron, the elements in the periodic table have been divided into four blocks viz s,p,d and f. The modern periodic table consists of 7 periods and 18 groups. Each period begins with the filling of a new energy shell. In accordance with the Aufbau principle, the seven periods (1 to 7) have 2, 8, 8, 18, 18, 32 and 32 elements respectively. The seventh period is still incomplete. To avoid the periodic table being too long, the two series of f-block elements, called lanthanoids and actinoids are placed at the bottom of the main

body of the periodic table.

Which of the elements whose atomic numbers are given below, cannot be accommodated in the present set up of the long form of the periodic table?

A. 107

B. 118

C. 126

D. 102

Answer: C







Watch Video Solution

29. In the modern periodic table, elements are arranged in order of increasing atomic numbers, which is related to the electronic configuration. Depending upon the type of orbitals receiving the last electron, the elements in the periodic table have been divided into four blocks, viz, p, d and f. The modern periodic table consists of 7 periods and

18 groups. Each period begins with the filling of a new energy shell. in accordance with the Aufbau principal, the seven periods (1 to 7) have 2,8,8,18,18,32 and 32 elements respectively. The seventh period is still incomplete. To avoid the periodic table being too long, the two series of f-block elements, called lanthanoids and actinoids, are placed at the bottom of the main body of the periodic table

The electronic configuration of the element which is just above the element with atomic number 43 in the same group is



Answer: A



Watch Video Solution

30. In the modern periodic table, elements are arranged in order of increasing atomic numbers, which is related to the electronic configuration. Depending upon the type of orbitals receiving the last electron, the elements in the periodic table have been divided into four blocks, viz, s, p, d and f. The modern periodic table consists of 7 periods and 18 groups. Each period begins with the filling of a new energy shell. in accordance with the Aufbau principle, the seven periods (1 to 7) have 2, 8, 8, 18, 18, 32 and 32 elements respectively. The seventh period is still incomplete. To avoid the periodic table being too long, the two series of f-block elements, called lanthanoids and actinoids, are placed at the bottom of the main body of the periodic table

The elements with atomic numbers 57, 89 and 103 are all

- A. noble gases
- B. halogens
- C. heavy metals
- D. light metals

Answer: B



Watch Video Solution

31. Ionisation energies three hypothetical elements are given below (in kJ/mole) :

	<i>I</i>	<i>II</i>	<i>III</i>
<i>X</i>	122	340	1890
<i>Y</i>	99	931	1100
<i>Z</i>	118	1220	16552

Which of the following pairs represents elements could be of the same group?

A. Y,Z

B. X,Y

C. X,Z

D. X,Y,Z

Answer: A



Watch Video Solution

32. Ionisation energies three hypothetical elements are given below (in kJ/mole) :

	<i>I</i>	<i>II</i>	<i>III</i>
<i>X</i>	122	340	1890
<i>Y</i>	99	931	1100
<i>Z</i>	118	1220	16552

What could be the value of the first electron affinity of Z^{++} in $KJmol^{-1}$?

- A. 118
- B. 1220
- C. 1652
- D. 734

Answer: B



Watch Video Solution

33. Ionisation energies three hypothetical elements are given below (in kJ/mole) :

	<i>I</i>	<i>II</i>	<i>III</i>
<i>X</i>	122	340	1890
<i>Y</i>	99	931	1100
<i>Z</i>	118	1220	16552

Energy (in kJ/mole) required for the process $Z \rightarrow Z^{2+} + 2e^{-}$ will be:

- A. 118
- B. 1220
- C. 1338
- D. 2872

Answer: C



Watch Video Solution

34. In 1931, Pauling defined the electronegativity of an atom as the tendency of the atom to attract electrons to itself when combined in a compound. The implication is that when a covalent bond is formed, the

electrons used for bonding need not be shared equally by both atoms. If the bonding electrons spend more time around one atom, that atom will have a δ^- charge, and consequently the other atom will have a δ^+ charge. In the extreme case, where the bonding electrons are round one atom all of the time, the bond is ionic. Pauling and others have attempted to relate the electronegativity difference between two atoms to the amount of ionic character in the bond between them.

Mulliken

In 1934, Mulliken suggested an alternative approach to electronegativity based on the ionization energy and electron affinity of an atom. Consider two atoms A and B. If an electron is transferred from A to B, forming ions A^+ and B^- , then the energy change is the ionization energy of atom A (I_A) minus the electron affinity of atom B (E_B), that is $I_A - E_B$. Alternatively, if the electron was transferred the other way to give B^+ and A^- ions, then the energy changed would be $I_B - I_A$. If A^+ and B^- are actually formed, then this process require less energy , and

$$(I_A - E_B) < (I_B - E_A)$$

$$\text{Rerranging } (I_A + E_A) < (I_A - E_B)$$

Now with respect to electronegativity for the same change,

$$E. N_A < E. N_B.$$

Thus Mulliken suggested that electronegativity is proportional to $I. E + E. A$ and could be regarded as the average of the ionization energy and the electron affinity of an atom.

$$\text{Electronegativity} = \frac{(I + E)}{2}$$

Mulliken used I and E values measured in electron volts, and the values were about 2.8 times larger than the Pauling values. It is to be noted that $I. E.$ and $E. A$ values are defined for singular gaseous atoms.

For the reaction $A(g) + e^- \rightarrow A^-(g)$ $\Delta H_r = \Delta H_{eg}$. The enthalpy change will be equal to (magnitude wise):

A. ΔH_{eg} of $A^-(g)$

B. $\Delta H_{I.E}$ of $A(g)$

C. $\Delta H_{E.N}$ of $A(g)$

D. $\Delta H_{I.E}$ of $A^-(g)$

Answer: D



Watch Video Solution

35. In 1931, Pauling defined the electronegativity of an atom as the tendency of the atom to attract electrons to itself when combined in a compound. The implication is that when a covalent bond is formed, the electrons used for bonding need not be shared equally by both atoms. If the bonding electrons spend more time around one atom, that atom will have a δ^- charge, and consequently the other atom will have a δ^+ charge. In the extreme case, where the bonding electrons are round one atom all of the time, the bond is ionic. Pauling and others have attempted to relate the electronegativity difference between two atoms to the amount of ionic character in the bond between them.

Mulliken

In 1934, Mulliken suggested an alternative approach to electronegativity based on the ionization energy and electron affinity of an atom. Consider two atoms A and B. If an electron is transferred from A to B, forming ions A^+ and B^- , then the energy change is the ionization energy of atom A (I_A) minus the electron affinity of atom B (E_B), that is $I_A - E_B$. Alternatively, if the electron was transferred the other way to give B^+ and A^- ions, then the energy changed would be $I_B - I_A$. If A^+ and B^-

are actually formed, then this process require less energy , and

$$(I_A - E_B) < (I_B - E_A)$$

Rerranging $(I_A + E_A) < (I_B - E_B)$

Now with respect to electronegativity for the same change,

$$E. N_A < E. N_B.$$

Thus Mulliken suggested that electronegativity is proportional to

$I. E + E. A$ and could be regarded as the average of the ionization

energy and the electron affinity of an atom.

$$\text{Electronegativity} = \frac{(I + E)}{2}$$

Mulliken used I and E values measured in electron volts, and the values

were about 2.8 times algrer than the Pauling values. It is to be noted that

$I. E.$ and $E. A$ values are defined for singular gaseous atoms.

For a reaction $A(g) + B(g) \rightarrow AB(S)$ the enthalpy change for the

recation will be [assuming $AB(s)$ to be an ionic compound] If

$$I. E_A + E. A_A < I. E_B + E. A_B$$

$$\text{A. } \Delta H_{I. E. A} + \Delta H_{eg_A} + \Delta H_{I. E_{AB(s)}}$$

$$\text{B. } \Delta H_{eg_A} + \Delta H_{I. E_B} + \Delta H_{L. E_{AB(s)}}$$


$$\text{C. } I. E_A + E. A_B + \Delta H_{L. E_{AB_s}}$$

$$D. I. E_B + E. A_B + \Delta_{L.E. AB(s)}$$

Answer: C



Watch Video Solution



36.

In above graph element F is:

- A. oxygen
- B. nitrogen
- C. fluorine
- D. neon

Answer: A



37. The amount of energy released when an electron is added to an isolated gaseous atom to produce a monovalent anion is called electron affinity of first electron affinity or electron gain enthalpy. The first electron affinity is given a negative sign as the addition of an electron to a neutral atom is an exoergic process. The addition of electron to A^- requires energy to overcome the force of repulsion. Thus, the second electron affinity is an endoergic process. The magnitude of electron affinity depends on a number of factors such as (i) atomic size (ii) effective nuclear charge (iii) screening effects (iv) half and fully filled orbitals and (v) shape of orbital. In general, electron affinity increase as the atomic radii decrease in a period. However, there are exceptions when the atoms have stable configuration. In a group, electron affinity decreases as the size increases. However, the members of 3rd period have somewhat higher values than the members in the 2nd period of the same subgroups.

Which of the following has least electron affinity?

A. Oxygen

B. Argon

C. Nitrogen

D. Boron

Answer: B



Watch Video Solution

38. The amount of energy released when an electron is added to an isolated gaseous atom to produce a monovalent anion is called electron affinity of first electron affinity or electron gain enthalpy. The first electron affinity is given a negative sign as the addition of an electron to a neutral atom is an exoergic process. The addition of electron to A^- requires energy to overcome the force of repulsion. Thus, the second electron affinity is an endoergic process. The magnitude of electron affinity depends on a number of factors such as (i) atomic size (ii) effective nuclear charge (iii) screening effects (iv) half and fully filled

orbitals and (v) shape of orbital. In general, electron affinity increase as the atomic radii decrease in a period. However, there are exceptions when the atoms have stable configuration. In a group, electron affinity decreases as the size increases. However, the members of 3rd period have somewhat higher values than the members in the 2nd period of the same subgroups.

The electron affinities of halogens are:

$$F = -332, Cl = -349, Br = -324, I = -295 \text{ kJ mol}^{-1} \quad \text{The}$$

higher value of Cl as compared to that of F^- is due to:

- A. higher atomic radius of F
- B. smaller electronegativity of F
- C. weaker electron repulsion in Cl
- D. more vacant p-subshell in Cl

Answer: C



Watch Video Solution

39. The amount of energy released when an electron is added to an isolated gaseous atom to produce a monovalent anion is called electron affinity of first electron affinity or electron gain enthalpy. The first electron affinity is given a negative sign as the addition of an electron to a neutral atom is an exoergic process. The addition of electron to A^- requires energy to overcome the force of repulsion. Thus, the second electron affinity is an endoergic process. The magnitude of electron affinity depends on a number of factors such as (i) atomic size (ii) effective nuclear charge (iii) screening effects (iv) half and fully filled orbitals and (v) shape of orbital. In general, electron affinity increase as the atomic radii decrease in a period. However, there are exceptions when the atoms have stable configuration. In a group, electron affinity decreases as the size increases. However, the members of 3rd period have somewhat higher values than the members in the 2nd period of the same subgroups.

Which of the following species has the highest electron affinity?



C. Na^+

D. O

Answer: D




Watch Video Solution

40. The properties of the elements (atomic/ionic radii, electron gain enthalpy, ionization enthalpy, electronegativity, valence, oxidising/reducing power, acid/base character, etc.) which are directly or indirectly related to their electronic configurations are called periodic properties. These properties show a regular gradation on moving from left to right in a period or from top to bottom in a group. Down a group, the atomic/ionic radii, metallic character and reducing character increase while ionization enthalpy and electronegativity decrease. Along a period from left to right, atomic/ionic and metallic character decrease while ionization enthalpy, electronegativity, non-metallic character and oxidizing power increase. However, electron gain enthalpy becomes less negative down a group but more negative along a period. In contrast, inert gases

have positive electron gain enthalpies which do not show any regular trend.

Which of the following isoelectronic ions has the lowest first ionization enthalpy?



Answer: D







Watch Video Solution

41. The properties of the elements (atomic/ionic radii, electron gain enthalpy, ionization enthalpy, electronegativity, valence, oxidising/reducing power, acid/base character, etc.) which are directly or indirectly related to their electronic configurations are called periodic properties. These properties show a regular gradation on moving from

left to right in a period or from top to bottom in a group. Down a group, the atomic/ionic radii, metallic character and reducing character increase while ionization enthalpy and electronegativity decrease. Along a period from left to right, atomic/ionic and metallic character decrease while ionization enthalpy, electronegativity, non-metallic character and oxidizing power increase. However, electron gain enthalpy becomes less negative down a group but more negative along a period. In contrast, inert gases have positive electron gain enthalpies which do not show any regular trend.

The outermost electronic configuration of the most electronegative elements is:




Answer: C






Watch Video Solution

42. The properties of the elements (atomic/ionic radii, electron gain enthalpy, ionization enthalpy, electronegativity, valence, oxidising/reducing power, acid/base character, etc.) which are directly or indirectly related to their electronic configurations are called periodic properties. These properties show a regular gradation on moving from left to right in a period or from top to bottom in a group. Down a group, the atomic/ionic radii, metallic character and reducing character increase while ionization enthalpy and electronegativity decrease. Along a period from left to right, atomic/ionic and metallic character decrease while ionization enthalpy, electronegativity, non-metallic character and oxidizing power increase. However, electron gain enthalpy becomes less negative down a group but more negative along a period. In contrast, inert gases have positive electron gain enthalpies which do not show any regular trend.

Amongst the following elements (whose electronic configurations are given below) the one having the highest ionization enthalpy is :





Answer: B



Watch Video Solution

43. The properties of the elements (atomic/ionic radii, electron gain enthalpy, ionization enthalpy, electronegativity, valence, oxidising/reducing power, acid/base character, etc.) which are directly or indirectly related to their electronic configurations are called periodic properties. These properties show a regular gradation on moving from left to right in a period or from top to bottom in a group. Down a group, the atomic/ionic radii, metallic character and reducing character increase while ionization enthalpy and electronegativity decrease. Along a period from left to right, atomic/ionic and metallic character decrease while ionization enthalpy, electronegativity, non-metallic character and oxidizing

power increase. However, electron gain enthalpy becomes less negative down a group but more negative along a period. In contrast, inert gases have positive electron gain enthalpies which do not show any regular trend.

Tick the correct order of second ionization enthalpy in the following:

A. $F > O > N > C$

B. $O > F > N > C$

C. $O > N > F > C$

D. $C > N > O > F$


Answer: B




Watch Video Solution

44. A quantitative measure of the tendency of an element to lose electron is given by its ionization enthalpy. It represents the energy required to remove an electron from an isolated gaseous atom (X) in its ground state. In other words, the first ionization enthalpy for an element X is the

enthalpy change ($\Delta_i H$) for the reaction depicted.



The ionization enthalpy is expressed in units of kJ mol^{-1} . We can define the second ionization enthalpy as the energy required to remove the second most loosely bound electron: it is the energy required to carry out the reaction,



Consider following statements for the two uncharged gaseous species

Species -1 = $1s^2, 2s^1$ Species -2 = $1s^2, 2p^1$

(P) Species-1 is ground state, species-2 is excited state of same atom

(Q) Species-1 is excited state, species-2 is ground state of same atom

(R) Ionization of species-1 is easier as compared to species-2

(S) Ionization of species-2 is easier as compared to species-1

Select correct statement(s)

A. P and R

B. Q and R

C. P and S


D. Q and S

Answer: C



Watch Video Solution

45. Born Haber cycle helps to determine lattice enthalpy of ionic compound as well as electron gain enthalpy of different elements. Based on the diagram drawn, answer the questions given below.



What is the value of IE_1 of $\text{Mg} + IE_1$ of Mg^{+} ?

A. 25 kJ

B. 5 kJ

C. 30 kJ


D. 20 kJ

Answer: C



Watch Video Solution

46. Born Haber cycle helps to determine lattice enthalpy of ionic compound as well as electron gain enthalpy of different elements. Based on the diagram drawn, answer the questions given below.



What is the electron gain enthalpy of $Cl(g)$?

- A. $-5kJ$
- B. $-25kJ$
- C. $-20kJ$
- D. $-10kJ$

Answer: D



Watch Video Solution

47. Nitrogen can form many oxides with oxygen, and thus is said to exhibit variable valency. Similarly, sulphur, phosphorus and carbon can exhibit variable valency.

Which sequence of compounds is according to the increasing order of the oxidation state of chlorine?

- A. $Cl_2, Cl_2O, Cl_2O_3, Cl_2O_5, Cl_2O_7, NH_4Cl$
- B. $Cl_2, Cl_2O, Cl_2O_3, Cl_2O_7, Cl_2O_5$
- C. $Cl_2, Cl_2O, Cl_2O_3, Cl_2O_5, Cl_2O_7$

D. CI_2 , CI_2O_3 , CI_2O_7 , CI_2

Answer: C



Watch Video Solution

48. The size of any species depends on various factors such as nature of charge, magnitude of charge/oxidation state, effective nuclear charge, electronic configuration etc.

It has been observed that size of Indium (In) $\approx$ Thallium (Tl). This can be attributed to:

- A. actinoid contraction
- B. lanthanoid contraction
- C. both (a) and (b)
- D. scandide contraction

Answer: B



Watch Video Solution

49. Ionisation energies of unknown elements are given below :

Element	IE(in kcal/mol)		
	I	II	III
M	209	548	919
N	100	735	1101
O	119	1092	1653
P	1500	2017	2320

Which amongst them is expected to be a noble gas?

A. M

B. N

C. O

D. P

Answer: D



Watch Video Solution

1. The amount of energy required to remove, an electron from the last orbit of an isolated (free) atom in gaseous state is known as ionisation energy or first ionisation of the element. Similarly the energy required for the removal of the electron from the unipositive ion (M^+ produced above) is referred to as second ionization energy and thus the third, fourth etc. The ionisation energy depends on various factors like nuclear charge, size of atom, type configurations, screening effect and penetration power of the electrons.

Which of the following statements is correct?

- A. Ionisation energies of elements decrease along the period.
- B. Ionisation energies of the II A group element are less than that of the corresponding IIIA group elements
- C. Ionisation energies of group 15 elements are less than that of the corresponding group 16 elements.
- D. Ionisation energy of Ga is greater than Al.

Answer: D



Comprehension 3

1. Numerous forms of the periodic table have been revised from time to time. A modern version, which is most convenient and widely used is the long or extended form of the periodic table. The aufbau principle (electrons are filled in the progressive order of their increasing energy i.e., by $n + l$ rule) and the electronic configuration of atom provides a theoretical foundation for the periodic classification. The horizontal rows are called periods. There are altogether seven periods. The first period consists of 2 elements. The subsequent periods consist of 2, 8, 8, 18 and 32 elements respectively. The seventh period is complete with 32 elements. Elements having similar outer electronic configurations in their atoms are grouped in vertical columns. These are referred to as groups or families. According to the recommendation of IUPAC, the groups are numbered 1 to 18 replacing the older notation of groups I, IA, IIA VIIA, VIII, IB, VII B. Each successive period in the periodic table is associated with the filling

up of next higher principal energy level following aufbau sequence. The number of elements in each period is twice the number of atomic orbitals available in the energy level that is being filled. All the elements are classified into four blocks, i.e., s-block, p-block, d-block and f-block depending on the type of atomic orbitals that are being filled with the last electron of the element.

The element with atomic number 56 is likely to have the same outer shell configuration as the element with atomic number.

A. 12

B. 18

C. 14

D. 20

Answer: A::D



Watch Video Solution

1. The reducing effect of the nuclear charge by the inner electrons for on outer electron is termed as shielding (or screening). As a result of shielding, the outer electrons in an atom always experience less nuclear charge than the actual nuclear charge Z . The effective nuclear charge (Z^*) as experienced by an electron is then obtained by subtracting the total shielding contributions from all the other electrons (i.e., except the one under consideration) from the actual nuclear charge.

$$Z^* = Z - \sigma$$

Where σ = sum of the shielding contributions. The rules for estimating contributions to σ are as follows (Slater's rule) Contribution to shielding by each electron is :

Electron Group	All Higher Group	Same Group	Group $n - 1$	Group $\leq$ $n - 2$
$1s$	0	0.30	—	—
(ns, sp)	0	0.35	0.85	1.00
(nd) or (nf)	0	0.35	1.00	1.00

According to Slater's treatment, the energy of an electron in n th shell of an atom having atomic number Z is given by the empirical equation

$$E = -13.6 \left(\frac{Z^*}{n} \right)^2 eV$$

Z^* = effective nuclear charge

Among the following, which electron of Fe atom experience minimum attraction from nucleus? (Atomic number of $Fe = 26$)

A. 3d

B. 4s

C. 2s

D. 2p

Answer: B




Watch Video Solution

Comprehension 5



1. The amount of energy required to remove the most loosely bound electron from an isolated gaseous atom is called as first ionization energy (IE_1). Similarly the amount of energies required to knock out second, third etc. electrons from the isolated and $IE_3 > IE_2 > IE_1$.

(i) Nuclear charge (ii) Atomic size (iii) penetration effect of the electrons

(iv) shielding effect of the inner electrons and (b) electronic configurations (exactly half filled and completely filled configurations are extra stable) are the important factors which affect the ionisation energies. Similarly, the amount of energy released when a neutral isolated gaseous atom accepts an extra electron to form gaseous anion is called electron affinity.



A positive electron affinity indicates that the ion X^{-} has a lower more negative energy than the neutral atom X. The second electron affinity for the addition of a second electron to an initially neutral atom is negative because the electron repulsion outweighs the nuclear attraction, e.g.,



The electron affinity of an element depends upon (i) atomic size (ii) nuclear charge and (iii) electronic configuration. In general, in a group, ionisation energy and electron affinity decrease as the atomic size increases. The members of third period have some higher (e.g., S and Cl) electron affinity values than the members of second period (e.g., O and F) because second period elements have very small atomic size. Hence, there

is tendency of electron-electron repulsion, which results in less evolution of energy in the formation of corresponding anion.

Identify the least stable ion amongst the following:



Answer: B



Watch Video Solution

Comprehension 6

1. The minimum amount of energy which is required to remove an outermost electron from any isolated neutral gaseous atom is known as first ionisation energy. These are the following factors which effect ionisation energy.

(i) Ionisation energy $\propto \frac{1}{\text{Principal quantum number}}$

(ii) Ionisation energy $\propto Z_{eff}$

(iii) If orbitals are fully filled or half filled so stability will be more and ionisation energy will be high.

(iv) If penultimate electron will effectively shield the nucleus, ionisation energy will be less and vice versa.

Choose the correct order of 1st ionisation energy

A. $Ne < F$

B. $O > N$

C. $Na > Al$

D. $Mg > Al$

Answer: D



Watch Video Solution

1. The ionisation energy (IE_1) of an atom is defined as the energy needed to remove an electron from gaseous atom in its ground state. The 2nd ionisation energy (IE_2) is the additional energy needed to remove the 2nd electron and so on. The successive ionisation energy of any species is increasing always.

Generally, the first ionisation energy increase along a period. But there are some exceptions. One which is not an exception is:

- A. N and O
- B. Na and Mg
- C. Mg and Al
- D. Be and B

Answer: B



Watch Video Solution

1. An atom is assumed to be spherical in shape and thus, the size of atom is generally given in terms of radius of the sphere and is called atomic radius. It is usually defined as the distance between the centre of the nucleus and outermost shell where electron are present. The exact measure of atomic radius is not easy due to following reasons:

- (i) The atom does not have well defined boundary. the probability of finding the electron is never zero even at large distance from the nucleus.
- (ii) It is not possible to get an isolated atom. the electron density around an atom is affected by the presence of neighbouring atoms, i.e., the size of the atom changes in going from one set of environment to another.
- (iii) the size of an atom is very small, of the order of about $1.2 \text{ \AA}$, i.e., $1.2 \times 10^{-10} m$.

An estimate of the size of the atom can, however, be made by knowing the distance between the atoms in the combined state. the distance between the atoms, i.e., bond length are generally measured by the application of techniques such as X-ray diffraction, electron diffraction, infrared spectroscopy, nuclear magnetic resonance spectroscopy, etc. However, bond lengths change with different type of bonding. Three types of radius are commonly used, i.e.,

(a) Covalent radius

(b) crystals radius

(c) Vander waal's radius

The correct order of effective nuclear charge Z_{eff} is

A. $B < C < N < O < F$

B. $B = C = N = O = F$

C. $B > C > N > O > F$

D. none of these

Answer: A



Watch Video Solution

Comprehension 9

1. Some of the properties which depend on electronic configuration elements are - atomic radii, ionisation potential, and electronegativity etc.

The element with maximum electronegativity belongs to :

A. period 2, group 17

B. period 3, group 18

C. period 4, group 17

D. period 2, group 16

Answer: A



Watch Video Solution

Comprehension 10

1. In the modern period table, elements are arranged in order of increasing atomic number which is related to the electric configuration. Depending upon the type of orbitals receiving the last electron, the elements in the periodic table have been divided into four blocks viz s,p,d and f. The modern periodic table consists of 7 periods and 18 groups. Each period begins with the filling of a new energy shell. In accordance with the Aufbau principle, the seven periods (1 to 7) have 2, 8, 8, 18, 18, 32 and 32 elements respectively. The seventh period is still incomplete. To avoid

the periodic table being too long, the two series of f-block elements, called lanthanoids and actinoids are placed at the bottom of the main body of the periodic table.

The element with atomic number 57 belongs to:

- A. s-block
- B. p-block
- C. d-block
- D. f-block

Answer: C



Watch Video Solution

Comprehension 11

1. Ionisation energies three hypothetical elements are given below (in kJ/mole) :

	<i>I</i>	<i>II</i>	<i>III</i>
<i>X</i>	122	340	1890
<i>Y</i>	99	931	1100
<i>Z</i>	118	1220	16552

Which of the following is likely to be 2nd group element?

- A. X
- B. Z
- C. Y
- D. X and Y

Answer: A



Watch Video Solution

Comprehension 12

1. In 1931, Pauling defined the electronegativity of an atom as the tendency of the atom to attract electrons to itself when combined in a compound. The implication is that when a covalent bond is formed, the

electrons used for bonding need not be shared equally by both atoms. If the bonding electrons spend more time around one atom, that atom will have a δ^- charge, and consequently the other atom will have a δ^+ charge. In the extreme case, where the bonding electrons are round one atom all of the time, the bond is ionic. Pauling and others have attempted to relate the electronegativity difference between two atoms to the amount of ionic character in the bond between them.

Mulliken

In 1934, Mulliken suggested an alternative approach to electronegativity based on the ionization energy and electron affinity of an atom. Consider two atoms A and B. If an electron is transferred from A to B, forming ions A^+ and B^- , then the energy change is the ionization energy of atom A (I_A) minus the electron affinity of atom B (E_B), that is $I_A - E_B$. Alternatively, if the electron was transferred the other way to give B^+ and A^- ions, then the energy changed would be $I_B - I_A$. If A^+ and B^- are actually formed, then this process require less energy , and

$$(I_A - E_B) < (I_B - E_A)$$

$$\text{Rerranging } (I_A + E_A) < (I_A - E_B)$$

Now with respect to electronegativity for the same change,

$$E. N_A < E. N_B.$$

Thus Mulliken suggested that electronegativity is proportional to $I. E + E. A$ and could be regarded as the average of the ionization energy and the electron affinity of an atom.

$$\text{Electronegativity} = \frac{(I + E)}{2}$$

Mulliken used I and E values measured in electron volts, and the values were about 2.8 times larger than the Pauling values. It is to be noted that $I. E.$ and $E. A$ values are defined for singular gaseous atoms.

We have X atoms of A. if all the atoms gain one electron each the energy released is aeV . If Y atoms of A lose one electron each, then the energy absorbed is beV . Then the electronegativity of A on the Mulliken scale will be:

A. $\frac{a + b}{2}$


B. $\frac{1}{2} \left(\frac{a}{X} + \frac{b}{Y} \right)$

C. $\frac{1}{2}(a + b) \times N_a$

D. $\frac{1}{2} \left(\frac{a}{X} + \frac{b}{Y} \right) \times N_a$

Answer: B

Comprehension 13



1.

Find the number of non metals in above graph, except noble gas. If A,B,C,E,F,G and H are 2nd period-elements.

A. 5

B. 6

C. 7

D. 8

Answer: A

Comprehension 14

1. The amount of energy released when an electron is added to an isolated gaseous atom to produce a monovalent anion is called electron affinity of first electron affinity or electron gain enthalpy. The first electron affinity is given a negative sign as the addition of an electron to a neutral atom is an exoergic process. The addition of electron to A^- requires energy to overcome the force of repulsion. Thus, the second electron affinity is an endoergic process. The magnitude of electron affinity depends on a number of factors such as (i) atomic size (ii) effective nuclear charge (iii) screening effects (iv) half and fully filled orbitals and (v) shape of orbital. In general, electron affinity increase as the atomic radii decrease in a period. However, there are exceptions when the atoms have stable configuration. In a group, electron affinity decreases as the size increases. However, the members of 3rd period have somewhat higher values than the members in the 2nd period of the same subgroups.

Which one of the following arrangements represents the correct order of electron gain enthalpy (with negative spin) of the given atomic species?

A. $F < Cl < O < S$

B. $S < O < Cl < F$

C. $O < S < F < Cl$

D. $Cl < F < S < O$

Answer: C



Watch Video Solution

Comprehension 15

1. The properties of the elements (atomic/ionic radii, electron gain enthalpy, ionization enthalpy, electronegativity, valence, oxidising/reducing power, acid/base character, etc.) which are directly or indirectly related to their electronic configurations are called periodic properties. These properties show a regular gradation on moving from

left to right in a period or from top to bottom in a group. Down a group, the atomic/ionic radii, metallic character and reducing character increase while ionization enthalpy and electronegativity decrease. Along a period from left to right, atomic/ionic and metallic character decrease while ionization enthalpy, electronegativity, non-metallic character and oxidizing power increase. However, electron gain enthalpy becomes less negative down a group but more negative along a period. In contrast, inert gases have positive electron gain enthalpies which do not show any regular trend.

If the ionic radii of K^+ and F^- are about $1.34\text{\AA}$ each, then the expected value of atomic radii of K and F should be respectively:

- A. 2.31 and $0.64\text{\AA}$
- B. 2.31 and $1.34\text{\AA}$
- C. 0.64 and $2.31\text{\AA}$
- D. 1.34 and $1.34\text{\AA}$


Answer: A




Watch Video Solution

Comprehension 16



1. A quantitative measure of the tendency of an element to lose electron is given by its ionization enthalpy. It represents the energy required to remove an electron from an isolated gaseous atom (X) in its ground state. In other words, the first ionization enthalpy for an element X is the enthalpy change ($\Delta_i H$) for the reaction depicted.





The ionization enthalpy is expressed in units of kJ mol^{-1} . We can define the second ionization enthalpy as the energy required to remove the second most loosely bound electron: it is the energy required to carry out the reaction,



Largest energy is required for which change?






Answer: D



Watch Video Solution

Comprehension 17

1. Born Haber cycle helps to determine lattice enthalpy of ionic compound as well as electron gain enthalpy of different elements. Based on the diagram drawn, answer the questions given below.



What is the lattice enthalpy for the formation of MgCl_2 ?

- A. -20kJ
- B. -40kJ
- C. -15kJ
- D. $+40\text{kJ}$



Answer: A



Watch Video Solution

1. Nitrogen can form many oxides with oxygen, and thus is said to exhibit variable valency. Similarly, sulphur, phosphorus and carbon can exhibit variable valency.

Amongst the following, select those which are most likely to be neutral.



Answer: A






Watch Video Solution

Comprehension 19

1. The size of any species depends on various factors such as nature of charge, magnitude of charge/oxidation state, effective nuclear charge,

electronic configuration etc.

Select the order of size which are correct?



Answer: C



Watch Video Solution

Comprehension 20

1. Ionisation energies of unknown elements are given below :

Element	IE(in kcal/mol)		
	I	II	III
M	209	548	919
N	100	735	1101
O	119	1092	1653
P	1500	2017	2320

Which pair of elements may represent same group?

A. M,N

B. N,O

C. O,P

D. M,P

Answer: B



Watch Video Solution

Match The Column Type

1. Match the following columns

Column-I		Column-II	
(a)	ns^2, np^5	(p)	Chromium
(b)	$(n-1)d^{10}, ns^1$	(q)	Copper
(c)	$(n-1)d^5, ns^1$	(r)	Krypton
(d)	$(n-1)d^{10}, ns^2, np^6$	(s)	Bromine

[Watch Video Solution](#)

2. Match the following columns

Column-I		Column-II	
(a)	Metalloid	(p)	Sulphur
(b)	Radioactive	(q)	Gold
(c)	Transition metal	(r)	Arsenic
(d)	Chalcogen	(s)	Uranium

[Watch Video Solution](#)

3. Match the following columns

Column-I		Column-II	
(a)	Representative element	(p)	Cerium
(b)	Lanthanide	(q)	Aluminium
(c)	Coinage metal	(r)	Thorium
(d)	Actinide	(s)	Gold

[Watch Video Solution](#)

4. Match the following columns

Column-I		Column-II	
(a)	Isoelectronic species	(p)	$A^+(g) + \text{energy} \rightarrow A^{++}(g) + e^-(g)$
(b)	Half filled orbital	(q)	Ar, K^+ , Ca^{++}
(c)	Second ionisation energy	(r)	Lutetium
(d)	Inner transition element	(s)	Antimony



Watch Video Solution

5. Match the following columns

Column-I		Column-II	
(a)	Highest 1 st ionisation energy	(p)	Technetium
(b)	Highest electronegativity	(q)	Lithium
(c)	Synthetic element	(r)	Helium
(d)	Strongest reducing agent	(s)	Fluorine



Watch Video Solution

6. Match the Column-I to Column-II :

Column-I (element / elements)		Column-II (group number)	
(a)	An element whose fourth shell contains two p -electrons	(p)	8th group
(b)	An element whose valence shell contains one unpaired p -electron	(q)	12th group
(c)	An element which receives last electron in $(n - 1) d$ -subshell	(r)	14th group
(d)	An element with the ground-state electron configuration $[\text{Ar}]4s^2 3d^{10}$	(s)	17th group



Watch Video Solution

7. The first (ΔH_1) and second (ΔH_2) ionisation enthalpies (in kJmol^{-1}) and the (ΔH_{eg}) electron gain enthalpy (in kJmol^{-1}) of a elements are given below:

	Element	ΔH_1	ΔH_2	$\Delta_{eg}H$
1.	P	520	7300	-60
2.	Q	419	3051	-48
3.	R	1681	3374	-328
4.	S	1008	1846	-295
5.	T	2372	5251	+48
6.	U	738	1451	-40

Based on the above information match the following columns.

Column-I		Column-II	
(a)	The least reactive element	(p)	R
(b)	The most reactive metal is	(q)	S
(c)	The most reactive non-metal is	(r)	T
(d)	The non-metal with least oxidising power (other than zero group element)	(s)	Q



Watch Video Solution

8. Match the following columns

Column-I		Column-II	
(a)	$1s^2$	(p)	shows highest negative oxidation state
(b)	$1s^2 2s^2 2p^5$	(q)	shows highest first ionisation energy
(c)	$1s^2 2s^1$	(r)	shows highest reducing power in aqueous solution
(d)	$1s^2 2s^2 2p^3$	(s)	shows highest electron affinity
		(t)	shows highest electronegativity



Watch Video Solution

9. Column-I contains some increasing orders of various species and column-II has the properties of the elements/ions.

Column-I		Column-II	
(a)	$\text{Na}^+ < \text{F}^- < \text{O}^{2-} < \text{N}^{3-}$	(p)	Electronegativity
(b)	$\text{Li}^+ < \text{Na}^+ < \text{K}^+ < \text{Rb}^+ < \text{Cs}^+$	(q)	Mobility of hydrated ions
(c)	$\text{O} < \text{S} < \text{F} < \text{Cl}$	(r)	Ionisation energy
(d)	$\text{Cl}^- < \text{K}^+ < \text{Ca}^{2+} < \text{Sc}^{3+}$	(s)	Electron affinity
		(t)	Ionic radii

 Watch Video Solution

10. Column-I contains the atomic number of few elements and column-II has the properties of the elements/salts.

Column-I		Column-II	
(a)	$Z = 19$	(p)	Diamagnetic and has higher atomic radius than $Z = 29$
(b)	$Z = 17$	(q)	Lower density than either side of the elements in the same group
(c)	$Z = 30$	(r)	Acts as metal as well as non-metal
(d)	$Z = 20$	(s)	Oxide is amphoteric
(e)	$Z = 51$	(t)	More negative electron gain enthalpy than $Z = 9$

 Watch Video Solution

11. Match the following columns

Column-I (Properties)		Column-II (Magnitude)	
(a)	Ionisation energy (IE_1)	(p)	Highest in halogens in their respective periods
(b)	Electron affinity (EA_1)	(q)	Highest in noble gas in their respective periods
(c)	Electronegativity	(r)	Highest in alkali metals in their respective periods
(d)	Electropositive character	(s)	Lowest in noble gases in their respective periods.

12. Match the values of 1st and 2nd ionisation energy



Watch Video Solution

12. Match the values of 1st and 2nd ionisation energy and electron gain enthalpy given in the column-I with types of elements/compounds given in column-II

Column-I				Column-II	
	$\Delta_1 H_1, \Delta_1 H_2, \Delta_{eg} H$ (in kJ mol^{-1})				
(a)	2372	5251	+48	(p)	Element which exists as a strong reducing agent
(b)	419	3051	-49	(q)	Element which exists as a monoatomic molecule
(c)	1681	3374	-333	(r)	Least reactive non-metal

(d)	1008	1846	-295	(s)	Element which acts as a strong oxidising agent
				(t)	Element whose oxide is strongly basic in nature



Watch Video Solution

13. Match the following columns

Column-I		Column-II	
(a)	$\text{Fe(III)} > \text{Fe(II)}$	(p)	Electronegativity
(b)	$\text{Al} > \text{Na}$	(q)	Basic character of their oxides
(c)	$\text{Cl} > \text{F}$	(r)	Electron gain enthalpy ($-\Delta_{eg} H$)
(d)	$\text{N} > \text{C}$	(s)	Degree of hydration



Watch Video Solution

14. Match the following columns

Column-I		Column-II	
(a)	NO_3^- , CO_3^{2-}	(p)	Semi-metals
(b)	Si, Ge, As, Sb	(q)	Isoelectronic species
(c)	He, Ne, Ar, Kr, Xe	(r)	van der Waals' radii
(d)	Na, Mg, Al, Si, P, S	(s)	Typical elements

 Watch Video Solution

15. Match the following columns

Column-I (Element)		Column-II (Property)	
(a)	Mg	(p)	s-block element
(b)	Be	(q)	Bridge element
(c)	Sc	(r)	Transition element
(d)	N	(s)	p-block element
		(t)	Alkaline earth elements

 Watch Video Solution

16. Match the following Columns

Column-I (Characteristic involved in the given process of Column II)		Column-II (Process described)	
(a)	Energy released	(p)	$S \longrightarrow S^{-}$
(b)	Energy absorbed	(q)	$O^{-} \longrightarrow O^{2-}$
(c)	Inert gas configuration is achieved	(r)	$Sr \longrightarrow Sr^{2+}$
(d)	Half filled configuration is achieved	(s)	$N^{-} \longrightarrow N$
		(t)	$Na^{+} \longrightarrow Na$



Watch Video Solution

17.

Column-I		Column-II	
(a)	Noble gas	(p)	Uranium
(b)	Transition element	(q)	Radon

(c)	Lanthanide	(r)	Cerium
		(s)	Copper



Watch Video Solution

18. Match the following columns

Column-I (Element)		Column-II [Atomic radius (pm)]	
(a)	Be	(p)	74
(b)	C	(q)	88
(c)	O	(r)	111
(d)	B	(s)	77
(e)	N	(t)	66



Watch Video Solution

19. Match the correct ionisation enthalpies electron gain enthalpies of the following elements.

Column-I		Column-II			
Elements			ΔH_1	ΔH_2	$\Delta_{eg}H$
(a)	Most reactive non-metal	(p)	419	3051	-
(b)	Most reactive metal	(q)	1681	3374	-328
(c)	Least reactive element	(r)	738	1451	-40
(d)	Metal forming binary halide	(s)	2372	5251	+48



Watch Video Solution

20. Match the following columns

Column-I (Electronic configuration)		Column-II (Electron gain enthalpy / kJ mol^{-1})	
(a)	$1s^2 2s^2 2p^6$	(p)	-53
(b)	$1s^2 2s^2 2p^6 3s^1$	(q)	-328
(c)	$1s^2 2s^2 2p^5$	(r)	-141
(d)	$1s^2 2s^2 2p^4$	(s)	+48



Watch Video Solution

21. Match the following columns

Column-I		Column-II	
(a)	Li, Na, K, Rb	(p)	Increase in radius
(b)	Li, B, C, N	(q)	Decrease in ionisation energy
(c)	Mg^{++} , Na^+ , F^- , O^{--}	(r)	Increase in Z_{eff}
(d)	F, Cl, Br, I	(s)	Decrease in Z_{eff}



Watch Video Solution

22. Match the order of species in column-I with properties in column-II.

Column-I		Column-II	
(a)	$\text{Na}^+ > \text{K}^+ > \text{Rb}^+$	(p)	Ionic mobility in aqueous solvent
(b)	$\text{B} < \text{C} < \text{O} < \text{N}$	(q)	Electronegativity of respective atom

(c)	$\text{Mg}^{2+} < \text{Li}^+ < \text{K}^+ < \text{Cs}^+$	(r)	Hydration energy
(d)	$\text{S} > \text{Se} > \text{O}$	(s)	1st Electron affinity
		(t)	1st Ionisation energy



Watch Video Solution

23. Match the following columns

Column-I (Correct characteristics)		Column-II (Elements)	
(a)	Radioactive lanthanide element	(p)	Beryllium
(b)	Shows diagonal relationship	(q)	Promethium
(c)	Transition element	(r)	Cadmium
(d)	Ground state electronic configuration $[\text{Kr}] 4d^{10}5s^2$	(s)	Osmium



Watch Video Solution

24. Match column-I (atomic number of elements) with column-II (position of element in periodic table).

Column-I		Column-II	
(a)	19	(p)	<i>p</i> -block
(b)	22	(q)	<i>f</i> -block
(c)	32	(r)	<i>d</i> -block
(d)	64	(s)	<i>s</i> -block



Watch Video Solution

25. Match the following columns

Column-I (Electronic configuration of element)		Column-II (Correct description)	
(a)	$1s^1, 2s^2 2p^2, 3s^1$	(p)	<i>s</i> or <i>p</i> block element
(b)	$1s^2, 2s^2 2p^6, 3s^2 3p^1$	(q)	III period element
(c)	$1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5, 4s^1$	(r)	Group number VI in the long form periodic table
(d)	$1s^2, 2s^1 2p^3, 3s^1 3p^3, 4s^1$	(s)	<i>d</i> -block element
		(t)	Valence shell electron ≥ 3



Watch Video Solution

26. Match the following columns

Column-I (Atomic Number)		Column-II	
(a)	52	(p)	s-block
(b)	56	(q)	p-block
(c)	57	(r)	d-block
(d)	60	(s)	f-block



Watch Video Solution

Subjective Type

1.0.5 moles of gaseous non-metallic X^- anions (having positive electron affinity) requires 806.4 kJ energy to get completely converted into gaseous X^+ ions. Calculate Pauling's electronegativity of the elements X.

Use Avogadro's no = 6×10^{23} and $1\text{eV} = 1.6 \times 10^{-19}\text{J}$.

[Use the fact that, Pauling's electronegativity

$$= \frac{\text{Mulliken's electronegativity}}{2.8} \quad \text{and} \quad \text{Mulliken's electronegativity} = \frac{\text{Ionisation energy} + \text{Electron affinity}}{2} \quad \Bigg]$$

 [View Text Solution](#)

2. $1.0g$ of Mg atom (atomic mass = 24.0amu) in the vapour phase absorbs $50.0kJ$ energy. Find the composition of the final magnesium, if the first and the second IE of Mg are $740kJmol^{-1}$ and $1450kJmol^{-1}$ respectively.

 [Watch Video Solution](#)

3. Find the sum of total number of 5f-electrons in Th and Ac .

 [Watch Video Solution](#)

4. Find the total number of species having two unpaired electrons from the following species Fe^{2+} , Cr , Cr^{3+} , Ti^{2+} , Mn^{2+} , V^{3+}

 [Watch Video Solution](#)

5. If internuclear distance between A atoms in A_2 is $10\text{\AA}$ and between B atoms in B_2 is $2\text{\AA}$, then calculate internuclear distance between A and B (Electronegativity difference between A and B has negligible value).



Watch Video Solution

6. Find the number of p-block elements from the following atomic numbers given below. 83 79 42 64 37 54 34



Watch Video Solution

7. Find the total number of 6^{th} period elements from the given atomic numbers? 81, 63, 80, 50, 54, 48, 86



Watch Video Solution

8. Find the total number of paramagnetic species among the following?

Sc^{3+} , Fe^{3+} , Mn^{2+} , Co^{4+} , Co^{3+} , Cr^{+} , Fe^{2+} , Mn^{3+} , Cr^{3+} , Zn^{2+} , Ti^{+4}



[Watch Video Solution](#)

9. What is the group number of Ba in periodic table?



[Watch Video Solution](#)

10. Select the number of elements which are called transition metals.

B, Sc, Al, Pd, Os, Zr, Rb, Ba, Fr



[Watch Video Solution](#)

11. Among the following species, how many have their ionic size greater than O^{2-} ?

Se²⁻, F⁻, N³⁻, P³⁻



[Watch Video Solution](#)

12. Among the following find out the total number of d-block elements.

La, Pd, Ni, Cd, Cs, Bi, Sn, Zr, K



Watch Video Solution

13. Number of oxides which is/are more basic as compared to Na_2O .

Li₂O, K₂O, Cs₂O, Rb₂O, MgO



Watch Video Solution

14. Bond length of A-A bond is 124 pm and bond length of B-B bond is 174 pm. Calculate the bond length (in pm) of A-B bond in AB molecule if percent ionic character of A-B bond is 19.5 %



Watch Video Solution

15. Find the number of chemical species in which outer shell 'd' orbital is having more than five electrons.

Ac, Cd, Zn, Sn, Cr, Pt, Ag



Watch Video Solution

16. Find the number of transition elements in the following:

Zn, Cd, Hg, Pt, U, Sn



Watch Video Solution

17. According to Hannay-Smith formula, if E.N difference between A and B is 2.1, then A-B molecule is expected to have x% ionic character, find $\frac{x}{10}$ value to the next integer.



Watch Video Solution

18. The element with the lowest atomic number that has a ground-state electronic configuration of $(n - 1)d^5ns^2$ is located in.....period.



Watch Video Solution

19. Find the total number of correct orders among the following.

(a) Order of electron affinity : $S > O^-$

(b) Order of ionic radius: $Al^{3+} < Mg^{2+} < O^{2-}$

(c) Order of electronegativity: $Si < P < C < N$

(d) Order of atomic radius: $O < C < S < Se$

(e) Order of second ionization energy: $C < N < F < O$

(f) Order of ionic radii : $S^{2-} < Cl^- < K^+ < Ca^{2+}$





Watch Video Solution

20. Period number of $Sc = x$





Modern periodic table group number of $Tl = x$ (according to 1 to 18 convention) Find the value of $y-x$



Watch Video Solution



Watch Video Solution



Watch Video Solution



Watch Video Solution

24. Among the following number of elements, which are transition elements are:

Ce, Nd, Sm, Yb, Ru, Th, U, Np, Am



Watch Video Solution

25. The number of different possible oxidation states of fluorine in the molecules $[F_2, OF_2, O_2F_2, HF] = t$ Find the value of t



Watch Video Solution

26. How many of the following species have lower IE_1 value compared to F^- ?

Cl^-, Br^-, O^-, I^-



Watch Video Solution

27. The number of oxides which are expected to be neutral amongst the oxides of nitrogen (viz. N_2O , NO , NO_2 , N_2O_4 , N_2O_5) = x The number of oxides which are expected to be more basic with respect to NiO amongst MgO , SrO , K_2O = y . Find the value of $x+y$.



Watch Video Solution

28. What is the value of $(n + l)$ for the unpaired e^- in an atom of an element which is present in the 3rd period and seventeenth group of the periodic table.



Watch Video Solution

29. A monoatomic anion of unit charge contain 45 neutrons and 36 electrons. What is the atomic mass number of element and in which group of periodic table does it lie? Write your answer as 1002 if answer is 10,2.



Watch Video Solution

[Watch Video Solution](#)

30. If $3e^-$ can be accommodated in each orbital then how many elements will be present in the 7th period of periodic table?



[Watch Video Solution](#)

31. How many elements will be present in the 12th period of periodic table.



[Watch Video Solution](#)

32. Identify the number of pairs which have higher electronegativity difference as compared to difference of Cs and F, Na and I, Li and F, H and F, I and Cl, Ca and O, Ba and S, C and Si.



[Watch Video Solution](#)