



CHEMISTRY

BOOKS - GRB CHEMISTRY (HINGLISH)

PRACTICAL ORGANIC CHEMISTRY

Exercise 1 Only One Correct Answer

1. Carbon and hydrogen are normally detected by strongly

heating the organic compound with

A. FeO

B. CaO

C. CuO

D. MnO


Answer: C


Watch Video Solution


Exercise 2 More Than One Correct Answer

1. Which of the following aromatic compounds will react

with $KMnO_4$?







Answer: A::B::C





Exercise 3 Passage 1


1. Compound (A) C_7H_8O is insoluble in aqueous $NaHCO_3$ and dissolves in aqueous NaOH and gives a characteristic colour with neutral $FeCl_3$. When treated

with $Br_2(A)$ forms compound $(B)C_7H_5OBr_3$.


The most probable structure of compound A is :







Β.



Answer: B




Passage 2

1. From the following sequence of reactions , [A] $(C_6H_{12}) \xrightarrow{HCl} (B)(C_6H_{13}Cl) + (C)(C_6H_{13}Cl)$ react with $AgNO_3$ to give white ppt. [B] $\xrightarrow{Alc.KOH}$ (D) (An isomer of A) gives positive test with Br_2/CCl_4 [D] $\xrightarrow{Ozonelysis}$ (E) gives positive iodoform test and negative Fehling's test . [A] $\xrightarrow{Ozonolysis}$ (F) + (G) , both F and G give positive Tollen's

test .

 $[F] + [G] \xrightarrow{\text{Conc.NaOH}} \text{HCOONa} + \text{alcohol}$


The structure A and B respectively are :



Answer: A

View Text Solution

Exercise 4 Matrix Match Type



View Text Solution



1. Lassigne's test is not used for the detection of :

B.S

C. Cl

D. 0

Answer: D



2. When an organic compound is present in aqueous medium and is less soluble in any organic solvent, then it is separated by :

A. continous extraction

B. distillation

C. chromatography

D. sublimation

Answer: A

Watch Video Solution

3. Ammonium molybdate is used for detection of which element in organic compound :

A. C

B. N

C. P

D. S

Answer: C

:




4. A white crystalline solid 'X' give following chemical test


(i) it liberates CO_2 with $NaHCO_3$


(ii) it forms a coloured dye on diazotisation and coupling with β -naphthol


(iii) with Br_2 water it forms white precipitate fo 2 ,4 ,6 tribromo aniline .

'X' can be identified as :











Answer: B

C.





5. Identify the reactant 'R'








Answer: A













Answer: C




7. A mixture of two organic compound gives red coloured precipitate with cuprous chloride and silver mirror on heating with Zn and NH_4Cl followed by $AgNO_3 + NH_4OH$ solution . The mixture contains :










Answer: A





which of the following reagents will not react with above

compound?

A. Na metal

B. $AgNO_3 + NH_4OH$


C. $Cu_2Cl_2 + NH_4OH$


D. $NaHCO_3$


Answer: D


> Watch Video Solution

9. Compound 'P', C_7H_8O is insolution in water , dilute when HCI and $NaHCO_3$ it disolves in dilite NaOH P is treated with $Br_2 - H_2O$ it converts rapidly into a compound of formula $C_7H_5OBr_3$ Idenity structure of P?










Answer: C





can be differentiated by :

A. carbylamine reaction


B. H_2SO_4

C. diazotisation followed by β -naphthol

D. mustard oil reaction

Answer: C





can be differentiated by :


A. $FeCl_3$

B. NaOH

- $\mathsf{C.} NaNO_2 + HCl$
- D. Fehling's solution

Answer: A





differentiated by :

A. NaOH

B. Na metal

C. $NaHCO_3$

D. $FeCl_3$

Answer: C





can be differentiated by :

A. carbylamine reaction

B. iodoform test

C. cold $KMnO_4$

D. $Br_2 - H_2O$

Answer: A



14.
$$CH_3 - \overset{O}{\overset{||}{C}} - H$$
 and $Ph - \overset{O}{\overset{||}{C}} - H$ can be

differentiated by:

A. Tollen's reagent

B. Fehling's solution

C. Lucas reagent


D. Victor meyer's test

Answer: B



15. Compound 'X' give positive test with 2,4- DNP and with

 $I_2\,/\,NaOH$ compound (X) may be :



D. all of these

Answer: D



16. An organic compound containing one oxygen gives red colour with cerric ammonium nitrate solution ,

decolourise alkaline $KMnO_4$, respond iodoform test and show geometrical isomerism . It should be :

A.
$$Ph - CH = CH - CH_2OH$$

B. $Ph - CH = CH - \overset{OH}{CH} - CH_3$
C. $Ph - CH = CH - O - CH_3$
(d) $Ph \longrightarrow OH$

Answer: B



17. Which of the following is true?

A. Alcohol give red colour with cerric ammonium

nitrate

B. Aldehyde and ketone give orange red colour with

2,4-DNP

C. RCOOH give CO_2 with $NaHCO_3$

D. All are true


Answer: D

Watch Video Solution

18. Compound (A) $C_{12}H_{20}$ discharges the colour of Br_2-H_2O and cold $KMnO_4$. On reduction with $H_2/{
m Pt}$

it gives compound (B) $C_{12}H_{22}$. A on ozonolysis give

cyclohexanone . Find structure of A :



D. None of these

Answer: A



19. Which of the following is true?

A. Tollen's reagent gives a positive test with all

aldehyde

B. Fehling's solution gives a positive test with all

C. Tollen's reagent gives a positive test with all

caroxylic acid

D. Tollen's reagent gives a positive test with lpha-methyl

keto


Answer: A



20. A monocarboxylic acid decolourise Br_2-H_2O , on heating with soda lime derivate of styrene is formed ,

with neutral $FeCl_3$, a buff coloured precipitate is formed

. Acid could be :






Answer: D



21. Which of the following compounds decolourise $Br_2 - H_2O$ and also give positive test with neutral $FeCl_3$:



Answer: C



22. Lassaigne's test for the detection of N fails in :

$$\stackrel{O}{\stackrel{\scriptstyle ||}{\scriptstyle \mid}}$$
A. $NH_2-\stackrel{O}{C}-NH-NH_2$

$$\mathsf{B.} NH_2 - NH_2$$

C.
$$NH_2 - \mathop{C}\limits_{\substack{||\\ O}} - NH_2$$

D.
$$C_6H_5 - NH - NH_2$$

Answer: B

Watch Video Solution

23. Which of the following compounds give positive test

with Tollen's reagent ?

$$A. H - \stackrel{O}{C} - OH$$

$$B. \stackrel{(b)}{\overbrace{}} \stackrel{O}{\overbrace{}} OH$$

$$C. CH_3 - CH - OC_2H_5$$

$$OH OH$$

$$OH O$$


$$D. CH_3 - \stackrel{||}{C} - H$$


Answer: A::B::C::D



24. Which of the following compound give negative test with Tollen's reagent ?

A.
$$Ph - \overset{O}{\overset{||}{C}} - H$$





$$\stackrel{O}{\stackrel{||}{\stackrel{}_{\scriptstyle \parallel}{\stackrel{}_{\scriptstyle \parallel}{\stackrel{}}_{\scriptstyle \parallel}{\stackrel{}_{\scriptstyle \parallel}{\stackrel{}}_{\scriptstyle \scriptstyle \scriptstyle}}{\stackrel{}}}}}}}$$
 D. CH_3 = CH_3

Answer: B::C::D



25. Which of the following reagents cannot be used for

differentiation between glucose and fructose ?

A. Lucas reagent

 $\mathsf{B.}\,Br_2-H_2O$

C. Tollen's reagent

D. 2,4- DNP

Answer: A::C



26. Which of the following reagents can be used to differentiate between Ph-C-H and $CH_3CH_2OH?$


A. NaOl

B. Fehling's solution

C. Tollen's reagent

D. $ZnCl_2$ / H


Answer: A::B




Answer: A::C::D

28. Which of the following compounds will react with $NaNH_2$?

A. $CH_3-\ \equiv CH$






C.

D.
$$Ph - egin{smallmatrix} O \ ert ert \ S \ ert \ O \ ert \ ert \ O \ ert \ e$$

Answer: A::B::C::D




Answer: A::C



30. Which of the following compounds may give reaction

with acidic $KMnO_4$?



D.

Answer: A::C::D



31. Which of the following reagents can be used to differentiate 1° and 3° alcohols ?

A. pcc

B. $K_2 Cr_2 O_7 \,/\, H^{\,\oplus}$

C. Jones reagent

D. $Br_2 - H_2O$

Answer: A::B::C

View Text Solution

32. Which of the following reagents cannot be used for differentiation between CH_3CHO and $CH_3 - C - Ph$?

A. NaOl

B. Tollen's agent

 $\mathsf{C}.\,H_2N-OH$


D. $Ph - NH - NH_2$

Answer: A::C::D

View Text Solution

33. Which of the following will not give white precipitate with ammoniacal silver nitrate solution ?

A. $CH_3 - C \equiv C - CH_3$





D. (d) Ph

Answer: A::B::C

View Text Solution

34. Which of the following tests can be used for differentiation among , 1° , 2° and 3° alcohol?

A. Lucas test

B. Victor meyer's test

C. Cu/ $300^{\circ}C$

D. Haloform reaction

Answer: A::B::C

D View Text Solution

35. Which of the following test can be used for identification of 1° amine ?

A. Carbylamine test

B. Hofmann mustard oil reaction


 $\mathsf{C}. NaNO_2 / HCl$

D. Fehling's solution

Answer: A::B::C



36. Unknown compound (A) $C_5H_{10}O$ gives positive test with 2,4-DNP but negative test with Tollen's reagent . It also give yellow precipitate with $I_2 / NaOH$. (A) is :








Answer: B::C




37. Compound (A) C_7H_8O is insoluble in aqueous $NaHCO_3$ and dissolves in aqueous NaOH and gives a characteristic colour with neutral $FeCl_3$. When treated with $Br_2(A)$ forms compound $(B)C_7H_5OBr_3$. The structure of compound (B) would be :











Answer: B





38. Compound (A) C_7H_8O is insoluble in aqueous $NaHCO_3$ and dissolves in aqueous NaOH and gives a characteristic colour with neutral $FeCl_3$. When treated with $Br_2(A)$ forms compound $(B)C_7H_5OBr_3$. What could be the structure of compound (A) if neither dissolves in aq. $NaHCO_3$ nor gives a characteristic colour with $FeCl_3$?











Answer: A::B



39. From the following sequence of reactions,

[A] $(C_6H_{12}) \stackrel{HCl}{\longrightarrow} (B)(C_6H_{13}Cl) + (C)(C_6H_{13}Cl)$ react

with $AgNO_3$ to give white ppt.

[B] $\xrightarrow{
m Alc.KOH}$ (D) (An isomer of A) gives positive test with Br_2/CCl_4

[D] $\xrightarrow{\text{Ozonelysis}}$ (E) gives positive iodoform test and


negative Fehling's test .


[A] $\xrightarrow{\mathrm{Ozonolysis}}$ (F) + (G) , both F and G give positive Tollen's


test.

 $[F] + [G] \xrightarrow{\text{Conc.NaOH}} \text{HCOONa + alcohol}$

The structure of C is :







Answer: C

D.



40. From the following sequence of reactions,

[A] $(C_6H_{12}) \stackrel{HCl}{\longrightarrow} (B)(C_6H_{13}Cl) + (C)(C_6H_{13}Cl)$ react

with $AgNO_3$ to give white ppt.

[B] $\xrightarrow{\text{Alc.KOH}}$ (D) (An isomer of A) gives positive test with Br_2/CCl_4

[D] $\xrightarrow{\text{Ozonelysis}}$ (E) gives positive iodoform test and

negative Fehling's test.


[A] $\xrightarrow{\mathrm{Ozonolysis}}$ (F) + (G) , both F and G give positive Tollen's

test .

 $[F] + [G] \xrightarrow{\text{Conc.NaOH}} \text{HCOONa + alcohol}$

The reaction involve in F and G with the NaOH is :

The structure of compound D is :



Answer: D



41. From the following sequence of reactions,

 $[\mathsf{A}] \ (C_6H_{12}) \stackrel{HCl}{\longrightarrow} (B)(C_6H_{13}Cl) + (C)(C_6H_{13}Cl) \text{ react}$

with $AgNO_3$ to give white ppt.

[B] $\stackrel{
m Alc.KOH}{
m \longrightarrow}$ (D) (An isomer of A) gives positive test with Br_2/CCl_4

[D] $\xrightarrow{\text{Ozonelysis}}$ (E) gives positive iodoform test and

negative Fehling's test .

[A] $\xrightarrow{\mathrm{Ozonolysis}}$ (F) + (G) , both F and G give positive Tollen's

test .

 $[F] + [G] \xrightarrow{\text{Conc.NaOH}} \text{HCOONa + alcohol}$

The reaction involve in F and G with the NaOH is :


The reaction involve in the F and G with NaOH is :

A. Reimer- Tiemann reaction


- B. Aldol condensation
- C. Cannizzaro reaction
- D. Perkin reaction

Answer: C









Column (II)

P. Decolourise Br₂ water

Q. Effervescence of CO_2 on reaction with NaHCO₃

R. Oxidation with alkaline $KMnO_4$

S. React with Na metal


View Text Solution



43.


(a) CH₃ —C≡CH O (b) H—C—OH **44.**

(c) $\sim NH_2$



- Column (II)
- P. Positive test with Fehling's solution
- Q. Positive test with Tollen's reagent
- R. Decolourise $Br_2 H_2O$
- S. Isocyanide test

View Text Solution





 $\operatorname{Column}(I)$ (a)Presence of halogen **46.** (b)Presence of sulphur $Q. Na_2[Fe(CN)_5NO]$ (c) Presence of nitrogen $R. Co(NO_3)_2$ (d)Presence of P and S

 $\operatorname{Column}(II)$ $P. HNO_3 / AgNO_3$ S. $FeCl_3$

