



## CHEMISTRY

### BOOKS - NARENDER AVASTHI CHEMISTRY (HINGLISH)

### CHEMICAL EQUILIBRIUM

#### Exercise

1. A reversible reaction is one which

- A. proceeds in one direction
- B. proceeds in both directions
- C. proceeds spontaneously
- D. all the statements are wrong

**Answer: b**



Watch Video Solution

2. The equilibrium constant  $K_c$  for the reaction



is 1.4 at  $400^\circ C$ . Suppose that 3 moles of  $P_4(g)$  and 2 moles of  $P_2(g)$  are mixed in 2 litre container at  $400^\circ C$ . What is the value of reaction quotient ( $Q_c$ )?

A.  $\frac{3}{2}$

B.  $\frac{2}{3}$

C. 1

D. none of these

**Answer: b**

[Watch Video Solution](#)

3. In a chemical reaction, equilibrium is said to have been established when the

- A. opposing reaction ceases
- B. concentrations of reactants and product are equal
- C. velocity of opposing reaction is the same as that of forward reaction
- D. reaction ceases to generate heat

Answer: bc



Watch Video Solution

4. The equilibrium constant for a reaction is  $K$ , and the reaction quotient is  $Q$ . For a particular reaction mixture, the ratio  $\frac{K}{Q}$  is 0.33. this means that:

- A. the reaction mixture will equilibrate to form more reactant species

- B. the reaction mixture will equilibrate to form more product species
- C. the equilibrium ratio of reactant to product concentration will be 3
- D. the equilibrium ratio of reactant to product concentrations will be 0.33

**Answer: b**

 [Watch Video Solution](#)

5. Consider the reaction  $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$  for which  $K_c = 278M^{-1}$ . 0.001 mole of each of the reagents  $SO_2(g)$ ,  $O_2(g)$  and  $SO_3(g)$  are mixed in a 1.0 L flask. Determine the reaction quotient of the system and the spontaneous direction of the system:

- A.  $Q_c = 1000$ , the equilibrium shifts to the right
- B.  $Q_c = 1000$ , the equilibrium shifts to the left
- C.  $Q_c = 0.001$ , the equilibrium shifts to the left

D.  $Q_c = 0.001$ , the equilibrium shifts to the right

**Answer: a**

 [Watch Video Solution](#)

6. In Q.No .5, if the mixture of gases was allowed to come to equilibrium .The volume of the reaction vessel was then rapidly increased by a factor of two .As a result of the change the reaction quotient ( $Q_c$ ) would:

- A. increase because of the pressure decrease
- B. decrease because of the pressure decrease
- C. remain the same because the equilibrium constant is independent of volume
- D. increase because the reaction is endothermic

**Answer: a**

 [Watch Video Solution](#)

7. For the reaction  $A(g) + 3B(g) \rightleftharpoons 2C(g)$  at  $27^\circ C$ , 2 moles of A, 4 moles of B and 6 moles of C are present in 2 litre vessel. If  $K_c$  for the reaction is 1.2, the reaction will proceed in :

- A. Forward direction
- B. backward direction
- C. neither direction
- D. none of these

**Answer: a**

 [Watch Video Solution](#)


8. For a reversible gaseous reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  at equilibrium , if some moles of  $H_2$  are replaced by same number of moles of  $T_2$  (T is tritium , isotope of H and assume isotopes do not have different chemical properties ) without affecting other parameters , then:

- A. the sample of ammonia obtained after something will be radioactive .
- B. moles of  $N_2$  after the change will be different as compared to moles of  $N_2$  present before the change
- C. the value of  $K_p$  or  $K_c$  will change
- D. the average molecular mass of new equilibrium will be same as that of old equilibrium


**Answer: a**

 **Watch Video Solution**


9. For the synthesis of ammonia by the reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  in the Haber's process ,the attainment of equilibrium is correctly predicated by the curve




A.



B.



C.



D.


**Answer: a**

 **Watch Video Solution**

10. The figure shows the change in concentration of species A and B as a function of time.



The equilibrium constant  $K_c$  for the reaction  $A(g) \rightleftharpoons 2B(g)$  is :




- A.  $K_c > 1$
- B.  $K < 1$
- C.  $K = 1$
- D. data insufficient

Answer: a



Watch Video Solution

11. Attainment of the equilibrium  $A(g) \rightleftharpoons 2C(g) + B(g)$  gave the following graph. Find the correct option. (% dissociation = Fraction dissociated  $\times 100$ )




- A. At  $t=5$  sec equilibrium has been reached and  $K_c = 40(\text{mol/litre})^2$
- B. At  $t=5$  sec equilibrium has been reached and % dissociation of A is 20%
- C. At  $t=5$  sec equilibrium has been reached and % dissociation of A is 30%
- D. none of these

**Answer: b**



Watch Video Solution

12. Using molar concentrations, what is the unit of  $K_c$  for the reaction ?



A.  $M^{-2}$

B.  $M^2$

C.  $M^{-1}$


D. M

Answer: b



Watch Video Solution

13. What is the unit of  $K_p$  for the reaction ?



A. atm

B.  $\text{atm}^{-2}$


C.  $\text{atm}^2$

D.  $\text{atm}^{-1}$

**Answer: b**

 [Watch Video Solution](#)

**14.** What is the equilibrium expression for the reaction



A.  $K_c = [O_2]^5$

B.  $K_c = [P_4O_{10}] / 5[P_4][O_2]$


C.  $K_c = [P_4O_{10}] / [P_4][O_2]^5$

D.  $K_c = 1 / [O_2]^5$


**Answer: d**

 [Watch Video Solution](#)

15. At  $527^\circ\text{C}$ , the reaction given below has  $K_c = 4$



what is the  $K_p$  for the reaction ?



A.  $16 \times (800R)^2$

B.  $\left(\frac{800R}{4}\right)^{-2}$

C.  $\left(\frac{1}{4 \times 800R}\right)^2$


D. none of these

**Answer: c**




**Watch Video Solution**

16. The equilibrium constant for the reaction



at temperature T is  $4 \times 10^{-4}$ .

The value of  $K_c$  for the reaction



at the same temperature is

A.  $4 \times 10^{-4}$

B. 50

C.  $2.5 \times 10^2$

D. 0.02


**Answer: b**



**Watch Video Solution**

17. The equilibrium constant  $K_c$  for the following reaction at  $842^\circ C$  is

$7.90 \times 10^{-3}$ . What is  $K_p$  at same temperature ?



A.  $8.64 \times 10^{-5}$

B.  $8.26 \times 10^{-4}$

C.  $7.90 \times 10^{-2}$


D.  $7.56 \times 10^{-2}$

**Answer: d**

 [Watch Video Solution](#)

18. The equilibrium constant  $K_p$  for the following reaction at  $191^\circ\text{C}$  is 1.24.

what is  $K_c$ ?



A. 6.7

B. 0.61

C. 8.30

D. 7.6

**Answer: d**

 [Watch Video Solution](#)

19. For the equilibrium  $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$ , what is the temperature at which  $\frac{K_p(atm)}{K_c(M)} = 3$ ?

A.  $0.027K$

B.  $0.36K$


C.  $36.54K$

D.  $273K$

Answer: c

 [Watch Video Solution](#)

20. For the reversible reaction



at  $500^\circ C$ , the value of  $K_p$  is  $1.44 \times 10^{-5}$  when the partial pressure is measured in atmosphere. The corresponding value of  $K_c$  with concentration in  $\text{mol } L^{-1}$  is



A.  $1.44 \times 10^{-5} / (0.082 \times 500)^{-2}$

B.  $1.44 \times 10^{-5} / (8.314 \times 773)^{-2}$

C.  $1.44 \times 10^{-5} / (0.082 \times 773)^2$

D.  $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$

Answer: d

 Watch Video Solution

21. For the reaction  $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$  the value of  $\left(\frac{K_c}{K_P}\right)$

is equal to :

A.  $\sqrt{RT}$

B.  $RT$

C.  $\frac{1}{RT}$

D. 1.0

Answer: b



[Watch Video Solution](#)

22. The concentration of a pure solid or liquid phase is not included in the expression of equilibrium constant because :

- A. density of solid and liquid are independent of their quantities .
- B. solids and liquids react slowly.
- C. solids and liquids at equilibrium do not interact with gaseous phase.
- D. the molecules of solids and liquids cannot migrate to the gaseous phase.

**Answer: a**



[Watch Video Solution](#)

23. A catalyst is a substance which

- A. increase the equilibrium concentration of the product.
- B. change the equilibrium constant of the reaction.
- C. shortens the time to reach equilibrium.
- D. supplies energy to the reaction.

**Answer: c**

 [Watch Video Solution](#)

**24.** What will be the effect of the equilibrium constant on increasing temperature. If the reaction neither absorbs heat nor releases heat?

- A. Equilibrium constant will remain constant.
- B. Equilibrium constant will decrease .
- C. Equilibrium constant will increase.
- D. Can not be predicted.

**Answer: a**

 [Watch Video Solution](#)

25. The equilibrium constant for a reaction

$N_2(g) + O_2(g) = 2NO(g)$  is  $4 \times 10^{-4}$  at  $2000K$ . In the presence of catalyst, the equilibrium constant is attained 10 times faster. The equilibrium constant in the presence of catalyst, at  $2000K$  is

A.  $40 \times 10^{-4}$

B.  $4 \times 10^{-4}$

C.  $4 \times 10^{-3}$

D. difficult to compute without more data

**Answer: a**

 [Watch Video Solution](#)

26. For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$

the equilibrium constant  $K_p$  changes with

A. total pressure

B. catalyst




C. concentration of  $H_2$  and  $I_2$

D. temperature

Answer: d

 Watch Video Solution

27. Consider the reaction :-



Which of the following relation is correct ?

A.  $K_3 = \frac{K_1}{K_2}$

B.  $K_3 = \frac{K_1^2}{K_2^2}$

C.  $K_3 = K_1 K_2$

$$D. K_3 = \sqrt{K_1 \cdot K_2}$$

Answer: d



Watch Video Solution

28. For the reaction  $2NO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons N_2O_5(g)$  if the equilibrium constant is  $K_p$ , then the equilibrium constant for the reaction  $2N_2O_5(g) \rightleftharpoons 4NO_2(g) + O_2(g)$  would be :

A.  $K_p^2$

B.  $\frac{2}{K_p}$

C.  $\frac{1}{K_p^2}$


D.  $\frac{1}{\sqrt{K_p}}$

Answer: c




Watch Video Solution

29. The equilibrium constant ( $K_c$ ) for the reaction



is  $4 \times 10^{-34}$  at  $25^\circ C$ . what is the equilibrium constant for the reaction ?



A.  $2 \times 10^{-17}$

B.  $2.5 \times 10^{33}$

C.  $5 \times 10^6$


D. none of these

Answer: d




Watch Video Solution

30. At a certain temperature , the following reactions have the equilibrium constants as shown below:



what is the equilibrium constant  $K_c$  for the reaction at the same

temperature?



A.  $2.5 \times 10^{76}$

B.  $4 \times 10^{23}$

C.  $4 \times 10^{-77}$




D. none of these

**Answer: c**




**Watch Video Solution**

**31. Given**



The equilibrium constant for



will be



A.  $K_1K_2K_3$

B.  $\frac{K_1K_2}{K_3}$

C.  $\frac{K_1K_3^2}{K_2}$

D.  $\frac{K_1K_3^2}{K_3}$

Answer: d



Watch Video Solution

32. In the reaction  $X(g) + Y(g) \rightleftharpoons 2Z(g)$ , 2 mole of X, 1 mole of Y and 1 mole of Z are placed in a 10 litre vessel and allowed to reach equilibrium. If final concentration of Z is 0.2 M, then  $K_c$  for the given reaction is :

A. 1.60

B.  $\frac{80}{3}$

C.  $\frac{16}{3}$

D. none of these

Answer: c



Watch Video Solution

33. An equilibrium mixture of the reaction  $2H_2S(g) \rightleftharpoons 2H_2(g) + S_2(g)$  had 0.5 mole  $H_2S$ , 0.10 mole  $H_2$  and 0.4 mole  $S_2$  in one litre vessel. The value of equilibrium constants (K) in mole  $litre^{-1}$  is

A. 0.0004

B. 0.008

C. 0.016

D. 0.160

Answer: c




Watch Video Solution

34.

Given

$[CS_2] = 0.120M$ ,  $[H_2] = 0.10$ ,  $[H_2S] = 0.10$ ,  $[H_2S] = 0.20$  and  $[CH_4] = 8$

for the following reaction at  $900^\circ C$  at eq. Calculate the equilibrium constant ( $K_c$ ).



A. 0.0120

B. 0.0980

C. 0.280

D. 0.120

**Answer: c**



[Watch Video Solution](#)

35. The equilibrium constant for the following reaction is 10.5 at 500 K .A


system

at

equilibrium

has

$[CO] = 0.250M$  and  $[H_2] = 0.120M$  what is the  $[CH_3OH]$ ?



A. 0.0378

B. 0.435

C. 0.546

D. 0.0499

**Answer: a**



[Watch Video Solution](#)

**36.** When sulphur ( in the form of  $S_B$ ) is heated at temperature  $T$ , at equilibrium , the pressure of  $S_B$  falls by 30 % from  $1.0atm$ , because  $S_B(g)$  is partially converted into  $S_2(g)$ .

Find the value of  $K_P$  for this reaction.

A. 2.96

B. 6.14

C. 204.8

D. none of these

**Answer: a**

 [Watch Video Solution](#)

37. 9.2 grams of  $N_2O_{4(g)}$  is taken in a closed one litre vessel and heated till the following equilibrium is reached  $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ . At equilibrium, 50 %  $N_2O_{4(g)}$  is dissociated. What is the equilibrium constant (in mol  $litre^{-1}$ ) (Molecular weight of  $N_2O_4 = 92$ ) ?

A. 0.1

B. 0.4

C. 0.2

D. 2

**Answer: c**

 [Watch Video Solution](#)

38. Two moles of  $NH_3$  when put into a previously evacuated vessel (one litre) partially dissociate into  $N_2$  and  $H_2$ . If at equilibrium one mole of  $NH_3$  is present, the equilibrium constant is

A.  $3/4 \text{ mol}^2 \text{ litre}^{-2}$

B.  $27/64 \text{ mol}^2 \text{ litre}^{-2}$

C.  $27/32 \text{ mol}^2 \text{ litre}^{-2}$


D.  $27/16 \text{ mol}^2 \text{ litre}^{-2}$

Answer: d



Watch Video Solution

39. In the presence of excess of anhydrous ( in torr) of water taken up is governed by  $K_p = 10^{12} \text{ atm}^{-4}$  for the following reaction at 273K



What is equilibrium vapour pressure ( in torr) of water in a closed vessel that contains  $SiCl_2 \cdot 2H_2O(s)$  ?

A. 0.001 torr

B.  $10^3$  torr

C. 0.76 torr


D. 1.31 → rr

Answer: c



Watch Video Solution

40.



If the vapour pressure of water is 38 torr then percentage of relative humidity is : (Assume all data at constant temperature)

A. 4


B. 10

C. 40

D. none of these

**Answer: c**

 [Watch Video Solution](#)



The equilibrium pressure at 25°C is 0.660 atm.  $\Delta G^\circ$  for the reaction ?

A. 0.109

B. 0.218

C. 1.89

D. 2.18

**Answer: a**

 [Watch Video Solution](#)



42. For the reaction  $2A(g) \rightleftharpoons B(g) + 3C(g)$ , at a given temperature,  $K_c = 16$ . What must be the volume of the flask, if a mixture of 2 mole each of A, B and C exist in equilibrium?

A.  $\frac{1}{4}$

B.  $\frac{1}{2}$

C. 1

D. none of these

Answer: b

 [Watch Video Solution](#)

43. One mole of pure ethyl alcohol was treated with one mole of pure acetic acid at  $25^\circ C$ . One-third of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be:

A.  $\frac{1}{4}$

B. 2


C. 3

D. 4

**Answer: a**



**Watch Video Solution**



This reaction is set-up in aqueous medium. We start with 1 mol of  $I_2$  and 0.5 mol of  $I^\ominus$  in 1L flask. After equilibrium reached, excess of  $AgNO_3$  gave 0.25 mol of yellow precipitate. Equilibrium constant is

A. 1.33

B. 2.66

C. 2.0


D. 3.0

**Answer: a**



**Watch Video Solution**

**45.** At  $87^\circ\text{C}$ , the following equilibrium is established.



If 0.3 mole hydrogen and 2 mole sulphur are heated to  $87^\circ\text{C}$  in a 2L vessel, what will be concentration of  $\text{H}_2\text{S}$  at equilibrium ?

A.  $0.11M$

B.  $0.022M$

C.  $0.044M$

D.  $0.08M$

**Answer: a**



**Watch Video Solution**

46. In the equilibrium  $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ , the partial pressure of  $SO_2$ ,  $O_2$  and  $SO_3$  are 0.662, 0.10 and 0.331 atm respectively. What should be the partial pressure of Oxygen so that the equilibrium concentrations of  $SO_3$  are equal?

A. 0.4atm

B. 1.0atm

C. 0.8atm

D. 0.25atm

**Answer: a**



**Watch Video Solution**

47. When heated, ammonium carbamate decomposes as follows:

$NH_4COOH(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$  At a certain temperature, the equilibrium pressure of the system is 0.318 atm.  $K_p$  for the reaction is:

A. 0.128

B. 0.426

C.  $4.76 \times 10^{-3}$

D. none of these

**Answer: c**

 [Watch Video Solution](#)

**48.** In the system  $A_{(s)} \rightleftharpoons 2B_{(g)} + 3C_{(g)}$ , if the concentration of  $C$  at equilibrium is increased by a factor of 2, it will cause the equilibrium concentration of  $B$  to change to:

A. Two times original value

B. One half of its original value

C.  $2\sqrt{2}$  times to the original value

D.  $\frac{1}{2\sqrt{2}}$  times the original value

**Answer: d**



**Watch Video Solution**

**49.**  $A + B \rightleftharpoons C + D$ . If finally the concentrations of A and B are both equal but at equilibrium concentration of D will be twice of that of A then what will be the equilibrium constant of reaction.

A.  $\frac{4}{9}$

B.  $\frac{9}{4}$

C.  $\frac{1}{9}$

D. 4

**Answer: d**



**Watch Video Solution**

50. The equilibrium  $K_c$  for the reaction  $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$  is 16. If 1 mole of each of all the four gases is taken in  $1\text{ dm}^3$  vessel, the equilibrium concentration of NO would be:

A.  $0.4\text{ M}$

B.  $0.6\text{ M}$

C.  $1.4\text{ M}$

D.  $1.6\text{ M}$

**Answer: d**



[Watch Video Solution](#)

51. On increasing the temperature, the rate of a reaction:

A. always increases

B. always decreases

C. first increases and then decreases

D. may increase or decrease depending upon the nature of the reaction

**Answer: a**

 [Watch Video Solution](#)

**52.** A catalyst increases the rate of a reaction by:

A. increasing the activation energy of a reaction

B. increasing the value of rate constant ( $K_f$  and  $K_b$ )

C. increasing the enthalpy change of the reaction


D. decreasing the enthalpy change of the reaction

**Answer: b**

 [Watch Video Solution](#)



53. At a certain temperature, only 50% HI is dissociated at equilibrium in the following reaction:



the equilibrium constant for this reaction is:

A. 0.25

B. 1.0

C. 3.0


D. 0.5

**Answer: a**



[Watch Video Solution](#)

54. The equilibrium constant  $K_p$  for the reaction



is 4.0 at  $1660^\circ C$ . Initially  $0.80 H_2$  and  $0.80 \text{ mol } CO_2$  are injected into a 5.0 litre flask. What is the equilibrium concentration of  $CO_2(g)$ ?

A. 0.533

B. 0.0534

C. 0.535

D. none of these

**Answer: b**



[Watch Video Solution](#)

55. At 273 K and 1atm , 10 litre of  $N_2O_4$  decompose to  $NO_2$  decomposes to  $NO_2$  according to equation



What is degree of dissociation ( $\alpha$ ) when the original volume is 25% less than that of existing volume?

A. 0.25

B. 0.33

C. 0.66

D. 0.5

Answer: b



Watch Video Solution

56. The equilibrium constant for the reaction  $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$  is 5. How many moles of  $CO_2$  must be added to 1 litre container already containing 3 moles each of CO and  $H_2O$  to make 2 M equilibrium concentration of CO?

A. 15

B. 19

C. 5

D. 20

Answer: b



Watch Video Solution

57. A nitrogen-hydrogen mixture initially in the molar ratio of 1:3 reached equilibrium to form ammonia when 25% of the  $N_2$  and  $H_2$  had reacted. If the pressure of the system was 21 atm, the partial pressure of ammonia at the equilibrium was:

A. 4.5 atm

B. 3.0 atm

C. 2.0 atm

D. 1.5 atm

**Answer: b**



[Watch Video Solution](#)

58.  $NH_3$  is heated at 15 atm, from  $25^\circ C$  to  $347^\circ C$  assuming volume constant. The new pressure becomes 50 atm at equilibrium of the reaction  $2NH_3 \rightleftharpoons N_2 + 3H_2$ . Calculate % moles of  $NH_3$  actually decomposed.

A. 65 %

B. 61.3 %

C. 62.5 %

D. 64 %

**Answer: b**



**Watch Video Solution**

59. 0.1 mole of  $N_2O_4(g)$  was sealed in a tube under one atmospheric conditions at  $25^\circ C$  Calculate the number of moles of  $NO_2(g)$  present, if the equilibrium  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$  ( $K_P = 0.14$ ) is reached after some time :

A.  $1.8 \times 10^2$

B.  $2.8 \times 10^2$

C. 0.034

D.  $2.8 \times 10^{-2}$

**Answer: c**



**Watch Video Solution**

60. 5 moles of  $SO_2$  and 5 moles of  $O_2$  are allowed to react. At equilibrium, it was found that 60% of  $SO_2$  is used up. If the pressure of the equilibrium mixture is one atmosphere, the partial pressure of  $O_2$  is :

A.  $0.52\text{atm}$

B.  $0.21\text{tm}$


C.  $0.41\text{aatm}$

D.  $0.82\text{atm}$

**Answer: c**



**Watch Video Solution**



For the reaction initially the mole ratio was 1:3 of  $N_2:H_2$ . At equilibrium 50% of each has reacted. If the equilibrium pressure is P, the partial pressure of  $NH_3$  at equilibrium is :

A.  $\frac{p}{3}$

B.  $\frac{P}{4}$

C.  $\frac{P}{6}$


D.  $\frac{p}{8}$

**Answer: a**



**Watch Video Solution**

62. 2.0 mole of  $PCl_5$  were introduced in a vessel of 5.0 L capacity of a particular temperature. At equilibrium,  $PCl_5$  was found to be 35 % dissociated into  $PCl_3$  and  $Cl_2$  the value of  $K_c$  for the reaction



A. 1.89

B. 0.377

C. 1.33


D. 13.3

**Answer: d**



**Watch Video Solution**

**63.** At certain temperature compound  $AB_2(g)$  dissociates according to the reaction



With degree of dissociation  $\alpha$  which is small compared with unity, the expression of  $K_p$  in terms of  $\alpha$  and initial pressure  $P$  is :

A.  $p \frac{\alpha^3}{2}$

B.  $\frac{P\alpha^2}{3}$

C.  $P \frac{\alpha^3}{3}$



D.  $\frac{P\alpha^2}{2}$

Answer: a



Watch Video Solution

64. For the reaction

$H_2(g) + CO(g) \rightleftharpoons CO(g) + H_2O(g)$ , if the initial concentration of  $[H_2] = [CO_2]$  and  $x$  moles /litres of hydrogen is consumed at equilibrium, the correct expression of  $K_p$  is :

A.  $\frac{x^2}{(1-x)^2}$

B.  $\frac{(1-x)^2}{(1-x)^2}$

C.  $\frac{x^2}{(2+x)^2}$

D.  $\frac{x^2}{(1-x)^2}$

Answer: a



Watch Video Solution

65. If  $D_T$  and  $D_o$  are the theoretical and observed vapour densities at a definite temperature and  $\alpha$  be the degree of dissociation of a substance, then  $\alpha$  in the terms of  $D_o$ ,  $D_T$  and  $n$  (number of moles of products formed from 1 mole reactant) is calculated by the formula :

A.  $\alpha = \frac{D_o - D_T}{(1 - n)D_T}$

B.  $\alpha = \frac{D_T - D_o}{(n - 1)D_T}$


C.  $\alpha = \frac{D_T - D_o}{(n - 1)D_o}$

D.  $\alpha = \frac{D - D_T}{(n - 1)D_T}$


Answer: c

 [Watch Video Solution](#)


66. For the dissociation of  $PCl_5$  into  $PCl_3$  and  $Cl_2$  in gaseous phase reaction, if  $d$  is the observed vapour density and  $D$  the theoretical vapour density with ' $\alpha$ ' as degree of dissociation, variation of  $D/d$  with ' $\alpha$ ' is given by ?



A.



B.



C.

D. none of these

**Answer: a**

 [Watch Video Solution](#)

67. At  $27^{\circ}C$  and 1 atm pressure,  $N_2O_4$  is 20% dissociation into  $NO_2$ .  
 .What is the density of equilibrium mixture of  $N_2O_4$  and  $NO_2$  at  $27^{\circ}C$  and 1 atm?

A.  $3.11\text{g/litre}$

B.  $2.11\text{g/litre}$

C.  $4.5\text{g/litre}$

D. none of these

**Answer: a**

 [Watch Video Solution](#)

68.  $\text{COCl}_2$  gas dissociates according to the equation,  $\text{COCl}_2 \rightleftharpoons \text{CO}(g) + \text{Cl}_2(g)$ . When heated to 700 K the density of the gas mixture at 1.16 atm and at equilibrium is  $1.16\text{g/litre}$ . The degree of dissociation of  $\text{COCl}_2$  at 700K is :

A. 0.28

B. 0.50

C. 0.72

D. 0.42

Answer: c



Watch Video Solution

69. The degree of dissociation of  $I_2$  molecule at  $1000^\circ C$  and under  $1.0\text{atm}$  is  $40\%$  by volume. If the dissociation is reduced to  $20\%$  at the same temperature, the total equilibrium pressure on the gas will be:

A.  $1.57\text{atm}$

B.  $2.57\text{atm}$

C.  $3.57\text{atm}$


D.  $4.57\text{atm}$

Answer: d



Watch Video Solution

70. Determine the value of equilibrium constant ( $K_C$ ) for the reaction



if 10 moles of  $A_2$ , 15 moles of  $B_2$  and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M:

A. 4.5

B. 1.5

C. 0.6


D. none of these

**Answer: a**



[Watch Video Solution](#)


71. At  $87^\circ C$ , the following equilibrium is established



If 0.50 mole of hydrogen and 1.0 mole of sulphur are heated to  $87^\circ C$

and 2.0 atm .the equilibrium gases mixture contains 40% chlorine by volume.

Calculate  $K_p$  at  $247^\circ C$  for the reaction



A. 0.966 atm

B. 1.38n atm

C. 0.0327 atm

D. 1atm


**Answer: a**



[Watch Video Solution](#)

**72.** Pure  $PCl_5$  is introduced into an evacuated chamber and to equilibrium at  $247^\circ C$  and 2.0 atm .The equilibrium gases mixure contains 40% choririne by volume .

Calculate  $K_p$  at  $247^\circ C$  for the reaction



A. 0.625 atm

B. 4atm

C. 1.6atm

D. none of these

**Answer: c**



**Watch Video Solution**

**73.** For the reaction

Calculate  $K_P$  at 900K where the equilibrium steam-hydrogen mixture was 45%  $H_2$  by volume :

A. 1.49

B. 1.22

C. 0.67

D. none of these



Answer: a

 Watch Video Solution

74. For the reaction

$XCO_3 \rightleftharpoons XO(s) + CO_2(g)$ ,  $K_p = 1.642 \text{ atm}$  at  $727^\circ C$  If 4 moles of  $XCO_3$

was put into a 50 litre container and heated to  $727^\circ C$

What mole percent of the  $XCO_3$  remains unreacted at equilibrium ?

A. 20

B. 25


C. 50

D. none of these

Answer: d

 Watch Video Solution

75.  $Fe_2O_3(s)$  may be converted to Fe by the reaction



What percentage of the  $H_2$  remains unreacted after the reaction has come to equilibrium?

A. ~22 %

B. ~34 %


C. ~66 %

D. ~78 %

**Answer: b**



[Watch Video Solution](#)



When the initial pressure of  $AB_3$  is 800 torr and the pressure developed at equilibrium is 900 torr, what fraction of  $AB_3(g)$  is dissociated?

A. 10 %

B. 20 %

C. 25 %


D. 30 %

**Answer: c**



**Watch Video Solution**

77. At 1000 K , a sample of pure  $NO_2$  gases decomposes as :



The equilibrium constant  $K_P$  is 156.25 atm .Analysis shows that the partial pressure of  $O_2$  is 0.25 atm at equilibrium .The parital pressure of  $NO_2$  at equilibrium is :

A. 0.01

B. 0.02

C. 0.04

D. none of these

Answer: b

 [Watch Video Solution](#)

78. pure nitrosyl chloride (NOCl) gas was heated to  $240^{\circ}\text{C}$  and the equilibrium pressure was  $1.0\text{ atm}$  and the  $K_p$ ?

A.  $1.02\text{ atm}$

B.  $16.875 \times 10^{-3}\text{ atm}$


C.  $16 \times 10^{-2}\text{ atm}$

D. none of these

Answer: b

 [Watch Video Solution](#)

79. At a certain temperature the equilibrium constant  $K_c$  is 0.25 for the reaction



If we take 1 mole of each of the four gases in a 10 litre container, what would be equilibrium concentration of  $A_2$  (g)?

A. 0.331 M

B. 0.033M

C. 0.133M

D. 1.33M

Answer: c



Watch Video Solution

80. At  $200^\circ C$   $PCl_5$  dissociates as follows :



It was found that the equilibrium vapours are 62 times as heavy as hydrogen. The degree of dissociation of  $PCl_5$  at  $200^\circ C$  is nearly :

A. 10 %

B. 42 %

C. 50 %

D. 68 %

**Answer: d**

 [Watch Video Solution](#)

**81.** For the dissociation reaction  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ , the degree of dissociation ( $\alpha$ ) in terms of  $K_p$  and total equilibrium pressure  $P$  is:

A.  $\alpha = \sqrt{\frac{4P + K_p}{K_p}}$

B.  $\alpha = \sqrt{\frac{K_p}{4P + K_p}}$

C.  $\alpha = \sqrt{\frac{K_p}{4P}}$


D. none of these

**Answer: b**


82. Consider the following equilibrium




Then select the correct graph, which shows the variation in concentrations of  $N_2O_4$  Against concentrations of  $N_2O_4$ :




A.



B.



C.



D.

**Answer: b**

[▶ Watch Video Solution](#)

83. The vapour pressure of mercury is 0.002 mm Hg at  $27^\circ C$ .  $K_c$  for the process  $Hg(l) \rightleftharpoons Hg(g)$  is :

A. 0.002

B.  $8.12 \times 10^{-5}$

C.  $6.48 \times 10^{-5}$

D.  $1.068 \times 10^{-7}$

**Answer: d**

[▶ Watch Video Solution](#)



84. Calculate the equilibrium constant ( $K_c$ ) for the reaction given below, if at equilibrium mixture contains 5.0 mole of  $A_2$ , 3 moles of  $B_2$  and 2 moles of  $AB_2$  at 8.21 atm and 300 K.

$$2AB_2(g) \rightleftharpoons A_2(g) + 2B_2(g) + \text{Heat}$$

A. 1.333

B. 2.66


C. 20

D. none of these

Answer: b

 [Watch Video Solution](#)

85. For the reaction (1) and (2)



Given,  $K_{p1} : K_{p2} = 9 : 1$

If the degree of dissociation of  $A(g)$  and  $X(g)$  be same then the total pressure at equilibrium

(1) and (2) are in the ratio:

A. 3 : 1

B. 36 : 1

C. 1 : 1

D. 0.5 : 1

**Answer: b**



[Watch Video Solution](#)

86. Given the following reaction at equilibrium



Some inert gas at constant pressure is added to the system. Predict which of the following facts:

A. more  $NH_3$  is produced

B. Less  $NH_3(g)$  is produced




C. No effect on the equilibrium

D.  $K_p$  of the reaction is decreased

**Answer: b**

 [Watch Video Solution](#)


**87.** In which of the following equilibrium, change in volume of the system does not alter the number of moles:



**Answer: a**

 [Watch Video Solution](#)

88. For the reaction



The number of moles of  $H_2$  at equilibrium will increase if:

- A. volume is increased
- B. volume is decreased
- C. argon gas is added at constant volume
- D.  $NH_3$  is removed

Answer: a



Watch Video Solution

89. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction



When equilibrium is re-established:

- A. The amount of  $Cl_2(g)$  remains unchanged

- B. the amount of  $Cl_2(g)$  increases
- C. The amount of  $SO_2(g)$  decreases
- D. The amount of  $SO_2(g)$  decreases

**Answer: b**

 [Watch Video Solution](#)

**90.** Some inert gas is added at constant volume to the following reaction at equilibrium

$N_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$  It is to be predicted the effect of adding the inert gas:

- A. The equilibrium shifts in the forward direction
- B. The equilibrium shifts in the backward direction
- C. The equilibrium remains unaffected
- D. The value of  $K_p$  is increased

Answer: c

 [Watch Video Solution](#)

91. Consider the reaction where  $K_p = 0.497$  at 500K




If the three gases are mixed in a rigid container so that the partial pressure of each gas is initially 1 atm, then which is the correct observation?

- A. More  $PCl_5$  will be produced
- B. More  $PCl_3$  will be produced
- C. Equilibrium will be established when 50% reaction is complete
- D. none of these

Answer: a

 [Watch Video Solution](#)

92. The preparation of  $SO_3(g)$  by reaction  $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$  is an exothermic reaction. If the preparation follows the following temperature-pressure relationship for % yield, then for temperatures  $T_1, T_2$  and  $T_3$  the correct option is:



A.  $T_3 > T_2 > T_1$

B.  $T_1 > T_2 > T_3$

$$C. T_1 = T_2 = T_3$$

D. Nothing could be predicated about temperature though given information

**Answer: b**



**Watch Video Solution**

93. In a vessel containing  $N_2$ ,  $H_2$  and  $NH_3$

at equilibrium, some helium gas is introduced.  $\xrightarrow{+}$  the pressure  $\in$  cause the dissociation of

$NH_3$ :

A. Increases

B. decreases

C. remains unaltered

D. changes unpredictably





**Answer: c**





Watch Video Solution

94. Le - Chatelier principle is not applicable to :






Answer: b



Watch Video Solution

95. Consider the following reactions .In which case the formation of product is favoured by decrease in pressure?





A. 2, 3

B. 3, 4

C. 2, 4



D. 1, 4

**Answer: d**



**Watch Video Solution**

**96.** In which of the following reactions, the formation of product is favoured by decrease in temperature ?



A. 1, 2

B. 2 only



C. 1,2,3

D. 3,4

**Answer: d**

 [Watch Video Solution](#)




97. For which of the following reaction is product formation favoured by low pressure and high temperature?



**Answer: b**

 [Watch Video Solution](#)



98. For which of the following reaction is product formation favoured by low pressure and high temperature?



Answer: c

 [Watch Video Solution](#)

99. Consider the following reaction at equilibrium and determine which of the indicated changes will cause the reaction to proceed to right.



(3)  $H_2(g) + F_2(g) \rightleftharpoons 2HF(g)$  (add  $F_2$ )(4)  $BaO(s) + SO_3(g)$

(g)  $H_2SO_4(s)$  (add  $BaO$ )`

A. 2, 3

B. 1,4

C. 2,4


D. 2,3,4

**Answer: a**



**Watch Video Solution**

**100.** If the pressure in a reaction vessel for the following reaction is increased by decreasing the volume ,what will happen to the concentrations of  $CO$  and  $CO_2$ ?



A. both the  $[CO]$  and  $[CO_2]$  will decrease

B. neither the  $[CO]$  nor the  $[CO_2]$  will change


C. the  $[CO]$  will decrease and the  $[CO_2]$  will increase

D. both the  $[CO]$  and  $[CO_2]$  will increase

**Answer: d**

 [Watch Video Solution](#)

**101.** Consider the following reaction and determine which of the conditions will shift the equilibrium position to the right ?



A. Increasing the temperature

B. increasing the pressure


C. adding a catalyst

D. none of above is correct

**Answer: d**

 [Watch Video Solution](#)

102. The conversion of ozone into oxygen is exothermic under what conditions is ozone is most stable?



- A. At low pressure and low temperature
- B. At high pressure and high temperature
- C. At high pressure and low temperature
- D. At low pressure and high temperature

Answer: b



Watch Video Solution

103. A System at equilibrium is described by the equation of fixed temperature T.



What effect will be the effect on equilibrium, if total pressure is resucing volume?

- A. Concentration of  $SO_2Cl_2(g)$  increases
- B. Concentrations of  $SO_2(g)$  increases
- C. Concentration of  $Cl_2(g)$  increases
- D. Concentration of all gases increaseses

**Answer: d**



[Watch Video Solution](#)

**104.** The reaction  $2NO_2(g) \rightleftharpoons N_2O_4(g)$  is an exothermic equilibrium .

This means that:

- A. equilibration of this gas mixture will be slower at high temperature
- B. A mole of  $N_2O_4$  will occupy twice the volume of a mole of  $NO_2$  at the same?




C. the equilibrium will move to the right if an equilibrium mixture is cooled

D. the position of equilibrium will move to the left with increasing gas pressure

**Answer: c**

 [Watch Video Solution](#)

105. Densities of diamond and graphite are  $\frac{3.5g}{mL}$  and  $\frac{2.3g}{mL}$ . 

$$\Delta_7 H = -1.9 \frac{kJ}{mole}$$

Favourable conditions for formation of diamond are:

A. high pressure and low temperature

B. low pressure and high temperature

C. high pressure and high temperature

D. low pressure and low temperature

**Answer: d**

 [Watch Video Solution](#)

**106.** For an equilibrium  $H_2O(s) \rightleftharpoons H_2O(l)$ , which of the following statements is true ?

- A. The pressure changes do not affect the equilibrium
- B. More of ice melts if pressure on the system is increased
- C. More of liquid freezes if pressure on the system is increased
- D. The pressure changes may increase or decrease the degree of advancement of the process

**Answer: b**

 [Watch Video Solution](#)

**107.** A pressure cooker reduces cooking time because

- A. the higher pressure inside the cooker crushes the food material
- B. cooking involves chemical change helped by a rise in temperature
- C. heat is more evenly distributed in the cooking space
- D. boiling point of water involved in cooking is increased

**Answer: d**

 [Watch Video Solution](#)

**108.** The vapour pressure of a liquid in a closed container depends upon

- A. 1 only
- B. 2 only
- C. 1 and 3 only
- D. 1,2,and3

**Answer: a**

 [Watch Video Solution](#)

**109.** The pressure on a sample of water at its triple point is reduced while the temperature is held constant. Which phase changes are favoured?

(1) melting of ice

(2) sublimation of ice

(3) vaporization of liquid water

A. 1 only

B. 3 only

C. 2 only


D. 2 and 3

**Answer: d**




**Watch Video Solution**


**110.** An exothermic reaction is represented by the graph :




A.



B.



C.




D.

**Answer: c**




**Watch Video Solution**


**111.** An endothermic reaction is represented by the graph :




A.



B.



C.




D.

**Answer: b**



**Watch Video Solution**

112. A schematic plot of  $\ln K_{eq}$  versus inverse of temperature for a reaction is shown below



the reaction must be:

- A. Exothermic
- B. Endothermic
- C. One with negligible enthalpy change
- D. Highly spontaneous at ordinary temperature

**Answer: a**

 [Watch Video Solution](#)

**113.** The correct relationship between free energy change in a reaction and the corresponding equilibrium constant  $K_c$  is:

A.  $\Delta G^\circ = RT \ln K$

B.  $\Delta G^\circ = -RT \ln K$


C.  $\Delta G = RT \ln K$

D.  $\Delta G = -RT \ln K$

**Answer: b**


 [Watch Video Solution](#)

**114.** For the chemical equilibrium,







$\Delta_r H^\ominus$  can be determined from which one of the following plots?




A.



B.



C.




D.

**Answer: a**



**Watch Video Solution**

115.  $K_p$  has the value of  $10^{-6} \text{ atm}^3$  and  $10^{-4} \text{ atm}^3$  at 298 K and 323 K respectively for the reaction



$\Delta_r H^\circ$  for the reaction is :

A.  $7.7\text{KJ/mol}$

B.  $-147.41\text{KJ/mol}$

C.  $147.41\text{KJ/mol}$

D. none of these

Answer: c



Watch Video Solution

**116.** Van's Hoff's equation shows the effect of temperature on equilibrium constants  $K_c$  and  $K_p$ . the  $K_P$  varies with tempertaure according to the realation:

A.  $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta H^\circ}{2.303R} \left( \frac{T_1 - T_2}{T_1 T_2} \right)$

B.  $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta H^\circ}{2.303R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$

C.

D.

**Answer: b**

 [Watch Video Solution](#)

117. For a reaction, the value of  $K_p$  increases with increase in temperature.

The  $\Delta H$  for the reaction would be :

A. positive

B. negative





C. zero

D. cannot be predicted

**Answer: A**

 [Watch Video Solution](#)

118. The most stable oxides of nitrogen will be :




Answer: A



Watch Video Solution

119. When 1 mole of pure ethyl alcohol ( $C_2H_5OH$ ) is mixed with 1 mole of acetic acid at  $25^\circ C$ . the equilibrium mixture contains  $2/3$  mole each of ester and water



The  $\Delta G^\circ$  for the reaction at  $298K$  is :

A. 3435 J

B. 4 J

C. - 3435 J

D. zero

**Answer: C**

 [Watch Video Solution](#)

**120.** The value of  $\Delta G^\circ$  for a reaction in aqueous phase having  $K_c = 1$ , would be :

A.  $-RT$

B.  $-1$

C. 0

D.  $+RT$

**Answer: C**

 [Watch Video Solution](#)


121. A plot of Gibbs energy of a reaction mixture against the extent of the reaction is :

- A. minimum at equilibrium
- B. zero at equilibrium
- C. maximum at equilibrium
- D. None of these

**Answer: A**

 [Watch Video Solution](#)

122. For the reaction at  $300K$



$$\Delta_r H^\circ = - kJ/mol, \Delta_r S^\circ = - 0.1K^{-1} \cdot mol^{-1}$$

What is the value of equilibrium constant ?

A. 0

B. 1


C. 10

D. None of these

**Answer: B**

 [Watch Video Solution](#)

**123.** Solid  $Ca(HCO_3)_2$  decomposes as



If the total pressure is 0.2 bar at 420K, what is the standard free energy change for the given reaction ( $\Delta_r G^\circ$ ) ?

A.  $840\text{kJ/mol}$

B.  $3.86\text{kJ/mol}$

C.  $6.98\text{kJ/mol}$

D.  $16.083\text{kJ/mol}$

**Answer: D**



**Watch Video Solution**

**124.** The standard free energy change of a reaction is  $\Delta G^\circ = -kJ/mol^{-1}$  at  $298K$ . Calculate the value of  $\log_{10} K_p$  ( $R = 8.314JK^{-1}mol^{-1}$ )

A. 20.16

B. 2.303

C. 2.016

D. 13.83



**Answer: A**




**Watch Video Solution**



125. The following equilibrium constants were determined at 1120K :



What is the equilibrium constant  $K_c$  for the foollowing reaction at 1120K :



A.  $3.31 \times 10^{11} M^{-1}$

B.  $5.5 \times 10^{10} M^{-1}$


C.  $5.51 \times 10^6 M^{-1}$

D. None of these


**Answer: A**

 [Watch Video Solution](#)

126. One mole of  $N_2$  (g) is mixed with 2 moles of  $H_2$ (g) in a 4 litre vessel If 50 % of  $N_2$ (g) is converted to  $NH_3$ (g) by the following reaction :



What will the value of  $K_c$  for the following equilibrium ?



A. 256

B. 16


C.  $\frac{1}{16}$

D. None of these

**Answer: C**



**Watch Video Solution**



127.


The gas  $A_2$  in the left flask allowed to react with gas  $B_2$  present in right flask as  $A_2(g) + B_2(g) \rightleftharpoons 2AB(g)$ ,  $K_c = 4$  at  $27^\circ\text{C}$ . What is the concentration of AB when equilibrium is established ?

- A. 1.33 M
- B. 2.66 M
- C. 0.66 M
- D. 0.33 M

Answer: C

 Watch Video Solution

**128.** Assume that the decomposition of  $HNO_3$  can be represented by the following equation



approaches equilibrium at  $400K$  temperature and  $30 \text{ atm}$  pressure. At equilibrium partial pressure of  $HNO_3$  is  $2 \text{ atm}$

Calculate  $K_c$  in  $(\text{mol}/L - K)$  at  $400K$

(Use:  $R = 0.08 \text{ atm} - L/\text{mol} - K$ )

A. 4

B. 8

C. 16

D. 32

**Answer: D**



**Watch Video Solution**

129. For the equilibrium:



at  $40^\circ C$ . A 5litre vessel contains 0.1 mole of  $LiCl \cdot NH_3$ . How many mole of  $NH_3$  should be added to the flask at this temperature to derive the backward reaction for completion?


- A. 0.2
- B. 0.59
- C. 0.69
- D. 0.79

**Answer: D**



[Watch Video Solution](#)

130. Solid Ammonium carbamate dissociates as:



In a closed vessel, solid ammonium carbonate is in equilibrium with its

dissociation products. At equilibrium, ammonia is added such that the partial pressure of  $NH_3$  at new equilibrium now equals the original total pressure. Calculate the ratio of total pressure at new equilibrium to that of original total pressure. Also find the partial pressure of ammonia gas added.

A. 4

B. 9

C.  $\frac{4}{9}$

D.  $\frac{2}{9}$

**Answer: C**



[Watch Video Solution](#)

**131.** For the reaction  $C_2H_6(g) \rightleftharpoons C_2H_4(g) + H_2(g)$

$K_p$  is  $5 \times 10^{-2}$  atm. Calculate the mole per cent of  $C_2H_6(g)$  at equilibrium if pure  $C_2H_6$  at 1 atm is passed over a suitable catalyst at  $900K$  :

A. 20

B. 33.33

C. 66.66

D. None of these

**Answer: C**



**Watch Video Solution**

**132.**  $2NOBr(g) \rightleftharpoons 2NO(g) + Br_2(g)$ . If nitrosyl bromide (NOBr) 40 % dissociated at certain temp. and a total pressure of 0.30 atm  $K_p$  for the reaction  $2NO(g) + Br_2(g) \rightleftharpoons 2NOBr(g)$  is

A. 45

B. 25

C. 0.022

D. 0.25

**Answer: A**



**Watch Video Solution**

**133.** Consider the partial decomposition of  $A$  as

$2A(g) \rightleftharpoons 2B(g) + C(g)$  At equilibrium 700mL gaseous mixture contains 100mL of gas C at 10 atm and 300K what is the value of  $K_p$  for the reaction ?

A.  $\frac{40}{7}$

B.  $\frac{1}{28}$

C.  $\frac{10}{28}$

D.  $\frac{28}{10}$


**Answer: C**



**Watch Video Solution**



134. At a certain temperature and 2 atm pressure equilibrium constant ( $K_p$ ) is 25 for the reaction




Initially if we take 2 moles of each of the four gases and 2 moles of inert gas, what would be the equilibrium partial pressure of  $NO_2$ ?

- A. 1.33 atm
- B. 0.1665 atm
- C. 0.133 atm
- D. None of these

**Answer: C**

 [Watch Video Solution](#)

135. 0.020 g of selenium vapour at equilibrium occupying a volume of 2.463 mL at 1 atm and  $27^\circ C$ . The selenium is in a state of equilibrium according to reaction



What is the degree of association of selenium ?

(At.mass of se = 79)

A. 0.205

B. 0.315

C. 0.14

D. None of these

**Answer: B**



[Watch Video Solution](#)

**136.** Determine the degree of association (polymerization) for the reaction in aqueous solution, if observed (mean) molar mass of HCHO and  $\text{C}_6\text{H}_{12}\text{O}_6$  is 150:  $6\text{HCHO} \rightleftharpoons \text{C}_6\text{H}_{12}\text{O}_6$

A. 0.50

B. 0.833

C. 0.90

D. 0.96

**Answer: D**



**Watch Video Solution**

**137.** A reaction system in equilibrium according to reaction  $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$  in one litre vessel at a given temperature was found to be 0.12 mole each of  $SO_2$  and  $SO_3$  and 5 mole of  $O_2$ . In another vessel of one litre contains 32 g of  $SO_2$  at the same temperature. What mass of  $O_2$  must be added to this vessel in order that at equilibrium 20 % of  $SO_2$  is oxidized to  $SO_3$ ?

A. 0.4125

B. 11.6 g

C. 1.6 g

D. None of these

**Answer: B**

 [Watch Video Solution](#)

**138.** The equilibrium constant  $K_p$  for the following reaction is 4.5

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$  What would be the average molar mass (in  $g/mol$ ) of an equilibrium mixture of  $N_2O_4$  and  $NO_2$  formed by the dissociation of pure  $N_2O_4$  at a total pressure of 2 atm ?

A. 69

B. 57.5


C. 80.5

D. 85.5

**Answer: B**

 [Watch Video Solution](#)

139. A flask containing 0.5 atm pressure of  $A_2(g)$ , some solid AB added into flask which undergoes dissociation according to




The total pressure (in atm) at equilibrium is :

- A. 0.70
- B. 0.6
- C. 0.10
- D. None of these

**Answer: A**

 [Watch Video Solution](#)

140. A vessel of 250 litre was filled with 0.01 mole of  $Sb_2S_3$  and 0.01 mole of  $H_2$  to attain the equilibrium at  $440^\circ C$  as



After equilibrium, the  $H_2S$  formed was analysed by dissolved it in water and treating

with excess of  $Pb^{20+}$  to give 1.19 g of PbS as precipitate. What is the value of  $K_c$  at  $440^\circ C$  ?

A. 1

B. 2

C. 4

D. 8

**Answer: A**



[Watch Video Solution](#)

**141.** For the reaction  $2A(g) + B(g) \rightleftharpoons C(g) + D(g)$ ,  $K_c = 10^{12}$ .if initially 4,2,6,2 moles of A,B,C,D respectively are taken in a 1 litre vessel, then the equilibrium concentration of A is :

A.  $4 \times 10^{-4}$

B.  $2 \times 10^{-4}$


C.  $10^{-4}$

$$D. 8 \times 10^{-4}$$

**Answer: A**

 [Watch Video Solution](#)

**142.** The equilibrium constant for the following reaction in aqueous solution is 0.90.



How many mole of glycerin should be added per litre of  $0.10M H_3BO_3$  so that 80 % of the  $H_3BO_3$  is converted to the boric-acid glycerin complex ?

A. 4.44

B. 4.52

C. 3.6

D. 0.08

**Answer: B**



[Watch Video Solution](#)

143. Rate of diffusion of ozonized oxygen is  $0.4\sqrt{5}$  times that of pure oxygen what is the per cent degree of association of oxygen assuming pure  $O_2$  in the sample initially ?

- A. 20
- B. 40
- C. 60
- D. None of these

**Answer: C**



[Watch Video Solution](#)

144. One mole of  $SO_3$  was placed in a two litre vessel at a certain temperature. The following equilibrium was established in the vessel





The equilibrium mixture reacts with 0.2 mole  $KMnO_4$  in acidic medium.

Hence,  $K_c$  is :

A. 0.50

B. 0.25

C. 0.125

D. None of these

**Answer: C**



[Watch Video Solution](#)

**145.** At  $800^\circ C$ , the following equilibrium is established as



The composition of equilibrium may be determined by measuring the rate of effusion of the mixture through a pin hole. It is found that at  $800^\circ C$  and 1 atm mixture effuses 1.6 times as fast as  $SO_2$  effuses under the similar conditions. (At. mass of F = 19) what is the value of  $K_p$  (in atm)

?

A. 0.315

B. 0.685

C. 0.46


D. 1.49

**Answer: D**



**Watch Video Solution**

**146.** The equilibrium constant for the ionization of  $RNH_2$  (g) in water as



is  $8 \times 10^{-6}$  at  $25^\circ C$ . find the pH of a solution at equilibrium when pressure of  $RNH_2(g)$  is 0.5 bar :

A.  $\approx 12.3$

B.  $\approx 11.3$

C.  $\approx 11.45$

D. None

**Answer: B**

 [Watch Video Solution](#)

**147.** When  $N_2O_5$  is heated at certain temperature, it dissociates as  $N_2O_5(g) \rightleftharpoons N_2O_3(g) + O_2(g)$ ,  $K_c = 2.5$ . At the same time  $N_2O_3$  also decomposes as :



$N_2O_3(g) \rightleftharpoons N_2O(g) + O_2(g)$ . "If initially" 4.0 moles of  $N_2O_5$  "are taken in" 1.0 litre flask and allowed to dissociate. Concentration of  $O_2$  at equilibrium is 2.5 M. "Equilibrium concentration of "  $N_2O_5$  is :

- A. 1.0 M
- B. 1.5M
- C. 2.166M
- D. 1.846 M

**Answer: D**

 [Watch Video Solution](#)

**148.** Two solid compounds X and Y dissociates at a certain temperature as follows



The total pressure of gases over a mixture of X and Y is :

- A. 4.5atm
- B. 0.45 atm
- C. 0.6 atm
- D. None of these

**Answer: B**



[Watch Video Solution](#)

**149.** For a gaseous reaction



equilibrium constants  $K_c$ ,  $K_p$  and  $K_x$  are represented by the following relation

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, K_p = \frac{P_C^c \cdot P_D^d}{P_A^a} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where  $[A]$  represents molar concentration of  $A$ ,  $p_A$  represents partial pressure of  $A$  and  $P$  represents total pressure,  $x_A$  represents mole fraction of  $A$

On the basis of above work-up select the write option

A.  $K_p = K_c(RT)^{\Delta ng}$ ,  $K_x = K_p(RT)^{\Delta ng}$

B.  $K_c = K_c(RT)^{\Delta ng}$ ,  $K_p = K_x P^{\Delta ng}$

C.  $K_c = K_x P^{\Delta ng}$ ,  $K_p = K_x P^{\Delta ng}$


D.  $K_c = K_p(RT)^{-\Delta ng}$ ,  $K_x = K_p(RT)^{\Delta ng}$

**Answer:**



**Watch Video Solution**

150. For a gaseous reaction



equilibrium constants  $K_c$ ,  $K_p$  and  $K_x$  are represented by the following relation

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, K_p = \frac{P^c \cdot P_D^d}{P_A^a} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where  $[A]$  represents molar concentration of  $A$ ,  $p_A$  represents partial pressure of  $A$  and  $P$  represents total pressure,  $x_A$  represents mole fraction of  $A$

For the reaction  $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$ ,  $K_p > K_x$  is obtained at :


- A. 0.5 atm
- B. 0.8 atm
- C. 1 atm
- D. 2atm

**Answer:**



**Watch Video Solution**


151. For a gaseous reaction



equilibrium constants  $K_c$ ,  $K_p$  and  $K_x$  are represented by the following relation

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, K_p = \frac{P^c \cdot P_D^d}{P_A^a} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where  $[A]$  represents molar concentration of  $A$ ,  $p_A$  represents partial pressure of  $A$  and  $P$  represents total pressure,  $x_A$  represents mole fraction of  $A$ . For the following equilibrium relation between  $K_c$  and  $K_x$  (in terms of mole fraction) is



A.  $K_c = K_x (RT)^{-1}$

B.  $K_c = K_x (RT)$

C.  $K_c = K_x \left( \frac{RT}{P} \right)$

D.  $K_c = K_x \left( \frac{P}{RT} \right)$

**Answer:**



152. Variation of equilibrium constant  $K$  with temperature is given by van't Hoff equation

$$\ln K = \frac{\Delta_r S^\circ}{R} - \frac{\Delta_r H^\circ}{RT}$$

for this equation,  $(\Delta_r H^\circ)$  can be evaluated if equilibrium constants  $K_1$  and  $K_2$  at two temperatures  $T_1$  and  $T_2$  are known.

$$\log\left(\frac{K_2}{K_1}\right) = \frac{\Delta_r H^\circ}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

For an isomerization  $X(g) \rightleftharpoons Y(g)$  the temperature dependency of equilibrium constant is given by:

$$\ln K = 2 - \frac{1000}{T}$$

The value of  $\Delta_r S^\circ$  at 300K is:

A.  $2R$

B.  $\frac{2}{R}$

C.  $1000R$

D. None of these



Answer:

 [Watch Video Solution](#)

**153.** Variation of equilibrium constant  $K$  with temperature is given by van't

Hoff equation

$$\ln K = \frac{\Delta_r S^\circ}{R} - \frac{\Delta_r H^\circ}{RT}$$

for this equation,  $(\Delta_r H^\circ)$  can be evaluated if equilibrium constants  $K_1$

and  $K_2$  at two temperatures  $T_1$  and  $T_2$  are known.

$$\log\left(\frac{K_2}{K_1}\right) = \frac{\Delta_r H^\circ}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

Select the correct statement :



- A. Value of  $K_{eq}$  always increases with increasing temperature
- B. For exothermic reaction value of  $K_{eq}$  increases with decreasing temperature
- C. For endothermic reaction value of  $K_{eq}$  increases with decreasing temperature
- D. For exothermic reaction slope is  $(\log K vs. 1/T)$  negative

**Answer:**



**Watch Video Solution**

**154.**  $N_2O_3$  is an unstable oxide of nitrogen and it decomposes into NO (g) and  $NO_2(g)$  where  $NO_2(g)$  is further dimerise into  $N_2O_4$  as



A flask is initially filled with pure  $N_2O_3(g)$  having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 bar.

The equilibrium partial pressure of  $N_2O_3(g)$  is :



- A. 0.5bar
- B. 1.0 bar
- C. 1.5 bar
- D. 0.1 bar

**Answer:**



**Watch Video Solution**

**155.**  $N_2O_3$  is an unstable oxide of nitrogen and it decomposes into NO (g) and  $NO_2(g)$  where  $NO_2(g)$  is further dimerise into  $N_2O_4$  as



A flask is initially filled with pure  $N_2O_3(g)$  having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 bar.

The equilibrium partial pressure of  $NO_2(g)$  is:

A. 0.066bar

B. 0.133 bar



C. 0.423 bar

D. 0.83 bar

**Answer:**

 [Watch Video Solution](#)

**156.**  $N_2O_3$  is an unstable oxide of nitrogen and it decomposes into NO (g) and  $NO_2(g)$  where  $NO_2(g)$  is further dimerise dimerise into  $N_2O_4$  as



A flask is initially filled with pure  $N_2O_3(g)$  having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 ber.

The value of  $K_{P2}$  is

A.  $0.16\text{bar}^{-1}$

B.  $0.32\text{bar}^{-1}$

C.  $0.48\text{bar}^{-1}$

D.  $0.64\text{bar}^{-1}$

**Answer:**



**Watch Video Solution**

**157.** If a system at equilibrium is subjected to a change of any one of the factors such as concentration, pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration become ' $Q$ ' (reaction quotient) and depending upon.

$Q < K$ : equilibrium will shift in forward direction

$Q > K$  equilibrium will shift in backward direction

Effect of change in pressure :

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is decrease in number of moles i.e., towards the direction in which there is decrease in volume.

Effect of change in pressure on melting point : There are two types of

solids :

Solids whose volume decreases on melting, e.g., ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume)  $\Leftrightarrow$  Liquid (higher volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

Solid whose volume increase on melting, e.g., Fe, Cu, Ag, Au, etc.

Solid (lower volume)  $\Leftrightarrow$  Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

Solubility of substances : When solid substance are dissolved in water, either heat is evolved.

for endothermic solubility process solubility increase with increase in temperature. For exothermic solubility decrease with increase in temperature.

Solubility of gases in liquids : when a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

Effect of temperature : For endothermic reaction as temperature

increases reaction shift in backward direction

'X'(g) solute when dissolved in water heat is evolved. Then solubility of 'X' will increase :

- A. high temperature, low pressure
- B. low temperature, high pressure
- C. high temperature, high pressure
- D. low temperature, high pressure

**Answer:**



[Watch Video Solution](#)

**158.** If a system at equilibrium is subjected to a change of any one of the factors such as concentration , pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration become 'Q' (reaction quotient) and depending upon.

$Q < K$ : equilibrium will shift in forward direction

$Q > K$  equilibrium will shift in backward direction

Effect of change in pressure :

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is decrease in number of moles i.e., towards the direction in which there is decrease in volume.

Effect of change in pressure on melting point : There are two types of solids :

Solids whose volume decreases on melting, e.g., ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume)  $\rightleftharpoons$  Liquid (higher volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

Solid whose volume increase on melting, e.g., Fe, Cu, Ag, Au, etc.

Solid (lower volume)  $\rightleftharpoons$  Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.



Solubility of substances : When solid substance are dissolved in water, either heat is evolved.

for endothermic solubility process solubility increase with increase in temperature. For exothermic solubility decrease with increase in temperature.

Solubility of gases in liquids : when a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

Effect of temperature : For endothermic reaction as temperature increases reaction shift in backward direction



Above equilibrium is favoured at :

- A. high pressure, low temperature
- B. high pressure, high temperature
- C. low pressure, high temperature
- D. low pressure, low temperature

**Answer:**



**159.** If a system at equilibrium is subjected to a change of any one of the factors such as concentration, pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration becomes 'Q' (reaction quotient) and depends upon.

$Q < K$ : equilibrium will shift in forward direction

$Q > K$  equilibrium will shift in backward direction

Effect of change in pressure :

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is a decrease in number of moles i.e., towards the direction in which there is a decrease in volume.

Effect of change in pressure on melting point : There are two types of solids :

Solids whose volume decreases on melting, e.g., ice, diamond,

carborundum, magnesium nitride and quartz.

Solid (higher volume)  $\Leftrightarrow$  Liquid (higher volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

Solid whose volume increase on melting, e.g., Fe, Cu, Ag, Au, etc.

Solid (lower volume)  $\Leftrightarrow$  Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.


Solubility of substances : When solid substance are dissolved in water, either heat is evolved.

for endothermic solubility process solubility increase with increase in temperature. For exothermic solubility decrease with increase in temperature.

Solubility of gases in liquids : when a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

Effect of temperature : For endothermic reaction as temperature increases reaction shift in backward direction

For the reaction



If pressure is increased by reducing the volume of the container then :

- A. total pressure at equilibrium will remain same
- B. concentration of all the component at equilibrium will change
- C. concentration of all the component at equilibrium will remain same
- D. equilibrium will shift in the backward direction

**Answer:**



[Watch Video Solution](#)

**160.** A catalyst :

- A. increase the average kinetic energy of reacting molecules
- B. decreases the activation energy
- C. can alter the reaction mechanism
- D. Can change pre-exponential factor

**Answer:**



[Watch Video Solution](#)

**161.** Which of the following is correct about the chemical equilibrium ?

A.  $(\Delta G)_{T,p} = 0$

B. Equilibrium constant is independent of initial concentration of reactants

C. Catalyst has no effect on equilibrium state


D. Reaction stops at equilibrium

**Answer:**



[Watch Video Solution](#)

**162.** For the reaction



If  $\alpha$  is negligible w.r.t 1 then degree of dissociation ( $\alpha$ ) of  $AB_2$  is proportional to :

A.  $\frac{1}{P}$

B.  $\frac{1}{V}$

C.  $\frac{1}{\sqrt{P}}$




D.  $\sqrt{V}$


**Answer:**



**Watch Video Solution**

**163.** Consider the reaction given below. In which cases will the reaction proceed toward right by increasing the pressure ?






**Answer:**

 [Watch Video Solution](#)

**164.** Ammonia is a weak base that reacts with water according to the equation



Select the correct option (s) that can increase the moles of ammonium ion in water:

- A. Addition of HCl
- B. Addition of NaOH
- C. Addition of  $NH_4Cl$
- D. Addition of  $H_2O$

**Answer:**

 [Watch Video Solution](#)

165. Consider the reaction  $2CO(g) + O_2(g) \rightleftharpoons 2CO_2(g) + \text{Heat}$

Under what conditions shift is undetminable ?

- A. Addition of  $O_2$  and decrease in volume
- B. Addition of CO and removal of  $CO_2$  at constant volume
- C. Increase in temperature and decrease in volume
- D. Addition of CO and increase in temperature at constant volume

**Answer:**

 [Watch Video Solution](#)

166. What will be the effect of addition of catalyst at constant temperature ?

- A. The equilibrium constant will remain constant
- B.  $\Delta H$  of the reaction will remain constant



C.  $K_f$  and  $K_b$  will increase upto same extent

D. equilibrium composition will change

**Answer:**

 [Watch Video Solution](#)

**167.** For the reaction  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ , the forward reaction at constant temperature favorrd by :

A. introducing an inert gas at constant volume

B. introducing chlorine gas at constant volume

C. introducing an inert gas at constant pressure

D. increasing the volume of the container

**Answer:**

 [Watch Video Solution](#)

168. For the reaction :  
 $Cl_2(g) + 3F_2(g) \rightleftharpoons 2ClF_3(g)$ ,  $\Delta H = -329kJ$ , dissociation of  $ClF_3(g)$  will be favoured by :

- A. increasing the temperature
- B. increasing the volume of the container
- C. adding of  $F_2$  gas
- D. adding of inert gas at constant pressure

**Answer:**



[Watch Video Solution](#)

169. Increase in the pressure for the following equilibrium results in the :



Equilibrium will shift left

- A. formation of more  $H_2O(l)$

B. formation of more  $H_2O(g)$

C. increase in b.p of  $H_2O(l)$

D. decrease in b.p. of  $H_2O(l)$

**Answer:**

 [Watch Video Solution](#)

**170.** Heating a II group metal carbonate leads to decomposition as :



Equilibrium will shift left

A. by addition of  $BaO(s)$

B. by addition of  $CO_2(g)$

C. by decreasing the temperature

D. by decreasing the volume of the vessel

**Answer:**



Watch Video Solution

171.  $N_2(g)$  and  $H_2(g)$  are allowed to react in a closed vessel at given temp. and pressure for the formation of  $NH_3(g)$ ,  $[N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g) + 22.4kcal]$  If he (g) is added at equilibrium at constant pressure than which is/are correct ?

- A. Concentration of  $N_2(g)$ ,  $H_2(g)$  and  $NH_3(g)$  decrease.
- B. Moles of  $NH_3(g)$  decreases.
- C. The extent of cooling depends on amount of he (g) added.
- D. Concentration of  $N_2$  and  $H_2$  increases and concentration of  $NH_3$  decreases.

Answer:



Watch Video Solution

172. Column-I and Column-II contains fore enteries each. Entries of Column-I are to be matched with, some entries of Column-II One or more than one entries of Column-I may have the mathching with the same entries of Column-II

Column-I	Column-II
(A) $\text{CaCO}_3(s) \rightleftharpoons \text{CaO}(s) + \text{CO}_2(g)$	(P) $K_p > K_c$ above room temperature
(B) $\text{CO}(g) + \text{Cl}_2(g) \rightleftharpoons \text{COCl}_2(g)$	(Q) $K_p = K_c$ above room temperature
(C) $\text{H}_2(g) + \text{I}_2(g) \rightleftharpoons 2\text{HI}(g)$	(R) $K_p < K_c$ above room temperature
(D) $\text{HCl}(g) \rightleftharpoons \text{H}^+(aq) + \text{Cl}^-(aq)$	(S) $K_p$ and $K_c$ not defined

 [Watch Video Solution](#)

173. Match the following columns

Column-I	Column-II
(A) $3\text{O}_2(g) \rightleftharpoons 2\text{O}_3(g)$	(P) no unit
(B) $\text{SO}_2(g) + \frac{1}{2}\text{O}_2(g) \rightleftharpoons \text{SO}_3(g)$	(Q) $\text{atm}^{-1/2}$
(C) $2\text{HF}(g) \rightleftharpoons \text{H}_2(g) + \text{F}_2(g)$	(R) $\text{atm}^{-1}$
(D) $\text{CO}(g) + 3\text{H}_2(g) \rightleftharpoons \text{CH}_4(g) + \text{H}_2\text{O}(g)$	(S) $\text{atm}^{-2}$

 [Watch Video Solution](#)

174. Match the following columns

**Column-I**

- (A)  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ ;  $\Delta H = -ve$   
(B)  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ ;  $\Delta H = +ve$   
(C)  $A(g) + B(g) \rightleftharpoons 2C(g) + D(g)$ ;  $\Delta H = +ve$   
(D)  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ ;  $\Delta H = +ve$

**Column-II**

- (P)  $K$  increases with increase in temperature  
(Q)  $K$  decreases with increase in temperature  
(R) Pressure has no effect  
(S) Moles of product increase due to addition of inert gas at constant pressure

 [Watch Video Solution](#)

175. Match the following columns

**Column-I**

- (A)  $\frac{K_{10+T^\circ C}}{K_{T^\circ C}} = 2$   
(B)  $\frac{K_{10+T^\circ C}}{K_{T^\circ C}} = \frac{1}{2}$   
(C)  $A(g) + B(g) \rightleftharpoons C(g)$   
(D)  $X(s) + Y(g) \rightleftharpoons Z(g)$

**Column-II**

- (P) Endothermic  
(Q) Not affected by pressure  
(R) Exothermic  
(S) Affected by volume

 [Watch Video Solution](#)

176. Match the following columns

**Column-I**

- (A) Pressure increased in  
 $2\text{NO}(g) \rightleftharpoons \text{N}_2(g) + \text{O}_2(g)$
- (B) Pressure increased in  
 $\text{CH}_4(g) + \text{H}_2\text{O}(g) \rightleftharpoons \text{CO}(g) + 3\text{H}_2(g)$
- (C) Temp. increased and pressure increased  
 $3\text{O}_2(g) \rightleftharpoons 2\text{O}_3(g); \quad \Delta H = 285 \text{ kJ}$
- (D) Pressure decreased and moles of  $\text{N}_2$  increased  
 $\text{N}_2(g) + 2\text{O}_2(g) \rightleftharpoons 2\text{NO}_2(g); \quad \Delta H = 66.4 \text{ kJ}$


**Column-II**

- (P) Equilibrium shifted in forward direction
- (Q) Equilibrium shifted in backward direction
- (R) Equilibrium remains unaffected
- (S) Theoretically we cannot predict




**Watch Video Solution**

(P)



(Q)



177.






(R)



(S)



[Watch Video Solution](#)

**178.** Assertion (A): The endothermic reactions are favoured at lower temperature and the exothermic reactions are favoured at higher temperature.

Reason (R) : when a system in equilibrium is disturbed by changing the

temperature, it will tend to adjust itself so as to overcome the effect of the change.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**

 [Watch Video Solution](#)

**179.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The melting point of ice decreases with increase of pressure.

STATEMENT-2: Ice contracts on melting .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



[Watch Video Solution](#)

**180.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according

to the instruction given below:

STATEMENT-1: The equilibrium of  $A(g) \rightleftharpoons B(g) + c(g)$  is not affected by changing the volume.

STATEMENT-2:  $K_c$  for the reaction does not depend on volume of the container.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**



**Watch Video Solution**

**181.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For a chemical reaction at initial stage rate of forward reaction ( $r_f$ ) is greater than rate of reversed reaction ( $r_b$ )

STATEMENT-2: When  $r_f = r_b$ , chemical reaction is at equilibrium.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



**Watch Video Solution**

**182.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the reaction  $A(g) \rightleftharpoons B(g) + C(g)$ ,  $K_p=1\text{atm}$ . If we start with equal moles of all gases at 9 tm of initial pressure, then at equilibrium partial pressure of A increases.

STATEMENT-2: Reaction quotient  $Q_p > K_p$  hence equilibrium shifts in backward direction.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**183.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The gas phase reaction  $PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$  shifts to the right on increasing pressure.

STATEMENT-2: When pressure increase, equilibrium shifts towards more number of moles.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**



**Watch Video Solution**

**184.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For a reaction at equilibrium, the Gibb's free energy of reaction is minimum at constant temp. and pressure.

STATEMENT-2: The Gibb's free energy of both reactants and products increases and become equal at equilibrium.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE



D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**

 [Watch Video Solution](#)

**185.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The physical equilibrium is not static but dynamic in nature.

STATEMENT-2: The physical equilibrium is a state in which two opposing process are proceeding at the same rate.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**

 [Watch Video Solution](#)

**186.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.

STATEMENT-2: Equilibrium constant depends upon the way in which the reaction is written.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**

 [Watch Video Solution](#)

**187.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: If  $Q_p < K_p$  reaction moves in direction of reactants.

STATEMENT-2: Reaction quotient is defined in the same way as equilibrium constant at any stage of the reaction.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**



[Watch Video Solution](#)

**188.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$  if the volume of vessel is reduced to half of its original volume, equilibrium concentration of all gases will be doubled.

STATEMENT-2: According to Le- Chatelier's principle, reaction shifts in a direction that tends to minimized the effect of the stess.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



[Watch Video Solution](#)

**189.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The equilibrium constant of the exothermic reaction at high temperature decreases.

STATEMENT-2: Since  $\ln \frac{K_2}{K_1} = \frac{\Delta H^\circ}{R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$  and for exothermic reaction ,

$$\Delta H^\circ = -ve \text{ and thereby, } \frac{K_2}{K_1} < 1$$

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**




**Watch Video Solution**

**190.** Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the reaction at certainn temperature



there will be no effect by addition of inert gas at constant volume.

STATEMENT-2: Molar concentration of all gases remains constant.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

191. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the physical equilibrium  $H_2O \rightleftharpoons H_2O(l)$  on increasing temperature and increasing pressure more water will form.

STATEMENT-2: Since forward reaction is endothermic in nature and volume of water is greater than that of the volume of ice.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**





192. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The catalyst does not alter the equilibrium constant.

STATEMENT-2: Because for the catalysed reaction and uncatalysed reaction  $\Delta H$  remains same and equilibrium constant depends of  $\Delta H$ .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**




Watch Video Solution

**193.** In the reaction  $C(s) + CO_2(g) \rightleftharpoons 2CO(g)$ , the equilibrium pressure is 12 atm. If 50 % of  $CO_2$  reacts, calculate  $K_p$ .



Watch Video Solution

**194.** Calculate partial pressure of B at equilibrium in the following equilibrium



Watch Video Solution

**195.** In a gaseous reaction  $A + 2B \rightleftharpoons 2C + D$  the initial concentration of B was 1.5 times that of A. At equilibrium the concentration of A and D were equal. Calculate the equilibrium constant  $K_C$ .



Watch Video Solution

196. For the reaction  $A(g) \rightleftharpoons B(g)$ ,  $K_C = 10$

$B(g) \rightleftharpoons C(g)$ ,  $K_C = 2$

$C(g) \rightleftharpoons D(g)$ ,  $K_C = 0.01$

Calculate  $K_C$  for the reaction  $D(g) \rightleftharpoons A(g)$ .

 [Watch Video Solution](#)

197. 5 litre vessel contains 2 moles of each of gases A and B at equilibrium.

If 1 mole each of A and B are removed. Calculate  $K_C$  for the reaction


$A(g) \rightleftharpoons B(g)$

 [Watch Video Solution](#)

198. Calculate  $K_P$  for the reaction  $A(g) \rightleftharpoons B(s) + 2C(g)$ ,  $K_C = 0.2$  at 305 K.

 [Watch Video Solution](#)

**199.** A mixture of 3 moles of  $SO_2$ , 4 moles of  $NO_2$ , 1 mole of  $SO_3$  and 4 moles of  $NO$  is placed in a 2.0L vessel.



At equilibrium, the vessel is found to contain 1 mole of  $SO_2$ . Calculate the value of  $K_C$ .

 [Watch Video Solution](#)

**200.** The density of an equilibrium mixture of  $N_2O_4$  and  $NO_2$  at 1 atm and 373.5K is 2.0 g/L.


Calculate  $K_C$  for the reaction  $N_2O_2(g) \rightleftharpoons 2NO_2(g)$

 [Watch Video Solution](#)

**201.** If chemical equilibrium is attained t standard states then what is the value of  $\Delta G^\circ$ ?

 [Watch Video Solution](#)

**202.** Calculate the equilibrium concentration ratio of C to A if equimolar ratio of A and B were allowed to come to equilibrium at 300K.




 [Watch Video Solution](#)

**203.** A definite amount of solid  $NH_4HS$  is placed in a flask already containing ammonia gas at a certain temperature and 0.1 atm pressure.

$NH_4HS$  decomposes to give  $NH_3$  and  $H_2S$  and at equilibrium total pressure in flask is 1.1 atm. If the equilibrium constant  $K_P$  for the reaction  $NH_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$  is represented as  $z \times 10^{-1}$  then find the value of z.

 [Watch Video Solution](#)

**204.** The gaseous reaction :  $A(g) + nB(g) \rightleftharpoons mC(g)$  is represented by following curves




What is the value of  $n+m$ ?


[Watch Video Solution](#)

Level 1 Q 93 To Q 122


1. An equilibrium mixture at 700 K of  $0.05M N_2(g)$  and  $0.2M NH_3(g)$  is present in a container. Now if this equilibrium is disturbed by adding  $N_2(g)$  so that its concentration becomes  $0.15M$  just after addition then which of the following graph represents the above situation more appropriately:




A.



B.



C.



D.

Answer: a



View Text Solution

Level 2

1. Calculate  $\Delta_r G$  for the reaction at  $27^\circ C$



Given :  $P_{H_2} = 0.5 \text{ bar}$ ,  $[Ag^+] = 10^{-5} M$ ,

$[H^+] = 10^{-3} M$ ,  $\Delta_r G^\circ [Ag^+(aq)] = 77.1 kJ/mol$

A.  $-154.2 kJ/mol$

B.  $-178.9 kJ/mol$

C.  $-129.5 kJ/mol$

D. None of these

**Answer: C**



**View Text Solution**