







CHEMISTRY

BOOKS - NARENDER AVASTHI CHEMISTRY (HINGLISH)

IONIC EQUILIBRIUM

Exercise

1. Morphine ($C_{17}H_{19}NO_3$), which is used medically to relieve pain is a base. What is its conjugate acid?






Answer:



Watch Video Solution

2. The conjugate base of $H_2PO_4^-$ is :



Answer:



Watch Video Solution

3. The strongest Bronsted base in the following anion is:



D. Br^-

Answer:

 [Watch Video Solution](#)

4. What salt can furnish H^+ in its aqueous solution?

A. NaH_2PO_2

B. Na_2HPO_3



C. Na_2HPO_4

D. All of these

Answer:

 [Watch Video Solution](#)

5. Which is the set of amphoteric species?



D. All of these

Answer:

 [Watch Video Solution](#)

6. The K_a values for HPO_4^{2-} and HSO_3^- are 4.8×10^{-13} and 6.3×10^{-8} respectively. Therefore, it follows the HPO_4^{2-} is ... acid than HSO_3^- and PO_4^{3-} is a base than SO_3^{2-}

A. weaker, stronger

B. stronger, weaker

C. weaker, weaker

D. stronger, stronger

Answer:



[Watch Video Solution](#)

7. Given the following K_a values, determine which species is the strongest base ?

$$HSO_4^- = 1.2 \times 10^{-2}, H_2PO_4^- = 6.3 \times 10^{-8}, HCO_3^- = 4.7 \times 10^{-11}$$

- A. CO_3^{2-}
- B. H_2SO_4
- C. SO_4^{2-}
- D. HPO_4^{2-}

Answer:



[Watch Video Solution](#)

8. Given that K_w for water is $10^{-13} M^2$ at $62^\circ C$, compute the sum of pOH and pH for a neutral aqueous solution at $62^\circ C$:

- A. 7.0
- B. 13.30
- C. 14.0
- D. 13.0

Answer:



[Watch Video Solution](#)

9. The value of the ion product constant for water, (K_w) at $60^\circ C$ is $9.6 \times 10^{-14} M^2$ what is the $[H_3O^+]$ of a neutral aqueous solution at $60^\circ C$ and an aqueous solution with a pH=7.0 at $60^\circ C$ are respectively?

- A. 3.1×10^{-8} acidic
- B. 3.1×10^{-7} , neutral

C. 3.1×10^{-8} , basic

D. 3.1×10^{-7} , basic

Answer:

 [Watch Video Solution](#)

10. For pure water:

A. pH increases while pOH decreases with rise in temperature

B. pH decreases while pOH increases with rise in temperature

C. both pH and pOH decreases with rise in temperature

D. both pH and pOH increases with rise in temperature

Answer:

 [Watch Video Solution](#)

11. A beer has a pH of 4.30. What is the $[H_3O^+]$?

A. 3.0×10^{-4}

B. 2.0×10^{-4}

C. 2.0×10^5

D. 5.0×10^{-5}

Answer:



[Watch Video Solution](#)

12. The hydrogen ion concentration of the oceans is about 2×10^{-9} M.

What is the pH?

A. 8.85

B. 9.3

C. 7.85

D. 8.7

Answer:



[Watch Video Solution](#)

13. The hydroxide ion concentration of a wine is 8×10^{-11} M. What is the pH of the wine?

A. 2.10

B. 2.9

C. 3.9

D. 4.9

Answer:



[Watch Video Solution](#)

14. The pH of a solution is 5. to this solution acid was added so that its pH value bcomes 2.0. The increase in H^+ concentration is :

A. 100 times

B. 5 times

C. 2.5 times

D. 1000 times

Answer:



Watch Video Solution

15. A solution has a $\text{pH}=9$. It is 1000 times more basic than the original solution. What was the pH of the original solution?

A. 12

B. 6

C. 9

D. 10

Answer:

 [Watch Video Solution](#)

16. Equal volumes of two HCl solutions of $pH = 3$ and $pH = 5$ were mixed. What is the pH of the resulting solution ?

A. 3.5

B. 4.0

C. 4.5

D. 3.3

Answer:

 [Watch Video Solution](#)

17. pOH of $0.002M HNO_3$ is :

A. $11 + \log 2$

B. $11 - \log 2$

C. $-3 + \log 2$

D. None of these

Answer:

 [Watch Video Solution](#)

18. Number of equivalents of HCl present in 100 mL of its solution whose pH is 4:

A. 10^{-4}

B. 10^{-3}

C. 10^{-2}

D. 10^{-5}

Answer:

 [Watch Video Solution](#)

19. To a 10mL of $10^{-3}\text{NH}_2\text{SO}_4$ solution water has been to make the total volume of one litre. Its pOH would be :

- A. 3
- B. 12
- C. 9
- D. 5

Answer:



[Watch Video Solution](#)

20. The pH of a solution of H_2SO_4 is 1. Assuming complete ionisation, find the molarity of H_2SO_4 solution :

- A. 0.1
- B. 0.2
- C. 0.05

D. 2.0

Answer:



[Watch Video Solution](#)

21. pH of a strong diprotic acid (H_2A) at concentrations:

(i) 10^{-4} M, (ii) 10^{-4} N

are respectively:

A. 3.7 and 4.0

B. 4 and 3.7

C. 4 and 4

D. 3.7 and 3.7

Answer:



[Watch Video Solution](#)

22. Calcium hydroxide is a strong base. Compute $[Ca^{2+}]$ and $[OH^-]$ "for" a solution that is prepared by dissolving 0.60g of $Ca(OH)_2$ in enough water to make a 1500 mL of solution.

[Atomic mass : $Ca = 40$, $O = 16$, $H = 1$]

A. 5.4×10^{-3} , 9.1×10^{-13}

B. 5.4×10^{-3} , 1.08×10^{-2}

C. 5.4×10^{-3} , 5.4×10^{-3}

D. 8.1×10^{-3} , 8.1×10^{-3}

Answer:



[Watch Video Solution](#)

23. pH of 10^{-6} M HCl (aq.) is :

A. just less than 6

B. exactly equal to 6

C. just greater than 6

D. just less than 7

Answer:

 [Watch Video Solution](#)

24. $10^{-5} M HCl$ solution at $25^{\circ} C$ is diluted 1000 times. The pH of the diluted solution will

A. be equal to 8

B. lie between 7 and 8

C. lie between 6 and 7

D. remain unchanged

Answer:

 [Watch Video Solution](#)

25. 4.0 g of NaOH and 4.9 g of H_2SO_4 are dissolved in water and volume is made upto 250 mL.

The pH of this solution is:

- A. 7.0
- B. 1.0
- C. 2.0
- D. 12.0

Answer:



[Watch Video Solution](#)

26. A 25.0 mL sample of 0.10 M HCl is titrated with 0.10 M NaOH. What is the pH of the solution at the points where 24.9 and 25.1 mL of NaOH have been added?

- A. 3.70, 10.70

B. 3.30, 10.30

C. 3.70, 10.30

D. 3.0, 11.0

Answer:

 [Watch Video Solution](#)

27. What is the pH of solution in which 25.0 mL of 0.1 M NaOH is added to 25 mL of 0.08M HCl and final solution is diluted to 500 mL?

A. 3

B. 11

C. 12

D. 13

Answer:

 [Watch Video Solution](#)

28. What is the pH of a solution in which 10.0 mL of 0.010 M $\text{Sr}(\text{OH})_2$ is added to 10.0 mL of 0.010 M HCl?

- A. 2.30
- B. 1.50
- C. 11.70
- D. 7.00

Answer:

 [Watch Video Solution](#)

29. At 90°C , pure water has $[\text{H}^+] = 10^{-6}$ M. If 100 mL of 0.2 M HCl is added to 200 mL of 0.1 M KOH at 90°C then pH of the resulting solution will be :

- A. 5

B. 6

C. 7


D. None of these

Answer:



[Watch Video Solution](#)

30. What change will occur for the following reaction if the hypochlorous acid solution is diluted from 0.1 to 0.01 M?



A. a decrease in the fraction of acid ionized

B. an increase in the fraction of acid ionized


C. no change in the fraction of acid ionized

D. we can not predict

Answer:

[Watch Video Solution](#)

31. Given K_a values of 5.76×10^{-10} and 4.8×10^{-10} for NH_4^+ and HCN respectively. What is the equilibrium constant for the following reaction?



A. 0.83

B. 1.2

C. 8.0×10^{-11}


D. 27.6×10^{-10}

Answer:

[Watch Video Solution](#)

32. Which is the strongest acid (pK_a value is given)?

A. $HCOOH$ [3.77]



Answer:



Watch Video Solution

33. Given : Enthalpy of ionization of two acids :

$$\Delta H^\circ (HCN) = 45.2 \text{ KJmol}^{-1}$$

$$\Delta H^\circ (CH_3COOH) = 2.1 \text{ KJmol}^{-1}$$

which relationship for the two acids is true ?

A. $pK_a(HCN) = pK_a(CH_3COOH)$

B. $pK_a(HCN) > pK_a(CH_3COOH)$

C. $pK_a(HCN) < pK_a(CH_3COOH)$

D. $pK_a(HCN) = \frac{45.2}{2.1} pK_a(CH_3COOH)$

Answer:

 [Watch Video Solution](#)

34. What is the hydronium ion concentration of a 0.25 M HA solution?

$$(K_a = 4 \times 10^{-8})$$

A. 10^{-4}

B. 10^{-5}

C. 10^{-7}

D. 10^{-10}

Answer:

 [Watch Video Solution](#)

35. What is the percent dissociation (α) of a 0.01 M HA solution?

$$(K_a = 10^{-4})$$

A. 9.5 %

B. 1 %

C. 10.5 %

D. 17 %

Answer:



[Watch Video Solution](#)

36. Given the two concentration of HCN ($K_a = 10^{-9}$) are 0.1 M and 0.001 M respectively. What will be the ratio of degree of dissociation?

A. 1

B. 0.1

C. 0.003

D. 0.01

Answer:

 [Watch Video Solution](#)

37. A 0.10 M solution of HF is 8.0 % dissociated. What is the K_a ?

A. 6.4×10^{-10}

B. 8.8×10^{-4}

C. 6.95×10^{-4}

D. 7.6×10^{-4}

Answer:

 [Watch Video Solution](#)

38. A weak base MOH of 0.1N concentration shows a pH value of 9 .

What is the percentage degree of ionization of the base ?

A. 0.01 %

B. 0.001 %

C. 0.1 %

D. 0.02 %

Answer:



Watch Video Solution

39. 0.01 M HA (aq.) is 2 % dissociated, $[OH^-]$ of solution is :

A. 2×10^{-4}

B. 10^{-8}

C. 5×10^{-11}

D. 5×10^{-12}

Answer:



Watch Video Solution

40. If degree of dissociation is 0.01 of decimolar solution of weak acid HA then pK_a of acid is :

- A. 2
- B. 3
- C. 5
- D. 7

Answer:



[Watch Video Solution](#)

41. What concentration of $HCOO^-$ is present in a solution of weak of 0.01 M $HCOOH$ ($K_a = 1.8 \times 10^{-4}$ and 0.01 M HCl)?

- A. 1.8×10^{-3}
- B. 10^{-2}
- C. 1.8×10^{-4}

D. 10^{-4}

Answer:

 Watch Video Solution

42. Chose the correct code

Column – I

(P) pK_b of X^- (K_a of $HX = 10^{-6}$)

(Q) pH of $10^{-8} M HCl$

(R) pH of $10^{-2} M$ acetic acid solution (Take K_a of acetic acid = 1.6×10^{-5})

(S) pOH of a solution obtained by mixing equal volumes of solution with

- A.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
1	2	4	3
- B.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
4	3	2	1
- C.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
2	1	4	3
- D.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
1	2	3	4

Answer:

 Watch Video Solution

43. How much water must be added to 300mL of a 0.2M solution of CH_3COOH for the degree of dissociation of the acid to double ? (Assume K_a of acetic is of order of 10^{-5}M)

- A. 600 mL
- B. 900 mL
- C. 1200 mL
- D. 1500 mL

Answer:

 [Watch Video Solution](#)

44. What is $[\text{NH}_4^+]$ in a solution that contain 0.02 M NH_3 ($K_b = 1.8 \times 10^{-5}$) and 0.01 M KOH ?

- A. 9×10^{-6}

B. 1.8×10^{-5}

C. 3.6×10^{-5}

D. None of these

Answer:



Watch Video Solution

45. A hand book states that the solubility of RNH_2 (g) in water at 1 atm and 0°C is 22.41 volumes of RNH_2 (g) per volume of water. (pK_b of $RNH_2 = 4$) Find the max. pOH that can be attained by dissolving RNH_2 in water:

A. 1

B. 2

C. 4

D. 6

Answer:



[Watch Video Solution](#)

46. The $[H^+]$ of a resulting solution that is 0.01 M acetic acid ($K_a = 1.8 \times 10^{-5}$) and 0.01 M in benzoic acid ($K_a = 6.3 \times 10^{-5}$):

A. 9×10^{-4}

B. 81×10^{-4}

C. 9×10^{-5}

D. 2.8×10^{-3}

Answer:



[Watch Video Solution](#)

47. 6.0 g weak acid HA (mol.mass=60 g/mol.) is dissolved in water and formed 10 m^3 solution. If $K_a(HA) = 10^{-9}$, then pOH of solution is :

[Given: $\log 4=0.6$]

- A. 7
- B. greater than 6.7 and less than 7.0
- C. greater than 7.0 and less than 7.3
- D. greater than 7.3

Answer:

 [Watch Video Solution](#)

48. Carbonic acid (H_2CO_3), a diprotic acid has $K_{a1} = 4.0 \times 10^{-7}$ and $K_{a2} = 7.0 \times 10^{-11}$. What is the $[HCO_3^-]$ of a 0.025 M solution of carbonic acid?

- A. 7.8×10^{-3}
- B. 6.6×10^{-4}
- C. 10^{-10}

D. 1.0×10^{-4}

Answer:

 [Watch Video Solution](#)

49. Carbonic acid (H_2CO_3), a diprotic acid has $K_{a1} = 4.0 \times 10^{-7}$ and $K_{a2} = 5.0 \times 10^{-11}$. What is the $[CO_3^{2-}]$ of a 0.025 M solution of carbonic acid?

A. 5.5×10^{-9}

B. 5.5×10^{-8}

C. 7.0×10^{-9}

D. 7.0×10^{-11}

Answer:

 [Watch Video Solution](#)

50. Selenious acid (H_2SeO_3), a diprotic acid has $K_{a1} = 3.0 \times 10^{-3}$ and $K_{a2} = 5.0 \times 10^{-8}$. What is the $[OH^-]$ of a 0.30 M solution of selenious acid?

A. 2.85×10^{-3}

B. 5.0×10^{-6}

C. 3.5×10^{-12}

D. 3.5×10^{-13}

Answer:



Watch Video Solution

51. Which of the hydrated species may exist?

I: $H_5O_2^+$, II: H_3O^+ , III: $H_3O_2^-$, IV: $H_7O_3^+$

A. II only

B. I and II

C. I, II and IV

D. I, II, III and IV

Answer:

 [Watch Video Solution](#)

52. Consider the following salts. Which one(s) when dissolved in water will produce an acidic solution?

1. NH_4Cl , 2. $KHSO_4$, 3. $NaCN$, 4. KNO_3

A. 2 and 3

B. 1 and 2

C. only 3

D. 2 and 4

Answer:

 [Watch Video Solution](#)

53. Consider the following salts. Which one(s) when dissolved in water will produce a basic solution?

1. $RbClO_4$, 2. $NaNO_2$, 3. NH_4Cl , 4. $NaCl$

A. 1 and 3

B. only 2

C. 1 and 2

D. 3 and 4

Answer:

 [Watch Video Solution](#)

54. At 25°C dissociation constants of acid HA and base BOH in aqueous solution are same. The pH of 0.01 M solution of HA is 5. The pOH of 10^{-4} M solution of BOH at the same temperature is :

A. 3.5

B. 4

C. 6

D. None of these

Answer:



[Watch Video Solution](#)

55. Which of the following solutions has the highest pH?

A. $0.2M HClO_4$

B. $0.20M CH_3COOH$

C. $0.020M HCl$

D. $0.2M NaCl$

Answer:



[Watch Video Solution](#)

56. pH of solutions of four sodium salts NaW, NaX, NaX, NaY and NaZ were found to be 7.0, 9.0, 10.0 and 11.0 respectively. If each solution has concentration 0.1 M, the weakest acid is :

- A. HW
- B. HX
- C. HY
- D. HZ

Answer:

 [Watch Video Solution](#)

57. The pH values 0.1 M solution of HCOONa (I), HCOOH (II), CH_3COONH_4 (III), NaOH (IV) HCl (V), will be in the order :

- A. IVgtIIlgtIgtIIgtV
- B. IVgtIgtIIlgtIIgtV

C. $\text{[H}^+\text{]} = 10^{-4} \text{ M}$

D. $\text{[H}^+\text{]} = 10^{-10} \text{ M}$

Answer:

 [Watch Video Solution](#)

58. pH of an aqueous NaCl solution at 50°C should be :

A. 7

B. > 7

C. < 7

D. 0

Answer:

 [Watch Video Solution](#)

59. Upon hydrolysis of sodium carbonate, the reaction takes place between:

A. Na^+ and water

B. Na^+ and OH^-

C. CO_3^{2-} and water

D. CO_3^{2-} and H^+

Answer:



[Watch Video Solution](#)

60. The solution of blue vitrol in water is acidic because:

A. $CuSO_4$ reacts with water

B. Cu^+ reacts with water

C. SO_4^{2-} reacts with water

D. $CuSO_4$ renives OH^- ions from water

Answer:



[Watch Video Solution](#)

61. 1 mL of 0.1 N HCl is added to 999 mL solution of NaCl. The pH of the resulting solution will be :

A. 7

B. 4

C. 2

D. 1

Answer:



[Watch Video Solution](#)

62. If a salt of strong acid and weak base hydrolyses appreciably ($\alpha = 0.1$), which of the following formula is to be used to calculate

degree of hydrolysis 'alpha'?

A. $\alpha = \frac{\sqrt{K_w}}{K_a \cdot a}$

B. $\alpha = \frac{\sqrt{K_w}}{K_b \cdot a}$

C. $\alpha = \frac{\sqrt{K_w}}{K_a \cdot K_b}$

D. None of these

Answer: b

 [Watch Video Solution](#)

63. The correct formula to calculate the hydroxyl ion concentration of an aqueous solution of NH_4NO_3 is:

A. $\sqrt{\frac{C \times K_w}{K_b}}$

B. $\sqrt{\frac{K_w \times K_b}{C}}$

C. $\sqrt{\frac{C \times K_w}{K_a}}$

D. $\sqrt{\frac{K_a \times K_w}{C}}$

Answer: B

 Watch Video Solution

64. $[H^+] = \sqrt{\frac{K_w K_a}{C}}$ is suitable for

A. $NaCl, NH_4Cl$

B. $CH_3COONa, NaCN$

C. $CH_3COONa, (NH_4)_2SO_4$

D. $CH_3COONH_4, (NH_4)_2CO_3$

Answer: b

 Watch Video Solution

65. What is the hydrolysis constant of the OCl^- ion? The ionization constant of $HOCl$ is 3.0×10^{-8} .

A. 3.33×10^{-8}

B. 3.33×10^{-7}

C. 3.0×10^{-7}

D. 3.33×10^{-6}

Answer:

 [Watch Video Solution](#)

66. What is the pH of a 0.10 M $C_6H_5O^-$ solution? The K_a of C_6H_5OH is

1.0×10^{-10}

A. 10.51

B. 11.04

C. 11.50

D. 12

Answer:

 [Watch Video Solution](#)

67. Calculate the $[OH^-]$ in $0.01M$ aqueous solution of $NaOCN$ (K_b for $OCN^- = 10^{-10}$):

A. $10^{-6} M$

B. $10^{-7} M$

C. $10^{-8} M$

D. None of these

Answer:

 [Watch Video Solution](#)

68. What is the ionization constant of an acid if the hydronium ion concentration of a $0.40 M$ solution is $1.40 \times 10^{-4} M$?

A. 1.96×10^{-8}

B. 1.22×10^{-9}

C. 4.90×10^{-8}

D. 1.40×10^{-6}

Answer:



Watch Video Solution

69. The degree of hydrolysis of 0.1 M RNH_3Cl solution is 1.0%. If the concentration of RNH_3Cl is made 0.4 M, what is the new degree of hydrolysis (in percentage)?

A. 0.01

B. 0.001

C. 0.2

D. 0.5

Answer:

[Watch Video Solution](#)

70. % hydrolysis of 0.1M CH_3COONH_4 , when $K_a(CH_3COOH) = K_b(NH_4OH) = 1.8 \times 10^{-5}$ is:

A. 0.55

B. 7.63

C. 0.55×10^{-2}

D. 7.63×10^{-3}

Answer:

[Watch Video Solution](#)

71. The enthalpy of neutralisation of four acids HA, HB, HC and HD with NaOH are 13, -12, -11, -10 Kcal//mol. Which salt has maximum degree of hydrolysis?

A. 1 M NaA

B. 1 M NaB

C. 1 M NaC

D. 1 M NaD

Answer:

 [Watch Video Solution](#)

72. Calculate $[H^+]$ at equivalent point between titration of 0.1 M, 25 mL of weak acid HA ($K_{a(HA)} = 10^{-5}$) with 0.05 M NaOH solution:

A. 3×10^{-9}

B. 1.732×10^{-9}

C. 8

D. 10

Answer:

 [Watch Video Solution](#)

73. When a salt of weak acid and weak base is dissolved in water, the pH of the resulting solution will be :

- A. be 7
- B. be greater than 7
- C. be less than 7
- D. depend upon K_a and K_b values

Answer:

 [Watch Video Solution](#)

74. What will be the pH of an aqueous solution of 1.0 M ammonium formate?

Given : $pK_a = 3.8$ and $pK_b = 4.8$

A. 7.5

B. 3.4

C. 6.5

D. 10.2

Answer:

 [Watch Video Solution](#)

75. What will be the pH and $\% \alpha$ (degree of hydrolysis) respectively for the salt BA of $0.1M$ concentration ? Given : K_a for $HA = 10^{-6}$ and K_b for $BOH = 10^{-6}$

A. 5, 1 %

B. 7, 10 %

C. 9, 0.01 %

D. 7, 0.01 %

Answer:



[Watch Video Solution](#)

76. The percentage degree of hydrolysis of a salt of weak acid (HA) and weak base (BOH) in its 0.1 M solution is found to be 10 %. If the molarity of the solution is 0.05 M, the percentage hydrolysis of the salt should be :

A. 5 %

B. 10 %

C. 20 %


D. None of these

Answer:



[Watch Video Solution](#)

77. What is the hydronium ion concentration of a 0.02 M solution of Cu^{2+} solution of copper(II) perchlorate? The acidity constant of the following reaction is 5×10^{-9} .



A. 1×10^{-5}

B. 7×10^{-4}


C. 5×10^{-4}

D. 1×10^{-4}

Answer:

 [Watch Video Solution](#)

78. What is the acidity constant for the following reaction given that the hydronium ion concentration of a 0.04 M solution of Ni^{2+} solution of nickel(II) perchlorate is 4.5×10^{-6} ?



A. 2×10^{-12}

B. 4×10^{-6}

C. 5×10^{-12}


D. 5×10^{-10}

Answer:

 [Watch Video Solution](#)

79. Calculate the pH at 25°C of a solution that is 0.10 M in $\text{Fe}(\text{NO}_3)_3$.

The acid dissociation constant for the reaction given below is 1.0×10^{-3} .



A. 2.00

B. 2.02

C. 2.30

D. 2.50

Answer:

 [Watch Video Solution](#)

80. Approximate pH of 0.01 M NaHA is calculated by :

($K_{a1} = 10^{-6}$ and $K_{a2} = 10^{-8}$ are ionization constants of H_2A)

A. $pH = 7 + \frac{pK_{a1}}{2} + \frac{\log C}{2}$

B. $pH = 7 - \frac{pK_{a1}}{2} - \frac{\log C}{2}$

C. $pH = \frac{pK_{a1} + pK_{a2}}{2}$

D. None of these

Answer:

 [Watch Video Solution](#)

81. H_3PO_4 is a weak triprotic acid, approximate pH 0.1 M $NaHPO_4(aq.)$ is calculated by:

A. $\frac{1}{2}[pK_{a1} + pK_{a2}]$

B. $\frac{1}{2}[pK_{a2} + pK_{a3}]$

C. $\frac{1}{2}[pK_{a1} + pK_{a3}]$

D. $pK_{a1} + pK_2$

Answer:

 [Watch Video Solution](#)

82. Which of the following is a buffer solution?

A. 500 mL of 0.1 N CH_3COOH + 500 mL of 0.1 N NaOH

B. 500 mL of 0.1 N CH_3COOH + 500 mL of 0.1 N HCl

C. 500 mL of 0.1 N CH_3COOH + 500 mL of 0.2 N NaOH

D. 500mL of 0.1N CH_3COOH + 500mL of 0.1N NaOH

Answer:

 [Watch Video Solution](#)

83. If 20 mL of 0.1 M NaOH is added to 30 mL of 0.2 M CH_3COOH ($pK_a=4.74$), the pH of the resulting solution is :

A. 4.44

B. 9.56

C. 8.96

D. 9.26




Answer:



[Watch Video Solution](#)

84. $H_2CO_3 + NaHCO_3$ found in blood helps in maintaining pH of the blood close to 7.4. An excess of acid entering the blood stream is removed by:

A. HCO_3^-



Answer:

 [Watch Video Solution](#)

85. 100mL of 0.02M benzoic acid ($pK_a = 4.2$) is titrated using 0.02M NaOH . pH values after 50mL and 100mL of NaOH have been added are

A. 3.50, 7

B. 4.2, 7

C. 4.2, 8.1

D. 4.2, 8.25

Answer:



Watch Video Solution

86. What is the pH of a solution of 0.28 M acid and 0.84 M of its conjugate base if the ionization constant of acid is 4×10^{-4} ?

- A. 3.88
- B. 3.34
- C. 7
- D. 10.12

Answer:



Watch Video Solution

87. The toxic compound 2,4-dinitrophenol has $K_a = 10^{-4}$. In an experiment, a buffer solution of 2,4-dinitrophenol was prepared with the pH adjusted to 5. Calculate the ratio of the concentrations of the dissociated ion to the undissociated acid:

A. 0.01

B. 0.1


C. 10

D. 100

Answer:

 [Watch Video Solution](#)

88. Equilibrium constant for the following reaction is 1×10^{-9} :



Determine the moles of pyridinium chloride (C_5H_5N) to obtain a buffer solution of pH=5 :

A. 0.1 mole

B. 0.2 mole

C. 0.3 mole

D. 0.4 mole

Answer:

 [Watch Video Solution](#)

89. Which one of the following mixture does not act as a buffer solution?

- A. Boric acid and borax
- B. Sodium phosphate & disodium hydrogen phosphate
- C. Sodium propionate and propionic acid
- D. Sod. Acetate and sodium propionate

Answer: d

 [Watch Video Solution](#)

90. The acid dissociation constant of uric acid is $K_a = 4.0 \times 10^{-6}$ M. The pH of a sample of urine is 6.0. What is the ratio of concentration of urate ion to uric acid in the urine?

A. 2.0

B. 4.0

C. 6.0

D. 0.25

Answer:

 [Watch Video Solution](#)

91. CH_3NH_2 (0.12 mole, $pK_b=3.3$) is added to 0.08 moles of HCl and the solution is diluted to one litre, resulting pH of solution is :

A. 10.7

B. 3.6

C. 10.4

D. 11.3

Answer:

 [Watch Video Solution](#)

92. An aqueous solution at room temperature contains 0.1 M NH_4Cl and 0.01M NH_4OH ($pK_b = 5$), the pH of the solution is :

A. 7.5

B. 6.8

C. 6.5

D. 8.0

Answer:

 [Watch Video Solution](#)

93. A 1L solution contains 0.2M NH_4OH and 0.2M NH_4Cl . If 1.0 mL of 0.001 M HCl is added to it what will be the $[OH^-]$ of the resulting solution ($K_b = 2 \times 10^{-5}$)

A. 2×10^{-5}

B. 5×10^{-10}

C. 2×10^{-3}

D. None of these

Answer:

 [Watch Video Solution](#)

94. 0.1 M formic acid solution is titrated against 0.1 M NaOH solution.

What would be the difference in pH between 1/5 and 4/5 stages of neutralization of acid?

A. $2 \log 3/4$

B. $2 \log 1/5$

C. $\log 1/3$

D. $2 \log 4$

Answer:



[Watch Video Solution](#)

95. The total number of different kind of buffers obtained during the titration of H_3PO_4 with $NaOH$ are:

- A. 3
- B. 1
- C. 2
- D. 4

Answer:



[Watch Video Solution](#)

96. A buffer solution is made up of acetic acid [$pK_a = 5$] having conc.=1.5M and sodium acetate having conc.=0.15 M. What is the number

OH^- ions present in 1 litre solution?

A. $10^{-10} N_A$

B. $10^{-4} N_A$

C. $10^{-3} N_A$



D. $10^{-6} N_A$

Answer:



[Watch Video Solution](#)

97. The pH of a solution of 0.10 M CH_3COOH increases when which of the following substances is added?






Answer:



[Watch Video Solution](#)

98. H^+ ion concentration of water does not change by adding:



Answer:



[Watch Video Solution](#)

99. pK_a of NH_4^+ is 9.26. Hence, effective range for $NH_4OH - NH_4Cl$ buffer is about pH:

A. 8.26 to 10.26

B. 4.74 to 5.74

C. 3.74 to 5.74

D. 8.26 to 9.26

Answer:



[Watch Video Solution](#)

100. 1.0 L solution is prepared by mixing 61 g benzoic acid ($pK_a = 4.2$) with 72 g of sodium benzoate and then 300 mL 1.0 M HBr solution was added. The pH of final solution is :

A. 3.6

B. 3.8

C. 4.2

D. 4.8

Answer:

 [Watch Video Solution](#)

101. The pH of a solution containing 0.4 M HCO_3^- and 0.2 M CO_3^{2-} is :

$$[K_{a1}(H_2CO_3) = 4 \times 10^{-7}, K_{a2}(HCO_3^-) = 4 \times 10^{-11}]$$

A. 10.4

B. 10.1

C. 6.1

D. 10.7

Answer:

 [Watch Video Solution](#)

102. The pH of the resultant solution of 20 mL of 0.1 M H_3PO_4 and 20 mL of 0.1 M Na_3PO_4 is :

A. $pK_{a1} + \log 2$


B. pK_{a1}

C. pK_{a2}

D. $\frac{pK_{a1} + pK_{a2}}{2}$





Answer:

[▶ Watch Video Solution](#)



103.





The graph represents the titration curve for :



Answer:

 [Watch Video Solution](#)

104. When 100 mL of 0.1 M NaCN solution is titrated with 0.1 M HCl solution the variation of pH of solution with volume of HCl added will be :



Answer:

 [Watch Video Solution](#)

105. The best indicator for the detection of the end point in the titration of a weak acid and a strong base is

- A. Methyl orange (3.1 to 4.4)
- B. Methyl red (4.2 to 6.3)
- C. Bromothymol blue (6 to 7.6)
- D. Phenolphthalein (8.2 to 10)

Answer:



[Watch Video Solution](#)

106. Select the best indicator from the given table for titration of 20 mL of 0.02 M CH_3COOH with 0.02 M NaOH. Given $pK_a(CH_3COOH) = 4.74$

Indicator	pH range
(I) Bromothymol blue	6.0-7.6
(II) Thymolphthalein	9.3-10.5
(III) Malachite green	11.4-13
(IV) M-Cresol purple	7.4-9.0

A. I

B. II

C. III

D. IV

Answer:



Watch Video Solution

107. Bromothymol blue is an indicator with a K_a value of 6×10^{-5} . What % of this indicator is in its basic form at a pH of 5 ?

A. 40

B. 85.7

C. 14.3

D. 60

Answer:

 [Watch Video Solution](#)

108. An acid-base indicator has a K_a of 3.0×10^{-5} . The acid form of the indicator is red and the basic form is blue. (a) By how much must the pH change in order to change the indicator from 75 % red to 75 % blue?

A. $8 \times 10^{-5} M$

B. $9 \times 10^{-5} M$

C. $1 \times 10^{-5} M$

D. $3 \times 10^{-4} M$

Answer:

 [Watch Video Solution](#)

109. An acid-base indicator which is a weak acid has a pK_{In} value =5.45. At what concentration ratio of sodium acetate to acetic acid would the indicator show a colour half-way between those of its acid and conjugate

base forms ?

[pK_a of acetic acid =4.75, $\log 2=0.3$]

A. 4: 1

B. 6: 1

C. 5: 1

D. 3: 1

Answer:



[Watch Video Solution](#)

110. A 20.0 mL sample of a 0.20 M solution of the weak diprotic acid H_2A is titrated with 0.250 M NaOH. The concentration of solution at the second equivalent point is:

A. 0.10 M NaHA

B. $0.153MN_{a_2}A$

C. $0.10MN_{a_2}A$

D. $0.0769MN_{a_2}A$

Answer:

 [Watch Video Solution](#)

111. During the titration of a weak diprotic acid (H_2A) against a strong base ($NaOH$), the pH of the solution half-way to the first equivalent point and that at the first equivalent point are given respectively by:

A. pK_{a1} and $pK_{a1} + pK_{a2}$

B. $\sqrt{K_{a1}C}$ and $\frac{pK_{a1} + pK_{a2}}{2}$

C. pK_{a1} and $\frac{pK_{a1} + pK_{a2}}{2}$

D. pK_{a1} and pK_{a2}

Answer:

 [Watch Video Solution](#)

112. In which of the following cases is the solution of $AgCl$ unsaturated?

A. $[Ag^+][Cl^-] < K_{sp}$

B. $[Ag^+][Cl^-] > K_{sp}$

C. $[Ag^+][Cl^-] = K_{sp}$

D. $[Ag^+][Cl^-] \leq K_{sp}$

Answer:



Watch Video Solution

113. When equal volumes of following solution are mixed, precipitation of $AgCl$?

($K_{sp} = 1.8 \times 10^{-10}$) will occur only with

A. $10^{-4}M(Ag^+)$ and $10^{-4}M(Cl^-)$

B. $10^{-5}M(Ag^+)$ and $10^{-5}M(Cl^-)$

C. $10^{-5}M(Ag^+)$ and $10^{-6}M(Cl^-)$

D. $10^{-10} M(Ag^+)$ and $10^{-10} M(Cl^-)$

Answer:

 [Watch Video Solution](#)

114. Choose the correct set of True/False for following statements:

(i) Silver chloride is more soluble in very concentrated sodium chloride solution than in pure water.

(ii) The pH of a buffer solution does not change on addition of small amount of an acid or a base.

(iii) Addition of NH_4Cl does not affect the pH of a solution of NH_4OH

(iv) Degree of hydrolysis of ammonium acetate does not depend upon the concentration of ammonium acetate solution.

(v) A mixture of acetic acid and sodium acetate can act as buffer solution.

A. TTF TT

B. FTTF F

C. TFTFT

D. FTTT

Answer:

 [Watch Video Solution](#)

115. A 1 litre solution containing NH_4Cl and NH_4OH has hydroxide ion ion concentration of 10^{-6} mol/litre. Which of the following hydroxides could be precipitated when the solution is added to 1 litre solution of 0.1 M metal ions?

- (I) $Ba(OH)_2$ ($K_{sp} = 5 \times 10^{-3}$), (II) $Ni(OH)_2$ ($K_{sp} = 1.6 \times 10^{-16}$)
(III) $Mn(OH)_2$ ($K_{sp} = 2 \times 10^{-13}$), (IV) $Fe(OH)_2$ ($K_{sp} = 8 \times 10^{-16}$)

A. I,II,IV

B. IV

C. II and IV

D. II,III,IV

Answer:



Watch Video Solution

116. 150 mL of 0.0008 M ammonium sulphate is mixed with 50 mL of 0.04 M calcium nitrate. The ionic product of $CaSO_4$ will be :

($K_{sp} = 2.4 \times 10^{-5}$ for $CaSO_4$)

A. $< K_{sp}$

B. $> K_{sp}$

C. $\approx K_{sp}$





D. None of these

Answer:



Watch Video Solution

117. In a saturated solution of $AgCl$, $NaCl$ is added gradually. The concentration of Ag^+ is plotted against the concentration of Cl^- . The graph appears as :



Answer:



Watch Video Solution

118. K_{sp} of $AgCl$ is 1×10^{-10} . Its solubility in $0.1\text{ M } KNO_3$ will be :

- A. 10^{-5} moles/litre
- B. $> 10^{-5}$ moles/litre
- C. $< 10^{-5}$ moles/litre
- D. None of these

Answer:



[Watch Video Solution](#)

119. 50 mL of a solution containing 10^{-3} mole of Ag^+ is mixed with 50 mL of a 0.1 M HCl solution. How much Ag^+ remains in solution ?

(K_{sp} of $AgCl = 1.0 \times 10^{-10}$)

- A. 2.5×10^{-9}
- B. 2.5×10^{-7}
- C. 2.5×10^{-8}

D. 2.5×10^{-10}

Answer:



Watch Video Solution

120. At a certain temperature, the solubility of the salt A_xB_y is S moles per litre. The general expression for the solubility product will be

A. S^2

B. $x^y y^x \cdot S^x + y$

C. $x^x y^y \cdot S^x + y$

D. $S^x + y$

Answer:



Watch Video Solution

121. What is the molarity of a saturated solution of $CaCO_3$?

$$(K_{sp} = 2.8 \times 10^{-9})$$

A. 2.6×10^{-5}

B. 2.8×10^{-9}

C. 5.2×10^{-5}

D. 5.6×10^{-9}

Answer:



Watch Video Solution

122. K_{sp} of $Zr_3(PO_4)_4$ in terms of solubility (S) is :

A. $108S^7$

B. $4S^3$

C. $6912S^7$

D. None of these

Answer:



Watch Video Solution

123. The solubility of electrolytes MX_1 , MX_2 and MX_3 is 1×10^{-3} moles per litre. Hence their respective solubility products are :

A. 10×10^{-6} , 4×10^{-9} , 27×10^{-12}

B. 10^{-9} , 4×10^{-9} , 32×10^{-12}

C. 10^{-9} , 8×10^{-8} , 32×10^{-12}

D. None of these

Answer:



Watch Video Solution

124. A saturated solution of $Ca_3(PO_4)_2$ has $[Ca^{2+}] = 2 \times 10^{-8}$ M and $[PO_4^{3-}] = 1.6 \times 10^{-5}$ M K_{sp} of $Ca_3(PO_4)_2$ is :

A. 3.2×10^{-13}

B. 3.2×10^{-34}





C. 2.048×10^{-33}

D. None of these

Answer:

 [Watch Video Solution](#)





125. Which of the following is most soluble in water?



Answer:

 [Watch Video Solution](#)

126. Silver ions are added to a solution with $[Br^-] = [Cl^-] = [CO_3^{2-}] = [AsO_4^{3-}] = 0.1M$. Which compound will precipitate with lowest $[Ag^+]$?



Answer:



[Watch Video Solution](#)

127. The solubility of different sparingly soluble salts are given as under :

S. No	Formula	Type	Solubility product
-------	---------	------	--------------------

(1)	AB		4.0×10^{-20}
-----	----	--	-----------------------

(2)	A_2B		3.2×10^{-11}
-----	--------	--	-----------------------

(3)	AB_3		2.7×10^{-31}
-----	--------	--	-----------------------

The correct increasing order of solubility is :

A. 1,2,3

B. 2,1,3

C. 1,2,3

D. 3,1,2

Answer:



[Watch Video Solution](#)

128. If K_{sp} for $HgSO_4$ is 6.4×10^{-5} , then solubility of this substance in mole per m^3 is

A. 8×10^{-3}

B. 6.4×10^{-5}

C. 8×10^{-6}

D. None of these

Answer:

 [Watch Video Solution](#)

129. The solubility of $Ba_3(AsO_4)_2$ (formula mass=690) is 6.9×10^{-2} g//100 mL. What is the K_{sp} ?

A. 1.08×10^{-11}

B. 1.08×10^{-13}

C. 1.0×10^{-15}

D. 6.0×10^{-13}

Answer:

 [Watch Video Solution](#)

130. The solubility of $AgBrO_3$ (formula mass=236) is 0.0072 g in 1000 mL. What is the K_{sp} ?

A. 2.2×10^{-8}

B. 3.0×10^{-10}

C. 3.0×10^{-5}

D. 9.3×10^{-10}

Answer:

 [Watch Video Solution](#)

131. The solubility of PbF_2 (formula mass =245) is 0.46 g/L. What is the solubility product?

A. 1.1×10^{-10}

B. 2.6×10^{-8}

C. 1.1×10^{-7}

D. 6.8×10^9

Answer:

 [Watch Video Solution](#)

132. How many grams of MgC_2O_4 (formula mass =122) will dissolve in 1.5 L of water?

$$(K_{sp} = 8.1 \times 10^{-5})$$

- A. 1.0
- B. 1.29
- C. 1.512
- D. 4.65

Answer:

 [Watch Video Solution](#)

133. What is the molarity of F^- ions in a saturated solution of BaF_2 ?

$$(K_{sp} = 1.0 \times 10^{-6})$$

A. 1.0×10^{-2}

B. 1.0×10^{-3}

C. 1.26×10^{-2}

D. 6.3×10^{-3}

Answer:

 [Watch Video Solution](#)

134. What is the molarity of F^- in a saturated solution of LnF_3 ?

$(K_{sp} = 7.9 \times 10^{-10})$

A. 2.3×10^{-3}

B. 8.3×10^{-3}

C. 1.0×10^{-3}

D. 7.0×10^{-3}

Answer:

 [Watch Video Solution](#)

135. What is the pH of a saturated solution of $Cu(OH)_2$?

$$(K_{sp} = 2.6 \times 10^{-19})$$

- A. 6.1
- B. 7.30
- C. 8.42
- D. 7.90

Answer:

 [Watch Video Solution](#)

136. The solubility product of $AgCl$ is $10^{-10} M^2$. The minimum volume (in m^3) of water required to dissolve $14.35mg$ of $AgCl$ is approximately :

- A. 0.01

B. 0.1

C. 100

D. 10

Answer:



[Watch Video Solution](#)

137. What is the molar solubility of $Fe(O)_2$ ($K_{sp}=8.0 \times 10^{-16}$) at pH 13.0?

A. 8.0×10^{-18}

B. 8.0×10^{-15}

C. 8.0×10^{-17}

D. 8.0×10^{-14}

Answer:



[Watch Video Solution](#)

138. What is the minimum pH necessary to cause a precipitate of $Pb(OH)_2$ ($K_{sp} = 1.2 \times 10^{-5}$) to form in a 0.12 M $PbCl_2$ solution?

- A. 12.4
- B. 10.8
- C. 12.0
- D. 11.1

Answer:

 [Watch Video Solution](#)

139. Which of the following would increase the solubility of $Pb(OH)_2$?

- A. Add hydrochloric acid
- B. Add a solution of $Pb(NO_3)_2$
- C. Add a solution of NaOH

D. None of the above-the solubility a compound is constant a constant temperature

Answer:

 [Watch Video Solution](#)

140. What is the molar solubility of Ag_2CO_3 ($K_{sp} = 4 \times 10^{-13}$) in $0.1MNa_2CO_3$ solution ?

A. 10^{-6}

B. 10^{-7}

C. 2×10^{-6}

D. 2×10^{-7}

Answer:

 [Watch Video Solution](#)

141. What is the concentration of Pb^{2+} when $PbSO_4$ ($K_{sp} = 1.8 \times 10^{-8}$) begins to precipitate from a solution that is $0.0045 M$ in SO_4^{2-} ?

- A. $4.0 \times 10^{-8} M$
- B. $1.0 \times 10^{-6} M$
- C. $2.0 \times 10^{-8} M$
- D. $4.0 \times 10^{-6} M$

Answer:



[Watch Video Solution](#)

142. What is the concentration of Ba^{2+} when BaF_2 ($K_{sp} = 1.0 \times 10^{-6}$) begins to precipitate from a solution that is $0.30 M F^-$?

- A. 9.0×10^{-7}
- B. 3.3×10^{-5}

C. 1.1×10^{-5}

D. 3.0×10^{-5}

Answer:

 [Watch Video Solution](#)

143. Solubility of AgCl in 0.2 M NaCl is x and that in 0.1 M AgNO_3 is y .

Then which of the following is correct?

A. $x = y$

B. $x > y$

C. $x < y$

D. We cannot predict

Answer:

 [Watch Video Solution](#)

144. What is the molarity of $Fe(CN)_6^{4-}$ in a saturated solution of $Ag_4[Fe(CN)_6]$?

$$(K_{sp} = 1.6 \times 10^{-41})$$

A. 1.6×10^{-8}

B. 5.2×10^{-8}

C. 2.0×10^{-8}

D. 2.3×10^{-9}

Answer:



Watch Video Solution

145. At 25°C , K_{sp} for $PbBr_2$ is equal to 8×10^{-5} . If the salt is 80 % dissociated, What is the solubility of $PbBr_2$ in mol//litre?

A. $\left[\frac{10^{-4}}{1.6 \times 1.6} \right]^{1/2}$

B. $\left[\frac{10^{-5}}{1.6 \times 1.6} \right]^{1/3}$

- C. $\left[\frac{10^{-4}}{0.8 \times 0.8} \right]^{1/3}$
- D. $\left[\frac{10^{-5}}{1.6 \times 1.6} \right]^{1/2}$

Answer:

 [Watch Video Solution](#)

146. What is the molar solubility of $Mn(OH)_2$ ($K_{sp} = 4.5 \times 10^{-14}$) in a buffer solution containing equal amounts of NH_4^+ and NH_3 ($K_b = 1.8 \times 10^{-5}$)?

- A. 3.0×10^{-4}
- B. 1.38×10^{-4}
- C. 1.38×10^{-3}
- D. 7.3×10^{-4}

Answer:

 [Watch Video Solution](#)

147. Find moles of NH_4Cl required to prevent $Mg(OH)_2$ from precipitating in a litre of solution which contains 0.02 mole NH_3 and 0.001 mole Mg^{2+} ions.

Given : $K_b(NH_3) = 10^{-5}$, $K_{sp}[Mg(OH)_2] = 10^{-11}$.

A. 10^{-4}

B. 2×10^{-3}

C. 0.02

D. 0.1

Answer:



[Watch Video Solution](#)

148. What mass of AgI will dissolve in 1.0 L of 1.0 M NH_3 ? Neglect change in conc. Of NH_3 .

[Given: $K_{sp}(AgI) = 1.5 \times 10^{-16}$), $K_f[Ag(NH_3)_2^+] = 1.6 \times 10^7$], (At.

Mass Ag=108, I=127)

A. 4.9×10^{-5} g

B. 0.0056 g

C. 0.035 g

D. 0.011 g

Answer:



Watch Video Solution

149. Consider the following statement and select correct option:

(I) K_{sp} of $Fe(OH)_3$ in aqueous solution is 3.8×10^{-38} at 298 K. The concentration of Fe^+ will increase when $[H^+]$ ion concentration decreases.

(II) In a mixture of NH_4Cl and NH_4OH in water, a further amount of NH_4Cl is added. The pH of the mixture will decrease.

(III) An aqueous solution of each of the following salt (NH_4I , $HCOOK$) will be basic, acidic respectively.

- A. only I is correct
- B. only II is correct
- C. only III is correct
- D. II and III are correct

Answer:

 [Watch Video Solution](#)

150. Equilibrium constants of T_2O (T or 3_1H isotope of 1_1H) and H_2O are different at 298 K. Let at 298 K pure T_2O has pT (like pH) is 7.62. The pT of a solution prepared by adding 10 mL of 0.2 M TCl to 15 mL of 0.25 M NaOT is:

- A. $2 - \log 7$
- B. $14 + \log 7$

C. $13.24 - \log 7$

D. $13.24 + \log 7$

Answer:

 [Watch Video Solution](#)

151. Liquid NH_3 dissociation to a slight extent, At a certain temp. its self dissociation constant $K_{SDC(NH_3)} = 10^{-30}$. The number of NH_4^+ ions are present per 100 cm^3 of pure liquid are :

A. 10^{-15}

B. 6.022×10^8

C. 6.022×10^7

D. 6.022×10^6

Answer:

 [Watch Video Solution](#)

152. To what volume of 10 litre of 0.5 M CH_3COOH ($K_a = 1.8 \times 10^{-5}$)

be diluted in order to double the hydroxide ion concentration :

A. 20 L

B. 30 L

C. 40 L

D. None of these

Answer:



[Watch Video Solution](#)

153. 20 mL of 0.1 M weak acid HA ($K_a = 10^{-5}$) is mixed with solution of

10 mL of 0.3 M HCl and 10 mL of 0.1 M NaOH. Find the value of $[A^-]$

$\frac{1}{[HA] + [A^-]}$ in the resulting solution :

A. 2×10^{-4}

B. 2×10^{-5}

C. 2×10^{-3}

D. 0.05

Answer:

 [Watch Video Solution](#)

154. What concentration of FCH_2COOH ($K_a = 2.6 \times 10^{-3}$) is needed so that $[H^+] = 2 \times 10^{-3}$?

A. 2×10^{-3} M

B. 2.6×10^{-3} M

C. 5.2×10^{-3} M

D. 3.53×10^{-3} M

Answer:

 [Watch Video Solution](#)

155. Calculate the ratio of $[HXOO^-]$ and $[F^-]$ in a mixture of 0.2 M $HCOOH$ ($K_a = 2 \times 10^{-4}$) and 0.1 M HF ($K_a = 6.6 \times 10^{-4}$):

A. 1:6.6

B. 1:3.3

C. 2:3.3

D. 3.3:2

Answer:

 [Watch Video Solution](#)

156. If first dissociation of $X(OH)_3$ is 100% where as second dissociation is 50% and third dissociation is negligible then the pH $4 \times 10^{-3} M X(OH)_3$ is :

A. 11.78

B. 10.78

C. 2.5

D. 2.22

Answer:



Watch Video Solution

157. H_3A is a weak triprotic acid

$$(K_{a1} = 10^{-5}, K_{a2} = 10^{-9}, K_{a3} = 10^{-13})$$

What is the value of pX of 0.1 M H_3A (aq.) solution ? Where $pX = -\log X$ and

$$X = \frac{[A^{3-}]}{[HA^{2-}]}$$

A. 7

B. 8

C. 9

D. 10

Answer:



[Watch Video Solution](#)

158. Calcium lactate is a salt of weak organic acid and strong base represented as $Ca(LaC)_2$. A saturated solution of $Ca(LaC)_2$ contains 0.6 mole in 2 litre solution. pOH of solution is 5.60. If 90 % dissociation of the salt takes place then what is pK_a of lactic acid?

A. $2.8 - \log(0.54)$

B. $2.8 + \log(0.54)$

C. $2.8 + \log(0.27)$

D. None the these

Answer:



[Watch Video Solution](#)

159. What is the concentration of $CH_3COOH(aq.)$ in a solution prepared by dissolving 0.01 mole of $NH_4^+CH_3COO^-$ in 1 L H_2O ?

$$\left[K_a(CH_3COOH) = 1.8 \times 10^{-5}, K_b(NH_4OH) = 1.8 \times 10^{-5} \right]$$

A. 5.55×10^{-5}

B. 0.10

C. 6.4×10^{-4}


D. 5.55×10^{-3}

Answer:



Watch Video Solution

160. K_a for the reaction,



what is the maximum pH value which could be used so that at least 80 %

of the total iron (*III*) in a dilute solution exists as Fe^{3+} ?

A. 2

B. 2.41

C. 2.79

D.

Answer:



Watch Video Solution

161. $Fe(OH)_2$ is diacidic base has $K_{b1} = 10^{-4}$ and $K_{b2} = 2.5 \times 10^{-6}$

What is the concentration of $Fe(OH)_2$ in 0.1 M $Fe(NO_3)_2$ solution?

A. 4×10^{-9}

B. 2.5×10^{-6}

C. 10^{-10}

D. 10^{-14}

Answer:

 [Watch Video Solution](#)

162. How many gm of solid KOH must be added to 100 mL of a buffer solution to make the pH of solution 6.0, if it is 0.1 M each w.r.t. acid HA and salt K A.

[Given: $pK_a(HA) = 5$]

A. 0.458

B. 0.327

C. 5.19

D. None of these

Answer:

 [Watch Video Solution](#)

163. Fixed volume of 0.1 M benzoic acid ($pK_a = 4.2$) solution is added into 0.2 M sodium benzoate solution and formed a 300 mL, resultant

acidic buffer solution. If pH of this buffer solution is 4.5 then find added volume of benzoic acid :

- A. 100 mL
- B. 150 mL
- C. 200 mL
- D. None of these

Answer:



[Watch Video Solution](#)

164. A 1.025 g sample containing a weak acid HX (mol. Mass=82) is dissolved in 60 mL water and titrated with 0.25 M NaOH. When half of the acid was neutralised the pH was found to be 5.0 and at the equivalence point the pH is 9.0. Calculate mass percentage of HX in sample :

- A. 50 %
- B. 75 %

C. 80 %

D. None of these

Answer:



Watch Video Solution

165. Which of the following expression for % dissociation of a monoacidic base (BOH) in aqueous solution at appreciable concentration is not correct?

A. $100 \times \sqrt{\frac{K_b}{c}}$

B. $\frac{1}{1 + 10^{(pK_b - pOH)}}$

C. $\frac{K_w[H^+]}{K_b + K_w}$

D. $\frac{K_b}{K_b + [OH^-]}$

Answer:



Watch Video Solution

166. A solution of weak acid HA was titrated with base NaOH. The equivalent point was reached when 40 mL of 0.1 M NaOH has been added. Now 20 mL of 0.1 M HCl were added to titrated solution, the pH was found to be 5.0. What will be the pH of the solution obtained by mixing 20 mL of 0.2 M NaOH and 20 mL of 0.2 M HA?

A. 7

B. 9

C. 11

D. None of these

Answer:



[Watch Video Solution](#)

167. A buffer solution 0.04 M in Na_2HPO_4 and 0.02 in Na_3PO_4 is prepared. The electrolytic oxidation of 1.0 milli-mole of the organic

compound RNHOH is carried out in 100 mL of the buffer. The reaction is $RNHOH + H_2O \rightarrow RNO_2 + 4H^+ + 4e^-$. The approximate pH of solution after the oxidation is complete is :

[Given: for H_3PO_4 , $pK_{a1} = 2.2$, $pK_{a2} = 7.20$, $pK_{a3} = 12$]

A. 6.90

B. 7.20

C. 7.5

D. None of these

Answer:



[Watch Video Solution](#)

168. When a 20 mL of 0.08 M weak base BOH is titrated with 0.08 M HCl, the pH of the solution at the end point is 5. What will be the pOH if 10 mL of 0.04 M NaOH is added to the resulting solution?

[Given: $\log 2 = 0.30$ and $\log 3 = 0.48$]

A. 5.40

B. 5.88

C. 4.92

D. None of these

Answer:



[Watch Video Solution](#)

169. Calculate approximate pH of the resultant solution formed by titration of 25 mL of 0.04 M Na_2CO_3 with 50 mL of 0.025 M HCl.

[Given: $pK_{a1} = 6.4$ and $pK_{a2} = 10.3$ for H_2CO_3]

A. 5.92

B. 6.88

C. 6.4

D. 5.88

Answer:

 [Watch Video Solution](#)

170. In the titration of solution of a weak acid HA and NaOH, the pH is 5.0 after 10 mL of NaOH solution has been added and 5.60 after 20 mL NaOH has been added.

What is the value of pK_a for HA?

A. 5.15

B. 5.3

C. 5.6

D. None of these

Answer:

 [Watch Video Solution](#)

171. A_3B_2 is a sparingly soluble salt with molar mass $M(\text{g mol}^{-1})$ and solubility $x \text{ gm litre}^{-1}$, the ratio of the molar concentration of B^{3-} to the solubility product of the salt is : —

A. $108 \frac{x^5}{m^5}$

B. $\frac{1}{108} \frac{M^4}{x^4}$

C. $\frac{1}{54} \frac{M^4}{x^4}$

D. None of these

Answer:



Watch Video Solution

172. A solution is $0.10 \text{ M } Ba(NO_3)_2$ and $0.10 \text{ M } Sr(NO_3)_2$. If solid Na_2CrO_4 is added to the solution, what is $[Ba^{2+}]$ when $SrCrO_4$ begins to precipitate?

$$[K_{sp}(BaCrO_4) = 1.2 \times 10^{-10}, K_{sp}(SrCrO_4) = 3.5 \times 10^{-5}]$$

A. 7.4×10^{-7}

B. 2.0×10^{-7}

C. 6.1×10^{-7}

D. 3.4×10^{-7}

Answer:

 [Watch Video Solution](#)

173. A solution is 0.01 M KI and 0.1 M KCl. If solid $AgNO_3$ is added to the solution, what is the $[I^-]$ when AgCl begins to precipitate?

$$[K_{SP}(AgI) = 1.5 \times 10^{-16}, K_{SP}(AgCl) = 1.8 \times 10^{-10}]$$

A. 3.5×10^{-7}

B. 6.1×10^{-8}





C. 2.2×10^{-7}

D. 8.3×10^{-8}

Answer:

 [Watch Video Solution](#)

174. Which of the following are conjugate acid-base pairs ?



Answer: A,B

 [Watch Video Solution](#)

175. If K_{a_1} , K_{a_2} and K_{a_3} be the first, second and third dissociation constant of H_3PO_4 and $K_{a_1} > K_{a_2} > K_{a_3}$ which is/are correct :

A. $[H^+] \approx \sqrt{K_{a1}[H_3PO_4]}$

B. $[H^+] \approx [HPO_4^{2-}]$

C. $K_{a2} \approx [HPO_4^{2-}]$

D. $[HPO_4^{2-}] = [PO_4^{3-}]$

Answer:

 [Watch Video Solution](#)

176. H_2A is a weak diprotic acid. If the pH of 0.1 M H_2A solution is 3 and concentration of A^{2-} is 10^{-12} at $25^\circ C$.

Select correct statement (s)

A. $[H^+]_{\text{total}} \approx [H^+]$ from first step of ionization of acid H_2A

B. Concentration of OH^- in solution is 10^{-3} M

C. The value of K_{a1} is nearly 10^{-5}

D. $pK_{a2} - pK_{a1} = 9$

Answer:



Watch Video Solution

177. Statement-1: pH value of acidic buffer solution changes , If buffer solution is diluted upto very large extent.

Statement-2: $[H^+]$ decreases due to change in concentration as well as α increases and decreases in concentration is more as compared to increases in α .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE AND STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer:



Watch Video Solution

178. Assertion : In a titration of weak monoprotic acid with strong base, the pH at the half equivalent point is pK_a .

Reason : At half equivalence point, it will form acidic buffer at its maximum capacity where $[acid] = [salt]$.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE AND STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer:



Watch Video Solution

179. Assertion: In the titration of Na_2CO_3 with HCl using methyl orange indicator, the volume of acid required is twice that of the acid required using phenolphthalein as indicator.

Reason: Two moles of HCl are required for the complete neutralisation of one mole of Na_2CO_3 .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE AND STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer:



[Watch Video Solution](#)

180. Assertion : Solubility of $AgCl$ in $NH_3(aq)$ is greater than in pure water.

Reason : When $AgCl$ dissolve in $NH_3(aq)$, complex ion $[Ag(NH_3)_2]^+$ formation takes place and solubility equilibrium of $AgCl$ shifted in forward direction.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE AND STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer:

 [Watch Video Solution](#)

181. Assertion (A): Solubility of $AgCN$ in acidic solutions is greater than in pure water.

Reason (R) : Solubility equilibrium of $AgCN$ is shifted in forward direction due to the formation of HCN .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE AND STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer:



Watch Video Solution

182. Calculate pOH of 0.1 M aq. Solution of weak base BOH ($K_b = 10^{-7}$) at $25^\circ C$.

 [Watch Video Solution](#)

183. pH of 0.01 M aq. Solution of HA is 4. Find the value of pK_a of HA at $25^\circ C$.

 [Watch Video Solution](#)

184. Calculate approximate pH of 10^{-10} M NaOH at $25^\circ C$.

 [Watch Video Solution](#)

185. Calculate pH of a resultant solution of 25 mL of 0.1 M HCl, 50 mL of 0.02 M HNO_3 and 25 mL of 0.1M NaOH

 [Watch Video Solution](#)

186. Calculate pH of a resultant solution of 0.1 M HA ($K_a = 10^{-6}$) and 0.5 M HB ($K_a = 2 \times 10^{-6}$) at 25°C .

 [Watch Video Solution](#)

187. 0.16g of N_2H_4 are dissolved in water and the total volume made upto 500 mL. Calculate the percentage of N_2H_4 that has reacted with water in this solution. (K_b of $\text{N}_2\text{H}_4 = 4.0 \times 10^{-6} <$)

 [Watch Video Solution](#)

188. Calculate pH of a buffer solution that contains 0.1M NH_4OH ($K_b = 10^{-5}$) and 0.1 M NH_4Cl .

 [Watch Video Solution](#)

189. Calculate the ratio of sodium formate and formic acid ($K_a = 2 \times 10^{-4}$) in a buffer solution of pH=4.3.

 [Watch Video Solution](#)

190. What is the pOH of 0.1 M KB (salt of weak acid and strong base) at $25^\circ C$? (Given : $pK_b \text{ of } B^- = 7$)

 [Watch Video Solution](#)

191. A certain weak acid has $K_a = 10^{-5}$. If the equilibrium constant for its reaction with a strong base is represented as 1×10^y then find the value of y.

 [Watch Video Solution](#)

192. If solubility of AgCl in 0.2 M solution of AgNO_3 is represented as $y \times 10^{-10}$ then find the value of y .

(Given: $K_{sp}(\text{AgCl}) = 10^{-10}$)

 [Watch Video Solution](#)

193. When one litre of a saturated solution of PbCl_2 (mol. Mass=278) is evaporated, the residue is found to weight 2.78g . If K_{sp} of PbCl_2 is represented as $y \times 10^{-6}$ then find the value of y .


 [Watch Video Solution](#)

194. A solution is saturated in SrCO_3 and SrF_2 . The CO_3^{2-} was found to be 10^{-3} mol/L . If the concentration of F^- in solution is represented as $y \times 10^{-2} \text{ M}$ then what is the value of y ?

[Given: $K_{sp}(\text{SrCO}_3) = 2.5 \times 10^{-10}$, $K_{sp}(\text{SrF}_2) = 10^{-10}$]

 [Watch Video Solution](#)

195. 10 mL of H_2A (weak diprotic acid) solution is titrated against 0.1M NaOH. pH of the solution is plotted against volume of strong base added and following observation is made




If pH of the solution at first equivalence point is pH_1 and at second equivalence point is pH_2 . Calculate the value of $(pH_2 - pH_1)$ at $25^\circ C$

Given for H_2A , $pK_{a_1} = 4.6$ and $pK_{a_2} = 8$, $\log 25 = 1.4$

[▶ Watch Video Solution](#)

196. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper blue is:



Given:

Acid	ka_1	ka_2	ka_3
H_3PO_4	10^{-3}	10^{-8}	10^{-12}
H_2CO_3	10^{-6}	10^{-11}	—
H_2CO_3	10^{-2}	10^{-5}	—



Watch Video Solution