



PHYSICS

NCERT - NCERT PHYSICS(GUJRATI)

MECHANICAL PROPERTIES OF FLUIDS

Example

1. The two thigh bones (femurs) each of cross-sectional area 10cm^2 support the upper part

of a human body of mass 40 kg . Estimate the average pressure sustained by the femurs.



[Watch Video Solution](#)

2. What is the pressure on a swimmer 10 m below the surface of a lake ? (Density of water

= 10^3 kgm^{-3} , Acceleration of gravity

$g = 10 \text{ ms}^{-2}$ and atm pressure

$P_a = 1.01 \times 10^5 \text{ Pa}$)



[Watch Video Solution](#)

3. The density of the atmosphere at sea level is $1.29\text{kg}/\text{m}^3$. Assume that it does not change with altitude. Then how high would the atmosphere extend? ($g = 9.8\text{ms}^{-2}$)



[Watch Video Solution](#)

4. At a depth of 1000m in an ocean (a) What is the absolute pressure? (b) what is the gauge pressure? (c) find the force acting on the window of area $20\text{cm} \times 20\text{cm}$ of a submarine at this depth, the interior of which is

maintained at sea level atmospheric pressure.

the density of sea water is $1.03 \times 10^3 \text{ kgm}^{-3}$.

$$g = 10 \text{ ms}^{-2}$$



Watch Video Solution

5. Two syringes of different cross sections (without needles) filled with water are connected with a tightly fitted rubber tube filled with water. Diameters of the smaller piston and larger piston 1.0 cm and 3.0 cm respectively. (a) Find the force exerted on the

large piston when a force of 10 N is applied to the smaller piston .(b) If the smaller piston is pushed in through 6.0 cm how much does the larger piston move out ?



[Watch Video Solution](#)

6. In a car lift compressed air exerts a force F on a small piston having a radius of 5.0 cm, This pressure is transmitted to a second piston of radius 15cm [Fig.10.7]. If the mass of the car to be lifted is 1350 kg. calculate F_1

What is the pressure necessary to accomplish this task?



[View Text Solution](#)

7. The flow of blood in a large artery of an anaesthetised dog is diverted through a Venturi Meter. The wider part of the meter has a cross-sectional area equal to that of the artery. $A = 8\text{mm}^2$. The narrow part has an area $a = 4\text{mm}^2$. The pressure drop in the

artery is 24Pa. What is the speed of the blood in the artery?



[View Text Solution](#)

8. A fully loaded boeing aircraft has a mass of $3.3 \times 10^5 \text{ Kg}$. Its total wing area is 500m^2 . It is in level flight with a speed of 960km/h (a) Estimate the pressure difference between the lower and upper surfaces of the wings (b) Estimate the fractional increase in the speed of the air on the surface of the wing relative to

the lower surface.[The density of air is $\rho = 1.2 \text{ kg m}^{-3}$]



Watch Video Solution

9. A metal block of area 0.10 m^2 is connected to a 0.010 kg mass via a string that passes over an ideal pulley (considered massless and frictionless) as in Fig 10.15. A liquid with a film thickness of 0.30 mm is placed between the block and the table. When released the block moves to the right with a constant speed of

0.085ms^{-1} . Find the coefficient of viscosity of the liquid.



[View Text Solution](#)

10. The terminal velocity of a copper ball of radius 2.0 mm falling through a tank of oil at 20°C is 6.5cm s^{-1} . Compute the viscosity of the oil at 20°C . Density of oil is $1.5 \times 10^3\text{kg m}^{-3}$. Density of copper is $8.9 \times 10^3\text{kg m}^{-3}$.



[View Text Solution](#)

11. The lower end of a capillary tube of diameter 2.00mm is dipped 8.00cm below the surface of water in a beaker. What is the pressure required in the tube in order to blow a hemispherical bubble at its end in water?

The surface tension of water at temperature of the experiments is $7.30 \times 10^{-2} \text{ Nm}^{-1}$.

atmospheric pressure = $1.01 \times 10^5 \text{ Pa}$

density of water

= 1000 kg/m^3 , $g = 9.80 \text{ ms}^{-2}$. Also calculate

the excess pressure.



[View Text Solution](#)

Exercise

1. Explain why

The angle of contact of mercury with glass is obtuse. While that of water with glass is acute

.



[Watch Video Solution](#)

2. Fill in the blanks using the words from the list appended with each statement

Surface tension of liquids generally.....with temperature (increases/decreases)



[View Text Solution](#)

3. Fill in the blanks using the words from the list appended with each statement

Viscosity of glass.....with temperature,whereas

viscosity of liquids....with temperature
(increases/decreases)



[View Text Solution](#)

4. Fill in the blanks using the words from the list appended with each statement

For solid of elastic modulus of rigidity, the shearing force is proportional to....while for fluids it is proportional to.....(shear strain/rate of shear strain).



[View Text Solution](#)

5. Fill in the blanks using the words from the list appended with each statement

For a fluid in a steady flow, the increase in flow speed at a constriction follows (conservation of mass/ Bernoulli's principle)



[View Text Solution](#)

6. A fluid flowing out of a small hole in a vessel results in a backward thrust on the vessel



[Watch Video Solution](#)

7. Explain why

A spinning cricket ball in air does not follow a parabolic trajectory.



[Watch Video Solution](#)

8. A 50kg girl wearing high heel shoes balances on a single heel. The heel is circular with a diameter 1.0 cm. What is the pressure exerted by the heel on the horizontal floor?



[View Text Solution](#)

9. A vertically-off shore structure is built to withstand a maximum stress of $10^9 Pa$ is the structure suitable for putting up on top of an oil well in the ocean? Take the depth of the ocean to be roughly 3km. And ignore ocean currents.



[View Text Solution](#)

10. A hydraulic automobile lift is designed to lift cars with a maximum mass of 3000 kg. The area of cross section of the piston carrying the load is 425cm^2 . What maximum pressure would the smaller piston have to exert?



View Text Solution

11. In the previous problem, if 15.0 cm of water and spirit each are further poured into the respective arms of the tube, what is the

difference in the levels of mercury in the two arms?



[Watch Video Solution](#)

12. Can Bernoulli's equations be used to describe the flow of water through a rapid in a river? Explain.



[View Text Solution](#)

13. Does it matter if one uses gauge instead of absolute pressures in applying Bernoulli's equation Explain?



Watch Video Solution

14. Glycerine flows steadily through a horizontal tube of length 1.5 m and radius 1.0 cm. IF the amount of glycerine collected per second at one end is $4.0 \times 10^{-3} \text{ kgs}^{-1}$. What is the pressure difference between the two

ends of the tube?(density of glycerine
= $1.3 \times 10^3 \text{ kg m}^{-3}$ and viscosity of glycerine
= 0.83 Pa s).[You may also like to check if the
assumption of laminar flow in the tube is
correct]




[Watch Video Solution](#)

15. In a test experiment on a model aeroplane
in a wind tunnel, the flow speeds on the upper
and lower surface of the wing are
 70 m s^{-1} and 63 m s^{-1} respectively. What is

the lift on the wing if its area is $2.5m^2$? Take the density of air to be $1.3kgm^{-3}$.

 [View Text Solution](#)

16. Figures 10.23(a) and (b) refer to the steady flow of a (non-viscous) liquid. Which of the two figures is incorrect? Why



 [View Text Solution](#)

17. The cylindrical tube of a spray pump has a cross section of 8.0cm^2 one end of which has 40 fine holes each of diameter 1.0 mm. If the liquid flow inside the tube is 1.5m min^{-1} . What is the speed of ejection of the liquid through the holes?



[View Text Solution](#)


18. A U-shaped wire is dipped in a soap solution, and removed. The thin soap film (which includes the small weight of the slider). The length of the slider is 30 cm. what is the surface tension of the film?



Watch Video Solution

19. Figure 10.24(a) shows a thin liquid film supporting a small weight $= 4.5 \times 10^{-2} N$
What is the weight supported by a film of the

same liquid at the same temperature in Fig.(b) and (c) ? Explain your answer physically.



अंकित 10.24



[View Text Solution](#)

20. What is the pressure inside the drop of mercury of radius 3.00mm at room temperature? Surface tension of mercury at that temperature ($20^{\circ}C$) is

$4.65 \times 10^{-1} Nm^{-1}$. The atmospheric pressure is $1.01 \times 10^5 Pa$. Also give the excess pressure inside the drop.



[View Text Solution](#)

21. What is the excess pressure inside a bubble of soap solution of radius 5.00 mm. given that the surface tension of soap solution at the temperature ($20^\circ C$) is $2.50 \times 10^{-2} Nm^{-1}$? If an air bubble of the same dimension were formed at depth of 40.0 cm inside a container

the soap solution (of relative density 1.20] what would be the pressure inside the bubble?[1 atmospheric pressure is $1.01 \times 10^5 Pa$).



[View Text Solution](#)

22. A tank with a square base of area $1.0m^2$ is divided by a vertical position in the middle. The bottom of the partition has a small hinged door of area $20cm^2$ The tank is filled with water in one compartment, and an acid

(of relative density 1.7) in the other. both to the height of 4.0 cm compute the force necessary to keep the door close.




[View Text Solution](#)

23. A manometer reads the pressure of a gas in an enclosure as shown in figure (a) When a pump removes some of the gas , the manometer reads as in figure (b). The liquid used in the manometers is mercury and the atmospheric pressure is 76 cm of mercury.

(a) Give the absolute and gauge pressure of the gas in the enclosure for cases (a) and (b), in units of cm of mercury.

(b) How would the levels change in case (b) if 13.6 cm of water (immiscible with mercury) are poured into the right limb of the manometer? (Ignore the small change in the volume of the gas).



24. Two vessels have the same area but different shapes. The first shape takes twice the volume of water that the second vessel requires to fill up to a particular common height? Is the force exerted by the water on the base of the vessel the same in the two cases? If so, why do the vessels filled with water to that same height give different readings on a weighing scale?



[View Text Solution](#)

25. In deriving Bernoulli's equation, we equated the work done on the fluid in the tube to its change in the potential and kinetic energy. (a) What is the largest average velocity of the blood flow in an artery of diameter $2 \times 10^{-3}m$ if the flow must remain laminar? (b) Do the dissipative forces become more important as the fluid velocity increases? Discuss qualitatively?



[View Text Solution](#)

26. What is the largest average velocity of blood flow in an artery of radius $2 \times 10^{-3}m$ if the flow must remain laminar?(b) What is the corresponding flow rate?(take viscosity of blood to be $2.084 \times 10^{-3}Pas$)



[View Text Solution](#)

27. A plane is in level flight at constant speed and each of its two wings has an area of $25m^2$. If the speed of the air is 180/h over the lower wing and 234km/h over the upper wing surface

, determine the plane's mass . (Take air density to be 1kgm^{-3}).



Watch Video Solution

28. In Millikan's oil drop experiment. What is the terminal speed of an uncharged drop of radius $2.0 \times 10^{-5}\text{m}$ and density $1.2 \times 10^3\text{kgm}^{-3}$. Take the viscosity of air at the temperature of the experiment to be $1.8 \times 10^{-5}\text{Pas}$. How much is the viscosity

force on the drop at that speed? Neglect buoyancy of the drop due to air.



[Watch Video Solution](#)

29. Mercury has an angle of contact equal to 140° with soda lime glass. A narrow tube of radius 1.00mm made of this glass is dipped in a trough containing mercury. By what amount does the mercury at the temperature of the experiment is 0.465 Nm^{-1} . Density of mercury $= 13.6 \times 10^3 \text{ kgm}^{-3}$



Watch Video Solution

30. Two narrow bores of diameters 3.00mm and 6.0mm are joined together to form a U-tube open at both ends. If the U-tube contains water, what is the difference in its levels in the two limbs of the tube? Surface tension of water at the temperature of the experiment is $7.3 \times 10^{-2} Nm^{-1}$. Take the angle of contact to be zero and density to water to be $1.0 \times 10^3 kgm^{-3}$



Watch Video Solution

31. It is known that density ρ of air decreases with height y as

$$\rho = \rho_0 e^{-y/y_0}$$

where $\rho_0 = 1.25 \text{kgm}^{-3}$ is the density at sea level. And y_0 is a constant . This density variation is called the law of atmosphere.

Obtain this law assuming that the temperature of atmosphere remains a constant (isothermal conditions). Also assume that the value of g remains constant.

A large He balloon of volume 1425m^3 is used

to lift a payload of 400kg. Assume that the balloon maintains constant radius as it rises.
How high does it rise?



Watch Video Solution