

# **BIOLOGY**

# **NEET & AIIMS**

# Mock test 29

**Example** 

**1.** One of the causes of variation known by early agriculturists, was hidden in

- A. Formation of deploid gametes
- B. Fusion of identical gametes
- C. sexual reproduction
- D. Both (1) & (2)

### **Answer: C**



**Watch Video Solution** 

**2.** Successfully bred domesticated varieties from while plans and elements are obtained through

- A. Artificial selection
- B. Selective crossing
- C. Fmasculation
- D. all except (3)

### **Answer: D**



**Watch Video Solution** 

3. A field of pea has plants bearing both yellow and green seeds green sheet colour is preferred, which of the following cross if performed repatedly has a higher probability

for obtaining pure line for green seed colour?

A. 
$$YY imes YY$$

B. 
$$Yy imes YY$$

$$\mathsf{C}.\, Yy imes yy$$

D. 
$$Yy imes Yy$$

### **Answer: C**



- **4.** By conducting\_\_\_\_ G. J. Mendel first demonstrate the scientific based of of\_\_\_\_.
  - A. selfing experiments, heredity and variation
  - B. Hybridisation experiments, chromosomal behaviour of gene
  - C. hybridisation experiment, inheritance and variation Himachal Bollywood

D. selfing experiments, one gene interaction

### **Answer: C**



**Watch Video Solution** 

**5.** mendelian experiments were infact the extension and development of hybridisation experiments on pea conducted by

A. Knight only

- B. Goss only
- C. Naudin only
- D. Both Knight and goss

#### **Answer: D**



**Watch Video Solution** 

**6.** select odd one out w.r.t recessive trait in pea plant

A. green seed colour

- B. green pod colour
- C. yellow pod colour
- D. white flower colour

### **Answer: B**



- **7.** Term 'Pure line' was coined by
  - A. Mendel
  - B. Johannsen

C. Bateson

D. T.H. Morgan

**Answer: B** 



**Watch Video Solution** 

**8.** which of the following trait of garden pea is not present on 4th chromosome?

A. flower position

B. stem height

C. pod shape

D. pod colour

### **Answer: D**



**Watch Video Solution** 

**9.** Mark the incorrect statement w.r.t law of segregation

A. Each gamete contains only one allele for

a gene

B. The two alleles present in the F\_1 generation segregate during random fusion of gametes, thus giving 3:1 ratio

- C. Each gamete is always pure for its/a trait
- D. The recessive trait which is not observed in the F\_1 generation reappears in F\_2 generation

Answer: B



**10.** Mendel failed to obtain the same results on

A. Hawkweed

B. Lablab

C. Cucurbita pepo

D. Both (1)&(2)

**Answer: D** 



<b>11.</b> Who	coined	the	term	'haterozygo	us'

- A. Bateson
- B. Saunders
- C. Jahannsen
- D. All except (3)

## **Answer: D**



<b>12.</b> who	gave	mendelian	inclusions	the	shape
of laws?					

- A. Carl Correns
- B. Hugo de Vries
- C. Tschermak
- D. All of these

### **Answer: A**



**13.** For hetrozygous tall plant of pea undergoing selfing, what would be the sum of phenotype and genotypes obtained in F\_1 generation?

- A. 2
- B. 5
- C. 3
- D. 4

**Answer: B** 



**14.** Percentage of inflated pod, wrinkled seed and axial flower obtained in F\_2 generation of Mendel's monohybrid cross are respectively

- A. 75%, 75%, 25%
- B. 25%, 50%,75%
- C. 75%,50%,25%
- D. 75%,25%,75%

### **Answer: D**



**15.** In a mendel's monohybrid cross for the shapes of pea seeds,1600 seeds were obtained in F\_2 generation. What will be the number of round seed and hybrid round seed, respectively?

A. 1200 & 800

B. 800 & 1200

C. 400 & 1200

D. 1200 & 400

### **Answer: A**



- **16.** When F<sub>1</sub> program shown intermediate phenotype between dominant and recessive phenotypes such as gene interaction is called
  - A. Complete dominance
  - B. Over dominance
  - C. pseudodominance
  - D. Incomplete dominance

### **Answer: D**



# **Watch Video Solution**

**17.** phenotypic and genotypic ratios are equal in

- A. incomplete dominance
- B. Test cross
- C. All monohybrid crosses
- D. Both (1)&(2)

### **Answer: D**



- **18.** white eye mutation leads to depigmentation in many parts of the body in Drosophila. it is referred as
  - A. Multiple alletes
  - B. Polymorphic effect
  - C. pleiotropic effect
  - D. Morphan syndrome

### **Answer: C**



- **19.** The \_\_\_\_ allele representing the original phenotype is \_\_\_\_ type and the allele \_\_\_\_ is generally the \_\_\_\_ type/allele.
  - A. Functional, dominant, modified, recessive
  - B. unmodified, wild, modified, mutant
  - C. modified, wild unmodified, recessive

D. Both (1) & (2)

**Answer: D** 



**Watch Video Solution** 

20. When a corss is made pink flowered and red flowered snapdragon plants, what proportion of phenotypein the offspring could be expected to be

A. 0.25

- B. 0.75
- C. 0.5
- D. 0.125

### **Answer: C**



**Watch Video Solution** 

**21.** Presence of more than two alleles for a gene in a population is referred as

A. Mutant alleles

- B. Multiple alleles
- C. isoalles
- D. Pseudoalleles

### **Answer: B**



**Watch Video Solution** 

**22.** select the set of parents that cannot produce child with blood group 'O'

A.  $I^A i imes I^A i$ 

B.  $I^B i imes I^A I^A$ 

C.  $ii imes I^Ai$ 

D.  $I^BI^B \times I^AI^B$ 

### **Answer: A**



Watch Video Solution

23. Human beings have \_\_\_\_\_ alleles for ABO blood grouping with \_\_\_\_\_ phenotypes and \_\_\_\_ genotypes.

- A. 4,3,6
- B. 4,6,3
- C. 3,4,6
- D. 6,3,4

### **Answer: C**



**Watch Video Solution** 

**24.** Theoretically, the modified allele could be responsible for the production of A. a non

functional enzyme B. normal or less effecient enzyme C. No enzyme at all

- A. A only
- B. A & B only
- C. B only
- D. A,B and C

#### **Answer: D**



**25.** For ABO system of blood group in humans, allele I^A produces \_\_\_\_ enzyme.

- A. N- acetylgalactosamyl transferase
- B. Glactosyl transferase
- C. N- muramic transferase
- D. NAG & NAM

**Answer: A** 



**26.** which of the following is correct regarding the location and event that occurs during capacitation of sperms?

A. Epididymis (location) & Maturation of sperm (Events)

B. Male urethra (Location) & Gain in sperm motility (Events)

C. female reproductive tract (Location) & chemical changes in sperms which prepare it to fertilize ovum (Events)

D. Vas deferens (Location) & Loss of sperm

tail (Events)

### **Answer: C**



**Watch Video Solution** 

**27.** which of the followin enzyme helps sperm to penetrate zone pellucida?

A. hyaluronidase

B. corona penetrating enzyme

- C. fertilizin
- D. acrosin

**Answer: D** 



**Watch Video Solution** 

**28.** which of the following event is/are included in fast block to prevent polyspermy?

- A. zona reactions
- B. depolarization of ovum membrane

C. cortical reaction

D. closing of Na^+ chanels of the ovum membrane

### **Answer: B**



Watch Video Solution

**29.** The event that triggers the formation of ootid is

A. binding of sperm head to receptors (

ZP3) on zona pollucida

B. Entry of sperm into secondary oocyte

C. Cortical reaction

D. Entry of sperm into fallopian tube

### Answer: B



**30.** A leads to mixing of genetic material of male and female which is known as B, which result is formation of C. Fill in the above blanks with suitable option

A. fertilization (A) & syngamy (B) & Synkaryon (C)

B. Fertilization (A) & Apomixis (B) & Embryo (C)

C. Syngamy (A) & Amphimixis (B) &

Synkaryon (C)

D. Syngamy (A) & Apomixis (B) & Synkaryon

(C)

### **Answer: C**



**Watch Video Solution** 

**31.** During embryonic development of human, second cleavage is completed after about

A. 30 hours of fertilization

B. 60 hours of fertilization

C. 72 hours of fertilization

D. 48 hours of fertilization

**Answer: B** 



**Watch Video Solution** 

**32.** which of the following is correct regarding cleavage division?

A. cleavage divisions are resulting in formation of blastomeres

B. cleavage divisions in mammals are slow and synchronous

C. during cleavage there is increase in DNA :cytoplasm ratio

D. zona pellucida breaks after 3rd cleavage division

Answer: C



33. in mammals embryo proper is formed from

A. trophoblast

B. inner cell mass

C. Cells of Rauber

D. Cytotrophoblast

**Answer: B** 



## 34. in humans implantation usually occurs at

- A. morula stage
- B. gastrula stage
- C. blastocyst stage
- D. Zygote Stage

## **Answer: C**



**35.** portion of endometrium lying between chorionic villi and myometrium is

- A. Decidua basalis
- B. Decidua capsularis
- C. Decidua vera
- D. Decidua parietalis

**Answer: A** 



**36.** which of the following events marks the beginning of gastrulation?

A. differentiation of inner cell mass into epiblast and hypoblast

B. formation of three germ layers

C. formation of primitive streak

D. formation of chorionic villi

## **Answer: C**



**37.** find the odd one w.r.t germ layers which form the following structures?

- A. Gonads
- B. Adrenal cortex
- C. kidneys
- D. Iris muscle

**Answer: D** 



**38.** presence of which hormone is diagnosed in

Gravidex test?

A. hCG

B. hPL

C. Chorionic thyrotropin

D. Estrogen

**Answer: A** 



**39.** which of the following statements is incorrect?

A. placenta is formed by structures contributes by both mother and fetus

B. TORCH infections are major cause of miscarriage in first trimester

C. Relaxin facilitates parturition by softening bones of public symphysis

D. there is no mixing of material and foetal

blood occur during foetal development

**Answer: C** 



**Watch Video Solution** 

**40.** which of the following is correct regarding embryonic development?

A. Heart sound of the foetus can be heard using stethoscope usually after four

week of gastation

B. First movement of foetus occurs just after 3rd month

C. Implantation occur at morula stage

D. Allantois serve as site of blood cell synthesis in foetal stage

Answer: A



**41.** Descending of testis into scrotal sac occur during

A. 3rd month of gestation

B. 5th month of gestation

C. 7th month of gestation

D. after birth of foetus

## **Answer: C**



**42.** Which extra embryonic membrane in human prevents descration of embryo inside uterus?

- A. Amnion
- B. Chorion
- C. Allantois
- D. Yock sac

**Answer: A** 

