



CHEMISTRY

JEE MAIN AND ADVANCED

ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

Examples


1. How can you perform the following conversions?

- (i) But-2-ene to Ethanol.
- (ii) Cyclohexanol to Cyclohexanone
- (iii) Butyne to Butan-2-one
- (iv) p-Nitrotoluene to p-Nitrobenzaldehyde
- (iv) Pent-3-en-2-ol to Pent-3-en-2-one





[Watch Video Solution](#)



2. Write the mechanism for following conversion:



 Watch Video Solution

3. Complete the following reactions:





[▶ Watch Video Solution](#)

4. How would you do the following conversions in not more than two steps?

(i) Propanone to Propene

(ii) Propanal to Butanone

(iii) Ethanol to 3-Hydroxybutanol

(iv) Benzaldehyde to Benzophenone

(v) Benzaldehyde to α -Hydroxyphenylacetic acid



(vi) Benzaldehyde to 3-Phenylpropan -1-ol





 **Watch Video Solution**

5. An organic compound A with molecular formula $C_9H_{10}O$ gives positive DNP and iodoform tests. It does not reduce. Tollen's reagent or Fehling reagent and does not decolourless bromine water or Bayer's reagent. On drastic oxidation with chromic acid, compound A gives a carboxylic acid having molecular formula $C_7H_6O_2$. Deduce the structure of organic compound A and write all the chemical reactions involved.

 **Watch Video Solution**

6. Identify A-D in the following reactions .





(3)

 **Watch Video Solution**

7. An unknown aldehyde A on reaction with alkali gives a β -hydroxy aldehyd, which loses water to form an unsaturated aldehyde, But-2-enal.

Another aldehyde 'B' disproportionation reaction in the presence of conc, alkali to form products C and D. C is an aryl alcohol wit the formula (C_2H_8)

(1) Identify A and B.



(ii) Write the sequence of reactions involved.

(iii) Name the product, when B reacts with Zn amalgam and hydrochloric acid.








Watch Video Solution

8. Complete the following reactions and identify A, B and C,



Watch Video Solution

9. Predict the main products in the following reactions:



 [Watch Video Solution](#)

10. How can you do the following- conversion?


(a) Pentan-1-ol to Pentanoic acid

(b) Propanal to Propanoic acid (C) 1-Phenylethanol to 3-Phenylpropanoic

acid



(d) Cyclopentene to Pentaanedioic acid

(e) 4-Nitrochlorobenzene to 4-Nitrobenzoic acid






 [Watch Video Solution](#)

11. Arrange the following compounds in order of decreasing acidic strength.



 [Watch Video Solution](#)

12. Complete the following reactions and form the products.



 [Watch Video Solution](#)

13. An organic compound A, C_8H_6 on reacting with dil. H_2SO_4 and $HgSO_4$ gives a compound 'B'-which can also be obtained from reaction of benzene with acid chloride in the presents of anh. $AlCl_3$ Compound B when reacted with iodine and aq. NaOH yields C and a yellow compound D. Identify A to D with proper justification.

 [Watch Video Solution](#)


14. How would you perform the following conversions?

- (i) Ethanol to 2-Hydroxybut-3-enoic acid
- (ii) Acetaldehyde to Butan-1-ol
- (iii) Acetophenone to 2-Phenylbutan-2-ol
- (iv) Ethanol to 3-Hydroxybutanol
- (v) Ethanol to Acetone



Watch Video Solution

15. Give the structure of the compounds in the following reactions



Watch Video Solution

16. Give the reasons for the following:






- (i) Iodoform is prepared by reacting acetone with hypoiodite and not with

iodine. Explain.

(ii) Halogen acids readily combine with alkene to form addition products but fail to react with carbonyl compounds. Discuss.


 [Watch Video Solution](#)

17. Identify compounds A to E in the following reactions.



 [Watch Video Solution](#)

18. Suggest a suitable oxidising agent for the given conversion.



 [Watch Video Solution](#)


19. An ester 'A' ($C_4H_8O_2$) on treatment with excess methyl magnesium chloride followed on acidification gives an alcohol 'B' as the sole organic product. Alcohol 'B' on oxidation with $NaOCl$ followed by acidification gives acetic acid. Deduce structure of A and B. Show the reactions involved.

 Watch Video Solution

20. Write mechanism for decarboxylation of β -keto acids.


 Watch Video Solution

21. Acid-catalyzed mechanism of given aldol reaction.




 Watch Video Solution

22. Propose mechanism of given reaction.



[▶ Watch Video Solution](#)



23.


Which type of reaction is use in above transformation step (A) and (B).

The structure of P and Q are?

[▶ Watch Video Solution](#)

24. In the following reaction sequence the correct structures of A, B and C




are



 **Watch Video Solution**

Illustrations



1. Predict the major products of the following reactions:



 **Watch Video Solution**

2. Write the structure of all possible products of reaction of $LiAlD_4$.

Followed by H_3O^+ with



 [Watch Video Solution](#)

3. Prepare





(a) Aniline from Benzoic acid.

(b) Ethyl amine from propane nitrile.

 [Watch Video Solution](#)

Assignment (Section A Competition Level Questions)


1. The general formula of both aldehyde & ketone is



Answer: B

 [Watch Video Solution](#)

2. The IUPAC name of following structure is



A. 1-Hydroxy-4-methylpentan-3-one

B. 2-Methyl-5-hydroxy-3-pentanone

C. 4-Methyl-3-oxo-1-pentanol

D. Hexanol-1-one-3

Answer: A

 [Watch Video Solution](#)

3. IUPAC name of $(C)Cl_3CHO$ is

- A. Chloral
- B. Trichloro acetaldehyde
- C. 1,1,1- Trichloroethanol
- D. 2,2,2- trichloroethanol

Answer: D

 [Watch Video Solution](#)

4. Which of the following is mixed ketone?

- A. Pentanone

B. Acetophenone

C. Benzaphenone

D. Butanone

Answer: B



[Watch Video Solution](#)

5. The reaction of acetaldehyde with HCN followed by hydrolysis gives a product which exhibits

A. Optical isomerism

B. Geometrical isomerism

C. Metamerism


D. Tautomerism

Answer: A



[Watch Video Solution](#)

6. Which factors will increase the reactivity of



group?

- (i) Presence of a group with positive inductive effect.
- (ii) Presence of a group with (-ve) inductive effect
- (iii) Presence of large alkyl group.

A. Only (i)

B. Only (ii)

C. Both (i) and (iii)

D. Both (ii) and (iii)

Answer: B



[Watch Video Solution](#)

7. Which of the following is optically active?

A. Ethylene glycol

B. Oxalic acid

C. Glycerol

D. Tartaric acid

Answer: D



[Watch Video Solution](#)

8. identify the wrong statement from the following:

A. Salicylic acid is a monobasic acid

B. Methyl salicylate is an ester

C. Salicylic acid gives violet colour with neutral ferric chloride as well as brisk effervescence with sodium bicarbonate

D. Methyl salicylate does not occur in natural oils.

Answer: D

 [Watch Video Solution](#)

9. The general formula of $C_nH_{2n}O_2$ could be for open chain

A. Diketones

B. Carboxylic acid

C. Diols

D. Dialdehyde

Answer: B

 [Watch Video Solution](#)

10. The IUPAC name of the phthalic acid-

A. Benzene-1,2-dicarboxylic acid

B. Benzene-1,4-dioic acid


C. Cyclo-1,3,5-trien-1,2-dioic acid

D. Benzene-1,3-dicarboxylic acid

Answer: A

 [Watch Video Solution](#)

11. In the following reaction, product P is



A. RCH_2OH

B. $RCOOH$

C. $RCHO$

D. RCH_3

Answer: C



Watch Video Solution

12. Dry distillation of calcium acetate gives :

- A. Acetaldehyde
- B. Ethane
- C. Acetic acid
- D. Acetone


Answer: D



Watch Video Solution

13. $CH_3 - CH_2 - C \equiv CH \xrightarrow[H_3O^{\oplus}]{R}$ Butanone, R is

- A. Hg^{+2}
- B. $KMnO_4$
- C. $KClO_3$



Answer: A



Watch Video Solution

14. Which of the following pathways produces 2-hexanone?

(i) 1-Hexyne is treated with $Hg^{2\oplus}$ / dil H_2SO_4

(ii) 3-methylhept-2-ene is treated with O_3 followed by hydrolysis.

(iii) n-butyl magnesium bromide reacts with acetaldehyde followed by hydrolysis and then oxidation with chromic acid.

(iv) Hydroboration-oxidation of 1-hexyne.

A. (i), (ii) and (iii)

B. (i) and (ii) only

C. (i), (ii) and (iv)

D. (i) and (iii) only

Answer: A



Watch Video Solution

15. An alkene of molecular formula C_9H_{18} on ozonolysis gives 2,2-dimethyl propanal and 2-butanone, then the alkene is

A. 2,2,4-trimethylhex-3-ene

B. 2,2,6-trimethylhex-3-ene


C. 2,3,4-trimethylhex-2-ene

D. 2,2,4-trimethylhex-2-ene

Answer: A




Watch Video Solution



In the above sequence of reactions, (A) and (B) are :




A. $(CH_3)_2C(OH)CH_3$, $(CH_3)_2C(OH)COOH$



Answer: A

 [Watch Video Solution](#)

17. A liquid was mixed with ethanol and a drop of concentrated H_2SO_4 was added. A compound with a fruity smell was formed. The liquid was



Answer: D

 [Watch Video Solution](#)

18. The major product obtained on interaction of phenol with sodium hydroxide and carbon dioxide is :

A. Benzoic acid

B. Salicylaldehyde

C. Salicylic acid gives violet colour with neutral ferric chloride as well as brisk effervescence with sodium bicarbonate

D. Phthalic acid

Answer: C



Watch Video Solution

19. Which of the following does not give benzoic acid on hydrolysis?

A. Phenyl cyanide


B. Benzoyl chloride

C. Benzyl chloride

D. Methyl benzoate

Answer: C

 Watch Video Solution



20.

The compound x is

A. CH_3COOH

B. $\text{BrCH}_2 - \text{COOH}$

C. $(\text{CH}_3\text{CO})_2\text{O}$

D. $\text{CHO} - \text{COOH}$

Answer: C

 [Watch Video Solution](#)


21. Acetic acid is obtained when:

- A. Methyl alcohol is oxidised with potassium permanganate
- B. Calcium acetate is distilled in the presence of calcium formate
- C. Acetaldehyde is oxidised with potassium dichromate and sulphuric acid
- D. Glycerol is heated with sulphuric acid


Answer: C

 [Watch Video Solution](#)


22. When m-chlorobenzaldehyde is treated with 50 % *KOH* solution, the product (s) obtained is (are)




A.



B.



C.



D.

Answer: C

 [Watch Video Solution](#)

23. Aldol condensation will not be observed in

A. Chloral

B. Phenyl acetaldehyde





C. Hexanal

D. Ethanal

Answer: A

 [Watch Video Solution](#)

24. Which of the following gives alcohol condensation reaction?



Answer: C

 [Watch Video Solution](#)

25. Which of the following organic compound exhibits positive Fehling test as well as iodoform test?

A. Methanol

B. Ethanol


C. Propanone


D. Ethanol

Answer: D


 Watch Video Solution

26. Predict the product







A.



B.



C.



D.

Answer: B

 [Watch Video Solution](#)

27. Which of the following will not undergo aldol condensation-

A. Acetaldehyde

B. Propanaldehyde

C. Benzaldehyde

D. Trideuteroacetaldehyde

Answer: C

 [Watch Video Solution](#)

28. Benzyl alcohol and sodium benzoate is obtained by the action of sodium hydroxide on benzaldehyde. This reaction is known as

A. Perkin reaction

B. Cannizzaro reaction

C. Sandmeyer reaction

D. Claisen condensation

Answer: B

 [Watch Video Solution](#)


29. Dimethyl ketones are usually characterised through

- A. Tollens reagent
- B. Iodoform test
- C. Schiff's test
- D. Benedict's reagent

Answer: B

 [Watch Video Solution](#)

30. In the Cannizzaro reaction given below:




- A. The attack of OH^- at the carbonyl group
- B. The transfer of hydride to the carbonyl group
- C. The abstraction of proton from the carboxylic group

D. The deprotonation of $PhCH_2OH$


Answer: B

 Watch Video Solution


31. Trichloroacetaldehyde, CCl_3CHO reacts with chlorobenzene in presence of sulphuric acid and produces.




A.



B.



C.






D.

Answer: D

 Watch Video Solution

32. Correct order of reactivity of



CH_3CHO , $C_2H_5COCH_3$ and CH_3COCH_3 is




Answer: A

 Watch Video Solution

33.




A.



B.





C.




D.




Answer: A

 [Watch Video Solution](#)



A.






Answer: B

 [Watch Video Solution](#)

35. Select the reagent for the following reaction:



A. SeO_2

B. $O_3, Zn / H_2O$

C. $O_3, H_2O_2 - (CH_3COOH)$

D. PCC

Answer: B

 [Watch Video Solution](#)

36. An aromatic compound X with molecular formula $C_9H_{10}O$ gives following chemical tests:


(i) Form 2,4-DNP derivative


(ii) Reduces Tollen's reagent

(iii) undergoes Cannizzaro's reaction


(iv) On vigorous oxidation 1, 2-Benzene dicarboxylic acid is obtained

The compound X is ,






B.



C.



D.

Answer: A

 [Watch Video Solution](#)

37. The increasing order of the rate of HCN addition to compound A-D is

(A) HCHO

(B) CH_3COOH_3

(C) $PhCOCH_3$

(D) $PhCOPh$

A. $A < B < C < D$

B. $D < B < C < A$


C. $D < C < B < A$

D. $C < D < B < A$

Answer: C

 Watch Video Solution

38. The correct order of reactivity of PhMgBr with



A. $(i) > (ii) > (iii)$


B. $(iii) > (ii) < (i)$

C. $(ii) < (iii) < (i)$

D. $(i) < (iii) < (ii)$





Answer: C

 Watch Video Solution



39. Identify 'X'


Identify 'X'




Answer: C

 Watch Video Solution


40. The diketone $CH_3 - \overset{\overset{O}{\parallel}}{C} - (CH_2)_2 - \overset{\overset{O}{\parallel}}{C} - CH_3$ on intermolecular aldol condensation gives the final product




A.



B.



C.




D.

Answer: C




Watch Video Solution


41. What will be the product, when carboxy phenol, obtained by Reimer Tiemann's process, is deoxidised with Zn powder?




A.



B.



C.







D.

Answer: D



Watch Video Solution

42. Which of the following aromatic acids is most acidic?






Answer: B



Watch Video Solution

43. $CH_3COOH \xrightarrow[P_2O_5]{\Delta} X$. Identify X





Answer: C



Watch Video Solution

44. Which one of the following orders of acidic strength is correct?




Answer: C




Watch Video Solution

45. Which of the following presents the correct order of the acidity in the given compounds?


A.




B.



C.



D.



Answer: D





Watch Video Solution

46. The correct order of acidic strength of carboxylic acid is

- A. Formic acid < Benzoic acid < Acetic acid
- B. Formic acid < Acetic Acid < Benzoic acid
- C. Acetic acid < Formic acid < Benzoic acid
- D. Acetic acid < Benzoic acid < Formic acid

Answer: D



Watch Video Solution

47. Treatment of benzoic acid with $Cl_2 / FeCl_3$ will give

- A. p-chlorobenzoic acid
- B. o-chlorobenzoic acid
- C. 2-4-dichlorobenzoic acid

D. m-chlorobenzoic acid

Answer: D

 [Watch Video Solution](#)

48. In esterification

- A. OH^- of acid is replaced by C_6H_5OH
- B. H^+ of acid is replaced by sodium metal
- C. OH^- of alcohol is replaced by chlorine
- D. OH^- of acid is replaced by RO^- group

Answer: D

 [Watch Video Solution](#)

49. Acetyl chloride is reduced with $LiAlH_4$, the product formed is

A. Methyl alcohol

B. Ethyl alcohol

C. Acetaldehyde

D. Acetone

Answer: B



Watch Video Solution

50. The reaction



A. Hell-Volhard-Zelinsky reaction

B. Birch reaction

C. Rosenmund reaction

D. Hunsdiecker reaction

Answer: A

 [Watch Video Solution](#)

51. Saponification of ethyl benzoate with caustic soda as alkali gives

- A. Benzyl alcohol and ethanoic acid
- B. Sodium benzoate and ethanol
- C. Benzoic acid and sodium ethoxide
- D. Phenol and ethanoic acid

Answer: B

 [Watch Video Solution](#)

52. Which of the following acids has the smallest dissociation constant?

- A. $CH_3CHF\text{COOH}$
- B. $FCH_2CH_2\text{COOH}$
- C. $BrCH_2CH_2\text{COOH}$



Answer: C



[Watch Video Solution](#)

53. Enol content is high test in

- A. Acetone
- B. Acetophenone
- C. Acetic acid
- D. Acetyl acetone

Answer: D



[Watch Video Solution](#)

54. Aliphatic aldehyde can be oxidised by

- A. Tollen's reagent
- B. Fehling solution
- C. Benedict solution
- D. All of these

Answer: D


 [Watch Video Solution](#)

55. Formaldehyde when treated with KOH (caustic potash) gives methanol and potassium formate, the reaction is known as

- A. Perkin reaction
- B. Claisen reaction
- C. Cannizzaro reaction
- D. Knoevengael reaction

Answer: C

 Watch Video Solution



56. Reactions

- A. Electrophilic substitution
- B. Nucleophilic substiution
- C. Nucleophilic substitution
- D. Electrophilic addition

Answer: B

 Watch Video Solution

57. Bakelite is a poylmer is

- A. HCHO + Phenol

B. HCHO + acetaldehyde

C. Phenol + H_2SO_4

D. HCHO + acetone

Answer: A



[Watch Video Solution](#)

58. Fehling test is positive for

A. Acetaldehyde

B. Benzaldehyde


C. Ether

D. Alcohol




Answer: A



[Watch Video Solution](#)




The similar reaction can take place with which of the following aldehyde?



Answer: C

 Watch Video Solution

60. Find the final product of the reaction



A. Benzene

B. Phthalimide


C. Benzamide

D. Acetaldehyde

Answer: B

 [Watch Video Solution](#)

Assignment (SECTION -B OBJECTIVE TYPES QUESTIONS (ONE OPTION IS CORRECT))



by α , β , γ and δ are in order:

A. $\alpha = \gamma < \beta = \delta$



B. $\alpha < \beta < \gamma = \delta$

C. $\alpha < \gamma = \delta < \beta$

$$D. \alpha = \beta = \gamma = \delta$$

Answer: C

 Watch Video Solution



Answer: B

 Watch Video Solution




3. $CH_3CHO + LiAlH_4 \rightarrow CH_3CH_2OH$ Nucleophile added in this reaction is





Answer: D

 Watch Video Solution

4. Complete the reaction





Answer: B

 **Watch Video Solution**

5. Consider the following sequence of reactions:




The final product [B] in the reaction would be:

- A. Meso
- B. Racemic
- C. Inversion
- D. All of these





Answer: B



6. Consider the following sequence of reaction



The final product [B] in the reaction would be:




Answer: D








Watch Video Solution



The product is





D. No reaction


Answer: A




Watch Video Solution




8.




A.



B.




C.





D.

Answer: A


 Watch Video Solution





9.




C. Both (1) and (2)



Answer: B




A.




B.

C. Both (1) and (2)




D.

Answer: A




11.


The product predominates is



A.




B.



C.

(4)



D.


Answer: A



Watch Video Solution


12. Which of the following carbonyl oxygen will form strongest hydrogen bond with H_2O molecule?

(1)




A.


(2)



B.



C.







D.

Answer: B

 [Watch Video Solution](#)

13. Which of the following would be the best synthesis of benzoic acid from bromobenzene?







Answer: C

 Watch Video Solution


14. Consider the following sequence of reactions:




Major product [B] of the given reaction would be




A.



B.



C.



Answer: C

[Watch Video Solution](#)


15. The compound which is not reduced by $LiAlH_4$ is

- A. Cyclohexanone
- B. 2-Methyl-1-butanol
- C. Ethyl benzoate
- D. ω -caprolactam






Answer: B

[Watch Video Solution](#)

16. Consider the following sequence of reactions.



The product [A] and [B] are respectively




Answer: A

Watch Video Solution

17.

The

compound




can be

compounded

oxidized

into



A. NaCN followed by hydrolysis

B. NaOI followed by H_3O^+





C. $KMnO_4$ hot followed by hydrolysis

D. $K_2Cr_2O_7$ followed H_3O^+

Answer: B

 Watch Video Solution


18. Which of the following conversion is known as Stefen's reduction?



Answer: B

 Watch Video Solution

19. Which reagent or sequence of reagents would best accomplish the following synthesis?




- A. (i) $LiAlH_4$ (ii) H^+ , Δ
- B. $NaBH_4$
- C. (i) Na / NH_3 (ii) $NaBH_4 / H^+$, Δ
- D. (i) Mg / Et_2O (ii) $LiAlH_6$, (iii) H^+ , Δ

Answer: B

 **Watch Video Solution**


20. An organic compound [X]. C_5H_8O reacts with hydroxylamine to form [Y]. In the presence of conc. H_2SO_4 gives δ -lactam. [X] neither give Benedicts test nor it respond positively towards haloform test. The compound [X] is

(1)




A.

(2)




B.

(3)



C.

(4)



D.



Answer: C



Watch Video Solution

21. Alanine can be obtained from acetaldehyde by the following sequence of reactions.

Alanine



Alanine


- A. Reactions with HCN, followed by NH_3 and finally acidic hydrolysis
- B. Reactions with HCN, acidic hydrolysis and finally reaction with NH_3
- C. Reaction with NH_3 , followed by HCN and finally acidic hydrolysis
- D. Reaction with NaHSO_3 , followed by NH_3 and finally acidic hydrolysis.

Answer: A




Watch Video Solution


22. Which of the following carboxylic acid is most reluctant to form ester with a given alcohol in the presence of a catalytic amount of concentrated H_2SO_4 ?




A.



B.



C.







D.

Answer: A

 [Watch Video Solution](#)

23. Among the given compounds



A. $III > IV > I > II$

B. $I > IV > II > III$


C. $III > I > IV > II$

D. $III > I > II > IV$

Answer: C







Watch Video Solution



24. A is

A is



Answer: C



View Text Solution

Assignment (SECTION -C Objective type questions more than one options are correct)

1. Which of the following reactions involve carbanion enolate as reactive intermediate?





- A. Kolbe-Schmidt reaction
- B. Reimer-Tiemann reaction
- C. Claisen condensation
- D. Aldol condensation

Answer: C::D



[Watch Video Solution](#)


2. Which of the following compounds can be synthesized by intramolecular aldol condensation in very good yield ?




Answer: B::D

 [Watch Video Solution](#)


3. Which of the following dicarboxylic acid will give cycle alkanone on heating?




A.



B.



C.







D.

Answer: A::B



Watch Video Solution

4. Which of the following compounds will give over all substitution product via. Addition/elimination mechanism with Ethylamine?



Answer: A::B

 [Watch Video Solution](#)

5. Which of the following reagents can be used to distinguish Benzaldehyde from acetophenone?

A. Tollen's reagent

B. Sodium hypoiodite



C. 2,4-Dinitrophenyl hydrazone

D. Benedicts solution


Answer: A::B

 [Watch Video Solution](#)

6. Which of the following compounds could liberate CO_2 with aqueous $NaHCO_3$?




(3)



C.

(4)




D.




Answer: A:D

 [Watch Video Solution](#)

7. Which of the following intermediates are involved in the acid catalyzed esterification of carboxylic acid?




A.







Answer: B::C::D

 [Watch Video Solution](#)

8. Consider the following sequence of reactions





Products of the given reaction would be





Answer: B::D

 [Watch Video Solution](#)

9. Which of the following compounds will give tertiary butanol as the major product when treated with excess of MeMgBr?






Answer: A::C

 Watch Video Solution

10. Which of the following statements are correct regarding given reaction?



[Labeled oxygen]

A. The given reaction primarily follows S_N2 -mechanism

B. In the given reaction condition inversion occurs at the chiral carbon


C. Labeled ^{18}O is present is formed carboxylic acid.

D. The given reaction primarily follows addition/elimination mechanism

Answer: A::C

 Watch Video Solution

11. Identify the set of reagents/ reaction conditions X and Y in the following of transformations



A. $X = Mg/Et_2O$. $Y = CO_2$, followed by H_3O^+

B. $X=AgCN$. $Y=(i) \xrightarrow{OH^-} / H_2$ (ii) H_3O^+

C. $X = NaCN$, $Y=(i) \xrightarrow{OH^-} / H_2O$ (ii) H_3O^+

D. X=alcoholic KOH, Y= (i) HCN, (ii) H_3O^+


Answer: A::C




Watch Video Solution

Assignment (SECTION -D Linked Comprehension type Questions)


1. Certain dicarboxylic acids spontaneously eliminate water when heated forming cyclic anhydrides. But for the reaction to be successful. The cyclic anhydrides product must normally have a ring size of five or six members. There are two important reasons, first, the second carboxyl group can serve as the acid catalyst (by intramolecular proton transfer), as well as the nucleophile. And second, the high temperature involved reduce the need for catalyst.




Which of the following dicarboxylic acid would you expect to form cyclic anhydride?




B.



C.



D.




Answer: C


 [View Text Solution](#)

2. Certain dicarboxylic acids spontaneously eliminate water when heated forming cyclic anhydrides. But for the reaction to be successful. The cyclic anhydrides product must normally have a ring size of five or six members. There are two important reasons, first, the second carboxyl


group can serve as the acid catalyst (by intramolecular proton transfer), as well as the nucleophile. And second, the high temperature involved reduce the need for catalyst.




Which of the following dicarboxylic acid will not form cyclic anhydrides?




A.



B.



C.




D.

Answer: C




Watch Video Solution


3. Certain dicarboxylic acids spontaneously eliminate water when heated forming cyclic anhydrides. But for the reaction to be successful. The cyclic anhydrides product must normally have a ring size of five or six members. There are two important reasons, first, the second carboxyl group can serve as the acid catalyst (by intramolecular proton transfer), as well as the nucleophile. And second, the high temperature involved reduce the need for catalyst.




Consider the following sequence of reaction,




The final product of the reaction would be




A.



B.



C.




D.

Answer: D

[▶ Watch Video Solution](#)


4. Both carbonyl compounds and acid derivatives though they contain



grouping yet


the reactions given by them are entirely different.

As Aldehydes and Ketones give addition product with a nucleophile, while carboxylic acid derivatives give nucleophilic acyl substitution through addition/elimination mechanism.



Where $X - = -\text{Cl} - \text{O} - \overset{\text{O}}{\parallel}{\text{C}} - \text{R} - \text{OR}, -\text{NR}(\text{R})$ etc.

where $X - = -\text{Cl}, -\text{O} - \overset{\text{O}}{\parallel}{\text{C}} - \text{R}, -\text{OR}', -\text{NRR}'$ etc.



Why aldehydes and Ketones give nucleophilic addition reaction while acid derivatives prefer nucleophilic acyl substitution reaction?

A. Carbonyl carbon of aldehyde and Ketones are more electron deficient than acid derivatives

B. Carbonyl carbon of acid derivatives are more electron deficient.

C. Acid derivatives also give addition reaction


D. In acid derivatives, acyl carbon is connected with a good leaving group

Answer: D



[Watch Video Solution](#)


5. Both carbonyl compounds and acid derivatives though they contain



grouping yet


the reactions given by them are entirely different.

As Aldehydes and Ketones give addition product with a nucleophile, while carboxylic acid derivatives give nucleophilic acyl substitution through addition/elimination mechanism.







Where $X = -Cl - O - \overset{O}{\parallel} C - R - OR, - NR(R)$ etc.

where X = Cl, -O-C(=O)-R, -OR', -NRR' etc.




Which of the following is most reactive towards a nucleophile?



Answer: A

 [View Text Solution](#)


6. Both carbonyl compounds and acid derivatives though they contain




grouping yet

the reactions given by them are entirely different.


As Aldehydes and Ketones give addition product with a nucleophile, while carboxylic acid derivatives give nucleophilic acyl substitution through addition/elimination mechanism.




Where $\text{X}^- = -\text{Cl}^- - \text{O}^- - \text{C}(=\text{O})^- - \text{R}^- - \text{OR}^-$, $-\text{N}(\text{R})\text{R}^-$ etc.




Carbonyl character is most suppressed in



A.




B.



C.

D.



Answer: A

 [Watch Video Solution](#)

Assignment SECTION - E Assertion - Reason Type Questions)

1. Statement-1: Benzaldehyde gives negative Benedict's test

and

Statement-2: Aldehydes do not respond positively with Benedicts reagent.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-1

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-1

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: C

 [Watch Video Solution](#)

2. Statement-1: When PhCHO is made to react in D_2O , deuterium is incorporated in product in Cannizzaro reaction.

and

Statement-2: In Cannizzaro reaction transfer of hydride takes place from one molecule to other.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-2

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-2

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: D



Watch Video Solution

3. Statement-1: Ketones in presence of highly electropositive elements such as Na, Mg, Yield 1,2 diol or pinacol.

and

Statement-2: Electrons released by electropositive elements convert C=O group into radical anion.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-3

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-3

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: A



[Watch Video Solution](#)

4. Statement -1: Cyclohexanone on reaction with secondary amines yields Schiff's base.

and

Statement-2: The initial adduct loses water to generate product.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-4

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-4

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: D



[Watch Video Solution](#)

5. Statement-1: Methanoic acid reduces mercuric chloride to mercurous chloride on heating while ethanoic acid does not.

and

Statement-2: Methanoic acid is stronger acid than ethanoic acid.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-1

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-1

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: B



[Watch Video Solution](#)

6. Statement-1: Acetic acid does not undergo haloform test.

and

Statement-2: Acetic acid does not contain any α -hydrogen.

- A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-6
- B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-6
- C. Statement-1 is true, statement -2 is False
- D. Statement-1 is False, Statement-2 is true

Answer: C

 [Watch Video Solution](#)

7. Statement-1: Decarboxylation of β - keto acid is very difficult

and

Statement-2: Decarboxylation takes place via a six membered cyclic transition state.

A. Statement-1 is True, Statement-2 is true, statement-2 is a correct explanation for statement-7

B. Statement -1 is true, Statement-2 is true, statement-2 is not a correct explanation for statement-7

C. Statement-1 is true, statement -2 is False

D. Statement-1 is False, Statement-2 is true

Answer: D

 [Watch Video Solution](#)

Assignment (SECTION - F Matrix Match type Questions)

1. Match the following

Column-I

- (A) Fries rearrangement
- (B) Claisen rearrangement
- (C) Bayer Villiger's rearrangement
- (D) Pinacol-pincolone rearrangement



Column-II

- (p) Acid-catalysed rearrangement
- (q) Concerned with ester
- (r) Involves electrophilic substitution
- (s) Intramolecular rearrangement



 [Watch Video Solution](#)

2. Match the following

Column-I




Column-II







Watch Video Solution

3. Match the following




4. Match the following:

Column I	Column II
(A) 	(p) Involves carbanion enolate
(B) 	(q) Involves nucleophilic addition
(C) 	(r) Molecular wt. of product more than that of reactant
(D) 	(s) Five membered ring formation
	(t) Generates resonance stabilized carbanion

 [Watch Video Solution](#)

Assignment (SECTION -G Integer Answer Type Questions)

1. Consider the following reactions.




Molecular weight of the product would increase by

[Assuming that mol. Wt. of C=12, O=16, H=1 and D=2]


 Watch Video Solution

2. What would be the maximum number of atoms involved in the formation of newly constructed ring in the given reaction?



 Watch Video Solution

3. What is the net negative charge on the major product of the given species when it is treated with excess NaOH?



Assignment (SECTION -H Multiple True-False Type Questions)

1. Statement -1: Acetophenone gives aldol condensation.

Statement-2: Benzaldehyde is more reactive than acetaldehyde towards nucleophilic addition.

Statement-3: Benzophenone has α -Hydrogens

A. TTT

B. FFF

C. TFF

D. TFT

Answer: C

2. Statement-1: Formation of cyclic acetal is entropically favourable.

Statement-2: PCC converts primary alcohol into aldehydes.

Statement-3: Aliphatic alcohols are practically insoluble in aqueous NaOH.

A. TTT

B. FFF

C. TFT

D. FTF



Answer: A



[Watch Video Solution](#)


Assignment (SECTION -I Subjective Type Questions)

1. Show how you would synthesize each compounds from starting materials containing no more than six carbon atoms.





 [Watch Video Solution](#)

2. Predict the products of the following reactions




(a)



(c)


 [Watch Video Solution](#)

3. Propose mechanism for the given reaction:




[▶ Watch Video Solution](#)

4. Supply the structures of A and B



[▶ Watch Video Solution](#)



5. An organic compound [A] C_9H_{12} gave (B) $C_8H_6O_4$ on oxidation by alkaline $KMnO_4$. [B] on heating does not form any anhydride. Also [B] reacts with Br_2 in the presence of iron to give only one monobromoderivative (



What are A,B and C?


 [Watch Video Solution](#)

6. Propose a synthesis of each of the following compounds from the indicated starting materials (s) and any other necessary reagents.





 [Watch Video Solution](#)

7. How many products are formed when compound [X] is decarboxylated?




 Watch Video Solution

8. Suggest a mechanism for each of the following reactions:




 Watch Video Solution

9. Suggest a synthesis of given product from the indicated starting material.



[▶ Watch Video Solution](#)


10. Identify the compounds [A] through [D] in the following sequence of reaction.







[▶ Watch Video Solution](#)

Assignment (SECTION -J Akash challengers Questions)

1. Consider the following sequence of reactions:




Intermediate [I] and corresponding major product [P] in the given reactions are




Answer: C


2. Consider the following sequences of reactions.




Major product would be




A.



B.



C.




D.

Answer: C


 Watch Video Solution

3. Draw a stepwise mechanism for the following reaction



 Watch Video Solution

4. Draw a mechanism for the following reaction.




 [Watch Video Solution](#)

5. Brevicomín, the aggregation pheromone of the western pine bark beetle, contains a bicyclic bridged ring system. Brevicomín is prepared by the acid catalyzed cyclization of 6,7-dihydroxy-2-nonanone.

(i) Suggest a structure for brevicomin.

(ii) Devise a synthesis of 6,7-dihydroxy-2-nonanone from 6-bromo-2-hexanone. You may also use three carbon alcohols and any required

organic or inorganic reagents.





[View Text Solution](#)

6. A compound [A] of molecular formula C_4H_9Br yields a compound [B] of molecular formula $C_4H_{10}O$ when reacted with aqueous $NaOH$. On oxidation [B] gives a ketone [C]. The vigorous oxidation of ketone gives ethanoic acid. Deduce the structure of A, B and C.

[Watch Video Solution](#)
















7. Suggest mechanisms for the following reactions which are catalyzed by strong base:



 [Watch Video Solution](#)

Try Yourself

1. Give common names and IUPAC names of the following compounds.



View Text Solution

2. Write the structures of the following compounds

- (i) 4-chloropentan-2-one
- (ii) p-Nitropropionophenone
- (iii) 3-Methylbutanal
- (iv) 4-Methylpent-3-en-2-one
- (v) p-Methoxy benzaldehyde
- (vi) o,o'-Dichlorobenzophenone
- (vii) Penta-1,4-dien-3-one
- (viii) 2-Methyl-4-oxohexanal
- (ix) Cyclohex-2-en-1-one
- (x) Pentan-2,4-dione



[Watch Video Solution](#)

3. Give chemical equations and name the main product formed when

- (i) Ethyl alcohol is mixed with air and passed over Ag catalyst at 520 K.
- (ii) Benzoyl chloride is treated with lithium tri-tert-butoxy aluminium hydride.
- (iii) Isopropylidene chloride is treated with caustic potash.

(iv) Methylmagnesium iodide is treated with HCN and the resultant product is hydrolysed.



 [Watch Video Solution](#)

4. What are the products formed by reductive ozonolysis of penta-1,3-diene?

 [Watch Video Solution](#)


5. Give chemical equations for following conversions:

(i) Propanal from allyl alcohol




 [Watch Video Solution](#)

6. Identify unknown compounds A to E in the following series of chemical reactions.





 Watch Video Solution

7. Complete the reaction



 Watch Video Solution

8. Complete the following reactions and identify A, B and C,



 [Watch Video Solution](#)

9. An organic compound [A] with molecular formula $C_5H_8O_2$ is reduced to n-pentane on treatment with $Zn - Hg/HCl$. The compound [A] forms a dioxime with hydroxyl amine and give a positive iodoform test and Tollen's test. Identify the compound [A] and deduce its structure?

 [Watch Video Solution](#)

10. Write reactions and conditions to bring about the following conversions.




(i) Toluene to benzaldehyde

(ii) Calcium formate to urotropin




Watch Video Solution

11. How can we distinguish chemically the following pairs of compounds?




Watch Video Solution









12. Write the various steps involved in the following conversion :



Watch Video Solution





13. Give IUPAC names of the following compounds.







Watch Video Solution

14. Complete the following reactions indicating the major product formed.





 [Watch Video Solution](#)

15. A compound [A] of molecular formula $\text{C}_4\text{H}_9\text{Br}$ yields a compound [B] of molecular formula $\text{C}_4\text{H}_{10}\text{O}$ when reacted with aqueous NaOH . On oxidation [B] gives a ketone [C]. The vigorous oxidation of ketone gives ethanoic acid. Deduce the structure of A, B and C.

 [Watch Video Solution](#)

16. How will you perform the following conversions?

(i) Acetaldehyde to acetone

(ii) Propanone to propene


(iii) Propene to propanone

(iv) Benzaldehyde to benzophenone

(v) Benzene to acetophenone


 [Watch Video Solution](#)

17. How will you convert benzene into



 [Watch Video Solution](#)


18. Arrange the following in decreasing order of nucleophilic addition



 [Watch Video Solution](#)

19. Give suitable explanation for the following:

(i) Chloral hydrate is a geminaldiol but still stable. How?



- (ii) Out of C_6H_5CHO and CH_3CH_2CHO which is more reactive towards nucleophilic addition?
- (iii) Why does pure HCN fail to react with aldehydes and ketones?
- (iv) Hydrazones of aldehydes and ketones are not prepared in strongly acidic medium. Why?
- (v) Sodium bisulphite is used to purify aldehydes and ketones. Explain.

[View Text Solution](#)

20. Hydrocarbon (X), C_7H_{12} , on reaction with boron hydride followed by treatment with CH_3COOH yields (A). On reductive ozonolysis (A) yields a mixture of two aldehydes, (B) and (C). Of these, only (B) can undergo Cannizzaro reaction. (A) exists in two geometrical isomers, (A-1) and

$(A - 2)$, of which $(A - 2)$ is more stable. Gives structures of (X), (A), (B), (C), $(A - 1)$, and $(A - 2)$ with proper reasoning.



Watch Video Solution