



CHEMISTRY

JEE MAIN AND ADVANCED

MOCK TEST 11

Example

1. in which of the following case entropy increases

- A. boiling of an egg
- B. crystallization of sugar from solution
- C. freezing of water
- D. stretching of rubber

Answer: A



View Text Solution

2. if the reaction is reversible all is at equilibrium ($\Delta S_{\text{universe}} = 0$), then the entropy of the system

A. will change abruptly

B. is greater than 1

C. will remain constant

D. is equal to 0

Answer: C



[View Text Solution](#)

3. the second law of thermodynamics states that

- A. in any spontaneous process entropy of the universe always increases
- B. energy can neither be created nor be destroyed
- C. energy of the universe remains constant
- D. $\Delta S_{\text{universe}} > 0$ for a spontaneous reaction

Answer: A



View Text Solution

4. in the reaction, ΔH and ΔS both are more than zero then in which of the following cases the reaction would be spontaneous

A. $\Delta H > T\Delta S$

B. $T\Delta S > \Delta H$

C. $\Delta H = T\Delta S$

D. $\Delta G > 0$

Answer: B



View Text Solution

5. the incorrect expression among the following is

A. $K = \frac{e^G}{R}T$

B. $\ln K = \frac{\Delta H^\circ - T\Delta S^\circ}{R}T$

C. $\Delta S_\Sigma = -\Delta H_s y \frac{s}{T}$

D. $\Delta S_s y s = q_r e \frac{y}{T}$

Answer: B



View Text Solution

6. $\Delta_t G^\circ$ for the following reaction



is, given that $\Delta_I G^\circ HI(g) = 1.8 \text{ kJ mol}^{-1}$.

$$\Delta_I G^\circ H_2S(g) = 33.8 \text{ kJ mol}^{-1}.$$

A. 30200 kJ

B. 30.2 kJ

C. -30200J

D. -302 J

Answer: C



View Text Solution

7. a process $A \rightarrow D$ is difficult to occur directly instead it takes place in three successive steps, $\Delta S(A \rightarrow B)=40$ e.u., $\Delta S(B \rightarrow C)=30$ e.u., $\Delta S(D \rightarrow C)=20$ e.u. where e.u. is entropy unit then the entropy change ΔS for the process $(A \rightarrow D)$ is

A. + 90 e.u.

B. + 50 e.u.

C. -90 e.u.

D. - 50 e.u.

Answer: B



View Text Solution

8. the entropy processes by certain substances at absolute zero is known as

A. Residual entropy

B. positive entropy

C. negative entropy




D. excess entropy

Answer: A




View Text Solution

9. the following equilibrium are given below,



constant of the reaction,



$K_1, K_2,$ and K_3 is

A. $K_1 \frac{K_2}{K_3}$

B. $K_1 \frac{K_3^2}{K_2}$

C. $K_2 \frac{K_3^3}{K_1}$


D. $K_1 K_2 K_3$

Answer: C



View Text Solution

10. consider the given reaction,



temperature if a mixture of 2 mol each of A, B

and C exist at equilibrium and $K_c = 9$ then

volume of the flask will be

A. 3L

B. 6L

C. 9L

D. 36L

Answer: B



View Text Solution

11. in a chemical reaction equilibrium is established when

A. backward and forward reaction ceases

B. concentration of reactants and products are equal

C. rate of backward reaction is equal to the rate of forward reaction

D. reaction ceases to generate heat

Answer: C

12. consider the reaction $2A_g + B_g \rightleftharpoons 2C_g$ for which $K_c = 350$. if 0.001 mole of each of the reactant and product are mix in a 2.0 L flux in the reaction quotient and spontaneous direction of the system will be

A. $Q_c = 0.002$, the equilibrium shifts to the left

B. $Q_c = 2000$, the equilibrium shifts to the left

C. $Q_c = 0.002$, the equilibrium shifts to right

D. $Q_c = 2000$, the equilibrium shifts to right

Answer: B



View Text Solution

13. for the homogeneous gaseous reaction

$A(g) + 2B(g) \rightleftharpoons C(g)$ at 300K. the value of

$K_c = 0.1$ When 2 mol of each of A and B are

mixed then what will be the approx

equilibrium pressure if 30% of A is converted

to C? [given that $\frac{(3.18)^1}{2} = 1.78$]

A. 90 atm

B. 100 atm

C. 178 atm

D. 1.78 atm

Answer: C



View Text Solution

14. for the reaction $P + Q \rightleftharpoons R + 2S$, initially the concentration of P is equal to that of Q (1 molar) but at equilibrium the concentration of R will be twice of that of P, then the equilibrium constant of the reaction is

A. $\frac{4}{3}$

B. $\frac{32}{3}$

C. $\frac{3}{10}$

D. $\frac{1}{10}$

Answer: B



[View Text Solution](#)

15. stage comes when no more sugar dissolves, instead it settles down at the bottom of the solution is now said to be

A. condensed

B. in a state of equilibrium

C. saturated

D. both (2) & (3)

Answer: D



View Text Solution