



CHEMISTRY

NCERT - NCERT CHEMISTRY(GUJRATI)

CHEMICAL EQUILIBRIUM - II

Self Evaluation A Choose The Correct Answer

1. State of chemical equilibrium is :

A. dynamic

B. stationery

C. none

D. both

Answer: A



Watch Video Solution

2. If the equilibrium constants of following reactions are $2A \rightleftharpoons B$ is K_1 and $B \rightleftharpoons 2A$ is K_2 , then

A. $K_1 = 2K_2$

B. $K_1 = 1/K_2$

C. $K_2 = (K_1)^2$


D. $K_1 = 1/K_2^2$

Answer: B



Watch Video Solution

3. For the equilibrium reaction



A. greater than K_c

B. less than K_c

C. Equal to K_c

D. Zero

Answer: C



Watch Video Solution

4. In the equilibrium $N_2 + 3H_2 \rightleftharpoons 2NH_3$, the maximum yield of ammonia will be obtained with the process having


- A. low pressure and high temperature
- B. low pressure and low temperature
- C. high temperature and high pressure
- D. high pressure and low temperature

Answer: C



Watch Video Solution

5. For the homogeneous gas reaction at 600 K



the equilibrium constant K_c has the unit

A. $\left(\text{mol dm}^{-3}\right)^{-1}$

B. $\left(\text{mol dm}^{-3}\right)$

C. $\left(\text{mol dm}^{-3}\right)^{10}$

D. $\left(\text{mol dm}^{-3}\right)^{-9}$

Answer: B



Watch Video Solution

6. Two moles of ammonia gas are introduced into a previously evacuated 1.0 dm^3 vessel in which it partially dissociates at high

temperature. At equilibrium 1.0 mole of ammonia remains. The equilibrium constant K_c for the dissociation is

A. $27 / 16 \left(\text{mole dm}^{-3} \right)^2$

B. $27 / 8 \left(\text{mole dm}^{-3} \right)^2$

C. $27 / 4 \left(\text{mole dm}^{-3} \right)^2$

D. None of these

Answer: A



Watch Video Solution

7. An equilibrium reaction is endothermic if K_1 and K_2 are the equilibrium constants at T_1 and T_2 temperatures respectively and if T_2 is greater than T_1 then

- A. K_1 is less than K_2
- B. K_1 is greater than K_2
- C. K_1 is equal to K_2
- D. None

Answer: A



Watch Video Solution


Self Evaluation B Answer In One Or Two Sentences



1. Dissociation of PCl_5 decreases in presence of increase in Cl_2 why ?



[Watch Video Solution](#)

2. Write the equilibrium constant for the following





[Watch Video Solution](#)

3. What is equilibrium constant ?



[Watch Video Solution](#)

4. What happens when



$\Delta n_g = 0$, $\Delta n_g = -ve$, $\Delta n_g = +ve$ in a

gaseous reaction.



Watch Video Solution

5. Calculate Δn_g , for the following reactions



Watch Video Solution

Self Evaluation C Answer Not Exceeding 60
Words


1. Write the K_p expression for



[Watch Video Solution](#)

Self Evaluation D Practice Problems

1. The equilibrium constant K_c for



constant of the forward reaction is

0.05 sec^{-1} . Calculate the rate constant of the reverse reaction.



[Watch Video Solution](#)

2. In the equilibrium $H_2 + I_2 \rightleftharpoons 2HI$ the number of moles of H_2 , I_2 and HI are 1,2,3 moles respectively. Total pressure of the reaction mixture is 60 atm. Calculate the partial pressures of H_2 , I_2 and HI in the mixture.



[Watch Video Solution](#)

3. In 1 litre volume reaction vessel, the equilibrium constant K_c of the reaction



What will be the degree of dissociation assuming only a small of 1 mole of PCl_5 has dissociated ?



[Watch Video Solution](#)

4. At temperature T_1 , the equilibrium constant of reaction is K_1 . At a higher temperature

T_2 , K_2 is 10% of K_1 . Predict whether the equilibrium is endothermic or exothermic.



[Watch Video Solution](#)

5. At 35°C , the value of K_p for the equilibrium reaction $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$ is 0.3174, Calculate the degree of dissociation when P is 0.2382 atm



[Watch Video Solution](#)

6. For the equilibrium $2\text{NOCl}_{(g)} \rightleftharpoons 2\text{NO}_{(g)} + \text{Cl}_{2(g)}$ the value of the equilibrium constant K_c is 3.75×10^{-6} at 790°C . Calculate K_p for this equilibrium at the same temperature.



[Watch Video Solution](#)

7. For the equilibrium $2\text{SO}_{3(g)} \rightleftharpoons 2\text{SO}_{2(g)} + \text{O}_{2(g)}$, the value of equilibrium constant is 4.8×10^{-3} at 700°C .

At equilibrium, if the concentration of SO_3 and SO_2 are $0.60M$ and $0.15M$ respectively. Calculate the concentration of O_2 in the equilibrium mixture.



[Watch Video Solution](#)

8. Hydrogen iodide is injected into a container at $458^\circ C$. Certain amount of HI dissociates to H_2 and I_2 . At equilibrium, concentration of HI is found to be $0.421M$ while $[H_2]$ and $[I_2]$ each equal to $6.04 \times 10^{-2}M$, at $458^\circ C$.

Calculate the value of the equilibrium constant of the dissociation of HI at the same temperature.



[Watch Video Solution](#)

9. Dissociation equilibrium constant of HI is 2.06×10^{-2} at $458^\circ C$. At equilibrium, concentrations of HI and I_2 are $0.36M$ and $0.15M$ respectively. What is the equilibrium concentration of H_2 at $458^\circ C$.




[Watch Video Solution](#)

10. The equilibrium constant for the reaction



K. Calculate the equilibrium constant for the



the same temperature.



[Watch Video Solution](#)

11. For the reaction $A + B \rightleftharpoons 3C$ at $25^\circ C$, a 3

litre volume reaction vessel contains 1,2 and 4

moles of A,B and C respectively at equilibrium,

calculate the equilibrium constant K_c of the reaction at $25^\circ C$.



Watch Video Solution

12. How much PCl_5 must be added to one litre volume reaction vessel at $250^\circ C$ in order to obtain a concentration of 0.1 mole of Cl_2 , K_c for $PCl_5 \rightleftharpoons PCl_3 + Cl_2$ is $0.0414 \text{ mol dm}^{-3}$ at $250^\circ C$.



Watch Video Solution

13. At 540, the equilibrium constant K_p for PCl_5 dissociation equilibrium at 1.0 atm 1.77 atm. Calculate equilibrium constant in molar concentration (K_c) at same temperature and pressure.



[Watch Video Solution](#)