



CHEMISTRY

JEE MAIN AND ADVANCED

THERMODYNAMICS

EXAMPLE

1. What kind of system does a water filled bucket represents?

 **Watch Video Solution**

2. On what values does the state functions depend?

 **Watch Video Solution**

3. Can adiabatic change be achieved by placing a system in thermostat?

 [Watch Video Solution](#)

4. A system gives out 20 J of heat and also does 40 J of work. What is the internal energy change?

 [Watch Video Solution](#)

5. An ideal gas enclosed in a cylinder fitted with a frictionless and weightless piston, pushes it outside. What would be the sign convention for the work ?

 [Watch Video Solution](#)

6. Calculate the amount of work done when one mole of an ideal gas contained in a bulb of 2 L capacity at 1 atm is allowed to enter in evacuated

bulb of 10 L capacity and in another case if is allowed to expand from 1 L to 5 L against a pressure of 1 atm.

 [Watch Video Solution](#)

7. If water vapour is assumed to be a perfect gas, molar enthalpy change for vaporization of 1 mol of water at 1 bar and 100°C is 41 kJ mol^{-1} . Calculate the internal energy, when 1 mol of water is vapourised at one bar pressure and 100°C .

 [Watch Video Solution](#)

8. A solution labelled as 1 molar NaCl is kept in the laboratory. What kind of property is this molarity, extensive or intensive ?

 [Watch Video Solution](#)

9. Calculate the amount of heat required to raise the temperature of 13.5 g aluminium from 300 K to 400 K, if specific heat of aluminium is $0.9 \text{ J } (^{\circ} \text{C})^{-1} \text{ g}^{-1}$.


 [Watch Video Solution](#)

10. A swimmer coming out from a pool is covered with a film of water weighing about 18 g. how much heat must be supplied to evaporate this water at 298 K ? Calculate the internal energy of vaporization at 100°C .

$\Delta_{vap}H^{\ominus}$ for water at 373 K = $40.66 \text{ kJ mol}^{-1}$

 [Watch Video Solution](#)

11. Calculate the enthalpy change where the standard heat of formation for gaseous NH_3 is -11.02 kcal/mol at 298 K. The reaction given is




 [Watch Video Solution](#)

12. If 150kJ of energy is needed for muscular work to walk a distance of one km, than how much of glucose one has to consume to walk a distance of five km, provided only 30 % energy is available for muscular work. The enthalpy of combustion of glucose is 3000kJmol^{-1}

 [Watch Video Solution](#)

13. Determine enthalpy change for the reaction,



Bond energies for $\text{C} - \text{H}$, $\text{C} - \text{Cl}$, $\text{Cl} - \text{Cl}$, $\text{H} - \text{Cl}$ are 412, 338, 242, 431kJmol^{-1} respectively.

 [Watch Video Solution](#)

14. When 1 mole of anhydrous CuSO_4 is dissolved in excess of water, -66.4 kJ heat is evolved. When one mole of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ is dissolved in water, the heat change is $+11.7\text{ kJ}$. Calculate enthalpy of hydration of CuSO_4 (anhydrous).



[Watch Video Solution](#)

15. The entropy change in the conversion of water to ice at 272 K for the system is $-21.85 \text{ JK}^{-1}\text{mol}^{-1}$ and that of surrounding is $+21.93 \text{ JK}^{-1}\text{mol}^{-1}$. State whether the process is spontaneous or not ?



[Watch Video Solution](#)

16. For a given reaction ΔG obtained was having positive sign convention. State whether the reaction was spontaneous or non-spontaneous.



[Watch Video Solution](#)

17. For the reaction $N_2(g) + 3H_2 \rightarrow 2NH_3(g)$

$\Delta H = -95.4 \text{ kJ}$ and $\Delta S = -198.3 \text{ JK}^{-1}$

Calculate the temperature at which Gibbs energy change (ΔG) is equal to zero. Predict the nature of the reaction at this temperature and above it.



[Watch Video Solution](#)

18. A monoatomic gas undergoes adiabatic process. Its volume and temperature are related as $TV^P = \text{constant}$. The value of p will be

- (1) 1.33 (2) 1.67
(3) 0.67 (4) 0.33



[Watch Video Solution](#)

19. An ideal gas with pressure P , volume V and temperature T is expanded isothermally to a volume $2V$ and a final pressure P_i . If the same gas is expanded adiabatically to a volume $2V$, the final pressure P_a . The ratio of the specific heats of the gas is 1.67. The ratio $\frac{P_a}{P_i}$ is



[Watch Video Solution](#)

20. Two moles of a perfect gas undergo the following processes:

- a. A reversible isobaric expansion from $(1.0\text{atm}, 20.0\text{L})$ to

(1.0atm, 40.0L).

b. A reversible isochoric change of state from (1.0atm, 40.0L) to (0.5atm, 40.0L)

c. A reversible isothermal expansion from (0.5atm, 40.0L) \rightarrow (1.0atm, 20.0L).

i. Sketch with labels each of the processes on the same $P - V$ diagram.

ii. Calculate the total work (w) and the total heat change (q) involved in the above process.

iii. What will be the values of ΔH for the overall process?



[Watch Video Solution](#)

21. At $0^\circ C$, ice and water are in equilibrium and $\Delta H^\circ = 6.00 \text{ kJ/mol}$ for the process



Value of ΔS° for the conversion of ice to liquid water is

- (1) $10.15 JK^{-1}mol^{-1}$ (2) $17.25 JK^{-1}mol^{-1}$
(3) $21.98 JK^{-1}mol^{-1}$ (4) $30.50 JK^{-1}mol^{-1}$



[Watch Video Solution](#)

22. The molar heat capacity of oxygen gas is given by the expression

$C_v = a + bT + cT^2$ where a, b and c are constants. What will be change in internal energy of 8 g of oxygen if it is heated from 200 K to 300 K at constant volume? Assume oxygen as an ideal gas. Given $a = 1.2 JK^{-1}mol^{-1}$, $b = 12.8 \times 10^{-2} JK^{-2}mol^{-1}$, $c = 3.3 \times 10^{-7} JK^{-3}mol^{-1}$.

- (1) 1000 J (2) 950.15 J
(3) 830.5 J (4) 315.5 J



[Watch Video Solution](#)

23. A balloon filled with helium gas having pressure 2 atm was kept at room temperature (of $27^\circ C$). What will be the resulting temperature if it bursts? (Assume that change occurs reversibly)

- (1) $-60^\circ C$ (2) $-45.65^\circ C$
(3) $-30.15^\circ C$ (4) $-10.5^\circ C$



[Watch Video Solution](#)

24. Find the entropy change when 90 g of H_2O at 10°C was converted into steam at 100°C . [Given

$$C_P(H_2O) = 75.29\text{JK}^{-1}\text{mol}^{-1} \text{ and } \Delta H_{\text{vap}} = 43.932\text{JK}^{-1}\text{mol}^{-1}]$$

 [Watch Video Solution](#)

25. What kind of system does a water filled bucket represents?

 [Watch Video Solution](#)

26. On what values does the state functions depend?

 [Watch Video Solution](#)

27. Can adiabatic change be achieved by placing a system in thermostal?

 [Watch Video Solution](#)

28. A system gives out 30 J of heat and also does 40 J of work. What is the internal energy change?

 [Watch Video Solution](#)

29. An ideal gas enclosed in a cylinder fitted with a frictionless and weightless piston, pushes it outside. What would be the sign convention for the work ?

 [Watch Video Solution](#)

30. Calculate the amount of work done when one mole of an ideal gas contained in a bulb of 2 L capacity at 1 atm is allowed to enter in evacuated bulb of 10 L capacity and in another case if is allowed to expand from 1 L to 5 L against a pressure of 1 atm.

 [Watch Video Solution](#)

31. If water vapour is assumed to be a perfect gas, molar enthalpy change for vaporization of 1 mol of water at 1 bar and $100^{\circ}C$ is $51kJmol^{-1}$. Calculate the internal energy, when 1 mol of water is vapourised at one bar pressure and $100^{\circ}C$.

 [Watch Video Solution](#)

32. A solution labelled as 1 molar NaCl is kept in the laboratory. What kind of property is this molarity, extensive or intensive ?

 [Watch Video Solution](#)

33. Calculate the amount of heat required to raise the temperature of 13.9g aluminium from $300K$ to $400K$, if specific heat of aluminium is $0.9J^{\circ}C^{-1}g^{-1}$.


 [Watch Video Solution](#)

34. A swimmer coming out from a pool is covered with a film of water weighing about 18 g. how much heat must be supplied to evaporate this water at 298 K ? Calculate the internal energy of vaporization at 100° C.

$$\Delta_{vap}H^{\ominus} \text{ for water at } 373 \text{ K} = 40.66 \text{ kJ mol}^{-1}$$

 [Watch Video Solution](#)

35. Calculate the enthalpy change where the standard heat of formation for gaseous NH_3 is -11.02 kcal/mol at 298 K. The reaction given is



 [Watch Video Solution](#)

36. If 150kJ of energy is needed for muscular work to walk a distance of one km, than how much of glucose one has to consume to walk a distance of five km, provided only 30 % energy is available for muscular work. The enthalpy of combustion of glucose is 3000 kJ mol^{-1}

 [Watch Video Solution](#)

37. What amount of heat will be released when (a) 0.1 mol of hydrochloric acid is neutralised by 0.1 mol of caustic soda solution (b) 0.2 mol of nitric acid is mixed with 0.2 mol of caustic soda solution?

 Watch Video Solution


38. Calculate amount of heat evolved when

(a) 500 cm^3 of 0.5 hydrochloric acid is mixed with 200 cm^3 of 0.8 M NaOH solution.

(b) 250 cm^3 of $0.4\text{ M H}_2\text{SO}_4$ is mixed with 500 cm^3 of 0.2 M NaOH solution.

 Watch Video Solution

39. Determine the enthalpy change for the given reaction.



Bond energies are given as follows:

$$C - H = 412 \text{ kJ mol}^{-1}$$

$$C - Cl = 338 \text{ kJ mol}^{-1}$$

$$Cl - Cl = 242 \text{ kJ mol}^{-1}$$

$$H - Cl = 431 \text{ kJ mol}^{-1}.$$

 [Watch Video Solution](#)

40. When 1 mole of anhydrous $CuSO_4$ is dissolved in excess of water, -66.4 kJ heat is evolved. When one mole of $CuSO_4 \cdot 5H_2O$ is dissolved in water, the heat change is +11.7 kJ. Calculate enthalpy of hydration of $CuSO_4$ (anhydrous).

 [Watch Video Solution](#)


41. The entropy change in the conversion of water to ice at 272 K for the system is $-21.85 \text{ JK}^{-1} \text{ mol}^{-1}$ and that of surrounding is $+21.93 \text{ JK}^{-1} \text{ mol}^{-1}$. State whether the process is spontaneous or not ?

 [Watch Video Solution](#)

42. For a given reaction ΔG obtained was having positive sign convention. State whether the reaction was spontaneous or non-spontaneous.

[Watch Video Solution](#)

43. For the reaction



$$\Delta H = -85.4 \text{ kJ} \text{ and } \Delta S = -188.3 \text{ Jk}^{-1}$$

Calculate the temperature at which Gibbs energy change (ΔG) is equal to zero. Predict the nature of the reaction at this temperature and above it.

[Watch Video Solution](#)

1. What kind of system does a thermos flask containing coffee represents?

 [Watch Video Solution](#)

2. A tree represents an _____ system.

 [Watch Video Solution](#)


3. In the equation $pV = nRT$, how many state variables are present?

 [Watch Video Solution](#)

4. Temperature is a state function or not?

 [Watch Video Solution](#)

5. Consider the following figure and tell the nature of work done.



[▶ Watch Video Solution](#)

6. A beaker containing water at 373 K is representing an open system, is placed in the room. Predict the nature of work done.

[▶ Watch Video Solution](#)

7. 160 J of work is done on the system and at the same time 100 J of heat is given out. What is the change in the internal energy?

[▶ Watch Video Solution](#)

8. If change in internal energy -80 J. The work done by system is $+40$ J.

Calculate the heat change.

 [Watch Video Solution](#)

9. If the external pressure is more than internal pressure then what will be the sign convention for the work done on a cylinder fitted with a frictionless and weightless piston containing one mole of an ideal gas?

 [Watch Video Solution](#)

10. What would be the sign convention for work done on a system for a cylinder containing one mole of an ideal gas where $P_{ext} = P_f$?

 [Watch Video Solution](#)

11. One mole of oxygen is allowed to expand isothermally and reversibly from 5 m^3 to 10 m^3 at 300 K . Calculate the work done in expansion of the gas.

 [Watch Video Solution](#)

12. How much heat would be absorbed by two litres of an ideal gas at 5 atm pressure if it expands isothermally into an evacuated flask of five litres?

 [Watch Video Solution](#)

13. The heat of combustion of naphthalene $\{C_{10}H_8(s)\}$ at constant volume was measured to be -5133 kJ mol^{-1} at 298 K . Calculate the value of enthalpy change (Given $R = 8.314\text{ JK}^{-1}\text{ mol}^{-1}$)

 [Watch Video Solution](#)

14. How it is possible to find the value of ΔU and not the value of U ?

 [Watch Video Solution](#)

15. Out of pressure, mass, moles and volume, which is an intensive property?

 [Watch Video Solution](#)

16. Pick the odd one out : (Intensive or extensive).

Refractive index, density, temperature and enthalpy.

 [Watch Video Solution](#)

17. Calculate the amount of heat required to raise the temperature of 5 g of iron from $25^{\circ}C$ to $75^{\circ}C$. The specific heat capacity of iron is 0.45 J/g .

 [Watch Video Solution](#)

18. Calculate the weight of gold having specific heat capacity $0.13 \text{ J } (\text{.}^\circ \text{C})^{-1} \text{g}^{-1}$ which is heated from 25°C to 300°C , where the heat absorbed is 1500 J .

 [Watch Video Solution](#)


19. 90 g of water spilled out from a vessel in the room on the floor. Assuming that water vapour behaving as an ideal gas, calculate the internal energy change when the spilled water undergoes complete evaporation at 100°C . (Given the molar enthalpy of vaporisation of water at 1 bar and $373 \text{ K} = 41 \text{ kJ mol}^{-1}$).

 [Watch Video Solution](#)

20. One mole of carbon undergoes incomplete combustion to produce carbon monoxide, Calculate $(\Delta H - \Delta U)$ for the formation of CO at 298 K . Given $R = 8.314 \text{ JK}^{-1} \text{mol}^{-1}$.

 [Watch Video Solution](#)

21. The standard heat of formation of at 298 K for $\text{Cl}_4(g)$, $\text{H}_2\text{O}(g)$, $\text{CO}_2(g)$ and $\text{HCl}(g)$ are -25.5, -57.8, -94.1 and -22.1 kcal mol^{-1} respectively. Calculate $\Delta_r H^\ominus$ for the reaction



Hint : $\Delta_r H^\ominus = \sum \Delta_f H_{\text{Products}}^\ominus - \sum \Delta_f H_{\text{Reactants}}^\ominus$

 Watch Video Solution

22. Why ΔH_f° of elements is zero?

 Watch Video Solution

23. An athlete is given 180 g of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$). He utilises 50% of the energy due to internal combustion in the body. In order to avoid storage of energy in the body, calculate the mass of water he would need to


perspire. Given enthalpy of combustion of glucose is $-2800 \text{ kJ mol}^{-1}$ and enthalpy of evaporation of water is 44 kJ mol^{-1}

 [Watch Video Solution](#)

24. A goobar gas plant produces methane and supplies to the families. If a family requires $15,000 \text{ kJ}$ of energy per day, how many days will the quote of 10 kg will last ? ($\Delta_c H$ of $\text{CH}_4 = -1665 \text{ kJ mol}^{-1}$)

 [Watch Video Solution](#)

25. Calculate the heat of combustion of eithene



The bond energy data are given below

$$\text{C} = \text{C} = 619 \text{ kJ mol}^{-1}$$

$$\text{C} - \text{H} = 414 \text{ kJ mol}^{-1}$$

$$\text{O} = \text{O} = 499 \text{ kJ mol}^{-1}$$


$$\text{C} = \text{O} = 724 \text{ kJ mol}^{-1}$$

$$\text{O} - \text{H} = 460 \text{ kJ mol}^{-1}$$



Watch Video Solution

26. Calculate the enthalpy change during the reaction :



Given, $e_{H-H} = 435 \text{kJmol}^{-1}$, $e_{Br-Br} = 192 \text{kJmol}^{-1}$ and

$e_{H-Br} = 368 \text{kJmol}^{-1}$.



Watch Video Solution

27. A metal chloride dissolves endothermally and when 7.45 g of its anhydrous form are dissolved in excess of water, the amount of heat absorbed is X kJ. Calculate the enthalpy of solution if molar mass of chloride is 74.5.



Watch Video Solution

28. Integral enthalpy of solution of KCl, when 1 mole of it is dissolved in 20 mole water is +15.90 kJ. When 1 mole of it is dissolved in 200 mole

water. ΔH is 15.58 kJ. Calculate enthalpy of hydration or dilution.

 [Watch Video Solution](#)

29. The entropy change in the conversion of water to ice at 273 K for the system is $-21.99 \text{ JK}^{-1}\text{mol}^{-1}$ and that of surrounding is $+21.99 \text{ JK}^{-1}\text{mol}^{-1}$. State whether the process is spontaneous or non-spontaneous.

 [Watch Video Solution](#)

30. The entropy change in the conversion of water to ice at 274 K for the system is $-22.13 \text{ JK}^{-1}\text{mol}^{-1}$ and for surrounding is $+22.05 \text{ JK}^{-1}\text{mol}^{-1}$. State whether the process is spontaneous or non-spontaneous.

 [Watch Video Solution](#)

31. For a process $\Delta S_{\text{total}} = 0$ and $\Delta G = 0$ was obtained. What does it mean?

 [Watch Video Solution](#)


32. If $\Delta G > 0$, then the nature of the process will be (spontaneous/non-spontaneous)?

 [Watch Video Solution](#)

33. A certain reaction is at equilibrium at 82°C and ΔH for this reaction is 21.3 kJ. What would be the value of ΔS (in $\text{JK}^{-1}\text{mol}^{-1}$) for the reaction?

 [Watch Video Solution](#)

34. Calculate the temperature above which the reaction of lead oxide to lead in the following reaction become spontaneous.



Given $\Delta H = 108.4 \text{ kJ mol}^{-1}$, $\Delta S = 190 \text{ JK}^{-1} \text{ mol}^{-1}$

 [Watch Video Solution](#)

35. What kind of system does a thermos flask containing coffee represents?

 [Watch Video Solution](#)

36. A tree represents an _____ system.

 [Watch Video Solution](#)


37. In the equation $pV = nRT$, how many state variables are present?

[▶ Watch Video Solution](#)

38. Temperature is a state function or not?

[▶ Watch Video Solution](#)

39. Consider the following figure and tell the nature of work done.



[▶ Watch Video Solution](#)

40. A beaker containing water at 373 K is representing an open system, is placed in the room. Predict the nature of work done.



[Watch Video Solution](#)

41. 180 J of work is done on the system and at the same time 120 J of heat is given out. What is the change in the internal energy?



[Watch Video Solution](#)

42. If change in internal energy -90 J. The work done by system is $+ 50$ J. Calculate the heat change.



[Watch Video Solution](#)

43. If the external pressure is more than internal pressure then what will be the sign convention for the work done on a cylinder fitted with a frictionless and weightless piston containing one mole of an ideal gas?



[Watch Video Solution](#)

44. What would be the sign convention for work done on a system for a cylinder containing one mole of an ideal gas where $P_{ext} = P_f$?

 [Watch Video Solution](#)

45. One mole of oxygen is allowed to expand isothermally and reversibly from 5 m^3 to 10 m^3 at 300 K. Calculate the work done in expansion of the gas.

 [Watch Video Solution](#)

46. How much heat would be absorbed by two litres of an ideal gas at 5 atm pressure if it expands isothermally into an evacuated flask of five litres?

 [Watch Video Solution](#)

47. The heat of combustion of naphthalene $\{C_{10}H_8(s)\}$ at constant volume was measured to be $-5133kJmol^{-1}$ at 298 K. Calculate the value of enthalpy change (Given $R = 8.314JK^{-1}mol^{-1}$)

 [Watch Video Solution](#)

48. How it is possible to find the value of ΔU and not the value of U ?

 [Watch Video Solution](#)

49. Out of pressure, mass , moles and volume, which is an intensive property?

 [Watch Video Solution](#)

50. Pick the odd one out : (Intensive or extensive) Refractive index, density, temperature and enthalpy.

 [Watch Video Solution](#)

51. Calculate the amount of heat required to raise the temperature of 5 g of iron from 25°C to 75°C . The specific heat capacity of iron is 0.45 J/g .

 [Watch Video Solution](#)

52. Calculate the weight of gold having specific heat capacity $0.13\text{ J} (\text{.}^{\circ}\text{C})^{-1}\text{g}^{-1}$ which is heated from 25°C to 300°C , where the heat absorbed is 1500 J .

 [Watch Video Solution](#)

53. 90 g of water spilled out from a vessel in the room on the floor. Assuming that water vapour behaving as an ideal gas, calculate the internal energy change when the spilled water undergoes complete evaporation at 100°C . (Given the molar enthalpy of vaporisation of water at 1 bar and $373\text{ K} = 41\text{ kJ mol}^{-1}$).




Watch Video Solution

54. One mole of carbon undergoes incomplete combustion to produce carbon monoxide, Calculate $(\Delta H - \Delta U)$ for the formation of CO at 298 K. Given $R = 8.314 \text{ JK}^{-1}\text{mol}^{-1}$.



Watch Video Solution

55. The standard heats of formation at 298K for $\text{CCl}_4(g)$, $\text{H}_2\text{O}(g)$, $\text{CO}_2(g)$ and $\text{HCl}(g)$ are -25.5 , -57.8 , -94.1 and $-22.1 \text{ kcal mol}^{-1}$ respectively. Calculate ΔH_{298}° for the reaction.



Watch Video Solution

56. An athlete is given 180 g of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$). He utilises 50% of the energy due to internal combustion in the body. In order to avoid storage


of energy in the body, calculate the mass of water he would need to perspire. Given enthalpy of combustion of glucose is $-2800 \text{ kJ mol}^{-1}$ and enthalpy of evaporation of water is 44 kJ mol^{-1}

 [Watch Video Solution](#)

57. A gobar gas plant produces methane and supplies to the families. If a family requires 15000 kJ of energy per day, how many days will the quota of 20 kg will last ($\Delta_c H \text{ of } CH_4 = -1665 \text{ kJ mol}^{-1}$).

 [Watch Video Solution](#)

58. Calculate the heat of combustion of ethene



The bond energy data are given below

$$C = C = 619 \text{ kJ mol}^{-1}$$

$$C - H = 414 \text{ kJ mol}^{-1}$$


$$O = O = 499 \text{ kJ mol}^{-1}$$

$$C = O = 724 \text{ kJ mol}^{-1}$$

$$O - H = 460 \text{ kJ mol}^{-1}$$

 [Watch Video Solution](#)

59. Calculate the enthalpy change during the reaction :



Given, $e_{H-H} = 435 \text{ kJ mol}^{-1}$, $e_{Br-Br} = 192 \text{ kJ mol}^{-1}$ and

$$e_{H-Br} = 368 \text{ kJ mol}^{-1}.$$

 [Watch Video Solution](#)

60. When 7.45 g of metal chloride dissolved in excess of water, the amount of heat absorbed is X kJ. Calculate the enthalpy of solution if molar mass of chloride is 74.5.

 [Watch Video Solution](#)

61. Enthalpy of solution for $BaCl_2 \cdot 2H_2O$ and $BaCl_2$ are 18.8 and $-30.6 kJ mol^{-1}$ respectively. Calculate the enthalpy of hydration of $BaCl_2$ to $BaCl_2 \cdot 2H_2O$.

 [Watch Video Solution](#)

62. The entropy change in the conversion of water to ice at 273 K for the system is $-21.99 JK^{-1} mol^{-1}$ and that of surrounding is $+21.99 JK^{-1} mol^{-1}$. State whether the process is spontaneous or non-spontaneous.

 [Watch Video Solution](#)

63. The entropy change in the conversion of water to ice at 274 K for the system is $-22.13 JK^{-1} mol^{-1}$ and for surrounding is $+22.05 JK^{-1} mol^{-1}$. State whether the process is spontaneous or non-spontaneous.

 [Watch Video Solution](#)

64. For a process $\Delta S_{\rightarrow tal} = 0$ and $\Delta G = 0$ was obtained. What does it mean?

 [Watch Video Solution](#)


65. If $\Delta G > 0$, then the nature of the process will be (spontaneous/non-spontaneous)?

 [Watch Video Solution](#)

66. Certain reaction is at equilibrium at $82^{\circ}C$ and ΔH for this reaction is 21.3 kJ. What would be the value of ΔS ($\in JK^{-1}mol^{-1}$) for the reaction.

 [Watch Video Solution](#)

67. Calculate the temperature above which the reduction of lead oxide to lead in the following reaction become spontaneous.



Given $\Delta H = 118.4 \text{ kJ mol}^{-1}$, $\Delta S = 200 \text{ JK}^{-1} \text{ mol}^{-1}$.

 [Watch Video Solution](#)

SECTION-A

1. Which one of the following is not an intensive property?

- A. Pressure
- B. Concentration
- C. Density
- D. Volume

Answer: D

 [Watch Video Solution](#)

2. A system in which state variables have constant values throughout the system is called in a state of

- A. Equilibrium
- B. Non-equilibrium
- C. Transition
- D. Zero entropy

Answer: A



[Watch Video Solution](#)

3. Which of the following conditions holds good for an adiabatic process ?

- A. $dq \leq 0$
- B. $dq > 0$

C. $dq = 0$

D. $dq = T$

Answer: C

 [Watch Video Solution](#)

4. A gas expands from 10 litres to 20 litres against a constant external pressure of 10 atm. The pressure-volume work done by the system is

A. 100 L atm

B. -100 L atm

C. 10 L atm

D. -10 L atm

Answer: B

 [Watch Video Solution](#)

5. Which of the following properties is not a state function ?

A. Work

B. Internal energy

C. Enthalpy

D. Entropy

Answer: A



[Watch Video Solution](#)

6. The heat capacity at constant pressure is related to heat capacity at constant volume by the relation

A. $C_p - C_v = R$

B. $C_v - R = C_p$

C. $C_v = -C_p - R$

D. $R - C_p = C_v$

Answer: A



[Watch Video Solution](#)

7. One mole of an ideal gas at 300 K is expanded isothermally from 1 litre volume to 10 litre volume. ΔU for this process is ($R = 2 \text{ cal } K^{-1} \text{ mol}^{-1}$)

A. -245.5 cal

B. 0

C. $+24 \text{ cal}$

D. -24 cal

Answer: B



[Watch Video Solution](#)

8. Calculate the work done when 1 mole of a gas expands reversibly and isothermally from 5 atm to 1 atm at 300 K. [Value of $\log 5 = 0.6989$].

A. -4015 J

B. $+4015\text{ J}$

C. Zero

D. -15 J

Answer: A

 [Watch Video Solution](#)

9. The change in enthalpy that takes place when one mole of the compound is formed from its elements is called

A. Heat of formation

B. Heat of fusion

C. Heat of combustion

D. Heat of sublimation

Answer: A

 [Watch Video Solution](#)

10. The change in enthalpy of a substance when one mole of the substance is completely burnt in excess of air is called

- A. Heat of sublimation
- B. Heat of combustion
- C. Heat of fusion
- D. Both (1) & (2)

Answer: B

 [Watch Video Solution](#)

11. Hess's law is used to determine

- A. Heat of formation of substances which are otherwise difficult to calculate

- B. Heat of transition
- C. Heat of dissociation
- D. All of these

Answer: D

 [Watch Video Solution](#)

12. A system absorbs 300 cal of heat. The work done by the system is 200 cal. ΔU for the above change is

- A. 100 cal
- B. 500 cal
- C. – 500 cal
- D. – 100 cal

Answer: A

 [Watch Video Solution](#)

13. In an isothermal process

- A. $q = 0$ and $\Delta U = 0$
- B. $q \neq 0$ and $\Delta U = 0$
- C. $q = 0$ and $\Delta U \neq 0$
- D. $q \neq 0$ and $\Delta U \neq 0$

Answer: B



[Watch Video Solution](#)

14. The work done during the expansion of a gas from a volume of 3 dm^3 against a constant external pressure of 3 atm is (1 L atm = 101.3 J)

- A. -608 J
- B. $+608 \text{ J}$
- C. -304 J

D. + 304 J

Answer: A



[Watch Video Solution](#)

15. 16 g of O_2 gas at STP is expanded so that volume is doubled. Hence, work done is

A. – 22.4 L atm

B. – 44.8 L atm

C. – 11.2 L atm

D. – 5.6 L atm

Answer: C



[Watch Video Solution](#)

16. In a process, a system does 140 J of work on the surroundings and only 40 J of heat is added to the system, hence change in internal energy is

- A. 180 J
- B. – 180 J
- C. 100 J
- D. – 100 J

Answer: D



[Watch Video Solution](#)

17. A system absorbs 800 J of heat and system works equivalent to 900 J by expanding. The value of ΔU for the system is

- A. – 100 J
- B. – 200 J
- C. – 300 J

D. -400 J

Answer: A



[Watch Video Solution](#)

18. 1 mole of an ideal gas undergoes an isothermal reversible expansion from 10 atm to 1 atm at 300 K. What will be the work done ?

A. $+57.44\text{ J}$

B. -57.44 J

C. -5744 J





D. $+5744\text{ J}$

Answer: C



[Watch Video Solution](#)

19. Which among the following represents standard enthalpy of formation?



Answer: A



Watch Video Solution

20. Which one of the following equations does not correctly represent the first law of thermodynamics for the given process?

A. Isothermal process, $q = -W$

B. Expansion of a gas in vacuum, $W = 0$

C. Adiabatic process, $\Delta U = q$

D. Process at constant volume, $\Delta U = q$

Answer: C



[Watch Video Solution](#)

21. For an isothermal process

A. $\Delta H = 0$

B. $\Delta U = 0$

C. $\Delta T = 0$

D. All of these

Answer: D



[Watch Video Solution](#)

22. Heat change of a process, in which volume remains constant, is represented by

A. ΔU

B. ΔH

C. ΔG

D. Δn

Answer: A



Watch Video Solution

23. If the total enthalpy of reactants and products is H_R and H_P respectively, then for exothermic reaction

A. $H_R = H_P$

B. $H_R < H_P$

C. $H_R > H_P$

D. $H_P \geq H_R$

Answer: C



Watch Video Solution

24. In which of the following change entropy decreases?

- A. Crystallization of sucrose from solution
- B. Dissolving sucrose in water
- C. Melting of ice
- D. Vaporization of camphor

Answer: A



Watch Video Solution

25. The value of entropy of the universe is always

- A. Constant
- B. Decreasing
- C. Increasing
- D. Zero

Answer: C

 [Watch Video Solution](#)

26. A reaction occurs spontaneously in forward direction if

- A. $T\Delta S > \Delta H$ where ΔH is positive and ΔS is negative.
- B. $T\Delta S = \Delta H$ where both ΔH and ΔS are positive
- C. $T\Delta S < \Delta H$ where both ΔH and ΔS are positive
- D. $T\Delta S > \Delta H$ where both ΔH and ΔS are positive

Answer: D

 [Watch Video Solution](#)

27. For an exothermic reaction to be spontaneous ($\Delta S = \text{negative}$)

- A. Temperature must be high
- B. Temperature must be zero
- C. Temperature may have any magnitude
- D. Temperature must be low

Answer: D



[Watch Video Solution](#)

28. The enthalpy change of a reaction does not depend on

- A. State of reactants and products
- B. Nature of reactants and products
- C. Different intermediate reactions

D. Initial and final enthalpy change of reaction

Answer: C

 [Watch Video Solution](#)

29. The free energy change for a reversible reaction at equilibrium is

A. Zero

B. Positive

C. Negative

D. Cannot say

Answer: A

 [Watch Video Solution](#)

30. In a chemical reaction, $\Delta H = 150\text{kJ}$ and $\Delta S = 100\text{JK}^{-1}$ at 300 K.

Therefore, ΔG will be

- A. Zero
- B. 300 kJ
- C. 330 kJ
- D. 120 kJ

Answer: D



[Watch Video Solution](#)

31. Pick out the correct option which represents for the work done by the system on the surroundings

- A. $p_{\text{in}} > p_{\text{ex}}$
- B. $p_{\text{in}} = p_{\text{ex}}$
- C. $p_{\text{in}} < p_{\text{ex}}$

D. $p_{\text{ex}} \geq 1 \text{ atm}$

Answer: A

 [Watch Video Solution](#)

32. A gas expands isothermally against a constant external pressure of 1 atm from a volume of 10 dm^3 to a volume of 20 dm^3 . It absorbs 800 J of thermal energy from its surroundings. The ΔU is

A. -312 J

B. $+123 \text{ J}$

C. $+312 \text{ J}$

D. -213 J

Answer: D

 [Watch Video Solution](#)

33. The free energy change, $\Delta G = 0$, when

- A. The reactants are completely consumed
- B. A catalyst is added
- C. The system is at equilibrium
- D. The reactants are initially start reacting

Answer: C



Watch Video Solution

34. If an insulated container containing liquid is stirred with a paddle to increase the temperature, which of the following is true?

- A. $\Delta U = W \neq 0, q = 0$
- B. $\Delta U = W = 0, q \neq 0$
- C. $\Delta U = 0, W = q \neq 0$
- D. $W = 0, \Delta U = q \neq 0$

Answer: A

 [Watch Video Solution](#)

35. The bond energies of $H - H$, $Br - Br$ and $H - Br$ are 433, 192 and 364 KJ mol^{-1} respectively. The ΔH° for the reaction $H_2(g) + Br_2(g) \rightarrow 2HBr(g)$ is

A. -261 kJ

B. $+103 \text{ kJ}$

C. $+261 \text{ kJ}$

D. -103 kJ

Answer: D

 [Watch Video Solution](#)

36. If one mole of ammonia and one mole of hydrogen chloride are mixed in a closed container to form ammonium chloride vapor, then

A. $\Delta H > \Delta U$

B. $\Delta H = \Delta U$

C. $\Delta H < \Delta U$

D. No relationship

Answer: C



Watch Video Solution

37. In thermodynamics, a process is called reversible when

A. Surrounding and system change into each other

B. There is no boundary between system and surrounding

C. The surrounding are always in equilibrium with the system

D. The system changes into surrounding spontaneously

Answer: C

 [Watch Video Solution](#)

38. $X(g) + 2Y(g) \rightarrow 2Z(g) + 3A(g)$ The change in enthalpy at $27^\circ C$ is 79.5 kJ. The value of ΔU is

A. 74.5 kJ

B. 4.99 kJ

C. 79.5 kJ

D. 75.9 kJ

Answer: A

 [Watch Video Solution](#)

39. In a spontaneous process the system undergoes

- A. No energy change
- B. Lowering of free energy
- C. Lowering of entropy of universe
- D. Increase in internal energy always

Answer: B

 [Watch Video Solution](#)

40. A quantity of an ideal gas at 20°C reversibly expands against a constant pressure of 2.0 atm from 1.0 L to 2.0 L. Calculate the work done

- A. -101.3 J
- B. -202.6 J
- C. -844 J
- D. -448 J

Answer: B

 [Watch Video Solution](#)

41. When the common salt dissolves in water, the entropy of the system increases. This means that the sign of ΔS is _____ and randomness of the system _____

- A. Undetermined, increases
- B. Positive, decreases
- C. Positive, increases
- D. Negative, decreases

Answer: C

 [Watch Video Solution](#)

42. Pick the extensive property from the given options

- A. Viscosity

B. Refractive index

C. Density

D. Heat capacity

Answer: D

 [Watch Video Solution](#)

43. If a process is both endothermic and spontaneous then

A. $\Delta S > 0$

B. $\Delta S < 0$

C. $\Delta H < 0$

D. $\Delta G > 0$

Answer: A

 [Watch Video Solution](#)

44. The specific heat of gas is found to be 0.075 calories at constant volume and its formula weight is 40. The atomicity of the gas would be

- A. One
- B. Two
- C. Three
- D. Four

Answer: A



[Watch Video Solution](#)

45. The standard entropies of $N_2(g)$, $H_2(g)$ and $NH_3(g)$ are 191.5, 130.5, $192.6 JK^{-1}mol^{-1}$. The value of ΔS^0 during the formation of 1 mole of ammonia is

- A. $-98.9 JK^{-1}mol^{-1}$
- B. Zero

C. $+129.4JK^{-1}mol^{-1}$

D. $-29.4JK^{-1}mol^{-1}$


Answer: A



Watch Video Solution

SECTION-B

1. Bond energies of H - H bond is 80 kJ/mol, I - I bond is 100 kJ/mol and for H - I bond is 200 kJ/mol, the enthalpy of the reaction :



A. -120 kJ

B. 200 kJ

C. $+100$ kJ

D. $+120$ kJ

Answer: B

 [Watch Video Solution](#)

2. The enthalpy of formation of $H_2O(l)$ is -280.70 kJ/mol and enthalpy of neutralisation of a strong acid and strong base is -56.70 kJ/mol. What is the enthalpy of formation of OH^- ions?

A. -22.9 kJ/mol

B. -224 kJ/mol

C. 58.7 kJ/mol

D. 214 kJ/mol

Answer: B

 [Watch Video Solution](#)

3. Heat of neutralisation of a strong dibasic acid in dilute solution by NaOH is nearly :

- A. -27.4 kcal/eq
- B. 13.7 kcal/mol
- C. -13.7 kcal/eq
- D. -13.7 kcal/mol

Answer: C



[Watch Video Solution](#)

4. The heat released in neutralisation of HCl and NaOH is 13.7 kcal/mol, the heat released on neutralisation of NaOH with CH_3COOH is 3.7 kcal/mol. The ΔH° of ionisation of CH_3COOH is

- A. 10.2 kcal
- B. 10 kcal





C. 3.7 kcal

D. 9.5 kcal

Answer: B

 [Watch Video Solution](#)

5. Which of the following reactions represents the enthalpy of formation of water?



Answer: B

 [Watch Video Solution](#)

6. The energy required to break 76 gm gaseous fluorine into free gaseous atom is 180 kcal at 25°C . The bond energy of F - F bond will be

- A. 180 kcal
- B. 90 kcal
- C. 45 kcal
- D. 104 kcal

Answer: B



[Watch Video Solution](#)

7. For the reaction $2\text{HgO}(s) \rightarrow 2\text{Hg}(l) + \text{O}_2(g)$

- A. $\Delta H > 0$ and $\Delta S < 0$
- B. $\Delta H > 0$ and $\Delta S > 0$
- C. $\Delta H < 0$ and $\Delta S < 0$
- D. $\Delta H < 0$ and $\Delta S > 0$

Answer: B

 [Watch Video Solution](#)

8. The heat of combustion of yellow phosphorus and red phosphorus are $-9.91 \text{ KJ mol}^{-1}$ and -8.78 KJ/mol respectively. The heat of transition from yellow phosphorus to red phosphorus is

A. -18.69 kJ

B. $+1.13 \text{ kJ}$

C. $+18.69 \text{ kJ}$

D. -1.13 kJ

Answer: D

 [Watch Video Solution](#)

9. If the entropy of vaporisation of a liquid is $110 \text{ JK}^{-1}\text{mol}^{-1}$ and its enthalpy of vaporisation is 50 kJ mol^{-1} . The boiling point of the liquid is

- A. 354.5 K
- B. 454.5 K
- C. 554.5 K
- D. 445.5 K

Answer: B



[Watch Video Solution](#)

10. Which of the following represents heat of formation (ΔH_f) ?

- A. $C(\text{diamond}) + O_2(g) \rightarrow CO_2(g)$
- B. $2CO(g) + O_2(g) \rightarrow 2CO_2(g)$
- C. $\frac{1}{2}H_2(g) + \frac{1}{2}F_2(g) \rightarrow HF(g)$
- D. $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$

Answer: C



Watch Video Solution

11. 1 mole of an ideal gas is expanded from an initial pressure of 1 bar to final pressure of 0.1 bar at constant temperature of 273 K. Predict which of the following is not true?

A. $\Delta E = 0$

B. $\Delta H = 0$

C. PV is constant


D. $\Delta S \leftarrow 0$

Answer: D



Watch Video Solution

12. Which of the following conditions should be satisfied for the given reaction to be spontaneous at $0^{\circ}C$ and 1 atm ?



A. $\Delta H = \Delta G$

B. $\Delta H < T\Delta S$

C. $\Delta H > T\Delta S$

D. $\Delta H = T\Delta S$

Answer: B



[Watch Video Solution](#)

13. The heat of formation of CO_2 is -407 kJ/mol. The energy required for the process $3CO_2(g) \rightarrow 3C(g) + 2O_3(g)$ is

A. Less than 1221 kJ

B. More than 1221 kJ

C. Is equal to 1221 kJ

D. Cannot be predicted

Answer: B

 [Watch Video Solution](#)

14. The free energy change due to a reaction is zero when

A. The reactants are initially mixed

B. A catalyst is added


C. The system is at equilibrium

D. The reactants are completely consumed

Answer: C

 [Watch Video Solution](#)

15. From the given graph



Which of the following statement is correct ?

- A. The point B represents the state of equilibrium
- B. The equilibrium composition strongly favours the reactant
- C. From the point B formation of product is equally spontaneous as of reactant
- D. From the point B formation of reactant is more spontaneous than that of product

Answer: A



Watch Video Solution

16. The heat of combustion of sucrose $C_{12}H_{22}O_{11}(s)$ at constant volume is $-1348.9 \text{ kcal mol}^{-1}$ at $25^\circ C$, then the heat of reaction at constant pressure, when steam is produced, is

- A. -1348.9 kcal
- B. 1342.344 kcal
- C. 1250 kcal
- D. -1250 kcal

Answer: B



Watch Video Solution

17. For the reaction, $C_7H_8(l) + 9O_2(g) \rightarrow 7CO_2(g) + 4H_2O(l)$, the calculated heat of reaction is 232 kJ mol^{-1} and observed heat of reaction is 50.4 kJ mol^{-1} , then the resonance energy is

A. $-182.2 \text{ kJ mol}^{-1}$

B. $+182.2 \text{ kJ mol}^{-1}$

C. 172 kJ mol^{-1}



D. -172 kJ mol^{-1}

Answer: A



Watch Video Solution

18. Given that ,



The heat of sublimation of A will be :

A. $x - y$

B. $x + y$

C. x or y

D. $-x + y$

Answer: B

 [Watch Video Solution](#)

19. AB , A_2 and B_2 are diatomic molecules. If the bond enthalpies of A_2 , AB and B_2 are in the ratio 1 : 1 : 0.5 and the enthalpy of formation of AB from A_2 and B_2 is -100 kJ mol^{-1} , what is the bond enthalpy of A_2 ?

A. 400 kJ mol^{-1}

B. 200 kJ mol^{-1}

C. 300 kJ mol^{-1}

D. 100 kJ mol^{-1}

Answer: A

 [Watch Video Solution](#)

20. One mole of a perfect gas expands isothermally to ten times its original volume. The change in entropy is

A. $0.1 R$

B. $10 R$

C. $2 R$

D. $2.303 R$

Answer: D



[Watch Video Solution](#)

SECTION-C

1. Which of the following is/are extensive properties?

A. Entropy

B. Density

C. Enthalpy

D. Boiling point

Answer: A::C

 [Watch Video Solution](#)

2. At 25° and 1 atm, which one(s) of the following has nonzero ΔH°_f ?

A. Fe

B. O

C. C(diamond)

D. Ne

Answer: B::C

 [Watch Video Solution](#)

3. The factors which influence the heat of reaction are

A. Physical state of reactants and products

B. The temperature of the reaction

C. The method by which the final products are obtained


D. Whether the reaction is carried out at constant pressure or constant temperature

Answer: A::B::D



Watch Video Solution

4. For the given graph, which parameters will be zero?



A. ΔQ

B. ΔH

C. ΔU

D. ΔS

Answer: B::C::D

[Watch Video Solution](#)

5. Which of the following is/are state function?

A. Enthalpy

B. Heat

C. Entropy

D. Gibb's free energy(G)

Answer: A::C::D



Watch Video Solution

6. Enthalpy of neutralization of strong acid with strong base is 13.7 kcal.

When an unknown acid (1 eq) is neutralized with (1 eq.) strong base, the enthalpy change is 10.7 kcal which of the following statements is/are correct regarding unknown acid?

A. Unknown acid is strong acid

B. Unknown acid is weak acid

C. 3.0 kcal heat utilised to dissociate the unknown acid

D. 10.7 kcal heat utilised to dissociate unknown acid

Answer: B::C

 [Watch Video Solution](#)

7. Which of the following statements is/are correct?

A. For spontaneous process, ΔG must be negative

B. At equilibrium, $\Delta G = 0$

C. At equilibrium, $\Delta G^\circ = 0$

D. At equilibrium, $\Delta G^\circ = -2.303RT \log K_P$

Answer: A::B::D

 [Watch Video Solution](#)

8. In which case of mixing of strong acid and strong base each of 1 N concentration temperature increase of solution will be same (assume

heat evolved in neutralisation is only used up to increase the temperature of solution)?

- A. 20 ml acid and 30 ml alkali
- B. 10 ml acid and 10 ml alkali
- C. 35 ml acid and 15 ml alkali
- D. 50 ml acid and 50 ml alkali

Answer: B::D



Watch Video Solution

9. A mole of steam is condensed at 100°C , the water is cooled to 0°C and frozen to i.e. . What is the difference in entropies of the stem and ice? The heat of vaporization and fusion are 540cal gm^{-1} and 80cal gm^{-1} respectively . Use the average heat capacity of liquid water as $1\text{cal gm}^{-1}\text{ degree}^{-1}$.

A. Entropy change during the condensation of steam is

$$-26.06 \text{ cal}/^\circ \text{C}$$

B. Entropy change during cooling of water from

$$100^\circ \text{C to } 0^\circ \text{C is } -5.62 \text{ cal}/^\circ \text{C}$$

C. Entropy change during freezing of water at 0°C is $-5.27 \text{ cal}/^\circ \text{C}$

D. Total entropy change is $-36.95 \text{ cal}/^\circ \text{C}$

Answer: A::B::C::D



Watch Video Solution

10. For a diatomic gas, which options is/are correct?

A. $\gamma = 1.40$

B. $C_P = \frac{7R}{2}$

C. $C_v = \frac{5R}{2}$

D. $\gamma = 1.67$

Answer: A::B::C



Watch Video Solution

11. When the gas is ideal and process is isothermal, then

A. $P_1V_1 = P_2V_2$

B. $\Delta U = 0$

C. $\Delta W = 0$

D. $\Delta H_1 = \Delta H_2$

Answer: A::B::D



Watch Video Solution

12. Which of the following relation is/are correct?

A. $\Delta G = \Delta H - T\Delta S$

$$B. \Delta G = \Delta H + T \left[\frac{\delta(\Delta G)}{\delta T} \right]_P$$

$$C. \Delta G = \Delta H + T\Delta S$$

$$D. \Delta G = \Delta H + \Delta nRT$$

Answer: A::B

 [Watch Video Solution](#)

13. Heat evolved during the combustion of 32 gm methanol is a bomb calorimeter was determined to be 470 kcal/mol at 25°C. The value of Δu of the reaction at the same temperature is

A. – 335.24 kcal

B. – 669.28kcal

C. – 470 kcal/mol

D. – 196.5×10^4 J

Answer: C::D

SECTION-D

1. Entropy is a measure of randomness of system. When a liquid is converted to the vapour state entropy of the system increases. Entropy in the phase transformation is calculated using $\Delta S = \frac{\Delta H}{T}$ but in reversible adiabatic process ΔS will be zero. The rise in temperature in isobaric or isochoric process increases the randomness of system, which is given by

$$\Delta S = 2.303 n C \log\left(\frac{T_2}{T_1}\right)$$

$$C = C_P \text{ or } C_V$$

The temperature at which liquid H_2O will be in equilibrium with its vapour is (ΔH and ΔS for vapourisation are 50 kJ mol^{-1} and $0.15 \text{ kJ mol}^{-1} \text{K}^{-1}$)

A. 77.30°C

B. 60.33°C

C. $50^{\circ}C$

D. $100^{\circ}C$

Answer: B

 [Watch Video Solution](#)

2. Entropy is a measure of randomness of system. When a liquid is converted to the vapour state entropy of the system increases. Entropy in the phase transformation is calculated using $\Delta S = \frac{\Delta H}{T}$ but in reversible adiabatic process ΔS will be zero. The rise in temperature in isobaric or isochoric process increases the randomness of system, which is given by

$$\Delta S = 2.303 n C \log\left(\frac{T_2}{T_1}\right)$$

$$C = C_P \text{ or } C_V$$

The change in entropy when 1 mole O_2 gas expands isothermally and reversibly from an initial volume 1 litre to a final volume 100 litre at $27^{\circ}C$

A. $20.5 JK^{-1} (mol^{-1})$

B. $38.29 JK^{-1} mol^{-1}$

C. $42.50 JK^{-1} mol^{-1}$

D. $50.65 JK^{-1} mol^{-1}$

Answer: B

 [Watch Video Solution](#)

3. Entropy is a measure of randomness of system. When a liquid is converted to the vapour state entropy of the system increases. Entropy in the phase transformation is calculated using $\Delta S = \frac{\Delta H}{T}$ but in reversible adiabatic process ΔS will be zero. The rise in temperature in isobaric or isochoric process increases the randomness of system, which is given by

$$\Delta S = 2.303 n C \log\left(\frac{T_2}{T_1}\right)$$

$$C = C_P \text{ or } C_V$$

Entropy change in a reversible adiabatic process is

A. Zero

B. Always positive

C. Always negative

D. Sometimes positive and sometimes negative

Answer: A

 [Watch Video Solution](#)

4. Heat of reaction is defined as the amount of heat absorbed or evolved at a given temperature when the reactants have combined to form the products is represented by a balanced chemical equation. If the heat is denoted by q then the numerical value of q depends on the manner in which the reaction is performed for the two methods of conducting chemical reactions in calorimeters.

Constant volume $W = 0$ and $q_v = \Delta E$ Bomb calorimeter

Constant pressure $W = -V \Delta P$, therefore $q_p = \Delta E + P \Delta V$

The heat capacity of a bomb calorimeter is 300 JK When 0.16 gm of

methane was burnt in this calorimeter the temperature rose by 3°C . The value of ΔU per mole will be

A. 100 kJ

B. 90 kJ

C. 900 kJ

D. 48 kJ

Answer: B



[Watch Video Solution](#)

5. Heat of reaction is defined as the amount of heat absorbed or evolved at a given temperature when the reactants have combined to form the products is represented by a balanced chemical equation. If the heat is denoted by q then the numerical value of q depends on the manner in which the reaction is performed for the two methods of conducting chemical reactions in calorimeters.

Constant volume $W = 0$ and $q_v = \Delta E$ Bomb calorimeter

Constant pressure $W = -V \Delta P$, therefore $q_P = \Delta E + P\Delta V$ (at constant pressure)

When maltose $C_{12}H_{22}O_{11}(s)$ burns in a calorimetric bomb at 298 K yielding carbon dioxide and water, the heat of combustion is -1350 kcal/mol, the heat of combustion of maltose at constant pressure will be

- A. - 2650 kcal/mol
- B. - 675 kcal/mol
- C. - 1350 kcal/mol
- D. - 1100 kcal/mol

Answer: C



[Watch Video Solution](#)

6. Heat of reaction is defined as the amount of heat absorbed or evolved at a given temperature when the reactants have combined to form the products is represented by a balanced chemical equation. If the heat is denoted by q then the numerical value of q depends on the manner in which the reaction is performed for the two methods of conducting

chemical reactions in calorimeters.

Constant volume $W = 0$ and $q_v = \Delta E$ Bomb calorimeter

Constant pressure $W = -V \Delta P$, therefore $q_P = \Delta E + P\Delta V$ or $(V \cdot \Delta P)$

The heat of combustion of maltose at constant volume in the above question if water vapour are not condensed

A. -1350 kcal/mol

B. -1343 kcal/mol

C. -1370 kcal/mol

D. -1400 kcal/mol

Answer: B



[Watch Video Solution](#)

7. In a fuel cell, methanol is used as a fuel and O_2 is used as oxidizer. The standard enthalpy of combustion of methanol is -726 kJ mol^{-1} . The standard free energies of formation of $CH_3OH(l)$, $CO_2(g)$ and $H_2O(l)$ are -166.3 , -394.4 and -237.1 kJ mol^{-1}

respectively.

The standard free energy change of the reaction will be

A. $-597.8 \text{ kJ mol}^{-1}$

B. $-298.9 \text{ kJ mol}^{-1}$

C. $-465.2 \text{ kJ mol}^{-1}$


D. $-702.3 \text{ kJ mol}^{-1}$

Answer: D



Watch Video Solution

8. In a fuel cell methanol is used as fuel and oxygen gas is used as an oxidizer. The reaction is :



At $298K$ standard Gibb's energies of formation for $CH_3OH(l)$, $H_2O(l)$ and $CO_2(g)$ are -166.2 , -237.2 and $-394.4 \text{ kJ mol}^{-1}$ respectively. If standard enthalpy of combustion of methanol is -726 kJ mol^{-1} , efficiency of the fuel cell will be :

A. 96.7 %

B. 66.2 %

C. 41.3 %

D. 85.1 %

Answer: A



Watch Video Solution

9. In a fuel cell, methanol is used as a fuel and O_2 is used as oxidizer. The standard enthalpy of combustion of methanol is -726 kJ mol^{-1} . The standard free energies of formation of $CH_3OH(l)$, $CO_2(g)$ and $H_2O(l)$ are -166.3 , -394.4 and $-237.1 \text{ kJ mol}^{-1}$ respectively.

The standard free energy change of the reaction will be

A. $-727.24 \text{ kJ mol}^{-1}$

B. $-724.76 \text{ kJ mol}^{-1}$

C. $-728.48 \text{ kJ mol}^{-1}$

D. $-723.42 \text{ kJ mol}^{-1}$

Answer: B

 [Watch Video Solution](#)

SECTION-E

1. Statement-1 : Chemical equation $H^+(aq) + OH^-(aq) \rightarrow H_2O(l), \Delta H = -X \text{ kJ}$ represents the enthalpy of formation of water

And

Statement-2 : $\Delta H_{\text{neutralisation}}$ of strong acid and strong base is constant.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: D

 [Watch Video Solution](#)

2. Statement-1 : In a cyclic process, both ΔH and ΔU are zero.

And

Statement-2 : ΔH and ΔU are path dependent functions.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: C

 [Watch Video Solution](#)

3. Statement-1 : Enthalpy (H) and entropy (S) both are extensive properties.

And

Statement-2 : Both are independent of mass.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: C



[Watch Video Solution](#)

4. Statement-1 : Work done in isothermal reversible process is more than irreversible process.

And

Statement-2 : Irreversible process is an infinitesimally slow process.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: C



[Watch Video Solution](#)

5. Statement-1 : ΔH_f° is zero for oxygen (O_2).

And

Statement-2 : ΔH_f° for all the elements at S.T.P. is zero.


- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: C



Watch Video Solution

6. Statement-1 : ΔH and ΔU are the same for the reaction



And

Statement-2 : All reactants and products are in gaseous form.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: B



Watch Video Solution

7. Statement-1 : All the exothermic reactions are spontaneous.

And

Statement-2 : For a spontaneous reaction, ΔG must be negative.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: D



[Watch Video Solution](#)

8. Statement-1 : Thermochemistry involves the calculation of heat change through balanced reactions.

And

Statement-2 : Enthalpy is a state function.

- A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1
- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: B



[Watch Video Solution](#)

9. Statement-1 : Entropy of egg increases after boiling.

And

Statement-2 : As boiling causes the denaturation of proteins.

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1

C. Statement-1 is True, Statement-2 is False

D. Statement-1 is False, Statement-2 is True

Answer: A



[Watch Video Solution](#)

10. Statement-1 : If same amounts are considered CO_2 requires more heat than O_2 , for the same rise in temperature.

And

Statement-2 : CO_2 being triatomic has higher heat capacity than O_2

A. Statement-1 is True, Statement-2 is True, Statement-2 is a correct explanation for Statement-1

- B. Statement-1 is True, Statement-2 is True, Statement-2 is Not a correct explanation for Statement-1
- C. Statement-1 is True, Statement-2 is False
- D. Statement-1 is False, Statement-2 is True

Answer: A

 [Watch Video Solution](#)

SECTION-F

1. Match the following

| Column-I | Column-II |
|------------------------|---------------------------------------|
| (A) Isothermal process | (p) Temperature changes |
| (B) Isobaric process | (q) Pressure changes |
| (C) Isochoric process | (r) Volume change |
| (D) Cyclic process | (s) Initial and final states are same |

 [Watch Video Solution](#)

2. Match the following

Column-I

(Enthalpy change in kcal)

(A) < 13.7 kcal






(B) $= 13.7$ kcal

(C) $gt 13.7$ kcal

(D) $= 27.4$ kcal

Column-II

(Neutralisation)



Assume heat of neutralisation of strong acid with strong base is 13.7 kcal.



Watch Video Solution

3. Match the following

Column-I

Column-II

(A) Enthalpy (p) Extensive property

(B) Entropy (q) $H - TS$

(C) Free energy (r) $E + PV$

(D) Pressure (s) Intensive property



Watch Video Solution

1. The maximum work done when pressure of n moles of H_2 was reduced from 20 atm to 1 atm at constant temperature of 273 K is found to be 8180 calories. What is the value of n ?

 [Watch Video Solution](#)

2. One mole of an ideal monoatomic gas expands reversibly and adiabatically from a volume of x litre to 14 litre at 27°C . Then value of x will be [Given, final temperature 189 K and $C_V = 3/2 R$].

 [Watch Video Solution](#)

3. For a reversible reaction $A \rightleftharpoons B$. Find $\frac{\log_{10} K}{10}$ at 2727°C temperature

Given


$$\Delta_r H^0 = -54.07 \text{ kJ mol}^{-1}$$

$$\Delta_r S^0 = 10 \text{ JK}^{-1}$$

$$R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$$

[▶ Watch Video Solution](#)

4. When a system is taken from A to C through path ABC, 10 J of heat flows to the system and 4 J of work is done by the system.



How much heat flows into the system in path ADC, if the work done by the system is 3 J?

[▶ Watch Video Solution](#)

5. The chemical reaction : $A \rightarrow P$, $\Delta H^\circ = 2.8 \text{ kJ}$ is spontaneous only above 400 K. Therefore ΔS of reaction must be at least (JK^{-1}).



Watch Video Solution

SECTION-H

1. Statement-1 : Neutralisation reaction is an Endothermic Process.

Statement-2 : Standard Enthalpy of Neutralisation for different pairs of strong acid and strong base are different.

Statement-3 : Standard Enthalpy of Neutralisation for a pair of strong acid and strong base is higher than that of weak acid and weak base.

A. F F T

B. F T T

C. T F T

D. F T F

Answer: A



[Watch Video Solution](#)

2. Statement-1 : Entropy of all elements is zero at zero Kelvin.

Statement-2 : Standard entropy of all elements is greater than zero.

Statement-3 : Entropy of all elements and compounds decreases with decrease of temperature and becomes zero at absolute zero temperature.

A. F F F

B. T T T

C. T F F

D. F T F

Answer: B



[Watch Video Solution](#)

3. Statement-1 : The third law of thermodynamics implies that absolute zero cannot be reached.

Statement-2 : ΔG° for an ideal gas reaction is a function of temperature.

Statement-3 : The adiabatic expansion of a gas into a vacuum is spontaneous.

A. F F F

B. T T T

C. T F F

D. F T F


Answer: B



[Watch Video Solution](#)

SECTION-I

1. Calculate enthalpy change of the following reaction :




The bond energy of C - H, C - C, C = C, H - H are 414, 615 and 436 kJ mol⁻¹ respectively.

 [Watch Video Solution](#)

2. At 300 K, 4 gm calcium is dissolved in hydrochloric acid in an open vessel at the atmosphere pressure 0.821 atm. Calculate work done by the system.

 [Watch Video Solution](#)

3. Heat of neutralisation of oxalic acid is -53.35 kJ/g.eq. using NaOH. Calculate the value of ΔH for the given reaction






 [Watch Video Solution](#)

4. If x gm of steam at 100°C is mixed with $5x$ gm of ice at 0°C , calculate the final temperature of resulting mixture. The heat of vapourisation and fusion are 540 cal gm^{-1} and 80 cal gm^{-1} respectively.

 [Watch Video Solution](#)

5. Calculate ΔH°_f for chloride ion from the following data :



 [Watch Video Solution](#)

6. The standard molar enthalpies of formation of cyclohexane (l) and benzene(l) at 25°C are -156 and $+49\text{KJmol}^{-1}$ respectively. The standard enthalpy of hydrogenation of cyclohexene (l) at 25° is -119KJmol^{-1} . Use these data to estimate the magnitude of the resonance energy of benzene.



[Watch Video Solution](#)

7. The specific heat at constant volume for a gas is 0.075 cal/g and at constant pressure is 0.125 cal/g . Calculate

- (i) Molecular weight of gas
- (ii) Atomicity of gas
- (iii) Number of atoms of gas in its 1 mole



[Watch Video Solution](#)

8. The enthalpy change involved in the oxidation of glucose is $-2880 \text{ kJ mol}^{-1}$. 25% of this energy is available for muscular work. If 100 kJ of muscular work is needed to walk one kilometer, what is the maximum distance that a person will be able to walk after consuming 120 gm of glucose?



[Watch Video Solution](#)

9. Calculate standard heat of formation of CS_2 . Given that standard heat of combustion of C, S and CS_2 are $-393.3, -293.72$ and $-1108.76 kJmol^{-1}$.




[Watch Video Solution](#)

SECTION-J

1. A heat engine carries one mole of an ideal monoatomic gas around the cycle as shown in the figure, the amount of heat added in the process AB

and heat removed in the process CA are :



- A. $q_{AB} = 750R$ and $q_{CA} = -450R$
- B. $q_{AB} = 650R$ and $q_{CA} = -350R$
- C. $q_{AB} = 750R$ and $q_{CA} = -350R$
- D. $q_{AB} = 650R$ and $q_{CA} = -450R$

Answer: C



Watch Video Solution

2. Calculate the average molar heat capacity at constant volume of a mixture containing 2 moles of monoatomic and 3 moles of diatomic ideal gas.

- A. R
- B. $2.1 R$
- C. $3.2 R$
- D. $4 R$

Answer: B



[Watch Video Solution](#)

3. What is the change in entropy when 2.5 mole of water is heated from $27^{\circ}C$ to $87^{\circ}C$?

Assume that the heat capacity is constant

$$(C_p)_m(H_2O) = 4.2J/g = k, \ln(1.2) = 0.18)$$

- A. $16.4 JK^{-1}$

B. 34.02 JK^{-1}

C. 2.89 JK^{-1}

D. 18.2 JK^{-1}

Answer: B

 [Watch Video Solution](#)

4. Which amount the following is most soluble in water?

A. Compound $\Delta_H(\text{hydration}) (\text{KJmol}^{-1})$ $\Delta H_{\text{lattice}} (\text{K})$
A -400 $+500$

B. Compound $\Delta_H(\text{hydration}) (\text{KJmol}^{-1})$ $\Delta H_{\text{lattice}} (\text{K})$
B -300 $+650$

C. Compound $\Delta_H(\text{hydration}) (\text{KJmol}^{-1})$ $\Delta H_{\text{lattice}} (\text{K})$
C $+200$ $+150$

D. Compound $\Delta_H(\text{hydration}) (\text{KJmol}^{-1})$ $\Delta H_{\text{lattice}} (\text{K})$
D -100 $+250$

Answer: C

 [Watch Video Solution](#)

5. A hungry man weighing 80 kg take quickly 20 g lunck, and then climbs up a mountain making it to a height of 200 m. If 60% of food energy was wasted as heat and the rest was used as climbing work. The fuel intake could have been any one of the following with given enthalpy of combustion?

- A. Glucose 16 kJ/g
- B. Wheat bread 20 kJ/g
- C. Fructose syrup 13 kg/g
- D. Olive oil 35 kg/g

Answer: B

 [Watch Video Solution](#)


6. According to second law of thermodynamics

- A. Heat can't flow spontaneously from a reservoir at lower temperature to a reservoir at higher temperature
- B. All spontaneous process leads to increase in entropy of universe
- C. Melting a solid increase entropy, therefore a spontaneous process
- D. All of these

Answer: D

 [Watch Video Solution](#)

7. Consider the reaction



The signs of ΔS and ΔG respectively are

- A. +, -, -
- B. -, +, +
- C. -, -, -
- D. -, +, -

Answer: D



[Watch Video Solution](#)

Exercise

1. Which of the following is an example of open system?

- A. Human body
- B. The earth
- C. Pond
- D. All of these

Answer: D



[Watch Video Solution](#)

2. Which of the following is not a state function?

A. H

B. U

C. q

D. G

Answer: C



Watch Video Solution

3. Which of the following is intensive property?

A. Temperature

B. Heat

C. Mass

D. Volume

Answer: A



Watch Video Solution

4. Closed system can exchange (with surroundings)

A. Heat

B. Matter

C. Heat and matter

D. Neither heat nor matter

Answer: A



[Watch Video Solution](#)

5. System and surroundings together constitute

A. The earth

B. The solar system

C. Galaxy

D. Universe

Answer: D



[Watch Video Solution](#)

6. Which of the following can be used to define the state of a system?

A. Pressure

B. Volume

C. Number of mole

D. All of these

Answer: D



[Watch Video Solution](#)

7. Which of the following is true about isolated system?

A. Heat passes into or out of the system

B. Matter enters or leaves the system

C. Work can be done on the system

D. All are correct

Answer: C



[Watch Video Solution](#)

8. Thermodynamics deals with

A. Macroscopic properties

B. Microscopic properties

C. Both Macroscopic & microscopic properties

D. Neither macroscopic nor microscopic properties

Answer: A



[Watch Video Solution](#)

9. The thermodynamic process in which temperature remains constant is

- A. Adiabatic process
- B. Isochoric process
- C. Isobaric process
- D. Isothermal process

Answer: D



[Watch Video Solution](#)

10. Which of the following is a state function?

- A. $q + w$
- B. Entropy
- C. Free Energy

D. All of these

Answer: D



[Watch Video Solution](#)

11. ΔE and ΔH both are zero in

A. Isobaric process

B. Adiabatic process

C. Isochoric process

D. Cyclic process

Answer: D



[Watch Video Solution](#)

12. The work done in open vessel at 300 K, when 56 g iron reacts with dilute HCl is

- A. 600 cal
- B. 300 cal
- C. 150 cal
- D. 100 cal

Answer: A



[Watch Video Solution](#)

13. Work done in reversible isothermal expansion is given by

- A. $-2.303RT \frac{\log V_2}{V_1}$
- B. $n \frac{R}{\lambda - 1} (T_2 - T_1)$
- C. $2.303RT \frac{\log V_2}{V_1}$
- D. None of these

Answer: A

 [Watch Video Solution](#)

14. When a gas undergoes adiabatic expansion, it gets cooler. This is because

- A. It is an exothermic process
- B. It is an endothermic process
- C. Internal energy of the system decreases
- D. Ideal gas becomes a real gas

Answer: C

 [Watch Video Solution](#)

15. Which of the following is correct expression for 1st law of thermodynamics under adiabatic condition?

A. $\Delta U = Q + W$

B. $\Delta U = Q - W$

C. $Q = -W$

D. $\Delta U = W$

Answer: D



[Watch Video Solution](#)

16. Expansion of a perfect gas into vacuum is related with

A. $\Delta H = 0$

B. $q = 0$

C. $W = 0$




D. All of these

Answer: D



[Watch Video Solution](#)

17. The following reactions carried in open vessel. The reaction for which $\Delta H = \Delta U$ will be



Answer: D

 [Watch Video Solution](#)

18. For the reaction $\frac{1}{8}S_8(s) + \frac{3}{2}O_2(g) \rightarrow SO_3(g)$, the difference of heat change at constant pressure and constant volume at $27^\circ C$ will be.

A. $+150R$

B. $-150R$

C. $+450R$

D. $-450R$

Answer: B



[Watch Video Solution](#)

19. Choose the correct statement(s)

A. Work done by the system is taken as negative

B. In expansion , there is work done by the system

C. $W_{\text{reversible}} > W_{\text{irreversible}}$

D. All of these

Answer: D



[Watch Video Solution](#)

20. Under which of the following condition $\Delta H = \Delta E = ?$

- A. When a process is carried out in a closed vessel
- B. When reaction involves only solids and liquids
- C. When $\Delta n_g = 0$
- D. All of these

Answer: D



[Watch Video Solution](#)

21. What is the change in internal energy if 10 J of heat is given to system at constant pressure and 20 J of work is done by the system?

- A. 10 J
- B. 30 J
- C. $-10J$
- D. $-30J$

Answer: C

 [Watch Video Solution](#)

22. For isothermal expansion , which is true?

A. $\Delta P = 0$

B. $\Delta S = 0$

C. $\Delta G = 0$

D. $\Delta U = 0$

Answer: D

 [Watch Video Solution](#)

23. For H_2 gas, $C_p - C_v = a$ and for O_2 gas $C_p - C_v = b$, so the relation between a and b is

A. $a = 16b$

B. $16a = b$

C. $a = 4b$

D. $a = b$

Answer: D

 [Watch Video Solution](#)

24. 550kJ cycle^{-1} work is done by 1 mol of an ideal gas in a cyclic process.

The amount of heat absorbed by the system in one cycle is

A. 550 kJ

B. -550kJ

C. 1100 kJ

D. Zero

Answer: A

 [Watch Video Solution](#)

25. An ideal gas undergoing expansion in vacuum shows

A. $\Delta E = 0$

B. $w = 0$

C. $q = 0$

D. All of these

Answer: D

 [Watch Video Solution](#)

26. 1 g H_2 gas expand at STP to occupy double of its original volume. The work done during the process is

A. 260 cal

B. 180 cal

C. 130 cal

D. 270 cal

Answer: D

 [Watch Video Solution](#)

27. An ideal gas is kept in a 5 litre cylinder at a pressure of 15 atm. In 30 minutes the gas is allowed to enter slowly into an evacuated vessel of 550 L capacity. Total work done during the process of energy is

A. 2500 L atm

B. 7500 L atm

C. 7425 L atm

D. 0

Answer: D

 [Watch Video Solution](#)

28. An ideal gas is allowed to expand from 5 L to 15 L once rapidly and once very slowly. The magnitude of work done in two processes are W_1 and W_2 , they are related as

A. $W_1 = W_2$

B. $W_1 > W_2$

C. $W_1 < W_2$

D. Data is insufficient

Answer: C



Watch Video Solution

29. Correct statement about *heat* and *w* or *k* is that

A. Heat appears at surface only and work in bulk

B. Heat appears in bulk and work only at surface

C. Both heat and work appear in bulk of sysem

D. Both heat and work appear on surface only

Answer: D

 [View Text Solution](#)

30. For one mole of a diatomic gas γ is equal to

A. $\frac{3}{5}$

B. $\frac{5}{3}$




C. $\frac{7}{5}$

D. $\frac{8}{6}$

Answer: C

 [Watch Video Solution](#)

31. The enthalpy of reaction is maximum in which of the following reaction?




D. All have same value

Answer: B



Watch Video Solution



32.

On the basis of Hess's law of constant heat summation, choose the correct relation:

A. $b = a - cd + c$

B. $c = a - b + d$



C. $d = a + b + c$

D. $a - c = b + d$

Answer: D

 [Watch Video Solution](#)

33. In which of the following $\Delta H = \Delta U$?




Answer: B



Watch Video Solution

34. Heat of combustion of CH_4 , C_2H_4 , C_2H_6 are -890 , -1411 and $-1560 kJ/mol$ respectively. Which has the lowest calorific fuel value ?



D. All of these

Answer: B

 [Watch Video Solution](#)

35. The heats of neutralisation of four acids A, B, C and D are $-13.7, -9.4, -11.2$ and -12.4 kcal respectively, when they are neutralised by a common base. The acidic character obeys the order :

A. $P > Q > R > S$

B. $P > S > R > Q$

C. $S > R > Q > P$

D. $S > Q > R > P$

Answer: B

 [Watch Video Solution](#)

36. Heat of neutralisation of NaOH and HCl is -57.46kJ / equivalent.

The heat of ionisation of water in kJ/mol is :

A. -57.46

B. $+57.46$

C. -114.92

D. $+114.92$

Answer: B



[Watch Video Solution](#)

37. Bond energy of $\text{N} - \text{H}$, $\text{H} - \text{H}$ and $\text{N} \equiv \text{N}$ are a , b , c respectively.

The Δh for the reaction,

$2\text{NH}_3 \rightarrow \text{N}_2 + 3\text{H}_2$ is :

A. $6a - 3b - c$

B. $6a + 3b + c$


C. $a + 6b - c$

D. $6a + b - 3c$

Answer: A

 [Watch Video Solution](#)

38. At 27°C , the combustion of ethane takes place according to the



$\Delta E - \Delta H$ for this reaction at 27°C will be

A. $+1347.1\text{J}$

B. -1247.1J

C. 6235.5J

D. $+6235.5\text{J}$

Answer: D

 [Watch Video Solution](#)

39. For the reaction



If the $\Delta H_f^\circ PCl_3$ is $-ykJ$, what is $\Delta H_f^\circ PCl_5$?


- A. $(x - y)kJ$
- B. $(y - x)kJ$
- C. $-(x + y)kJ$
- D. $(x + y)kJ$



Answer: C



Watch Video Solution

40. Heat of which of the following reaction gives enthalpy of formation directly?





D. All of these

Answer: C

 [Watch Video Solution](#)

41. When ΔH and $T\Delta S$ both are negative , then for spontaneous process which option is true?

A. $\Delta H > T\delta S$

B. $\Delta H < T\delta S$

C. $\Delta H = T\delta S$

D. $\Delta H = \Delta U$

Answer: A

 [Watch Video Solution](#)

42. Which of the following relation is false?

A. $\Delta G = \Delta H - T\Delta S$

B. $\Delta S = q + T$

C. $\Delta H = \Delta U + \Delta n_g RT$

D. $\Delta H = \Delta U + P\Delta V$

Answer: B



Watch Video Solution

43. In a reversible process, the value of $\Delta S_{sys} + \Delta S_{surr}$ is

A. > 0

B. < 0

C. $= 0$

D. All of these

Answer: C

 [Watch Video Solution](#)

44. A boiled egg shows a/an In entropy

- A. Increase
- B. Decrease
- C. No change
- D. None of these

Answer: A

 [Watch Video Solution](#)

45. The ΔG in the process of melting of Ice at $-15^{\circ}C$ is

- A. $\Delta G < 0$

B. $\Delta G > 0$

C. $\Delta G = 0$

D. All of these

Answer: B

 [Watch Video Solution](#)

46. The spontaneous nature of a reaction is impossible if

A. $\Delta H = +ve, \Delta S = +ve$

B. $\Delta H = -ve, \Delta S = -ve$

C. $\Delta H = -ve, \Delta S = +ve$

D. $\Delta H = +ve, \Delta S = -ve$

Answer: D

 [Watch Video Solution](#)

47. ΔG° of reversible reaction at its equilibrium is

- A. Positive
- B. Negative
- C. Always zero
- D. May be (1) or (2)

Answer: D

 [Watch Video Solution](#)

48. Latent heat of fusion of ice is $0.333kJg^{-1}$. The increase in entropy when 1 mole water melts at $0^\circ C$ will be

- A. $21.96kJ^{-1}mol^{-1}$
- B. $21.98kcalK^{-1}mol^{-1}$
- C. $21.96JK^{-1}mol^{-1}$
- D. $21.98calK^{-1}mol^{-1}$

Answer: C

 [Watch Video Solution](#)

49. The value for ΔH_{vap} and ΔS_{vap} for ethanol are respectively $38.594 \text{ kJ mol}^{-1}$ and 109.8 JK^{-1} . The boiling point of ethanol will be

- A. 78.5 K
- B. 431.5 K
- C. 351.5 K
- D. 331.5 K

Answer: C

 [Watch Video Solution](#)

50. 3RD LAW OF THERMODYNAMICS

- A. Entropy change
- B. Enthalpy
- C. Free Energy
- D. Absolute entropy

Answer: D

 [Watch Video Solution](#)

ASSIGNMENT (Section - A) Objective Type Questions

1. Tea placed in thermos flask is an example of

- A. Open system
- B. Close system
- C. Isolated system
- D. It can't act as system

Answer: C



[Watch Video Solution](#)

2. Gaseous system is placed with pressure P_1 , volume V_1 and temperature T_1 , it has undergone thermodynamic changes where temperature is remaining constant, it is

- A. Adiabatic process
- B. Isothermal process
- C. Isobaric process
- D. Isochoric process

Answer: B



[Watch Video Solution](#)

3. The respective examples of extensive and intensive properties are

A. Enthalpy, Entropy

B. Entropy, Enthalpy

C. Entropy, Temperature

D. Temperature, Entropy

Answer: C

 [Watch Video Solution](#)

4. A thermally isolated gaseous system can exchange energy with the surroundings. The mode of energy may be

A. Heat

B. Work

C. Heat and radiation

D. Internal energy

Answer: B

 [Watch Video Solution](#)

5. Which of the following is a state function?

- A. q
- B. Work
- C. $q + w$
- D. All of these

Answer: C

 [Watch Video Solution](#)

6. For the reaction $PCI_5(g) \rightarrow PCI_3(g) + CI_2(g)$

- A. $\Delta H = \Delta E$
- B. $\Delta H > \Delta E$
- C. $\Delta H < \Delta E$

D. Can't predicted

Answer: B



[Watch Video Solution](#)

7. If ' r ' is the work done on the system and ' s ' is heat evolved by the system then,

A. $\Delta E = r + s$

B. $\Delta E = r - s$

C. $\Delta E = r$


D. $\Delta E = s$

Answer: A



[Watch Video Solution](#)

8. For the reaction



A. $\Delta H - \Delta E = (b - d)RT$

B. $\Delta H - \Delta E = (c - b)RT$

C. $\Delta H - \Delta E = (a + b) - (c + d)RT$

D. $\Delta H - \Delta E = (a - d)RT$

Answer: B



[Watch Video Solution](#)

9. A system absorbs 10 kJ of heat and does 4 kJ of work. The internal energy of the system

A. Decrease by 6 kJ

B. Increase by 6 kJ

C. Decrease by 14 kJ

D. Increase by 14 kJ

Answer: B

 [Watch Video Solution](#)

10. In a reaction, all reactant and products are liquid, then

A. $\Delta H > \Delta E$

B. $\Delta H < \Delta E$

C. $\Delta H = \Delta E$

D. Can't predicted

Answer: C

 [Watch Video Solution](#)

11. Regarding the internal energy of the molecule, which of the following statement is correct?



- A. Its absolute value can be successfully calculated.
- B. Its absolute value cannot be determined.
- C. It is the sum of vibrational and rotational energies.
- D. Both (1) & (3)

Answer: B



[Watch Video Solution](#)

12. The heat of transition (ΔH_t) of graphite into diamond would be, where



- A. $x_1 + x_2$

B. $x_2 - x_1$

C. $x_1 - x_2$

D. x_1x_2

Answer: B



Watch Video Solution

13. For the given reactions, $A \rightarrow D$, $\Delta H = x$. Steps involved are

$$A \rightarrow B, \Delta H_1 = x_1$$

$$B \rightarrow C, \Delta H_2 = ?$$

$$C \rightarrow D, \Delta H_3 = x_3$$

A. $\Delta H_2 = x - (x_1 + x_3)$

B. $\Delta H_2 = x + x_1 + x_3$

C. $\Delta H_2 = x_1 - x_3 - x$

D. $\Delta H_2 = (x_1 + x) - x_3$

Answer: A



Watch Video Solution

14. The heats of combustion of yellow P and red P are $-9.91kJ$ and $-8.78kJ$ respectively. The heat of transition of yellow to red phosphorus is

A. $-18.69kJ$

B. $+1.13kJ$

C. $+18.69kJ$

D. $-1.13kJ$

Answer: D



Watch Video Solution

15. If the heat of formation of NO_2 is 'x' $\left[\frac{1}{2}N_2(g) + O_2(g) \rightarrow NO_2(g) \right]$
the heat of reaction $N_2(g) + O_2(g) \rightarrow 2NO_g$ is y and the heat of
reaction $2NO_g + O_2(g) \rightarrow 2NO_2(g)$ is z, then

A. $2x + z = y$

B. $2y + z = x$

C. $2x - z = y$

D. $2z + x = y$

Answer: C



Watch Video Solution

16. In the reactions



A. $x = y$

B. $x = 2y$

C. $x = \frac{y}{2}$

D. $x = \sqrt{y}$

Answer: C



Watch Video Solution

17. $\Delta H_1 C_2 H_4 = 12.5 kcal$

Heat of atomisation of C = 171 Kcal

Bond energy of $H_2 = 104.3 kcal$

Bond energy C - H = 99.3 kcal

What is C = C bond energy?

A. $140.9 kcal$

B. $49 kcal$


C. $40 kcal$

D. $76 kcal$

Answer: A

 [Watch Video Solution](#)

18. The difference between ΔH and ΔE for the reaction



A. $-7.43kJ$


B. $+3.72kJ$

C. $-3.72kJ$

D. $+7.43kJ$

Answer: A

 [Watch Video Solution](#)





The data can predict that -

- A. Rhombic sulphur is yellow in colour
- B. Monoclinic sulphur has metallic lustre
- C. Monoclinic sulphur is more stable
- D. $\Delta H_{Transition} \text{ of } S(R) \rightarrow S(M)$ is endothermic process

Answer: D

 [Watch Video Solution](#)



the enthalpy of formation of H_2SO_4 at 298 K will be

A. -814.4 kJ

B. -650.3 kJ

C. $-320.5kJ$

D. $-433.5kJ$

Answer: A



[Watch Video Solution](#)

21. The volume of a gas expands by $0.25m^3$ at a constant pressure of 10^3Nm^2 . The work done is equal to

A. $2.5erg$

B. $250J$

C. $250wa$

D. $250 \neq w \rightarrow n$

Answer: B



[Watch Video Solution](#)

22. When 1 g of anhydrous oxalic acid is burnt at $25^{\circ}C$, the amount of heat liberated is $2.835kJ$. ΔH combustion is (oxalic acid : $C_2H_2O_4$)

A. $-255.15kJ$

B. $-445.65kJ$

C. $-295.24kJ$

D. $-155.16kJ$

Answer: A



Watch Video Solution

23. The heat of neutralization of LiOH and HCl at $25^{\circ}C$ is $34.868kJmol^{-1}$

. The heat of ionisation of LiOH will be

A. $44.674kJ$

B. $22.232kJ$

C. $32.684kJ$

D. 96.464kJ

Answer: B

 [Watch Video Solution](#)

24. Which compound will absorb the maximum amount of heat when dissolved in the same amount of water ? (Integral heats of solution at 25°C in kcal/mol of each solute are given in brackets)

A. $\text{HCl}(\Delta H = -17.74)$

B. $\text{HNO}_3(\Delta H = -7.85)$

C. $\text{NH}_4\text{NO}_3(\Delta H = +16.8)$

D. $\text{NaCl}(\Delta H = +1.02)$

Answer: C

 [Watch Video Solution](#)

25. $HA + OH \rightarrow H_2O + A^{-} + q_1 kJ$ $H^{+} + OH^{-} \rightarrow H_2O + q_2 kJ$ the

enthalpy of ionisation of HA is

- A. $(q_1 + q_2)$
- B. $(q_1 - q_2)$
- C. $(q_2 - q_1)$
- D. $-(q_1 + q_2)$

Answer: C



[Watch Video Solution](#)

26. An athlete takes 100 g of glucose of energy equivalent to 1560 kJ. How much amount of energy is uptaken by 1 g molecule of glucose?


- A. $15.6 kJ$
- B. $2808 kJ$
- C. $1560 kJ$

D. $28.08kJ$

Answer: B

 [Watch Video Solution](#)

27. For the reaction



Which of the following is true ?

A. $-936.9 = \Delta E - (2 \times 10^3 \times 298 \times 3)kcal$

B. $+936.9 = \Delta E + (2 \times 10^3 \times 298 \times 3)kcal$

C. $-936.9 = \Delta E - (2 \times 10^3 \times 298 \times 2)kcal$

D. $-936.9 = \Delta E + (2 \times 10^3 \times 298 \times 2)kcal$

Answer: A

 [Watch Video Solution](#)

28. For strong acid strong base neutralisation energy for 1 mole H_2O formation is $-57.1kJ$. If 0.25 mole of strong monoprotic acid is reacted with 0.5 mole of strong base then enthalpy of neutralisation is

A. $-(0.25 \times 57.1)$

B. 0.5×57.1

C. 57.1

D. $-(0.5 \times 57.1)$

Answer: A



[Watch Video Solution](#)

29. The heat of combustion of solid benzoic acid at constant volume is $-321.3kJ$ at 27° degree Celsius. The heat of combustion at constant pressure is

A. $-321.3 - 300R$

B. $-321.30 + 300R$

C. $-321.3 - 150R$


D. $-321.3 + 900R$

Answer: C



Watch Video Solution

30.



Latent heat of vaporization of liquid water into water vapour = x_4 then

ΔH_f (heat of formation of liquid water) is:

A. $x_1 + \frac{x_2}{2} - x_3 + x_4$

B. $2x_3 - x_1 - \frac{x_2}{2} - x_4$

C. $x_1 + \frac{x_2}{2} - 2x_3 - x_4$

D. $x_1 + \frac{x_2}{2} - 2x_3 + x_4$

Answer: D



Watch Video Solution



Watch Video Solution

31. A cylinder contains either ethylene or propylene 12 ml of gas required 54 ml of oxygen for complete combustion. The gas is

- A. Ethylene
- B. Propylene
- C. 1 : 1 mixture of two gases
- D. 1 : 2 mixture

Answer: B



Watch Video Solution

32. The specific heat of gas is found to be 0.075 calories at constant volume and its formula weight is 40. The atomicity of the gas would be

- A. One

B. Two

C. Three

D. Four

Answer: A



[Watch Video Solution](#)

33. $H(g) + O(g) \rightarrow O - H(g)$, ΔH for this reaction is

A. Heat of formation of O - H

B. Bond energy of O - H

C. Heat of combustion of H_2

D. Zero at all temperatures

Answer: B



[Watch Video Solution](#)

34. Energy required to dissociate 4 g of gaseous H_2 into free gaseous atoms is 872 kJ at $25^\circ C$. The bond energy of H-H bond will be

A. 8.72kJ

B. 4.36kJ

C. 436kJ

D. 43.6kJ

Answer: C



Watch Video Solution

35. The dissociation energy of $CH_4(g)$ is $360 \text{ kcal mol}^{-1}$ and that of $C_2H_6(g)$ is $620 \text{ kcal mol}^{-1}$. The C-C bond energy

A. $260 \text{ kcal mol}^{-1}$

B. $180 \text{ kcal mol}^{-1}$

C. $130 \text{ kcal mol}^{-1}$

D. 80kcalmol^{-1}

Answer: D

 [Watch Video Solution](#)

36. The enthalpy of reaction, $2\text{HC} = \text{CH} + 5\text{O}_2 \rightarrow 4\text{CO}_2 + 2\text{H}_2\text{O}$

If the bond energies of $\text{C} - \text{H}$, $\text{C} = \text{C}$, $\text{O} = \text{O}$, $\text{C} = \text{O}$ and $\text{O} - \text{H}$ bonds are p, q, r, s, t respectively

A. $[8s + 4t] - [4p + q + 5r]$

B. $[4p + 2q + 5r] - [8s + 4t]$


C. $[4p + 2q + 5r + 8s + 4t]$

D. $[2p + q + 5r] - [8s + 4t]$

Answer: B

 [Watch Video Solution](#)

37. Calculate heat of formation of isoprene using bond energy data.



$$\text{Given } C - H = 98.8 \text{ kcal}$$

$$H - H = 104 \text{ kcal}$$

$$C - C = 83 \text{ kcal}$$

$$C = C = 147 \text{ kcal and } C(s) \rightarrow C(g) = 171 \text{ kcal}$$

A. -21 kcal

B. 21 kcal

C. 40 kcal

D. 50 kcal

Answer: B



Watch Video Solution

38. In a flask, colourless N_2O_4 is in equilibrium with brown-coloured NO_2 .

At equilibrium, when the flask is heated to $100^\circ C$ the brown colour

deepens and on cooling, the brown colour became less coloured. The change in enthalpy ΔH for the system is

- A. Negative
- B. Positive
- C. Zero
- D. Undefined

Answer: B



Watch Video Solution

39. For which of these reactions will there be ΔS positive?

- A. $H_2O(g) \rightarrow H_2O(l)$
- B. $H_2(g) + I_2(g) \rightarrow 2HI(g)$
- C. $CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$
- D. $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$

Answer: C

 [Watch Video Solution](#)

40. For stretched rubber, Entropy

- A. Increases
- B. First increases then decreases
- C. Decreases
- D. First decreases then increases

Answer: C

 [Watch Video Solution](#)

41. The most random state of H_2O system is

- A. ice

B. Liquid water

C. Steam

D. Randomness is same in all

Answer: A



[Watch Video Solution](#)

42. ΔS for the reaction ,

$MgCO_3(s) \rightarrow MgO(s) + CO_2(g)$ will be :

A. Zero

B. $-ve$

C. $+ve$

D. ∞

Answer: C



[Watch Video Solution](#)

43. The standard entropies of $N_2(g)$, $H_2(g)$ and $NH_3(g)$ are 191.5, 130.5, $192.6 J K^{-1} mol^{-1}$. The value of ΔS° of formation of ammonia is

A. $-98.9 J K^{-1} mol^{-1}$

B. Zero

C. $+129.4 J K^{-1} mol^{-1}$

D. $-29.4 J K^{-1} mol^{-1}$

Answer: A

 [Watch Video Solution](#)

44. What is the increase in entropy when $11.2 L$ of O_2 are mixed with $11.2 L$ of H_2 at STP?

A. $0.576 \frac{J}{K}$

B. $5.76 \frac{J}{K}$

C. $7.56 \frac{J}{K}$

D. $2.76 \frac{J}{K}$

Answer: B




Watch Video Solution

45. Given $S_{C_2H_6}^\circ = 225 J mol^{-1} K^{-1}$,

$S_{C_2H_4}^\circ = 220 J mol^{-1} K^{-1}$, $S_{H_2}^\circ = 130 J mol^{-1} K^{-1}$

Then ΔS° for the process



A. $+25J$

B. $-125J$

C. $135J$

D. $315J$

Answer: B

 [Watch Video Solution](#)

46. For the melting of NaCl heat required is $7.26 \text{ kcal mol}^{-1}$ and ΔS increases by $6.73 \text{ cal mol}^{-1} \text{ K}^{-1}$. The melting point of the salt is

A. 805.75°C

B. 500K


C. 1.77K

D. 1.77°C

Answer: A

 [Watch Video Solution](#)

47. The ΔS for the reaction



$$S_{H_2}^\circ(g) = 126.6, S_{O_2}^\circ(g) = 201.20,$$

$S_{H_2O}(l) = 68.0 JK^{-1}mol^{-1}$ respectively is

A. $-318.4 JK^{-1}mol^{-1}$

B. $318.4 JK^{-1}mol^{-1}$

C. $31.84 JK^{-1}mol^{-1}$

D. $3.184 JK^{-1}mol^{-1}$

Answer: A



Watch Video Solution

48. Which of the following is correct ?

A. ΔH ΔS Nature of reaction
(+) (-) Nonspontaneous regardless of temperature

B. ΔH ΔS Nature of reaction
(-) (+) Spontaneous only at high temperature

C. ΔH ΔS Nature of reaction
(+) (+) Spontaneous only at low temperature

D. ΔH ΔS Nature of reaction
(+) (-) Spontaneous at all temperatures

Answer: B

 [Watch Video Solution](#)

49. Entropy of vaporisation of water at 100°C , if molar heat of vaporisation is 8710calmol^{-1} will be

A. $20\text{calmol}^{-1}\text{K}^{-1}$

B. $23.36\text{calmol}^{-1}\text{K}^{-1}$

C. $24\text{calmol}^{-1}\text{K}^{-1}$

D. $28.0\text{calmol}^{-1}\text{K}^{-1}$

Answer: B

 [Watch Video Solution](#)

50. A particular reaction at 27°C for which $\Delta H > 0$ and $\Delta S > 0$ is found to be non-spontaneous. The reaction may proceed spontaneously

if

- A. The temperature is decreased
- B. The temperature is increased
- C. The temperature is kept constant
- D. It is carried in open vessel at $27^{\circ}C$

Answer: B



Watch Video Solution

51. It is impossible for a reaction to take place if

- A. $\Delta H_{is} + ve$ and $\Delta S_{is} + ve$
- B. $\Delta H_{is} - ve$ and $\Delta S_{is} + ve$
- C. $\Delta H_{is} + ve$ and $\Delta S_{is} - ve$
- D. $\Delta H_{is} - ve$ and $\Delta S_{is} - ve$

Answer: C



Watch Video Solution

52. The standard free energy change ΔG° is related to K (equilibrium constant) as`

A. $\Delta G^\circ = -2.303RT \log K$

B. $\Delta G^\circ = 2.303RT \log K$

C. $\Delta G^\circ = RT \log K$

D. $\Delta G^\circ = -RT \log K$

Answer: A



Watch Video Solution

53. The sole criterion for the spontaneity of a process is

A. Tendency to acquire minimum energy

B. Tendency to acquire maximum randomness

C. Tendency to acquire minimum energy and maximum randomness

D. Tendency to acquire maximum stability

Answer: D

 [Watch Video Solution](#)

54. For an endothermic reaction to be spontaneous

A. $\Delta G = 0$

B. $\Delta > 0$


C. $\Delta G < 0$

D. ΔG maybe +ve or -ve

Answer: C

 [Watch Video Solution](#)

55. At 27°C for reaction,



proceeds spontaneously because the magnitude of

- A. $\Delta H = T\Delta S$
- B. $\Delta H > T\Delta S$
- C. $\Delta H < T\Delta S$
- D. $\Delta H > 0$ and $T\delta S < 0$

Answer: B



[Watch Video Solution](#)

ASSIGNMENT (Section - B) Objective Type Questions

1. For one mole of an ideal gas (C_p and C_v are molar heat capacities at constant pressure and constant volume respectively)

A. $C_p - C_v = R$

B. $C_p - C_v = \frac{R}{2}$

C. $C_p - C_v = -2R$

D. $C_p - C_v = 0$

Answer: A

 [Watch Video Solution](#)

2. When an ideal gas is compressed adiabatically and reversibly, the final temperature is:

A. Higher than the initial temperature

B. Lower than the initial temperature





C. The same as the initial temperature

D. Dependent on the rate of compression

Answer: A

 [Watch Video Solution](#)

3. ΔS° will be highest for the reaction



Answer: B

 [Watch Video Solution](#)

4. In a reversible process,

$\Delta S_{sys} + \Delta S_{surr}$ is

A. $+ve$

B. $-ve$

C. Zero

D. All of these

Answer: A



[Watch Video Solution](#)

5. A closed flask contains a substance in all its three states, solids, liquids and vapour at its triple point. In this situation the average KE of the water molecule will be

A. Maximum in vapour state

B. Maximum in solid state

C. Greater in the liquid than in vapour state

D. Same in all the three states

Answer: D



[Watch Video Solution](#)

6. In thermodynamics, a process is called reversible when

- A. System and surrounding change into each other
- B. There is no boundary between system and surrounding
- C. The surroundings are always in equilibrium with the system
- D. The system changes into the surroundings spontaneously

Answer: C



[Watch Video Solution](#)

7. The molar heat capacity of water at constant pressure P is $75JK^{-1}mol^{-1}$. When $1.0kJ$ of heat is supplied to 1000 g of water, which is free to expand, the increase in temperature of water is

- A. $1.2K$
- B. $2.4K$

C. $4.8K$

D. $0.24K$

Answer: D



[Watch Video Solution](#)

8. $16g$ oxygen gas expands at STP to occupy double of its oxygen volume. The work done during the process is:

A. 260 kcal

B. 180 kcal

C. 130 kcal

D. 271 kcal

Answer: D



[Watch Video Solution](#)

9. When enthalpy and entropy change for a chemical reaction are -2.5×10^3 cal and 7.4 cal deg^{-1} respectively. Predict that reaction at 298 K is


- A. Spontaneous
- B. Reversible
- C. Irreversible
- D. Non-spontaneous

Answer: A



[Watch Video Solution](#)

10. The temperature at which the given reaction is at equilibrium



$$\Delta H = 40.5 \text{ kJ mol}^{-1} \text{ and } \Delta S = 0.086 \text{ kJ mol}^{-1} \text{ K}^{-1}$$

- A. 470.93 K

B. 362.12 K

C. 262.12 K

D. 562.12 K

Answer: A



Watch Video Solution

11. One mole of a non-ideal gas undergoes a change of state $(2.0\text{atm}, 3.0\text{L}, 95\text{K}) \rightarrow (4.0\text{atm}, 5.0\text{L}, 245\text{K})$ with a change in internal energy, $\Delta U = 30.0\text{L. atm}$. Calculate change in enthalpy of the process in L. atm .

A. 40.0

B. 42.3

C. 44.0

D. 56.0

Answer: C

 [Watch Video Solution](#)

12. Which of the following can be zero for isothermal reversible expansion?

A. ΔE

B. ΔH

C. ΔT

D. All of these

Answer: D

 [Watch Video Solution](#)

13. In an isolated container water is stirred with a rod to increase the temperature. Which of the following is true?

A. $\Delta U = W \neq 0, q = 0$

B. $\Delta U = W = q \neq 0$

C. $\Delta U = 0, W = q \neq 0$

D. $W = 0, \Delta U = q \neq 0$

Answer: A

 [Watch Video Solution](#)

14. Two atoms of hydrogen combine to form a molecule of hydrogen gas, the energy of the H_2 molecule is :

A. Greater than that of separate atoms

B. Equal to that of separate atoms

C. Lower than that of separate atoms

D. Sometimes lower and sometimes higher

Answer: C

 [Watch Video Solution](#)

15. The temperature of 15 ml of a strong acid increases by $2^{\circ}C$ when 15 ml of a strong base is added to it. If 5 ml of each are mixed, temperature should increase by

A. $0.6^{\circ}C$

B. $0.3^{\circ}C$

C. $2^{\circ}C$

D. $6^{\circ}C$

Answer: C

 [Watch Video Solution](#)

16. The standard heat of formation of $NO_2(g)$ and $N_2O_4(g)$ are 8.0 and $4.0kcalmol^{-1}$ respectively. The heat of dimerisation of NO_2 in kcal is`

A. $-12kcal$

B. $12kcal$

C. $4kcal$

D. $16kcal$

Answer: A



Watch Video Solution

17. If $\frac{1}{2}X_2O_{(s)} \rightarrow X_{(s)} + \frac{1}{4}O_{2(g)}$, $\Delta H = 90kJ$ then heat change during reaction of metal X with $1moleO_2$ to form oxide to maximum extent is :

A. $360kJ$

B. $-360kJ$


C. $-180kJ$

D. $+180kJ$

Answer: B

 [Watch Video Solution](#)

18. For a gaseous reaction



ΔH is 27 kcal at $37^\circ C$. Assuming $R = 2\text{calK}^{-1}\text{mol}^{-1}$ the value of ΔH for the above reaction will be

- A. 15.8kcal
- B. 28.2kcal
- C. 25.7kcal
- D. 20.0kcal

Answer: B

 [Watch Video Solution](#)

19. A mixture of 2 moles of CO and 1 mole of O_2 , in a closed vessel is ignited to convert the CO to CO_2 if ΔH is the enthalpy change and ΔU is the change in internal energy then:

A. $\Delta H = \Delta U$

B. $\Delta H > \Delta U$

C. $\Delta H < \Delta U$

D. The relationship depends upon the capacity of vessel

Answer: C



Watch Video Solution

20. If the bond dissociation energies of XY , X_2 and Y_2 (all diatomic molecules) are in the ratio 1:1:0.5 and $\Delta_f H$ of XY is $-200 kJ mol^{-1}$.

The bond dissociation energy of X_2 will be :

A. $800 kJ mol^{-1}$

B. 200kJmol^{-1}

C. 300kJmol^{-1}

D. 400kJmol^{-1}

Answer: A



[Watch Video Solution](#)

21. Vapour density of a gas is 8. Its molecular mass will be

A. 8

B. 16

C. 32

D. 64

Answer: B



[Watch Video Solution](#)

22. If x mole of ideal gas at 27°C expands isothermally and reversibly from a volume of y to $10y$, then the work done is

A. $w = xR300\ln y$

B. $w = -300xR\ln\frac{y}{10}$

C. $w = -300xR\ln 10$

D. $w = 100xR\ln\frac{1}{y}$

Answer: C



Watch Video Solution

23. Enthalpy of formation of NH_3 is $-X$ kJ and $\Delta H_{\text{H}-\text{H}}$, $\Delta H_{\text{N}-\text{H}}$ are respectively Y kJmol^{-1} and Z kJmol^{-1} . The value of $\Delta H_{\text{N}=\text{N}}$ is

A. $Y - 6Z + \frac{X}{3}$

B. $-3Y + 6Z + 2X$

C. $3Y + 6Z + X$


$$D. Y + 6X + Z$$

Answer: B



[Watch Video Solution](#)

24. A system X undergoes following changes



The overall process may be called as

- A. Reversible process
- B. Cyclic process
- C. Cyclic reversible process
- D. Isochoric process

Answer: B



[Watch Video Solution](#)

25. The heat of neutralisation for strong acid and strong base forming 2 moles of water is

A. $-2 \times 57.1 \text{ kJ}$

B. -57.1 kJ

C. $-\frac{57.1}{2} \text{ kJ}$


D. Strong acid and strong base will not undergo neutralisation

Answer: A



Watch Video Solution

26. The value of ΔH° in kJ for the reaction will be



$$\Delta H_f^\circ(\text{CS}_2) = -X$$

$$\Delta H_f^\circ(\text{NOCl}) = -y$$

$$\Delta H_f^\circ(\text{CCL}_4) = +z$$

$$\Delta H_f^\circ(\text{SO}_2) = -r$$

A. $x + 4y - z - 2r$

B. $r + z + 4y - x$

C. $2r + z + 4y + x$

D. $x + 4y + z - 2r$

Answer: D

 [Watch Video Solution](#)

27. The heat liberated on complete combustion of 1 mole of CH_4 gas to $CO_2(g)$ and $H_2O(l)$ is 890 kJ. Calculate the heat evolved by 2.4L of CH_4 on complete combustion.

A. $95.3kJ$

B. $8900kJ$

C. $890kJ$

D. $8.9kJ$

Answer: A



Watch Video Solution

28. The work done in an open vessel at $300K$, when $112g$ iron reacts with dil. HCL is:

A. $1.1kcal$

B. $0.6kcal$

C. $0.3kcal$

D. $0.2kcal$

Answer: A



Watch Video Solution

29. Which statement is correct?


- A. $\frac{dh}{(dT)_P} < \frac{dE}{(dT)_V}$
- B. $\frac{dH}{(dT)_P} + \frac{dE}{(dT)_V} = R$
- C. $\frac{dE}{(dV)_T}$ for ideal gas is zero
- D. All of these

Answer: C

 [View Text Solution](#)

30. A schematic representation of enthalpy changes for the reaction,

$C_{\text{graphite}} + \frac{1}{2}O_2(g) \rightarrow CO(g)$ is given below. The missing value is



A. $+10.5kJ$

B. -11.05kJ




C. -110.5kJ

D. -10.5J

Answer: C

 [Watch Video Solution](#)


31. Which of the following equations represents standard heat of formation of CH_4 ?



Answer: B

 [Watch Video Solution](#)

32. Different types of systems are given below



The A and B systems respectively are

- A. Open system, Closed system
- B. Isolated system, Closed system
- C. Adiabatic system, Isolated system
- D. Closed system, Isolated system

Answer: D



[Watch Video Solution](#)

33. Set of intensive properties is shown by

- A. Mole fraction, standard electrode potential, heat capacity
- B. Viscosity, refractive index, specific heat
- C. Density, Gibbs free energy, internal energy
- D. Number of moles, molarity, electrode potential

Answer: B

 [Watch Video Solution](#)


34. For the expansion occurring from initial to final stage in finite time, which is incorrect?

- A. Equilibrium exist in initial and final stage
- B. Work obtained is maximum
- C. Driving force is much greater than the opposing force
- D. Both (1) & (2)

Answer: B

 [Watch Video Solution](#)

35. Calorific value of ethane, in k J/g if for the reaction



A. - 12.4

B. - 52

C. - 24.8

D. - 104

Answer: B

 [Watch Video Solution](#)

ASSIGNMENT (Section - C) Previous Years Questions

1. A gas is allowed to expand in a well insulated container against a constant external pressure of 2.5 atm from an initial volume of 2.50 L to a

final volume of 4.50 L. the change in internal energy ΔU of the gas in joules will be-

A. $1136.25J$

B. $-500J$

C. $-505J$

D. $+505J$

Answer: C



Watch Video Solution

2. For a given reaction, $\Delta H = 35.5kJmol^{-1}$ and $\Delta S = 83.6JK^{-1}mol^{-1}$. The reaction is spontaneous at : (Assume that ΔH and ΔS do not vary with temperature)

A. $T < 425K$

B. $T > 425K$

C. All temperature

$$D. T > 298K$$

Answer: B

 [Watch Video Solution](#)

3. For a sample of perfect gas when its pressure is changed isothermally from $p_i \rightarrow p_f$ the entropy change is given by

A. $\Delta S = nR \ln \left(\frac{p_f}{p_i} \right)$

B. $\Delta S = nR \ln \left(\frac{p_i}{p_f} \right)$

C. $\Delta S = nRT \ln \left(\frac{p_f}{p_i} \right)$

D. $\Delta S = nRT \ln \left(\frac{p_i}{p_f} \right)$

Answer: B

 [Watch Video Solution](#)

4. The correct thermodynamic conditions for the spontaneous reaction at all temperature is:

- A. $\Delta H < 0$ and $\Delta S < 0$
- B. $\Delta H < 0$ and $\Delta S = 0$
- C. $\Delta H > 0$ and $\Delta S < 0$
- D. $\Delta H < 0$ and $\Delta S > 0$

Answer: B::D

 [Watch Video Solution](#)

5. Heat of combustion of carbon to CO_2 is -393.5 kJ/mol. The heat released upon the formation of 35.2 kg of CO_2 from carbon and oxygen gas is-

- A. -630 kJ
- B. -3.15 kJ

C. $-315kJ$


D. $+315kJ$

Answer: C



Watch Video Solution

6. For the reaction:



$$\Delta U = 2.1\text{cal}, \Delta S = 20\text{calK}^{-1}\text{at } 300\text{K}$$

Hence ΔG is

A. 2.7kcal

B. -3.9kcal

C. 9.3kcal





D. -9.3kcal

Answer: B



Watch Video Solution

7. In which of the following reactions, standard reaction entropy change (ΔS°) is positive and standard Gibbs energy change (ΔG°) decreases sharply with increasing temperature?



Answer: C



Watch Video Solution

8. Standard enthalpy of vaporisation $\Delta V_{vap} \cdot H^\ominus$ for water at $100^\circ C$ is $40.66 kJ mol^{-1}$. The internal energy of Vaporization of water at $100^\circ C$ (in $kJ mol^{-1}$) is

A. + 43.76

B. + 40.66

C. + 37.56

D. - 43.76

Answer: C

 [Watch Video Solution](#)

9. The enthalpy of fusion of water is 1.435 kcal/mol. The molar entropy change for the melting of ice at $0^{\circ}C$ is :

A. $5.260ca \frac{l}{molK}$

B. $0.526ca \frac{l}{molK}$

C. $10.52ca \frac{l}{molK}$

D. $21.04ca \frac{l}{molK}$

Answer: A

 [Watch Video Solution](#)

10. Equal volumes of two monoatomic gases, A , B , at the same temperature and pressure are mixed. The ratio of specific heats (C_p / C_v) of the mixture will be

A. 0.83

B. 1.50

C. 3.3

D. 1.67

Answer: D

 [Watch Video Solution](#)

11. Which of the following is correct option for the free expansion of an ideal gas under adiabatic condition ?

A. $q = 0, \Delta T < 0, w \neq 0$

B. $q = 0, \Delta T \neq 0, w = 0$

C. $q \neq 0, \Delta T = 0, w = 0$

D. $q = 0, \Delta T = 0, w = 0$

Answer: D

 [Watch Video Solution](#)

12. If the enthalpy change for the transition of liquid water to steam is 30kJmol^{-1} at 27°C , the entropy change for the process would be

A. $100\text{Jmol}^{-1}\text{K}^{-1}$

B. $10\text{Jmol}^{-1}\text{K}^{-1}$

C. $1.0\text{Jmol}^{-1}\text{K}^{-1}$

D. $0.1\text{Jmol}^{-1}\text{K}^{-1}$

Answer: A

 [Watch Video Solution](#)

13. Enthalpy change for the reaction, $4H_{(g)} \rightarrow 2H_{2(g)}$ is -869.6 kJ

The dissociation energy of H-H bond is :

A. $+217.4 \text{ kJ}$

B. -434.8 kJ

C. -869.6 kJ

D. $+434.8 \text{ kJ}$

Answer: D

 [Watch Video Solution](#)

14. Consider the following processes :-



For $B + D \rightarrow E + 2C$, ΔH will be

A. $-325k \frac{J}{m} ol$

B. $325k \frac{J}{m} ol$

C. $525k \frac{J}{m} ol$

D. $-175k \frac{J}{m} ol$

Answer: D

 [Watch Video Solution](#)

15. Standard entropies of X_2 , Y_2 and XY_3 are 60, 40 and $50 \text{ JK}^{-1} \text{ mol}^{-1}$ respectively. For the reaction



to be at equilibrium, the temperature should be

A. 750 K

B. 1000 K

C. 1250 K

D. 500 K

Answer: A

 Watch Video Solution

16. Match List I (Equations) with List II (Type of processes) and select the correct option.

| List I (Equations) | List II (Type of processes) |
|------------------------------------|--------------------------------|
| A. $K_p > Q$ | 1. Non-spontaneous |
| B. $\Delta G^\circ < RT \ln Q$ | 2. Equilibrium |
| C. $K_p = Q$ | 3. Spontaneous and endothermic |
| D. $T > \frac{\Delta H}{\Delta S}$ | 4. Spontaneous |

A. $a(i), b(ii), c(iii), d(iv)$

B. $a(iii), b(iv), c(ii), d(i)$

C. $a(iv), b(i), c(ii), d(iii)$

D. $a(ii), b(i), c(iv), d(iii)$

Answer: C



[Watch Video Solution](#)

17. 1 moles of an ideal gas expanded spontaneously into vacuum. The work done will be

- A. Infinite
- B. 3 Joules
- C. 9 Joules
- D. Zero

Answer: D



[Watch Video Solution](#)

18. For vaporization of water at 1 atmospheric pressure the values of ΔH and ΔS are $50.63 \text{ kJ mol}^{-1}$ and $118.8 \text{ JK}^{-1} \text{ mol}^{-1}$ respectively. The temperature when Gibbs energy change (ΔG) for this transformation will be zero, is

A. 273.4K

B. 393.4K


C. 426.1K

D. 293.4K


Answer: C

 [Watch Video Solution](#)

19. The following two reactions are known :




$$\Delta H = - 26.8kJ$$



$$\Delta H = - 16.5kJ$$

Correct target equation is



A. +10.3kJ

B. $-43.3kJ$

C. $-10.3kJ$


D. $+6.2kJ$

Answer: D



Watch Video Solution

20. The values of ΔH and ΔS for the reaction,



are $170KJ$ and $170JK^{-1}$ respectively. This reaction will be spontaneous at

A. 910 K

B. 1110 K

C. 510 K

D. 710 K

Answer: B

 Watch Video Solution

21. From the following bond energies


$H - H$ bond energy $431.37 \text{ kJ mol}^{-1}$

$C = C$ bond energy $606.10 \text{ kJ mol}^{-1}$

$C - C$ bond energy $336.49 \text{ kJ mol}^{-1}$

$C - H$ bond energy $410.5 \text{ kJ mol}^{-1}$

Enthalpy for the reaction



will be

A. $-243.6 \text{ kJ mol}^{-1}$

B. $-120.0 \text{ kJ mol}^{-1}$

C. $553.0 \text{ kJ mol}^{-1}$

D. $1523.6 \text{ kJ mol}^{-1}$

Answer: B

 [Watch Video Solution](#)

22. Bond dissociation enthalpy of H_2 , Cl_2 and HCl are 434, 242 and 431 kJ mol^{-1} respectively. Enthalpy of formation of HCl is

A. 245 kJ mol^{-1}

B. 93 kJ mol^{-1}

C. -245 kJ mol^{-1}

D. -93 kJ mol^{-1}

Answer: D

 [Watch Video Solution](#)

23. For the gas-phase decomposition,



A. $\Delta H > 0$ and $\Delta S < 0$

B. $\Delta H = 0$ and $\Delta S < 0$

C. $\Delta H > 0$ and $\Delta S > 0$

D. $\Delta H < 0$ and $\Delta S < 0$

Answer: C



Watch Video Solution

24. Which of the following are not state functions?

(i) $q + w$

(ii) q

(iii) w

(iv) $H - TS$

A. (II) and(III)

B. (I) and(IV)


C. (II) , (III) and(IV)

D. (I), (II) and(III)



Answer: A

 Watch Video Solution


25. Consider the following reactions:



$$\Delta H = -X_1 \text{KJmol}^{-1}$$



$$\Delta H = -X_3 \text{KJmol}^{-1}$$



$$\Delta H = +X_4 \text{KJmol}^{-1}$$

Enthalpy of formation of $H_2O(l)$ is

A. $+X_1 \text{kJmol}^{-1}$

B. $+X_2 \text{kJmol}^{-1}$

C. $+X_3 \text{kJmol}^{-1}$

$$D. + X_4 kJmol^{-1}$$

Answer: B

 [Watch Video Solution](#)

26. Given the bond energies of $H - H$ and $Cl - Cl$ are $430 kJmol^{-1}$ and $240 kJmol^{-1}$, respectively, and $\Delta_f H^\circ$ for HCl is $-90 kJmol^{-1}$. Bond enthalpy of HCl is

A. $245 kJmol^{-1}$

B. $2909 kJmol^{-1}$

C. $380 kJmol^{-1}$

D. $425 kJmol^{-1}$

Answer: D

 [Watch Video Solution](#)

27. Identify the correct statement for change of Gibbs energy for a system (ΔG_{system}) at constant temperature and pressure.


- A. If $\Delta G_{\text{system}} > 0$, the process is spontaneous
- B. If $\Delta G_{\text{system}} = 0$ the system has attained equilibrium
- C. If $\Delta G_{\text{system}} = 0$, the system is still moving in a particular direction
- D. If $\Delta G_{\text{system}} < 0$, the system is not spontaneous

Answer: B

 [Watch Video Solution](#)

28. Assume each reaction is carried out in an open container. For which reaction will $\Delta H = \Delta E$?


- A. $H_2(g) + Br_2(g) \rightarrow 2HBr(g)$
- B. $C(s) + 2H_2O(g) \rightarrow 2H_2(g) + CO_2(g)$
- C. $PCl_5(g) \rightarrow PCl_3(g) + Cl_2(g)$



Answer: A

 [Watch Video Solution](#)

29. The enthalpy and entropy change for the reaction



respectively. The temperature at which the reaction will be in equilibrium

is

A. $363.64K$

B. $273K$

C. $450K$

D. $300K$

Answer: A

 [Watch Video Solution](#)

30. At 25°C , standard enthalpy of combustion of combustion of hydrogen, cyclohexene (C_6H_{10}) and cyclohexane (C_6H_{12}) are -241 , -3800 and -3920 kJ mol^{-1} respectively. Determine the heat of hydrogenation of cyclohexene.

A. -121kJpermol

B. $+121\text{kJpermol}$

C. $+242\text{kJpermol}$

D. -242kJpermol

Answer: A

 [Watch Video Solution](#)

31. A reaction occurs spontaneously if

A. $T\Delta S < \Delta H$ and both ΔH and ΔS are $+ve$

B. $T\Delta S > \Delta H$ and both ΔH and ΔS are $+ve$

C. $T\Delta S = \Delta H$ and both ΔH and ΔS are +ve

D. $T\Delta S > \Delta H$ and ΔH is +ve and ΔS is -ve

Answer: B

 [Watch Video Solution](#)

32. Which of the following pairs of a chemical reaction is certain to result in a spontaneous reaction ?

A. Exothermic and decreasing disorder

B. Endothermic and increasing disorder

C. Exothermic and increasing disorder

D. Endothermic and decreasing disorder

Answer: C

 [Watch Video Solution](#)

33. The absolute enthalpy of neutralization of the reaction,

$MgO(s) + 2HCl(aq.) + H_2O(l)$ will be

- A. Less than $-57.33kJmol^{-1}$
- B. $-57.33kJmol^{-1}$
- C. Greater than $-57.33kJmol^{-1}$
- D. $57.33kJmol^{-1}$

Answer: A



Watch Video Solution

34. Which reaction, with the following values of ΔH , ΔS , at 400 K is spontaneous and endothermic?




- A. $\Delta H = -48kJ, \Delta S = +135\frac{J}{K}$
- B. $\Delta H = -48kJ, \Delta S = -135\frac{J}{K}$
- C. $\Delta H = +48kJ, \Delta S = +135\frac{J}{K}$

$$D. \Delta H = +48 \text{ kJ}, \Delta S = -135 \frac{\text{J}}{\text{K}}$$

Answer: C

 Watch Video Solution

35. Given :



Based on the above thermochemical equations, find out which one of the following algebraic relationships is correct ?

A. $x = y - z$

B. $z = x + y$

C. $x = y + z$


D. $y = 2z - x$

Answer: C



Watch Video Solution

36. Following reaction occurring in an automobile



ΔH , ΔS and ΔG would be-

A. -, +, +

B. +, +, -

C. +, -, +

D. -, +, -

Answer: D



Watch Video Solution

37. When 5 litres of a gas mixture of methane and propane is perfectly combusted at $0^\circ C$ and 1 atmosphere, 16 litre of oxygen at the same temperature and pressure is consumed. The amount of heat released

from this combustion in kJ

$(\Delta H_{comb}(CH_4) = 890 \text{ kJ mol}^{-1}, \Delta H_{comb}(C_3H_8) = 2220 \text{ kJ mol}^{-1})$ is

A. 32

B. 38

C. 317

D. 477

Answer: C



Watch Video Solution

38. The enthalpy of formation for $C_2H_4(g)$, $CO_2(g)$ and $H_2O(l)$ at $25^\circ C$ and 1 atm. Pressure be 52, -394 and -286 kJ mol^{-1} respectively. The enthalpy of combustion of $C_2H_4(g)$ will be

A. $+14.2 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

B. $+1412 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

C. $-141.2 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

D. $-1412k\frac{J}{m}ol$

Answer: D

 [Watch Video Solution](#)

39. For a reaction to occur spontaneously

- A. ΔH must be negative
- B. ΔS must be negative
- C. $(\Delta H - T\Delta S)$ must be negative
- D. $(\Delta H + T\Delta S)$ must be negative

Answer: C

 [Watch Video Solution](#)

40. Given that $C + O_2 \rightarrow CO_2$, $\Delta H^\circ = -xkJ$

$2CO + O_2 \rightarrow 2CO_2$, $\Delta H^\circ = -ykJ$

What is heat of formation of CO?

A. $\frac{y - 2x}{2}$

B. $2x - y$

C. $y - 2x$

D. $\frac{2x - y}{2}$

Answer: A



Watch Video Solution

41. Identify the correct statement regarding entropy

A. At absolute zero of temperature, the entropy of all crystalline substances is taken to be zero

- B. At absolute zero of temperature, the entropy of a perfectly crystalline substance is $+ve$
- C. At absolute zero of temperature, entropy of a perfectly crystalline substance is taken to be zero
- D. At $0^{\circ}C$, the entropy of a perfectly crystalline substance is taken to be zero.

Answer: C

 [Watch Video Solution](#)

42. One mole of an ideal gas at 250 K is expanded isothermally from an initial volume of 5 litre to 10 litres. The ΔE for this process is $(R = 2cal. Mol^{-1}K^{-1})$



- A. 1381.1cal
- B. Zero
- C. 163.7cal

D. 9Latm

Answer: B



Watch Video Solution



The heat of formation of SO_2 is : –

A. $(2x + y)$

B. $(x - y)$

C. $(x + y)$

D. $(y - 2x)$

Answer: D



Watch Video Solution

44. At $37^{\circ}C$ latent heat of fusion of a compound is 3930 J/mol . Entropy change is

A. $13.10 \frac{\text{J}}{\text{m}} \text{olK}$

B. $10.77 \frac{\text{J}}{\text{m}} \text{olK}$

C. $9.07 \frac{\text{J}}{\text{m}} \text{olK}$

D. $0.977 \frac{\text{J}}{\text{m}} \text{olK}$

Answer: A



Watch Video Solution

45. For the reaction $C_2H_5OH(l) + 3O_2(g) \rightarrow 2CO_2(g) + 3H_2O(l)$, which one is true ?

A. $\Delta H = \Delta E - RT$

B. $\Delta H = \Delta E + RT$

C. $\Delta H = \Delta E + 2RT$

$$D. \Delta H = \Delta E - 2RT$$

Answer: A

 [Watch Video Solution](#)

46. Change in enthalpy for reaction $2H_2O_2(l) \rightarrow 2H_2O(l) + O_2(g)$

if heat of formation of $H_2O_2(l)$ and $H_2O(l)$ are -188 and -286 kJ/mol respectively is

A. $-196 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

B. $+196 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

C. $+948 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

D. $-948 \text{ k} \frac{\text{J}}{\text{m}} \text{ol}$

Answer: A

 [Watch Video Solution](#)

47. When 1 mol of gas is heated at constant volume temperature is raised from 298 to 308 K. Heat supplied to the gas is 500 J. Then which statement is correct ?

A. $q = \Delta U = -500J, w = 0$

B. $q = \Delta U = +500J, w = 0$

C. $q = w - 500J, \Delta U = 0$

D. $\Delta U = 0, q = w = -500J$

Answer: B

 [Watch Video Solution](#)

48. Enthalpy of $CH_4 + \frac{1}{2}O_2 \rightarrow CH_3OH$ is negative. If enthalpy of combustion of CH_4 and CH_3OH are x and y respectively. Then which relation is correct ?

A. $x > y$

B. $x < y$

C. $x = y$

D. $x \geq y$

Answer: A



Watch Video Solution

49. The unit of entropy is

A. $JK^{-1}mol^{-1}$

B. $Jmol^{-1}$

C. $J^{-1}K^{-1}mol^{-1}$

D. $JKmol^{-1}$

Answer: A



Watch Video Solution

50. In a closed insulated container, a liquid is stirred with a paddle to increase the temperature, which of the following is true?

A. $\Delta E = W \neq 0, q = 0$

B. $\Delta E = W = q \neq 0$

C. $\Delta E = 0, W = q \neq 0$

D. $W = 0, \Delta E = q \neq 0$

Answer: A

 [Watch Video Solution](#)

51. Two moles of ideal gas at $27^\circ C$ temperature is expanded reversibly from 2 litre to 20 liter. Find entropy change ($R = 2cal/molK$).

A. 92.1

B. 0

C. 4

D. 9.2

Answer: D

 [Watch Video Solution](#)

52. Heat of combustion ΔH° for $C(s)$, $H_2(g)$ and $CH_4(g)$ are 94, -68 and -213Kcal/mol . Then ΔH° for $C(s) + 2H_2(g) \rightarrow \Delta CH_4(g)$ is

A. -17kcal

B. -111kcal


C. -170kcal

D. -85kcal

Answer: A

 [Watch Video Solution](#)

53. For the reaction



$\Delta H - \Delta E$ is`

A. $+RT$

B. $-4RT$

C. $+3RT$

D. $-RT$

Answer: B



Watch Video Solution

54. What is the entropy change ($\in JK^{-1}mol^{-1}$) when one mole of ice is converted into water at $0^\circ C$? (The enthalpy change for the conversion of ice to liquid water is $6.0kJmol^{-1}$ at $0^\circ C$)

A. 20.13

B. 2.013





C. 2.198

D. 21.98

Answer: D

 [Watch Video Solution](#)

55. For which one of the following equations is ΔH_{react}° equal to ΔH_r° for the product?



Answer: C

 [Watch Video Solution](#)

56. Standard enthalpy and standard entropy changes for the oxidation of ammonia at 298 K are $-582.64 \text{ kJ mol}^{-1}$ and $-150.6 \text{ J mol}^{-1}$, respectively. Standard Gibbs energy change for the same reaction at 298 K is

A. $-221.1 \text{ kJ mol}^{-1}$

B. $-537.7 \text{ kJ mol}^{-1}$

C. $-439.3 \text{ kJ mol}^{-1}$

D. $-523.2 \text{ kJ mol}^{-1}$

Answer: B



Watch Video Solution

57. Considering entropy (S) as a thermodynamics parameter, the criterion for the spontaneity of any process is

A. $\Delta S_{\text{system}} + \Delta S_{\text{surroundings}} > 0$

B. $\Delta S_{\text{system}} - \Delta S_{\text{surroundings}} > 0$

C. $\Delta S_{\text{system}} > 0$ only

D. $\Delta S_{\text{surroundings}} > 0$ only

Answer: A

 [Watch Video Solution](#)

58. The work done during the expansion of a gas from a volume of 4dm^3 to 6dm^3 against a constant external pressure of 3 atm is ($1\text{Latm} = 101.32\text{J}$)

A. -6J

B. -608J

C. $+304\text{J}$

D. -304J

Answer: B



Watch Video Solution

59. The enthalpy of hydrogenation of cyclohexene is $-119.5 \text{ kJ mol}^{-1}$. If resonance energy of benzene is $-150.4 \text{ kJ mol}^{-1}$, its enthalpy of hydrogenation would be :

A. $-358.5 \text{ kJ mol}^{-1}$

B. $-508.9 \text{ kJ mol}^{-1}$




C. $-208.1 \text{ kJ mol}^{-1}$

D. $-269.9 \text{ kJ mol}^{-1}$


Answer: C



Watch Video Solution



ΔG° for the following reaction



A. $-1462J$

B. $-1317J$

C. $-501J$


D. $-731J$

Answer: D



Watch Video Solution

ASSIGNMENT (Section -D) Assertion-Reason Type Questions



ΔH and ΔU are same for this reaction.

R: Entropy increases during the conversion of diamond to graphite.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: B



Watch Video Solution

2. A: Specific heat is an intensive property

R: Heat capacity is an extensive property.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion, then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: B



[Watch Video Solution](#)

3. A: All reactions which are exothermic are spontaneous

R: All reactions in which entropy increases are spontaneous

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)
- B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)
- C. If Assertion is true statement but Reason is false, then mark (3)
- D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: D

 [Watch Video Solution](#)

4. A: Enthalpy of neutralisation of 1 equivalent each of HCl and H_2SO_4 with $NaOH$ is same.

R : Enthalpy of neutralisation is always the heat evolved when 1 mole acid is neutralised by a base.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)
- B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)
- C. If Assertion is true statement but Reason is false, then mark (3)
- D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: C

 [Watch Video Solution](#)

5. A: q and w are path function

R: $q + w$ is a state function

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: B

 [Watch Video Solution](#)

6. A : Dissolution of sugar in water proceed via increases in entropy

R: Entropy decreases when egg is hard boiled.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark

(3)

Answer: C



Watch Video Solution

7. A : For an isolated system $\Delta G = - T \Delta S_{\text{total}}$

R: For an isolated system $q = 0$.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark

(3)

Answer: B



Watch Video Solution

8. A: Combustion is an exothermic process.

R: Combustion is a spontaneous process.

A. If both Assertion & Reason are true and the reason is the correct

explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the

correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark

(3)

Answer: B



Watch Video Solution

9. A: Total enthalpy change of a multistep process is sum of $\Delta H_1 + \Delta H_2 + \Delta H_3 + \dots$

R: When heat is absorbed by the system, the sign of q is taken to be negative.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: C



[Watch Video Solution](#)

10. A : Bond energy is equal to enthalpy of formation with negative sign.

R: Bond energy is energy required to dissociate 1 mole single bond.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)
- B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)
- C. If Assertion is true statement but Reason is false, then mark (3)
- D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: C



[Watch Video Solution](#)

11. A: ΔH is positive for endothermic reactions.

R : If total enthalpies of reactants and products are H_R & H_P respectively then for an endothermic reaction $H_R < H_P$

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: A



Watch Video Solution

12. A : The energy of the universe is constant, whereas the entropy of the universe is continuously increasing.

R : For spontaneous process $\Delta S > 0$.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: B



Watch Video Solution

13. A: A non- spontaneous process becomes spontaneous when coupled with a suitable spontaneous reaction.

R: The overall free energy of coupled spontaneous reaction is negative.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: A



[Watch Video Solution](#)

14. A: An ideal crystal has more entropy than a real crystal.

R: An ideal crystals has more disorder.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion and Reason is false, then mark (4)

Answer: D



[Watch Video Solution](#)

15. A: Work done in an irreversible isothermal process at constant volume is zero.

R: Work is assigned negative sign during expansion and is assigned positive sign during compression.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1)

B. If both Assertion & Reason are true and the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3)

D. If both Assertion is true statement but Reason is false, then mark (3)

Answer: B



[Watch Video Solution](#)