



## MATHS

# BOOKS - MHTCET PREVIOUS YEAR PAPERS AND PRACTICE PAPERS

## SOLVED PAPER 2017

### Mcqs

1. The number of principal solutions of  $\tan 2\theta = 1$  is

A. one

B. two

C. three

D. four

**Answer: B**

 [Watch Video Solution](#)

2. The objective function,  $z = 4x_1 + 5x_2$ , subject to  $2x_1 + x_2 \geq 7$ ,  $2x_1 + 3x_2 \leq 15$ ,  $x_2 \leq 3$ ,  $x_1, x_2 \geq 0$  has minimum value at the point

A. On X-axis

B. on Y-axis

C. at the origin

D. on the line parallel to X-axis

**Answer: A**



**Watch Video Solution**

**3.** If  $z_1$  and  $z_2$  are z-coordinates of the points of trisection of the segment joining the points  $A(2,1,4)$ ,  $B(-1,3,6)$ , then

$$z_1 + z_2 +$$

A. 1

B. 4

C. 5

D. 10

**Answer: D**



**Watch Video Solution**

4. The maximum value of  $f(x) = \frac{\log x}{x}$  ( $x \neq 0, x \neq 1$ ) is

A.  $e$

B.  $\frac{1}{e}$

C.  $e^2$

D.  $\frac{1}{e^2}$

**Answer: B**



**Watch Video Solution**

5.  $\int_0^1 x \tan^{-1} x dx =$

A.  $\frac{\pi}{4} + \frac{1}{2}$

B.  $\frac{\pi}{4} - \frac{1}{2}$

C.  $\frac{1}{2} - \frac{\pi}{4}$

D.  $-\frac{\pi}{4} - \frac{1}{2}$

**Answer: B**



**Watch Video Solution**

**6. The statement pattern  $(\sim p \wedge q)$  is logically equivalent to**

A.  $(P \vee q) \vee \sim p$

B.  $(p \vee q) \wedge \sim p$

C.  $(p \wedge q) \rightarrow p$

D.  $(p \vee q) \rightarrow p$

**Answer: B**

 **Watch Video Solution**

7. If  $g(x)$  is the inverse function of  $f(x)$  and

$f'(x) = \frac{1}{1+x^4}$ , then  $g'(x)$  is

A.  $1 + [g(x)]^4$

B.  $1 - [g(x)]^4$

C.  $1 + [f(x)]^4$

D.  $\frac{1}{1 + [g(x)]^4}$

**Answer: A**



Watch Video Solution

8. The inverse of the matrix  $\begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix}$  is

A.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 2 & -3 \end{bmatrix}$

B.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

C.  $-\frac{1}{3} \begin{bmatrix} 3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

D.  $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ -3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

**Answer: B**



Watch Video Solution

9. If  $\int \frac{1}{\sqrt{9-16x^2}} dx = \alpha \sin^{-1}(\beta x) + c$ , then  $\alpha + \frac{1}{\beta} =$

A. 1

B.  $\frac{7}{12}$

C.  $\frac{19}{12}$

D.  $\frac{9}{12}$

**Answer: A**



**Watch Video Solution**

10.  $O(0,0)$ ,  $A(1,2)$ ,  $B(3,4)$  are the vertices of  $\triangle OAB$ . The joint equation of the altitude and median drawn from  $O$  is



A.  $x^2 + 7xy - y^2 = 0$

B.  $x^2 + 7xy + y^2 = 0$

C.  $3x^2 - xy - 2y^2 = 0$

D.  $3x^2 + xy - 2y^2 = 0$

**Answer: D**



**Watch Video Solution**

11.  $f(x) = \left[ \tan\left(\frac{\pi}{4} + x\right) \right]^{\frac{1}{x}}$ ,  $x \neq 0$  and  $f(x) = k$ ,  $x = 0$

is continuous at  $x=0$  then  $k=$

A.  $e$

B.  $e^{-1}$

C.  $e^2$

D.  $e^{-2}$

**Answer: C**

 [Watch Video Solution](#)

12. For an invertible matrix  $A$  if  $A(\text{adj}A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$ , then

$|A| =$

A. 100

B.  $-100$

C. 10

D.  $-10$

**Answer: C**

 [Watch Video Solution](#)

**13.** The solution of the differential equation

$$\frac{dy}{dx} = \tan\left(\frac{y}{x}\right) + \frac{y}{x} \text{ is}$$

A.  $\cos\left(\frac{y}{x}\right) = cx$

B.  $\sin\left(\frac{y}{x}\right) = cx$

C.  $\cos\left(\frac{y}{x}\right) = cy$

D.  $\sin\left(\frac{y}{x}\right) = cy$

**Answer: B**

 [Watch Video Solution](#)

14. In  $\Delta ABC$ , if  $\sin^2 A + \sin^2 B = \sin^2 C$  and  $l(AB) = 10$ , then the maximum value of the area of  $\Delta ABC$  is

A. 50

B.  $10\sqrt{2}$

C. 25

D.  $25\sqrt{2}$

**Answer: C**



**Watch Video Solution**

15. If  $x = f(t)$  and  $y = g(t)$ , then  $\frac{d^2y}{dx^2}$  is equal to

- A.  $\frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^3}$
- B.  $\frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^2}$
- C.  $\frac{g'(t) \cdot f''(t) - f'(t) \cdot g''(t)}{[f'(t)]^3}$
- D.  $\frac{g'(t) \cdot f''(t) + f'(t) \cdot g''(t)}{[f'(t)]^3}$

**Answer: A**



**Watch Video Solution**

**16.** The equation of line equally inclined to coordinate axes and passing through  $(-3, 2, -5)$  is

A.  $\frac{x + 3}{1} = \frac{y - 2}{1} = \frac{z + 5}{1}$

B.  $\frac{x + 3}{-1} = \frac{y - 2}{1} = \frac{5 + z}{-1}$

$$C. \frac{x+3}{-1} = \frac{y-2}{1} = \frac{z+5}{1}$$

$$D. \frac{x+3}{-1} = \frac{2-y}{1} = \frac{z+5}{-1}$$

**Answer: B**

 [Watch Video Solution](#)

17. If  $\int_0^{\frac{\pi}{2}} \log \cos x dx = \frac{\pi}{2} \log \left( \frac{1}{2} \right)$ , then

$$\int_0^{\frac{\pi}{2}} \log \sec x dx =$$

A.  $\frac{\pi}{2} \log \left( \frac{1}{2} \right)$

B.  $1 - \frac{\pi}{2} \log \left( \frac{1}{2} \right)$

C.  $1 + \frac{\pi}{2} \log \left( \frac{1}{2} \right)$

D.  $\frac{\pi}{2} \log 2$

**Answer: D**



**Watch Video Solution**

**18.** A boy tosses fair coin 3 times. If he gets Rs  $2X$  for  $X$  heads, then his expected gain equals to Rs.....

A. 1

B.  $\frac{3}{2}$

C. 3

D. 4

**Answer: C**



**Watch Video Solution**

19. Which of the following statement pattern is a tautology?

A.  $p \vee (q \rightarrow p)$

B.  $\sim q \rightarrow \sim p$

C.  $(q \rightarrow p) \vee (\sim p \leftrightarrow q)$

D.  $p \wedge \sim p$

**Answer: C**

 [Watch Video Solution](#)

20. If the angle between the planes

$$r \cdot (m\hat{i} - \hat{j} + 2\hat{k}) + 3 = 0 \text{ and } r \cdot (2\hat{i} - m\hat{j} - \hat{k}) - 5 = 0$$



is  $\frac{\pi}{3}$ , then  $m =$

A. 2

B.  $\pm 3$

C. 3

D.  $-2$

**Answer: C**



**Watch Video Solution**

**21.** If the origin and the points  $P(2,3,4)$ ,  $Q(1,2,3)$  and  $R(x,y,z)$  are coplanar, then

A.  $x - 2y - z = 0$

B.  $x + 2y + z = 0$

C.  $x - 2y + z = 0$

D.  $2x - 2y + z = 0$

**Answer: C**



**Watch Video Solution**

22. if lines represented by equation  $px^2 - qy^2 = 0$  are distinct, then

A.  $pq > 0$

B.  $pq < 0$

C.  $pq = 0$

D.  $p + q = 0$

**Answer: A**

 [Watch Video Solution](#)

23. Let  $\square PQRS$  be a quadrilateral. If M and N are the mid-points of the sides PQ and RS respectively, then  $PS+QR=$

A.  $3\overline{MN}$

B.  $4\overline{MN}$

C.  $2\overline{MN}$

D.  $2\overline{NM}$

**Answer: C**



**Watch Video Solution**

**24.** If slopes of lines represented by  $kx^2 + 5xy + y^2 = 0$  differ by 1, then k=

A. 2

B. 3

C. 6

D. 8

**Answer: C**



**Watch Video Solution**

25. If vector  $r$  with direction cosines  $l, m, n$  is equally inclined to the coordinate axes, then the total number of such vectors is

A. 4

B. 6

C. 8

D. 2

**Answer: C**



**Watch Video Solution**

26.

If

$$\int \frac{1}{(x^2 + 4)(x^2 + 9)} dx = A \tan^{-1} \frac{x}{2} + B \tan^{-1} \left( \frac{x}{3} \right) + C$$

, then  $A-B=$

A.  $\frac{1}{6}$

B.  $\frac{1}{30}$

C.  $-\frac{1}{30}$

D.  $-\frac{1}{6}$

**Answer: A**



**Watch Video Solution**

**27.** If  $\alpha$  and  $\beta$  are roots of the equation  $x^2 + 5|x| - 6 = 0$ , then the value of  $|\tan^{-1} \alpha - \tan^{-1} \beta|$  is

A.  $\frac{\pi}{2}$

B. 0

C.  $\pi$

D.  $\frac{\pi}{4}$

**Answer: A**



**Watch Video Solution**

28. If  $x = a\left(t - \frac{1}{t}\right)$ ,  $y = a\left(t + \frac{1}{t}\right)$ , where  $t$  be the parameter, then  $\frac{dy}{dx} = ?$

A.  $\frac{y}{x}$

B.  $\frac{-x}{y}$

C.  $\frac{x}{y}$

D.  $\frac{-y}{x}$

**Answer: C**



**Watch Video Solution**

**29.** The point on the curve  $y = \sqrt{x-1}$ , where the tangent is perpendicular to the line  $2x+y-5=0$  is

A. (2,-1)

B. (10,3)

C. (2,1)

D. (5,-2)



**Answer: C**



**Watch Video Solution**

30. If  $\int \sqrt{\frac{x-5}{x-7}} dx = A\sqrt{x^2 - 12x + 35}$

$+ \log|x - 6 + \sqrt{x^2 - 12x + 35}| + C$ , then  $A =$

A.  $-1$

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D.  $1$

**Answer: D**



**View Text Solution**

31. At random variable  $X \sim B(n, p)$  if values of mean and variance of  $X$  are 18 and 12 respectively, then total number of possible values of  $X$  are

A. 54

B. 55

C. 12

D. 18

**Answer: B**



**Watch Video Solution**

32. The area of the region bounded by the lines  $y=2x+1$ ,  $y=3x+1$  and  $x=4$  is

A. 16 sq unit

B.  $\frac{121}{3}$  sq unit

C.  $\frac{121}{6}$  sq unit

D. 8 sq unit

**Answer: D**



**Watch Video Solution**

33. A box contains 6 pens, 2 of which are defective, two pens are taken randomly from the box. If r.v.X: number of

defective pens obtained, then standard deviation of  $X=$

A.  $\pm \frac{4}{3\sqrt{5}}$

B.  $\frac{8}{3}$

C.  $\frac{16}{45}$

D.  $\frac{4}{3\sqrt{5}}$

**Answer: D**



[View Text Solution](#)

**34.** If the volume of spherical ball is increasing at the rate of  $4\pi$  cc/s, then the rate of change of its surface area when the volume is  $288\pi$  cc is

A.  $\frac{4}{3}\pi \text{ cm}^2 / s$

B.  $\frac{2}{3}\pi \text{ cm}^2 / s$

C.  $4\pi \text{ cm}^2 / s$

D.  $2\pi \text{ cm}^2 / s$

**Answer: A**



**Watch Video Solution**

**35.** If  $f(x) = \log(\sec^2 x)^{\cot^2 x}$  for  $x \neq 0$  and  $k$  for  $x = 0$  is continuous at  $x = 0$ , then  $k$  is

A.  $e^{-1}$

B. 1

C. e

D. 0

**Answer: B**

 [Watch Video Solution](#)

**36.** If  $c$  denotes the contradiction, then dual of the compound statement  $\sim p \wedge (q \vee c)$  is

A.  $\sim p \vee (q \wedge t)$

B.  $\sim p \wedge (q \vee t)$

C.  $p \vee (\sim q \vee t)$

D.  $\sim p \vee (q \wedge c)$

**Answer: A**



**Watch Video Solution**

**37.** Form the differential equation of the family of all the parabolas whose vertex is at origin and axis is y-axis.

A.  $x \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

B.  $x \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$

C.  $\frac{d^2y}{dx^2} - y = 0$

D.  $\frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$

**Answer: A**



**Watch Video Solution**

38.  $\int_0^3 [x] dx = . . .$ , where  $[x]$  is greatest integer function.

A. 3

B. 0

C. 2

D. 1

**Answer: A**

 [Watch Video Solution](#)

39. The objective function off LPP defined over the convex set attains it optimum value at



A. least two off the corner value at

B. all the corner points

C. atleast one of the corner points

D. none of the corner points

**Answer: C**



**Watch Video Solution**

40. If the inverse of the matrix  $\begin{bmatrix} \alpha & 14 & -1 \\ 2 & 3 & 1 \\ 6 & 2 & 3 \end{bmatrix}$  does not

exist, then the value of  $\alpha$  is

A. 1

B.  $-1$

C. 0

D. -2

**Answer: D**

 [Watch Video Solution](#)

41. If  $f(x) = x$  for  $x \leq 0$  and  $f(x) = 0$  for  $x > 0$ , the  $f(x)$  at  $x=0$  is

A. continuous but not differentiable

B. not continuous but differentiable

C. continuous and differentiable

D. not continuous and not differentiable

**Answer: A**



Watch Video Solution

42. The equation of the plane through  $(-1,1,2)$ , whose normal makes equal acute angles with coordinate axes is

A.  $r \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$

B.  $r \cdot (\hat{i} + \hat{j} + \hat{k}) = 6$

C.  $r \cdot (3\hat{i} - 3\hat{j} + 3\hat{k}) = 2$

D.  $r \cdot (\hat{i} - \hat{j} + \hat{k}) = 3$

**Answer: A**



Watch Video Solution

43. Probability that a person will develop immunity after vaccinations is 0.8. if 8 people are given the vaccine, then probability that all develop immunity is=

A.  $(0.2)^8$

B.  $(0.8)^8$

C. 1

D.  ${}^8C_6(0.2)^6(0.8)^2$

**Answer: B**



**Watch Video Solution**

44. If the distance of points  $2\hat{i} + 3\hat{j} + \lambda\hat{k}$  from the plane  $r \cdot (3\hat{i} + 2\hat{j} + 6\hat{k}) = 13$  is 5 units, then  $\lambda =$

A.  $6, -\frac{17}{3}$

B.  $6, \frac{17}{3}$

C.  $-6, -\frac{17}{3}$

D.  $-6, \frac{17}{3}$

**Answer: A**



**Watch Video Solution**

45. The value of  $\cos^{-1}\left(\cot\left(\frac{\pi}{2}\right)\right) + \cos^{-1}\left(\sin\left(\frac{2\pi}{3}\right)\right)$

is

A.  $\frac{2\pi}{3}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{2}$

D.  $\pi$

**Answer: A**



**Watch Video Solution**

**46.** The particular solution of the differential equation  $x \, dy + 2y \, dx = 0$ , at  $x=2, y=1$  is

A.  $xy=4$

B.  $x^2y = 4$

C.  $xy^2 = 4$

D.  $x^2y^2 = 4$

**Answer: B**

 [Watch Video Solution](#)

47.  $ABC$  is a triangle and  $A = (2, 3, 5)$ ,  $B = (-1, 3, 2)$  and  $C = (\lambda, 5, \mu)$ . If the median through  $A$  is equally inclined to the axes, then find the value of  $\lambda$  and  $\mu$ .

A. 10, 7

B. 9, 10

C. 7, 9

D. 7,10

**Answer: D**

 [Watch Video Solution](#)

**48.** For the following distribution function  $F(x)$  of a rv.X.

$x$	1	2	3	4	5	6
$F(x)$	0.2	0.37	0.48	0.62	0.85	1

$$P(3 < x \leq 5) =$$

A. 0.48

B. 0.37

C. 0.27

D. 1.47



**Answer: B**

 [Watch Video Solution](#)

49. The lines

$$\frac{x - 1}{2} = \frac{y + 1}{2} = \frac{z - 1}{4} \text{ and } \frac{x - 3}{1} = \frac{y - 6}{2} = \frac{z}{1}$$

intersect each other at point

A. (-2,-4,5)

B. (-2,-4,-5)

C. (2,4,-5)

D. (2,-4,-5)

**Answer: B**

 [Watch Video Solution](#)

50.  $\int \frac{\sec^8 x}{\operatorname{cosec} x} dx =$

A.  $\frac{\sec^8 x}{8} + c$

B.  $\frac{\sec^7 x}{7} + c$

C.  $\frac{\sec^6 x}{6} + c$

D.  $\frac{\sec^9 x}{9} + c$

**Answer: B**



**Watch Video Solution**