



## MATHS

# BOOKS - MHTCET PREVIOUS YEAR PAPERS AND PRACTICE PAPERS

## SOLVED PAPER 2018

### Mcqs

1. If  $\int_0^k \frac{dx}{2 + 18x^2} = \frac{\pi}{24}$ , then the value of k is

A. 3

B. 4

C.  $\frac{1}{3}$

D.  $\frac{1}{4}$

**Answer: C**



**Watch Video Solution**

2. The cartesian coordinates of the point on the parabola

$y^2 = -16x$ , whose parameter is  $\frac{1}{2}$ , are

A. (-2,4)

B. (4,-1)

C. ( - 1, - 4)

D. ( - 1, 4)

**Answer: C**

 [Watch Video Solution](#)

3.  $\int \frac{1}{\sin x \cdot \cos^2 x} dx =$

A.  $\sec x + \log|\sec x + \tan x| + C$

B.  $\sec x \cdot \tan x + C$

C.  $\sec x + \log|\sec x - \tan x| + C$

D.  $\sec x + \log|\operatorname{cosec} x - \cot x| + C$

**Answer: D**

 [Watch Video Solution](#)

4. If  $\log_{10} \left( \frac{x^3 - y^3}{x^3 + y^3} \right) = 2$ , then  $\frac{dy}{dx} =$

A.  $\frac{x}{y}$

B.  $-\frac{y}{x}$

C.  $-\frac{x}{y}$

D.  $\frac{y}{x}$

**Answer: D**



**Watch Video Solution**

5. If  $f: R - \{2\} \rightarrow R$  is a function defined by

$f(x) = \frac{x^2 - 4}{x - 2}$ , then its range is

A.  $R$

B.  $R - \{2\}$

C.  $R - \{4\}$

D.  $R - \{-2, 2\}$

**Answer: C**

 [Watch Video Solution](#)

6. If  $f(x) = \begin{cases} x^2 + \alpha & \text{for } x \geq 0 \\ 2\sqrt{x^2 + 1} + \beta & \text{for } x < 0 \end{cases}$  is continuous at  $x=0$  and  $f\left(\frac{1}{2}\right) = 2$ , then  $\alpha^2 + \beta^2$  is

A. 3

B.  $\frac{8}{25}$

C.  $\frac{25}{8}$

D.  $\frac{1}{3}$

**Answer: C**

 [Watch Video Solution](#)

7. If  $y = (\tan^{-1} x)^2$ , then

$$(x^2 + 1)^2 \frac{d^2y}{dx^2} + 2x(x^2 + 1) \frac{dy}{dx} =$$

A. 4

B. 2

C. 1

D. 0

**Answer: A**

 [Watch Video Solution](#)

8. The line  $5x + y - 1 = 0$  coincides with one of the lines given by  $5x^2 + xy - kx - 2y + 2 = 0$ , then the value of  $k$  is

A.  $-11$

B.  $31$

C.  $11$

D.  $-31$

**Answer: C**

 [Watch Video Solution](#)

9. If  $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}$ , then  $(A^2 - 5A)A^{-1} =$

A.  $\begin{bmatrix} 4 & 2 & 3 \\ -1 & 4 & 2 \\ 1 & 2 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} -4 & 2 & 3 \\ -1 & -4 & 2 \\ 1 & 2 & -1 \end{bmatrix}$

C.  $\begin{bmatrix} -4 & -1 & 1 \\ 2 & -4 & 2 \\ 3 & 2 & -1 \end{bmatrix}$

D.  $\begin{bmatrix} -1 & -2 & 1 \\ 4 & -2 & -3 \\ 1 & 4 & -2 \end{bmatrix}$

**Answer: B**



**Watch Video Solution**



10. The equation of line passing through (3,-1,2) and perpendicular to the lines

$$\vec{r} = (\hat{i} + \hat{j} - \hat{k}) + \lambda(2\hat{i} - 2\hat{j} + \hat{k})$$

and  $\vec{r} = (2\hat{i} + \hat{j} - 3\hat{k}) + \mu(\hat{i} - 2\hat{j} + 2\hat{k})$  is

A.  $\frac{x+3}{2} = \frac{y+1}{3} = \frac{z-2}{2}$

B.  $\frac{x-3}{3} = \frac{y+1}{2} = \frac{z-2}{2}$

C.  $\frac{x-3}{2} = \frac{y+1}{3} = \frac{z-2}{2}$

D.  $\frac{x-3}{2} = \frac{y+1}{2} = \frac{z-2}{3}$

**Answer: C**



**Watch Video Solution**

11. Letters in the word HULULULU are rearranged. The probability of all three L being together is

A.  $\frac{3}{20}$

B.  $\frac{2}{5}$

C.  $\frac{3}{28}$

D.  $\frac{5}{23}$

**Answer: C**



**Watch Video Solution**

12. The sum of the first 10 terms of the series  $9+99+999+ \dots$

Is

A.  $\frac{9}{8}(9^{10} - 1)$

B.  $\frac{100}{9}(10^9 - 1)$

C.  $10^9 - 1$

D.  $\frac{100}{9}(10^{10} - 1)$

**Answer: B**



**Watch Video Solution**

13. If  $A, B, C$  are the angles of  $\triangle ABC$ , then  
 $\cot A \cdot \cot B + \cot B \cdot \cot C + \cot C \cdot \cot A =$

A. 0

B. 1

C. 2

D. - 1

**Answer: B**

 [Watch Video Solution](#)

14. if  $\int \frac{dx}{\sqrt{16 - 9x^2}} = A \sin^{-1}(Bx) + C$ , then  $A+B=$

A.  $\frac{9}{4}$

B.  $\frac{19}{4}$

C.  $\frac{3}{4}$

D.  $\frac{13}{12}$

**Answer: D**

 [Watch Video Solution](#)

15. Evaluate:  $\int e^x \left( \frac{2 + \sin 2x}{1 + \cos 2x} \right) dx$

A.  $e^x \tan x + C$

B.  $e^x + \tan x + C$

C.  $2e^x \tan x + C$

D.  $e^x \tan 2x + C$

**Answer: A**

 [Watch Video Solution](#)

16. A coin is tossed three times. If  $X$  denotes the absolute difference between the number of heads and the number of tails, then  $P(X=1)=$

A.  $\frac{1}{2}$

B.  $\frac{2}{3}$

C.  $\frac{1}{6}$

D.  $\frac{3}{4}$

**Answer: D**



**Watch Video Solution**

17. If  $2 \sin\left(\theta + \frac{\pi}{3}\right) = \cos\left(\theta - \frac{\pi}{6}\right)$ , then  $\tan \theta =$

A.  $\sqrt{3}$

B.  $-\frac{1}{\sqrt{3}}$

C.  $\frac{1}{\sqrt{3}}$

D.  $-\sqrt{3}$

**Answer: D**

 [Watch Video Solution](#)

**18.** The area of the region bounded by  $x^2 = 4y$ ,  $y = 1$ ,  $y = 4$  and the Y-axis lying in the first quadrant is ..... Square units.

A.  $\frac{22}{3}$

B.  $\frac{28}{3}$

C. 30

D.  $\frac{27}{4}$

**Answer: B**



**Watch Video Solution**

19. if  $f(x) = \frac{e^{x^2} - \cos x}{x^2}$ , for  $x \neq 0$  is continuous at  $x = 0$

, then value of  $f(0)$  is

A.  $\frac{2}{3}$

B.  $\frac{5}{2}$

C. 1

D.  $\frac{3}{2}$

**Answer: D**



**Watch Video Solution**



20. The maximum value of  $Z = 2x + y$  subject to  $3x + 5y \leq 26$  and  $5x + 3y \leq 30, x \geq 0, y \geq 0$  is

A. 12

B. 11.5

C. 10

D. 17.33

**Answer: A**



[View Text Solution](#)

21. If  $\vec{a}, \vec{b}, \vec{c}$  are mutually perpendicular vectors having magnitudes 1, 2, 3 respectively, then

$$\left[ \vec{a} + \vec{b} + \vec{c} \quad \vec{b} - \vec{a} \quad \vec{c} \right] =$$

A. 0

B. 6

C. 12

D. 18

**Answer: C**



[Watch Video Solution](#)

22. If points  $P(4, 5, x)$ ,  $Q(3, y, 4)$  and  $R(5, 8, 0)$  are collinear, then the value of  $x+y$  is

A.  $-4$

B. 3

C. 5

D. 4

**Answer: D**



**Watch Video Solution**

**23.** If the slope of one of the lines given by  $ax^2 + 2hxy + by^2 = 0$  is two times the other, then

A.  $8h^2 = 9ab$

B.  $8h^2 = 9ab^2$

C.  $8h = 9ab$

D.  $8h = 9ab^2$

**Answer: A**



**Watch Video Solution**

**24.** The equation of the line passing through the point  $(-3,1)$  and bisecting the angle between coordinate axes is

A.  $x+y+2=0$

B.  $-x + y + 2 = 0$

C.  $x - y + 4 = 0$

D.  $2x + y + 5 = 0$

**Answer: A**



**Watch Video Solution**

25. The negation of the statement: "Getting above 95% marks is necessary condition for Hema to get the admission in good college"

A. Hema gets above 95% marks but she does not get the admission in good college

B. Hema does not get above 95% marks and she gets admission in good college

C. If hema does not get above 95% marks then she will not get the admission in good college

D. Hema does not get above 95% marks or she gets the admission in good college

**Answer: B**



Watch Video Solution

26.  $\cos 1^\circ \cdot \cos 2^\circ \cdot \cos 3^\circ \dots \dots \cos 179^\circ$  is equal to :

A. 0

B. 1

C.  $-\frac{1}{2}$

D.  $-1$

**Answer: A**



Watch Video Solution

27. If the planes

$$x - cy - bz = 0, cx = y + az = 0 \text{ and } bx + ay - z = 0$$

pass through a straight line, then find the value of

$$a^2 + b^2 + c^2 + 2ab.$$

A.  $1 - abc$

B.  $abc - 1$

C.  $1 - 2abc$

D.  $2abc - 1$

**Answer: C**



**Watch Video Solution**

28. The point of intersection of lines represented by

$$x^2 - y^2 + x + 3y - 2 = 0$$
 is

A. (1,0)

B. (0,2)

C.  $\left(-\frac{1}{3}, \frac{3}{2}\right)$

D.  $\left(\frac{1}{2}, \frac{1}{2}\right)$

**Answer: C**



**Watch Video Solution**

29. A die is rolled. If X denotes the number of positive divisors of the outcome o the outcome, then the range of the random variable X is



A. {1,2,3}

B. {1,2,3,4}

C. {1,2,3,4,5,6}

D. {1,3,5}

**Answer: C**



**Watch Video Solution**

**30.** A die is thrown four times. The probability of getting perfect square in at least one throw is

A.  $\frac{16}{81}$

B.  $\frac{65}{81}$

C.  $\frac{23}{81}$

D.  $\frac{58}{81}$

**Answer: B**

 [Watch Video Solution](#)

31.  $\int_0^{\pi/4} x \cdot \sec^2 x dx =$

A.  $\frac{\pi}{4} \log \sqrt{2}$

B.  $\frac{\pi}{4} - \log \sqrt{2}$

C.  $1 + \log \sqrt{2}$

D.  $1 - \frac{1}{2} \log 2$

**Answer: B**

 [Watch Video Solution](#)

32. In  $\triangle ABC$ , with usual notations, if  $a, b, c$  are in AP then a

$$\cos^2\left(\frac{C}{2}\right) + c \cos^2\left(\frac{A}{2}\right) =$$

A.  $\frac{3a}{2}$

B.  $\frac{3c}{2}$

C.  $\frac{3b}{2}$

D.  $\frac{3abc}{2}$

**Answer: C**



**Watch Video Solution**

33. If  $x = e^\theta(\sin \theta - \cos \theta)$ ,  $y = e^\theta(\sin \theta + \cos \theta)$ , then  $\frac{dy}{dx}$  at  $\theta = \frac{\pi}{4}$  is

A. 1

B. 0

C.  $\frac{1}{\sqrt{2}}$

D.  $\sqrt{2}$

**Answer: A**



**Watch Video Solution**

34. The number of solutions of  $\sin x + \sin 3x + \sin 5x = 0$  in the interval  $\left[\frac{\pi}{2}, 3\frac{\pi}{2}\right]$  is

A. 2

B. 3

C. 4

D. 5

**Answer: B**



**Watch Video Solution**

35. Solve :  $\tan^{-1} 2x + \tan^{-1} 3x = \frac{\pi}{4}$

A.  $-1$

B.  $\frac{1}{3}$

C.  $\frac{1}{6}$

D.  $\frac{1}{2}$

**Answer: C**

 [Watch Video Solution](#)

36. Matrix  $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ , then the value of  $a_{31}A_{31} + a_{32}A_{32} + a_{33}A_{33}$  is

A. 1

B. 13

C. -1

D. -13

**Answer: C**



37. The contrapositive of the statement: "If the weather is fine then my friends will come and we go for a picnic".

A. The weather is fine but my friends will not come or we do not go for a picnic

B. if my friends do not come or we do not go for picnic then weather will not be fine

C. If the weather is not fine then my friends will not come or we do not go for a picnic

D. The weather is not fine but my friends will come and we go for a picnic

**Answer: B**



**Watch Video Solution**

**38.** If  $f(x) = \frac{x}{x^2 + 1}$  is increasing function, then the value

of  $x$  lies in

A.  $\mathbb{R}$

B.  $(-\infty, -1)$

C.  $(1, \infty)$

D.  $(-1, 1)$

**Answer: D**



**Watch Video Solution**



39. If  $X = \{4^n - 3n - 1 : n \in \mathbb{N}\}$  and  $\{9(n - 1) : n \in \mathbb{N}\}$ ,  
the prove that  $X \subset Y$ .

A.  $X$

B.  $Y$

C.  $\phi$

D.  $\{0\}$

**Answer: A**

 [Watch Video Solution](#)

40. The statement pattern  $p \wedge (\sim p \wedge q)$  is

A. a tautology

B. a contradiction

C. equivalent to  $p \wedge q$

D. equivalent to  $p \vee q$

**Answer: B**



**Watch Video Solution**

**41.** If the line  $y = 4x - 5$  touches to the curve  $y^2 = ax^3 + b$  at the point  $(2,3)$ , then  $7a+2b=$

A. 0

B. 1

C.  $-1$

D. 2

**Answer: A**

 [Watch Video Solution](#)

**42.** The sides of a rectangle are given by  $x = \pm a$  and  $y = \pm b$ . The equation of the circle passing through the vertices of the rectangle is

A.  $x^2 + y^2 = a^2$

B.  $x^2 + y^2 = a^2 + b^2$

C.  $x^2 + y^2 = a^2 - b^2$

D.  $(x - a)^2 + (y - b)^2 = a^2 + b^2$

**Answer: B**

 [Watch Video Solution](#)

43. The minimum value of the function  $f(x)=x \log x$  is

A.  $-\frac{1}{e}$

B.  $-e$

C.  $\frac{1}{e}$

D.  $e$

**Answer: A**



**Watch Video Solution**

44. if  $X \sim B(n, p)$  with  $n = 10, p = 0.4$ , then  $E(X^2) =$

A. 4

B. 2.4

C. 3.6

D. 18.4

**Answer: D**



**Watch Video Solution**

**45.** The general solution of differential equation

$$\frac{dx}{dy} = \cos(x + y) \text{ is}$$

A.  $\tan\left(\frac{x + y}{2}\right) = y + C$

B.  $\tan\left(\frac{x + y}{2}\right) = x + C$

C.  $\cot\left(\frac{x + y}{2}\right) = y + C$

D.  $\cot\left(\frac{x + y}{2}\right) = x + C$

**Answer: A**

 [Watch Video Solution](#)

**46.** If planes

$$\vec{r} \cdot (p\hat{i} - \hat{j} + 2\hat{k}) + 3 = 0 \text{ and } \vec{r} \cdot (2\hat{i} - p\hat{j} - \hat{k}) - 5 = 0$$

include angle  $\frac{\pi}{3}$ , then the value of  $p$

A. 1,-3

B. -1, 3

C. -3

D. 3

**Answer: D**

 [Watch Video Solution](#)

47. Find the differential equation of all the parabolas with latus rectum ' $4a$ ' and whose axes are parallel to x-axis.

- A. one
- B. four
- C. three
- D. two

**Answer: D**



**Watch Video Solution**

48. If lines

$$\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4} \text{ and } x-3 = \frac{y-k}{2} = z$$

intersect, then the value of  $k$  is

A.  $\frac{9}{2}$

B.  $\frac{1}{2}$

C.  $\frac{5}{2}$

D.  $\frac{7}{2}$

**Answer: A**



**Watch Video Solution**



49. If a line makes angle  $120^\circ$  and  $60^\circ$  with the positive directions of X and Z-axes respectively, then the angle made by the line with positive Y-axis is

A.  $150^\circ$

B.  $60^\circ$

C.  $135^\circ$

D.  $120^\circ$

**Answer: C**



**Watch Video Solution**

50. L and M are two points with position vectors  $2\vec{a} - \vec{b}$  and  $\vec{a} + 2\vec{b}$ , respectively. The position vector of the point N which divides the line segment LM in the ratio 2:1 externally is

A.  $3\vec{b}$

B.  $4\vec{b}$

C.  $5\vec{b}$

D.  $3\vec{a} + 4\vec{b}$

**Answer: C**



**Watch Video Solution**