



# PHYSICS

## BOOKS - MHTCET PREVIOUS YEAR PAPERS AND PRACTICE PAPERS

### MHTCET 2017

#### Physics

1. Let  $v_1$  be the frequency of series limit of Lyman series,  $v_2$  the frequency of the first line

of Lyman series and  $\nu_3$  the frequency of series limit of Balmer series. Then which of the following is correct ?

A.  $\nu_1 - \nu_2 = \nu_3$

B.  $\nu_1 \nu_3 = \nu_2$

C.  $\nu_1 + \nu_2 = \nu_3$

D.  $\nu_1 - \nu_3 = 2\nu_1$

**Answer: A**



**Watch Video Solution**

2. When three capacitors of equal capacities are connected in parallel and one of the same capacity is connected in series with its combination . The resultant capacity is  $3.75\mu F$  . The capacity of each capacitor is

A.  $5\mu F$

B.  $6\mu F$

C.  $7\mu F$

D.  $8\mu F$

**Answer: C**



3. Sensitivity of moving coil galvanometer is 'S'.

If a shunt of  $\left(\frac{1}{6}\right)$  th of the resistant of galvanometer is connected to moving coil galvanometer, its sensitivity becomes.

A.  $\frac{S}{3}$

B.  $\frac{S}{7}$

C.  $\frac{S}{9}$

D.  $\frac{S}{12}$

**Answer: C**



**Watch Video Solution**

4. Two resistances are connected in the two gaps of a meter bridge. The balance point is  $20\text{cm}$  from the zero end. When a resistance  $15\Omega$  is connected in series with the smaller of two resistance, the null point shifts to  $40\text{cm}$ . The smaller of the two resistance has the value.

A. 12

B. 24

C. 36

D. 48

**Answer: B**



**Watch Video Solution**

5. In Fraunhofer diffraction pattern, slit width is  $0.2\text{mm}$  and screen is at 2 m away from the lens. If wavelength of light used is  $5000\text{\AA}$ , then

the distance between the first minimum on either side of the central maximum is ( $\theta$  is small and measured in radian)

A.  $10^{-1}m$

B.  $10^{-2}m$

C.  $2 \times 10^{-2}m$

D.  $2 \times 10^{-1}m$

**Answer: B**



**Watch Video Solution**

6. In series LCR circuit  $R = 18\Omega$  and impedance is  $33\Omega$ . An rms voltage  $220V$  is applied across the circuit. The true power consumed in AC circuit is

A.  $220W$

B.  $400W$

C.  $600W$

D.  $800W$

**Answer: D**



**Watch Video Solution**



7. Two parallel plate air capacitance of same capacity  $C$  are connected in series to a battery of emf  $E$ . Then one of the capacitors is completely filled with dielectric material of constant  $K$ . The change in the effective capacity of the series combination is

A.  $\frac{C}{2} \left[ \frac{K - 1}{K + 1} \right]$

B.  $\frac{2}{C} \left[ \frac{K - 1}{K + 1} \right]$

C.  $\frac{C}{2} \left[ \frac{K + 1}{K - 1} \right]$

D.  $\frac{C}{2} \left[ \frac{K - 1}{K + 1} \right]^2$

**Answer: A**



**Watch Video Solution**

8. The polarising angle the transparent medium is ' $\theta$ ' and ' $v$ ' is the speed of light in that medium. Then the relation between ' $\theta$ ' and ' $v$ ' is

( $c$  = velocity of light in air)

A.  $\theta = \tan^{-1} \left( \frac{v}{c} \right)$

B.  $\theta = \cot^{-1} \left( \frac{v}{c} \right)$

$$C. \theta = \sin^{-1}\left(\frac{v}{c}\right)$$

$$D. \theta = \cos^{-1}\left(\frac{v}{c}\right)$$

**Answer: B**



**Watch Video Solution**

9. Two identical light waves having phase difference ' $\phi$ ' propagate in same direction. When they superpose, the intensity of resultant wave is proportional to

A.  $\cos^2 \phi$

B.  $\cos^2, \frac{\phi}{2}$

C.  $\cos^2, \frac{\phi}{3}$

D.  $\cos^2, \frac{\theta}{4}$

**Answer: B**



**Watch Video Solution**

**10.** For a transistor ,  $\alpha_{dc}$  and  $\beta_{dc}$  are the current ratios, then the value of  $\frac{\beta_{dc} - \delta_{dc}}{\alpha_{dc} \cdot \beta_{dc}}$

A. 1

B. 1.5

C. 2

D. 2.5

**Answer: A**



**Watch Video Solution**

**11.** A radioactive element has rate of disintegration 10,000 disintegrations per minute at a particular instant. After four

minutes it becomes 2500 disintegrations per minute. The decay constant per minute is

A.  $0.2 \log_e 2$

B.  $0.5 \log_e 2$

C.  $0.6 \log_e 2$

D.  $0.8 \log_e 2$

**Answer: B**



**Watch Video Solution**

12. When the same monochromatic ray of light travels through glass slab and through water, the number of waves in glass slab of thickness 6 cm is same as in water column of height 7 cm. If refractive index of glass is 1.5, then refractive index of water is

A. 1.258

B. 1.269

C. 1.286

D. 1.310

**Answer: C**



**Watch Video Solution**

**13.** If the electron in hydrogen atom jumps from second Bohr orbit to ground state and difference between energies of the two states is radiated in the form of photons. If the work function of the material is  $4.2eV$ , then stopping potential is

[Energy of electron in  $n$ th orbit  
 $= -\frac{13.6}{n^2}eV$ ]



A. 2V

B. 4V

C. 6 V

D. 8 V

**Answer: C**



**Watch Video Solution**

**14.** The magnetic moment of electron due to orbital motion is proportional to  
( $n$  = principle quantum numbers)

A.  $\frac{1}{n^2}$

B.  $\frac{1}{n}$

C.  $n^2$

D.  $n$

**Answer: D**



**Watch Video Solution**

**15. Photodiode is a device**

A. which is always operated in reverse bias

- B. which is always operated in forward bias
- C. in which photo current is independent of intensity of incident radiation
- D. which may be operated in forward or reverse bias.

**Answer: A**



**Watch Video Solution**

16. A wheel of moment of inertia  $2\text{kgm}^2$  is rotating about an axis passing through centre and perpendicular to its plane at a speed  $60\text{rad/s}$ . Due to friction, it comes to rest in 5 minutes. The angular momentum of the wheel three minutes before it stops rotating is

A.  $24\text{kgm}^2 / \text{s}$

B.  $48\text{kgm}^2 / \text{s}$

C.  $72\text{kgm}^2 / \text{s}$

D.  $96\text{kgm}^2 / \text{s}$

**Answer: C**



**Watch Video Solution**

**17.** The equation of the progressive wave is

$$y = 3 \sin \left[ \pi \left( \frac{t}{3} - \frac{x}{5} \right) + \frac{\pi}{4} \right], \text{ where } x \text{ and } y$$

are in metre and time in second. Which of the following is correct.

A. Velocity  $v = 1.5 \text{ m/s}$

B. Amplitude  $A = 3 \text{ cm}$

C. Frequency  $f = 0.2 \text{ Hz}$

D. Wavelength  $\lambda = 10\text{m}$

**Answer: D**



**Watch Video Solution**

**18.** Two spherical black bodies of radii  $R_1$  and  $R_2$  and with surface temperature  $T_1$  and  $T_2$  respectively radiate the same power.  $R_1 / R_2$  must be equal to

A.  $\frac{T_1}{T_2}$

B.  $\frac{T_2}{T_1}$

C.  $\left(\frac{T_1}{T_2}\right)^3$

D.  $\left(\frac{T_2}{T_1}\right)^2$

**Answer: C**



**Watch Video Solution**

**19.** The closed and open organ pipes have same length. When they are vibrating simultaneously in first overtone, produce three beats. The length of open pipe is made

$\frac{1}{3}$  rd and closed pipe is made thre time the original , the number of beats produced will be

A. 8

B. 14

C. 17

D. 20

**Answer: C**



**Watch Video Solution**



20. A lift of mass 'm' is connected to a rope which is moving upward with maximum acceleration 'a'. For maximum safe stress, the elastic limit of the rope is 'T'. The minimum diameter of the rope is

(g = gravitational acceleration)

A.  $\left[ \frac{2m(g + a)}{\pi T} \right]^{\frac{1}{2}}$

B.  $\left[ \frac{4m(g + a)}{\pi T} \right]^{\frac{1}{2}}$

C.  $\left[ \frac{m(g + a)}{\pi T} \right]^{\frac{1}{2}}$

D.  $\left[ \frac{m(g + a)}{2\pi T} \right]^{\frac{1}{2}}$

**Answer: B**



**Watch Video Solution**

21. A solid sphere of mass 2 kg is rolling on a frictionless horizontal surface with velocity  $6m/s$ . It collides on the free end of an ideal spring whose other end is fixed. The maximum compression produced in the spring will be (Force constant of the spring = 36 N/m)

A.  $\sqrt{14}m$

B.  $\sqrt{2.8}m$

C.  $\sqrt{14}m$

D.  $\sqrt{0.7}m$

**Answer: B**



**Watch Video Solution**

22. A flywheel at rest is to reach an angular velocity of 24 rad/s in 8 second with constant angular acceleration. The total angle turned through during this interval is

A. 24 rad

B. 48 rad

C. 72 rad

D. 96 rad

**Answer: D**



**Watch Video Solution**

**23.** Two uniform wires of a the same material are vibrating under the same tension. If the first overtone of the first wire is equal to the

second overtone of the second wire and radius of the first wire is twice the radius of the second wire, then the ratio of the lengths of the first wire to second wire is

A.  $\frac{1}{3}$

B.  $\frac{1}{4}$

C.  $\frac{1}{5}$

D.  $\frac{1}{6}$

**Answer: A**



**Watch Video Solution**

24. When one end of the capillary is dipped in water, the height of water column is ' $h$ '. The upward force of 105 dyne due to surface tension is balanced by the force due to the weight of water column . The inner circumference of the capillary is

(Surface tension of water =  $7 \times 10^{-2} N/m$ )

A. 1.5 cm

B. 2 cm

C. 2.5 cm

D. 3 cm

**Answer: A**



**Watch Video Solution**

**25.** For a rigid diatomic molecule, universal gas constant  $R = nC_p$ , where ' $C_p$ ' is the molar specific heat at constant pressure and 'n' is a number. Hence n is equal to

**A. 0.2257**

B. 0.4

C. 0.2857

D. 0.3557

**Answer: C**



**Watch Video Solution**

**26.** A ideal gas has pressure ' $p$ ', volume ' $V$ ' and absolute temperature ' $T$ '. It ' $m$ ' is the mass of each molecules and ' $K$ ' is the Boltzmann constant , the density of the gas is



A.  $\frac{\text{pm}}{MT}$

B.  $\frac{KT}{\text{pm}}$

C.  $\frac{Km}{\text{pT}}$

D.  $\frac{pK}{Tm}$

**Answer: A**



**Watch Video Solution**

**27.** A big water drop is formed by the combination of 'n' small water drops of equal

radii. The ratio of the surface energy of 'n' drops to the surface energy of big drop is

A.  $n^2 : 1$

B.  $n : 1$

C.  $\sqrt{n} : 1$

D.  ${}^3\sqrt{n} : 1$

**Answer: D**



**Watch Video Solution**

28. The ratio of binding energy of a satellite at rest on earth's surface to the binding energy of a satellite of same mass revolving around of the earth at a height  $h$  above the earth's surface is ( $R$  = radius of the earth).

A.  $\frac{2(R + h)}{R}$

B.  $\frac{R + h}{2}$

C.  $\frac{R + h}{R}$

D.  $\frac{R}{R + h}$

**Answer: A**



Watch Video Solution

29. A particle performing SHM starts equilibrium position and its time period is 16 seconds. After 2 seconds its velocity is  $\pi m / s$ .

Amplitude of oscillation is

$$\left( \cos 45^\circ = \frac{1}{\sqrt{2}} \right)$$

A.  $2\sqrt{2}m$

B.  $4\sqrt{2}m$

C.  $6\sqrt{2}m$

D.  $8\sqrt{2}m$

**Answer: D**



**Watch Video Solution**

**30.** In sonometer experiment , the string of length 'L' under tension vibrates in second overtone between two bridges. The amplitude of vibration is maximum at

A.  $\frac{L}{3}, \frac{2L}{3}, \frac{5L}{6}$

B.  $\frac{L}{8}, \frac{L}{4}, \frac{L}{2}$

C.  $\frac{L}{2}, \frac{L}{4}, \frac{L}{6}$

D.  $\frac{L}{6}, \frac{L}{2}, \frac{5L}{6}$

**Answer: D**



**Watch Video Solution**

**31.** The depth at which the value of acceleration due to gravity is  $\frac{1}{n}$  times the value at the surface, is (R=radius of the earth)

$$A. d = R \left( \frac{n}{n-1} \right)$$

$$B. d = R \left( \frac{n-1}{2n} \right)$$

$$C. d = R \left( \frac{n-1}{n} \right)$$

$$D. d = R^2 \left( \frac{n-1}{n} \right)$$

**Answer: C**



**Watch Video Solution**

**32.** A particle performing SHM starting extreme position. Graphical representation

shows that, between displacement and acceleration, there is a phase difference of

A. 0 rad

B.  $\frac{\pi}{4}$  rad

C.  $\frac{\pi}{2}$  rad

D.  $\pi$  rad

**Answer: D**



**Watch Video Solution**



**33.** The fundamental frequency of an air column in a pipe closed at one end is 100 Hz. If the same pipe is open at both the ends, the frequencies produced in Hz are

- A. 100, 200, 300, 400, . . . . .
- B. 100, 300, 500, 700, . . . . .
- C. 200, 300, 400, 500, . . . . .
- D. 200, 400, 600, 800, . . . . .

**Answer: D**



**Watch Video Solution**

34. For a particle moving in vertical circle, the total energy at different positions along the path

- A. is conserved
- B. increases
- C. decreases
- D. may increases or decreases

**Answer: A**





**35.** A simple pendulum of length 'L' has mass 'M' and it oscillates freely with amplitude energy is

(g = acceleration due to gravity)

A.  $\frac{MgA^2}{2L}$

B.  $\frac{MgA}{2L}$

C.  $\frac{MgA^2}{L}$

D.  $\frac{2MgA^2}{L}$

**Answer: A**



**Watch Video Solution**

**36.** On a photosensitive material, when frequency of incident radiation is increased by 30% kinetic energy of emitted photoelectrons increases from  $0.4eV$  to  $0.9eV$ . The work function of the surface is

A.  $1 eV$

B.  $1.267eV$

C.  $1.4eV$

D.  $1.8eV$

**Answer: B**



**Watch Video Solution**

**37.** Out of the following graphs, which graph shows the correct relation (graphical representation) for LC parallel resonant circuit ?

A. 

B. 

C. 

D. 

**Answer: D**



[View Text Solution](#)

**38.** According to de-Broglie hypothesis, the wavelength associated with moving electron of mass 'm' is ' $\lambda_e$ '. Using mass energy relation

and Planck's quantum theory, the wavelength associated with photon is ' $\lambda_p$ '. If the energy (E) of electron and photon is same, then relation between  $\lambda_e$  and ' $\lambda_p$ ' is

A.  $\lambda_p \propto \lambda_e$

B.  $\lambda_p \propto \lambda_e^2$

C.  $\lambda_p \propto \sqrt{\lambda_e}$

D.  $\lambda_p \propto \frac{1}{\lambda_e}$

**Answer: A**



**Watch Video Solution**

39. A parallel plate air capacity ' $C$ ' farad, potential ' $V$ ' volt and energy ' $E$ ' joule . When the gap between the plates is completely filled with dielectric

- A. both  $V$  and  $E$  increase
- B. both  $V$  and  $E$  decrease
- C.  $V$  decrease,  $E$  increases
- D.  $V$  increases.  $E$  decrease

**Answer: B**





40. The resistivity of a potentiometer wire is  $40 \times 10^{-8} \Omega - m$  and its area of cross section is  $8 \times 10^{-6} m^2$ . If 0.2 A current is flowing through the wire the potential gradient will be

A.  $10^{-1} v / m$

B.  $10^{-2} V / m$

C.  $10^{-3} V / m$

D.  $10^{-4} V / m$

**Answer: B**



**Watch Video Solution**

**41.** A ceiling fan rotates about its own axis with some angular velocity. When the fan is switched off, the angular velocity becomes  $\left(\frac{1}{4}\right)$ th of the original in time 't' and 'n' revolutions are made in that time. The number of revolutions made by the fan during the time interval between switch off and rest are (Angular retardation is uniform)

A.  $\frac{4n}{15}$

B.  $\frac{8n}{15}$

C.  $\frac{16n}{15}$

D.  $\frac{32n}{15}$

**Answer: C**



**Watch Video Solution**

**42.** A disc of the moment of inertia ' $I_1$ ' is rotating in horizontal plane about an axis passing through a centre and perpendicular

to its plane with constant angular speed ' $\omega_1$ '.  
. Another disc of moment of inertia ' $I_2$ ',  
having zero angular speed is placed discs are  
rotating disc. Now, both the discs are rotating  
with constant angular speed ' $\omega_2$ '. The energy  
lost by the initial rotating disc is

A.  $\frac{1}{2} \left[ \frac{l_1 + l_2}{l_1 l_2} \right] \omega_1^2$

B.  $\frac{1}{2} \left[ \frac{l_1 l_2}{l_1 - l_2} \right] \omega_1^2$

C.  $\frac{1}{2} \left[ \frac{l_1 - l_2}{l_1 l_2} \right] \omega_1^2$

D.  $\frac{1}{2} \left[ \frac{l_2 l_2}{l_1 + l_2} \right] \omega_1^2$

**Answer: D**



Watch Video Solution

43. A particle executes *SHM* on a straight line. At two positions, its velocities are  $u$  and  $v$  while accelerations are  $\alpha$  and  $\beta$  respectively [ $\beta > \alpha > 0$ ]. The distance between these two positions is

A.  $\frac{u^2 - v^2}{\alpha + \beta}$

B.  $\frac{u^2 + v^2}{\alpha + \beta}$

C.  $\frac{u^2 - v^2}{\alpha - \beta}$

D.  $\frac{u^2 - v^2}{\alpha - \beta}$

**Answer: A**



**Watch Video Solution**

**44.** The observer is moving with velocity  $v'_0$  towards the stationary source of sound and then after crossing moves away from the source with velocity  $v'_0$ . Assume that the medium through which the sound waves travel is at rest. If  $v$  is the velocity of sound

and  $n$  is the frequency emitted by the source, then the difference between apparent frequencies heard by the observer is

A.  $\frac{2nv_0}{v}$

B.  $\frac{nv_0}{v}$

C.  $\frac{v}{2nv_0}$

D.  $\frac{v}{nv_0}$

**Answer: A**



**Watch Video Solution**

45. A metal rod of length 'L' and cross-sectional area 'A' is heated through ' $T$ '° C. What is the force required to prevent the expansion of the rod lengthwise ?

A.  $\frac{Y A \alpha T}{(1 - \alpha T)}$

B.  $\frac{Y A \alpha T}{(1 + \alpha T)}$

C.  $\frac{(1 - \alpha T)}{Y A \alpha T}$

D.  $\frac{(1 + \alpha T)}{Y A \alpha T}$

**Answer: B**



**Watch Video Solution**



**46.** Two coils P and Q are kept near each other. When no current flows through coil P and current increases in coil Q at the rate  $10A/s$ , the emf in coil P is 15 mV. When coil Q carries no current and current of  $1.8A$  flows through coil P, the magnetic flux linked with the coil Q is

A.  $1.4mWb$

B.  $2.2mWb$

C.  $2.7mWb$

D.  $2.9mWb$

**Answer: C**



**Watch Video Solution**

47. In Young's double experiment , in air interference pattern second minimum is observed exactly in front of one slit. The distance between the two coherent source is 'd' and the distance between source and screen 'D'. The wavelength of light source used is

A.  $\frac{d^2}{D}$

B.  $\frac{d^2}{2D}$

C.  $\frac{d^2}{3D}$

D.  $\frac{d^2}{4D}$

**Answer: C**



**Watch Video Solution**

**48.** In communication system, the process of superimposing a low frequency signal on a high frequency wave is known as

A. repeater

B. attenuation

C. modulation

D. demodulation

**Answer: C**



**Watch Video Solution**

**49.** A bar magnet has length 3 cm, cross-sectional area  $2\text{cm}^3$  and magnetic moment

$3Am^2$ . The intensity of magnetisation of bar magnet is

A.  $2 \times 10^5 A / m$

B.  $3 \times 10^5 A / m$

C.  $4 \times 10^5 A / m$

D.  $5 \times 10^7 A / m$

**Answer: D**



**Watch Video Solution**

50. The magnetic flux near the axis and inside the air core solenoid of length  $60\text{cm}$  carrying current 'f' is  $1.57 \times 10^{-6}\text{Wb}$ . Its magnetic moment will be (cross-sectional area of a solenoid is very small as compared to its length.

$$\mu_0 = 4\pi \times 10^{-7}\text{SI unit )}$$

A. 0.25 A

B. 0.50A

C. 0.75A

D. 1A

**Answer: C**



**Watch Video Solution**