



MATHS

BOOKS - CENGAGE MATHS (ENGLISH)

AREA

Illustration

1. Find the area of the closed figure bounded by the curves

$$y = \sqrt{x}, y = \sqrt{4 - 3x} \text{ and } y = 0$$



[Watch Video Solution](#)

2. Find the area, lying above the x -axis and included between the circle

$$x^2 + y^2 = 8x \text{ and the parabola } y^2 = 4x.$$



[Watch Video Solution](#)

3. Find the area bounded by

(i) $y = \log_e|x|$ and $y = 0$

(ii) $y = |\log_e|x||$ and $y = 0$



Watch Video Solution

4. Find the area bounded by $y = \frac{1}{x^2 - 2x + 2}$ and x-axis.



Watch Video Solution

5. find the area of $y = 3 - 2x - x^2$



Watch Video Solution

6. If the area of bounded between the x-axis and the graph of $y = 6x - 3x^2$ between the ordinates $x = 1$ and $x = a$ is 19 units, then a

can take the value 4 or -2 two value are in (2,3) and one in $(-1, 0)$

two value are in (3,4) and one in $(-2, -1)$ none of these

 [Watch Video Solution](#)

7. Prove that area common to ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and its auxiliary circle

$x^2 + y^2 = a^2$ is equal to the area of another ellipse of semi-axis a and $a - b$.

b.

 [Watch Video Solution](#)

8. Let $f(x) = \text{maximum} \{x^2, (1-x)^2, 2x(1-x)\}$ where $x \in [0, 1]$.

Determine the area of the region bounded by the curve $y = f(x)$ and the

lines $y = 0, x = 0, x = 1$.

 [Watch Video Solution](#)

9. Consider the region formed by the lines $x = 0$, $y = 0$, $x = 2$, $y = 2$. If the area enclosed by the curves $y = e^x$ and $y = 1nx$, within this region, is being removed, then find the area of the remaining region.

 [Watch Video Solution](#)

10. Draw a rough sketch of the curve $y = \frac{x^2 + 3x + 2}{x^2 - 3x + 2}$ and find the area of the bounded region between the curve and the x-axis.

 [Watch Video Solution](#)

11. Find the area bounded by the curve $y = (x - 1)(x - 2)(x - 3)$ lying between the ordinates $x = 0$ and $x = 3$.

 [Watch Video Solution](#)

12. Find the area bounded by the curve $x = \begin{cases} -2 - y, & y < -1 \\ y^3, & -1 \leq y \leq 1 \\ 2 - y, & y > 1 \end{cases}$

and $x=0$ is

 [Watch Video Solution](#)

13. Find the area enclosed by the graph of $y = \log_e(x + 1)$, y -axis, and the line $y=1$

 [Watch Video Solution](#)

14. Find the area bounded by the curve $y = \sin^{-1} x$ and the line $x = 0, |y| = \frac{\pi}{2}$.

 [Watch Video Solution](#)

15. Find the area of the region bounded by the curves $y = \sqrt{x + 2}$ and $y = \frac{1}{x + 1}$ between the lines $x=0$ and $x=2$.



Watch Video Solution

16. Find the area bounded by $y = \sin^{-1} x$, $y = \cos^{-1} x$, and the X -axis.



Watch Video Solution

17. Find the area bounded by the parabola $y = x^2 + 1$ and the straight line $x + y = 3$.



Watch Video Solution

18. Find the area bounded by the curves $y = \sin x$ and $y = \cos x$ between two consecutive points of the intersection.



Watch Video Solution

19. Find the ratio in which the area bounded by the curves $y^2 = 12x$ and $x^2 = 12y$ is divided by the line $x = 3$.

 [Watch Video Solution](#)

20. Find the area of the figure bounded by the parabolas $x = -2y^2$, $x = 1 - 3y^2$.

 [Watch Video Solution](#)

21. The area common to regions $x^2 + y^2 - 2x \leq 0$ and $y \geq \sin(\pi x/2)$

 [Watch Video Solution](#)

22. Find the area of the region enclosed by the curves $y = x \log x$ and $y = 2x - 2x^2$.

 [Watch Video Solution](#)

23. Find the area bounded by $y^2 \leq 4x$, $x^2 + y^2 \geq 2x$, and $x \leq y + 2$ in the first quadrant.

 [Watch Video Solution](#)

24. Sketch the region bounded by the curves $y = x^2$ and $y = \frac{2}{1 + x^2}$. Find the area.

 [Watch Video Solution](#)

25. Find the area bounded by the curves $y = x^3 - x$ and $y = x^2 + x$.

 [Watch Video Solution](#)

26. Find the area bounded by $y = -x^3 + x^2 + 16x$ and $y = 4x$

 [Watch Video Solution](#)

27. Find the area of the region enclosed by $y = -5x - x^2$ and $y = x$ on interval $[-1, 5]$



Watch Video Solution


28. If the area enclosed by curve $y = f(x)$ and $y = x^2 + 2$ between the abscissa $x = 2$ and $x = \alpha$, $\alpha > 2$, is $(\alpha^3 - 4\alpha^2 + 8)$ sq. unit. It is known that curve $y = f(x)$ lies below the parabola $y = x^2 + 2$.



Watch Video Solution

29. Let C_1 and C_2 be the graphs of the functions $y = x^2$ and $y = 2x$, respectively, where $0 \leq x \leq 1$. Let C_3 be the graph of a function $y = f(x)$, where $0 \leq x \leq 1$, $f(0) = 0$. For a point P on C_1 , let the lines through P , parallel to the axes, meet C_2 and C_3 at Q and R , respectively (see figure). If for every position of P (on C_1), the areas of the shaded

regions OPQ and ORP are equal, determine the function $f(x)$.



[Watch Video Solution](#)

30. If the area bounded by $f(x) = \sqrt{\tan x}$, $y = f(c)$, $x = 0$ and $x = a$, $0 < c < a < \frac{\pi}{2}$ is minimum then find the value of c .

[Watch Video Solution](#)

31. Find the area of the region R which is enclosed by the curve $y \geq \sqrt{1 - x^2}$ and $\max\{|x|, |y|\} \leq 4$.

 [Watch Video Solution](#)

32. Plot the region in the first quadrant in which points are nearer to the origin than to the line $x = 3$.

 [Watch Video Solution](#)

33. Consider a square with vertices at $(1, 1)$, $(-1, 1)$, $(-1, -1)$ and $(1, -1)$. Let S be the region consisting of all points inside the square which are nearer to the origin than to any edge. Sketch the region S and find its area.

 [Watch Video Solution](#)

34. Let $O(0, 0)$, $A(2, 0)$, and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let

R be the region consisting of all those points P inside OAB which satisfy $d(P, OA) \leq \min [d(P, OB), d(P, AB)]$, where d denotes the distance from the point to the corresponding line. Sketch the region R and find its area.

 [Watch Video Solution](#)

35. Find the area enclosed by $y = g(x)$, x -axis, $x=1$ and $x=37$, where $g(x)$ is inverse of $f(x) = x^3 + 3x + 1$.

 [Watch Video Solution](#)

36. Find the area bounded by the curve $f(x) = x + \sin x$ and its inverse function between the ordinates $x = 0$ to $x = 2\pi$.

 [Watch Video Solution](#)

1. Find the area bounded by the curve $x^2 = y$, $x^2 = -y$ and $y^2 = 4x - 3$

 [Watch Video Solution](#)

2. Find the area of the region enclosed by the curve $y = \left| x - \frac{1}{x} \right| (x > 0)$ and the line $y=2$

 [Watch Video Solution](#)

3. Find the area of the region bounded by the curves $y = x^2$, $y = |2 - x^2|$, and $y = 2$, which lies to the right of the line $x = 1$.

 [Watch Video Solution](#)

4. The ratio in which the line $x - 1 = 0$ divides the area bounded by the curves $2x + 1 = \sqrt{4y + 1}$, $y = x$ and $y = 2$ is

 [Watch Video Solution](#)

5. If S_0, S_1, S_2, \dots are areas bounded by the x-axis and half-wave of the curve $y = \sin \pi \sqrt{x}$, then prove that S_0, S_1, S_2, \dots are in A.P..

 [Watch Video Solution](#)

6. Find the area of the figure enclosed by the curve $5x^2 + 6xy + 2y^2 + 7x + 6y + 6 = 0$.

 [Watch Video Solution](#)

7. Find the area bounded by the curves $x^2 + y^2 = 4$, $x^2 = -\sqrt{2}y$ and $x = y$

 [Watch Video Solution](#)

8. Find the area of the region bounded by the curve $C: y = \tan x$, $\tan \geq n$ drawn $\rightarrow C$ at $x = \frac{\pi}{4}$, and the x-axis.

 [Watch Video Solution](#)

9. Compute the area of the region bounded by the curves $y = ex(\log)_e x$ and $y = \frac{\log x}{ex}$

 [Watch Video Solution](#)

10. If A_n be the area bounded by the curve $y = (\tan x)^n$ and the lines $x = 0$, $y = 0$, $x = \pi/4$, then for $n > 2$.

 [Watch Video Solution](#)

1. Find the area of the smaller part of the circle $x^2 + y^2 = a^2$ cut off by the line $x = \frac{a}{\sqrt{2}}$

 [Watch Video Solution](#)

2. Find the area enclosed by the curves $x^2 = y$, $y = x + 2$,

 [Watch Video Solution](#)

3. A curve is given by $y = \begin{cases} \sqrt{4 - x^2}, & 0 \leq x < 1 \\ \sqrt{3x}, & 1 \leq x \leq 3 \end{cases}$. Find the area lying between the curve and x-axis.

 [Watch Video Solution](#)

4. Find the area bounded by $x = 2y - y^2$

 [Watch Video Solution](#)

5. Find the area bounded by the x-axis, part of the curve $y = \left(1 - \frac{8}{x^2}\right)$, and the ordinates at $x = 2$ and $x = 4$. If the ordinate at $x = a$ divides the area into two equal parts, then find a .

 [Watch Video Solution](#)

6. Find the area of the region bounded by the x-axis and the curves defined by $y = \tan x$, (where $-\frac{\pi}{3} \leq x \leq \frac{\pi}{3}$) and $y = \cot x$ (where $\frac{\pi}{6} \leq x \leq \frac{2\pi}{3}$)

 [Watch Video Solution](#)

7. Find the area bounded by $y = \left|\sin x - \frac{1}{2}\right|$ and $y = 1$ for $x \in [0, \pi]$

 [Watch Video Solution](#)

8. If the area bounded by the graph of $y = xe^{-ax}$ ($a > 0$) and the x-axis is $1/9$ then find the value of a .

 [Watch Video Solution](#)

9. The area bounded by the curve $xy^2 = a^2(a - x)$ and y-axis is

 [Watch Video Solution](#)

Concept Application Exercise 9 2

1. Find the area lying in the first quadrant and bounded by the curve $y = x^3$ and the line $y = 4x$.

 [Watch Video Solution](#)

2. Find the area bounded by the curve $x^2 = 4y$ and the straight line $x = 4y - 2$.

 [Watch Video Solution](#)

3. Find the area enclosed by the figure described by the equation $x^4 + 1 = 2x^2 + y^2$.

 [Watch Video Solution](#)

4. In what ratio does the x-axis divide the area of the region bounded by the parabolas $y = 4x - x^2$ and $y = x^2 - x$?

 [Watch Video Solution](#)

5. Find the area of the circle $x^2 + y^2 = 16$ which is exterior to the parabola $y^2 = 6x$ by using

integration.

 [Watch Video Solution](#)

6. Find the area of the region bounded by the curves $y = x^2 + 2$, $y = x$, $x = 0$, and $x = 3$.

 [Watch Video Solution](#)

7. Find the area of the region bounded by the limits $x = 0$, $x = \frac{\pi}{2}$, and $f(x) = \sin x$, $g(x) = \cos x$.

 [Watch Video Solution](#)

8. Find the area bounded by $y = \tan^{-1} x$, $y = \cot^{-1} x$, and y -axis in the first quadrant.

 [Watch Video Solution](#)

9. Find the area bounded by $y = -(\log)_e x$, $y = (\log)_e x$, $y = (\log)_e(-x)$, and $y = -(\log)_e(-x)$.

 [Watch Video Solution](#)

10. Find the area of the region $\{(x, y) : y^2 \leq 4x, 4x^2 + 4y^2 \leq 9\}$

 [Watch Video Solution](#)

11. Sketch the region bounded by the curves $y = \sqrt{5 - x^2}$ and $y = |x - 1|$ and find its area.

 [Watch Video Solution](#)

12. Sketch the curves and identify the region bounded by the curves $x = \frac{1}{2}$, $x = 2$, $y = \log x$ and $y = 2^x$. Find the area of this region.

 [Watch Video Solution](#)

13. Find the area bounded by $y = x^2$ and $y = x^{1/3}$ for $x \in [-1, 1]$.



Watch Video Solution

14. Find the smallest area bounded by the curves $y = x - \sin x$, $y = x + \cos x$.



Watch Video Solution

Concept Application Exercise 9 3

1. Find the continuous function f where $(x^4 - 4x^2) \leq f(x) \leq (2x^2 - x^3)$ such that the area bounded by $y = f(x)$, $y = x^4 - 4x^2$, then y -axis, and the line $x = t$, where $(0 \leq t \leq 2)$ is k times the area bounded by $y = f(x)$, $y = 2x^2 - x^3$, $y = a\xi s$, and line $x = t$ (where $0 \leq t \leq 2$).



Watch Video Solution

2. If the area bounded by the x-axis, the curve $y = f(x)$, ($f(x) > 0$) and the lines $x = 1, x = b$ is equal to $\sqrt{b^2 + 1} - 1$ then find $f(x)$.

 [Watch Video Solution](#)

3. The area bounded by the graph of $y = f(x)$, $f(x) > 0$ on $[0, a]$ and x-axis is $\frac{a^2}{2} + \frac{a}{2} \sin a + \frac{\pi}{2} \cos a$ then find the value of $f\left(\frac{\pi}{2}\right)$.

 [Watch Video Solution](#)

4. A curve $y = f(x)$ is such that $f(x) \geq 0$ and $f(0) = 0$ and bounds a curvilinear triangle with the base $[0, x]$ whose area is proportional to $(n + 1)^{th}$ power of $f(x)$. If $f(1) = 1$ then find $f(x)$.

 [Watch Video Solution](#)

5. Find the area of curve enclosed by :

$$|x + y| + |x - y| \leq 4, |x| \leq 1, y \geq \sqrt{x^2 - 2x + 1}.$$

 [Watch Video Solution](#)

6. Consider two regions

R_1 : points P are nearer to $(1,0)$ than to $x = -1$.

R_2 : Points P are nearer to $(0,0)$ than to $(8,0)$ Find the area of the region common to R_1 and R_2 .

 [Watch Video Solution](#)

7. If $f: [-1, 1] \rightarrow \left[-\frac{1}{2}, \frac{1}{2}\right]: f(x) = \frac{x}{1+x^2}$, then find the area bounded by $y = f^{-1}(x)$, the x -axis and the lines $x = \frac{1}{2}, x = -\frac{1}{2}$.

 [Watch Video Solution](#)

1. Area enclosed by the curve $y = f(x)$ defined parametrically as

$$x = \frac{1-t^2}{1+t^2}, y = \frac{2t}{1+t^2} \text{ is equal to } \pi \text{ sq units} \quad (\text{b}) \quad \frac{\pi}{2} \text{ sq units}$$

$\frac{3\pi}{4} \text{ sq units}$ (d) $\frac{3\pi}{2} \text{ sq units}$

A. π sq. units

B. $\pi/2$ sq. units

C. $\frac{3\pi}{4}$ sq. units

D. $\frac{3\pi}{2}$ sq. units

Answer: A



Watch Video Solution

2. Let $f(x) = \text{minimum}(x+1, \sqrt{1-x})$ for all $x \leq 1$. Then the area

bounded by $y = f(x)$ and the x-axis is k units then $6k$ is equal to

A. $\frac{7}{3}$ sq. units

B. $\frac{1}{6}$ sq. units

C. $\frac{11}{6}$ sq. units

D. $\frac{7}{6}$ sq. units

Answer: D



Watch Video Solution

3. The area of the closed figure bounded by $x = -1$, $y = 0$, $y = x^2 + x + 1$, and the tangent to the curve $y = x^2 + x + 1$ at $A(1,3)$ is

A. (a) $4/3$ sq. units

B. (b) $7/3$ sq. units

C. (c) $7/6$ sq. units

D. (d) None of these

Answer: C



Watch Video Solution

4. The area bounded by the curve $a^2y = x^2(x + a)$ and the x-axis is

- A. $a^2 / 3$ sq. units
- B. $a^2 / 4$ sq. units
- C. $3a^2 / 4$ sq. units
- D. $a^2 / 12$ sq. units

Answer: D



[Watch Video Solution](#)

5. The area between the curve $y = 2x^4 - x^2$, the x-axis, and the ordinates of the two minima of the curve is

- A. $11 / 60$ sq. units
- B. $7 / 120$ sq. units
- C. $1 / 30$ sq. units

D. $7/90$ sq. units

Answer: B



Watch Video Solution

6. The area of the closed figure bounded by $x = -1$, $x = 2$, and $y = \begin{cases} -x^2 + 2, & x \leq 1 \\ 2x - 1, & x > 1 \end{cases}$ and the abscissa axis is (a) $\frac{16}{3}$ sq units (b) $\frac{10}{3}$ sq units (c) $\frac{13}{3}$ sq units (d) $\frac{7}{3}$ sq units

A. $16/3$ sq. units

B. $10/3$ sq. units

C. $13/3$ sq. units

D. $7/3$ sq. units

Answer: A



Watch Video Solution

7. The value of the parameter a such that the area bounded by $y = a^2x^2 + ax + 1$, coordinate axes, and the line $x=1$ attains its least value is equal to

- A. $\frac{1}{4}$ sq. units
- B. $-\frac{1}{2}$ sq. units
- C. $\frac{3}{4}$ sq. units
- D. -1 sq. units

Answer: C

 [Watch Video Solution](#)

8. The positive value of the parameter ' k ' for which the area of the figure bounded by the curve $y = \sin(kx)$, $x = \frac{2\pi}{3k}$, $x = \frac{5\pi}{3k}$ and x-axis is less than 2 can be

- A. $\frac{1}{8} < k < \frac{3}{8}$
- B. $0 < k < \frac{1}{8}$

C. $1 < k < 2$

D. $\frac{3}{8} < k < \frac{5}{8}$

Answer: C

 [Watch Video Solution](#)

9. The area bounded by the curve $y = x(1 - \log_e x)$ and x-axis is

A. $\frac{e^2}{4}$

B. $\frac{e^2}{2}$

C. $\frac{e^2 - e}{2}$

D. $\frac{e^2 - e}{4}$

Answer: A

 [Watch Video Solution](#)

10. The area inside the parabola $5x^2 - y = 0$ but outside the parabola $2x^2 - y + 9 = 0$ is (a) $12\sqrt{3}$ sq units (b) $6\sqrt{3}$ sq units (c) $8\sqrt{3}$ sq units (d) $4\sqrt{3}$ sq units

A. $12\sqrt{3}$ sq. units

B. $6\sqrt{3}$ sq. units

C. $8\sqrt{3}$ sq. units

D. $4\sqrt{3}$ sq. units

Answer: A



[Watch Video Solution](#)

11. Area enclosed between the curves $|y| = 1 - x^2$ and $x^2 + y^2 = 1$ is (a)

$\frac{3\pi - 8}{3}$ (b) $\frac{\pi - 8}{3}$ (c) $\frac{2\pi - 8}{3}$ (d) None of these

A. $\frac{3\pi - 8}{3}$ sq. units

B. $\frac{\pi - 8}{3}$

C. $\frac{2\pi - 8}{3}$ sq. units

D. None of these

Answer: A



Watch Video Solution

12. If A_n is the area bounded by $y=x$ and $y = x^n, n \in N$, then

$$A_2 \cdot A_3 \dots A_n =$$

A. $\frac{1}{n(n+1)}$

B. $\frac{1}{2^n n(n+1)}$

C. $\frac{1}{2^{n-1} n(n+1)}$

D. $\frac{1}{2^{n-2} n(n+1)}$

Answer: D



Watch Video Solution

13. The area of the region in 1st quadrant bounded by the y-axis,

$$y = \frac{x}{4}, y = 1 + \sqrt{x}, \text{ and } y = \frac{2}{\sqrt{x}} \text{ is}$$

- A. $2/3$ sq. units
- B. $8/3$ sq. units
- C. $11/3$ sq. units
- D. $13/6$ sq. units

Answer: C



Watch Video Solution

14. The area of the closed figure bounded by $y = \frac{x^2}{2} - 2x + 2$ and the tangents to it at $\left(1, \frac{1}{2}\right)$ and $(4, 2)$ is $\frac{9}{8}$ sq units (b) $\frac{3}{8}$ sq units $\frac{3}{2}$ sq units

(d) $\frac{9}{4}$ sq units

- A. $9/8$ sq. units
- B. $3/8$ sq. units

C. $3/2$ sq. units

D. $9/4$ sq. units

Answer: A



[Watch Video Solution](#)

15. The area of the region bounded by

$x^2 + y^2 - 2x - 3 = 0$ and $y = |x| + 1$ is

A. $\frac{\pi}{2} - 1$ sq. units

B. 2π sq. units

C. 4π sq. units

D. $\pi/2$ sq. units

Answer: A



[Watch Video Solution](#)

16. The area enclosed by the curve $y = \sqrt{4 - x^2}$, $y \geq \sqrt{2} \sin\left(\frac{x\pi}{2\sqrt{2}}\right)$,

and the x-axis is divided by the y-axis in the ratio.

(a) $\frac{\pi^2 - 8}{\pi^2 + 8}$

(b) $\frac{\pi^2 - 4}{\pi^2 + 4}$

(c) $\frac{\pi - 4}{\pi - 4}$

(d) $\frac{2\pi^2}{2\pi + \pi^2 - 8}$

A. $\frac{\pi^2 - 8}{\pi^2 + 8}$

B. $\frac{\pi^2 - 4}{\pi^2 + 4}$

C. $\frac{\pi - 4}{\pi - 4}$

D. $\frac{2\pi^2}{2\pi + \pi^2 - 8}$

Answer: D



Watch Video Solution

17. The area bounded by the curve $y^2 = 1 - x$ and the lines

$y = \frac{[x]}{x}$, $x = -1$, and $x = \frac{1}{2}$ is

A. $\frac{3}{\sqrt{2}} - \frac{11}{6}$ sq. units

B. $3\sqrt{2} - \frac{11}{4}$ sq. units

C. $\frac{6}{\sqrt{2}} - \frac{11}{5}$ sq. units

D. None of these

Answer: A



[Watch Video Solution](#)

18. The area bounded by the curves $y = (\log)_e x$ and $y = ((\log)_e x)^2$ is

A. $e - 2$ sq. units

B. $3 - e$ sq. units

C. e sq. units

D. $e - 1$ sq. units

Answer: B



[Watch Video Solution](#)

19. The area bounded by $y = 3 - |3 - x|$ and $y = \frac{6}{|x + 1|}$ is

A. $\frac{15}{2} - 6$ In 2 sq. units

B. $\frac{13}{2} - 3$ In 2 sq. units

C. $\frac{13}{2} - 6$ In 2 sq. units

D. None of these

Answer: C



[Watch Video Solution](#)

20. Find the area enclosed between the curves:

$y = \log_e(x + e)$, $x = \log_e\left(\frac{1}{y}\right)$ & the x-axis.

A. 2 sq. units

B. 1 sq. units

C. 4 sq. units

D. None of these

Answer: A



[Watch Video Solution](#)

21. Find the area enclosed the curve $y = \sin x$ and the X-axis between $x = 0$ and $x = \pi$.

A. $\frac{7}{2}$ sq. units

B. $\frac{7}{4} + \sqrt{3}$ sq. units

C. $\frac{7\sqrt{3}}{4}$ sq. units

D. $7 - \frac{\sqrt{3}}{4}$ sq. units

Answer: A



[Watch Video Solution](#)

22. The area bounded by $y = x^2$, $y = [x + 1]$, $0 \leq x \leq 2$ and the y-axis is where $[.]$ is greatest integer function.

A. $\frac{1}{3}$

B. $\frac{\sqrt{2}}{3}$

C. 1

D. $\frac{7}{3}$

Answer: B



[Watch Video Solution](#)

23. The area of the region bounded by the parabola $(y - 2)^2 = x - 1$, the tangent to the parabola at the point (2,3) and the X-axis is

A. 7 sq. units

B. 6 sq. units

C. 9 sq. units

D. None of these

Answer: C

 [Watch Video Solution](#)

24. The area bounded by the curves $y = xe^x$, $y = xe^{-x}$ and the line $x = 1$ is $\frac{2}{e}$ sq units (b) $1 - \frac{2}{e}$ sq units $\frac{1}{e}$ sq units (d) $1 - \frac{1}{e}$ sq units

A. $\frac{2}{e}$ sq. units

B. $1 - \frac{2}{e}$ sq. units

C. $\frac{1}{e}$ sq. units

D. $1 - \frac{1}{e}$ sq. units

Answer: A

 [Watch Video Solution](#)

25. The area of the region whose boundaries are defined by the curves $y=2 \cos x$, $y=3 \tan x$, and the y-axis is

A. $1 + 3 \ln \left(\frac{2}{\sqrt{3}} \right)$ sq. units

B. $1 + \frac{3}{2} \ln 3 - 3 \ln 2$ sq. units

C. $1 + \frac{3}{2} \ln 3 - \ln 2$ sq. units

D. $\ln 3 - \ln 2$ sq. units

Answer: B



[Watch Video Solution](#)

26. Area bounded by $y = \sec^{-1} x$, $y = \cot^{-1} x$ and line $x=1$ is given by

A. $\log(3 + 2\sqrt{2}) - \frac{\pi}{2}$ sq. units

B. $\frac{\pi}{2} - \log(3 + 2\sqrt{2})$ sq. units

C. $\pi - \log_e 3$ sq. units

D. None of these

Answer: A

 [Watch Video Solution](#)

27. The area bounded by the curve $y = \frac{3}{|x|}$ and $y + |2 - x| = 2$ is

 [Watch Video Solution](#)

28. The area enclosed by $y = x^2 + \cos x$ and its normal at $x = \frac{\pi}{2}$ in the first quadrant is

A. $\frac{\pi^5}{32} - \frac{\pi^4}{64} + \frac{\pi^3}{32} + 1$

B. $\frac{\pi^5}{16} - \frac{\pi^4}{32} + \frac{\pi^3}{24} - 1$

C. $\frac{\pi^5}{32} - \frac{\pi^4}{32} + \frac{\pi^3}{16}$

D. $\frac{\pi^5}{32} - \frac{\pi^4}{32} + \frac{\pi^3}{24} + 1$

Answer: D



Watch Video Solution

29. Given $f(x) = \int_0^x e^t (\log_e \sec t - \sec^2 t) dt$, $g(x) = -2e^x \tan x$, then the area bounded by the curves $y = f(x)$ and $y = g(x)$ between the ordinates $x = 0$ and $x = \frac{\pi}{3}$, is (in sq. units)

A. $\frac{1}{2} e^{\frac{\pi}{3}} \log_e 2$

B. $e^{\frac{\pi}{3}} \log_e 2$

C. $\frac{1}{4} e^{\frac{\pi}{3}} \log_e 2$

D. $e^{\frac{\pi}{3}} \log_e 3$

Answer: B



Watch Video Solution

30. The area of the loop of the curve $ay^2 = x^2(a - x)$ is $4a^2$ sq units (b)

$\frac{8a^2}{15}$ sq units $\frac{16a^2}{9}$ sq units (d) None of these

A. $4a^2$ sq. units

B. $\frac{8a^2}{15}$ sq. units

C. $\frac{16a^2}{9}$ sq. units

D. None of these

Answer: B



Watch Video Solution

31. Area of the region enclosed between the curves $x = y^2 - 1$ and

$x = |y|\sqrt{1 - y^2}$ is

A. 1 sq. units

B. $4/3$ sq. units

C. $2/3$ sq. units

D. 2 sq. units

Answer: D



Watch Video Solution

32. The area bounded by the loop of the curve $4y^2 = x^2(4 - x^2)$ is given by (1) $\frac{7}{3}$ (2) $\frac{8}{3}$ (3) $\frac{11}{3}$ (4) $\frac{16}{3}$

A. $\frac{7}{3}$ sq. units

B. $\frac{8}{3}$ sq. units

C. $\frac{11}{3}$ sq. units

D. $\frac{16}{3}$ sq. units

Answer: D



Watch Video Solution

33. The area enclosed by the curves $xy^2 = a^2(a - x)$ and $(a - x)y^2 = a^2x$ is

A. $(\pi - 2)a^2$ sq. units

B. $(4 - \pi)a^2$ sq. units

C. $\pi a^2 / 3$ sq. units

D. None of these

Answer: A



Watch Video Solution

34. The area bounded by the two branches of curve $(y - x)^2 = x^3$ and the straight line $x = 1$ is $\frac{1}{5}$ sq units (b) $\frac{3}{5}$ sq units $\frac{4}{5}$ sq units (d) $\frac{8}{4}$ sq units

A. $1/5$ sq. units

B. $3/5$ sq. units

C. $4/5$ sq. units

D. $8/4$ sq. units

Answer: C



Watch Video Solution

35. The area bounded by the curves

$y = \sin^{-1}|\sin x|$ and $y = (\sin^{-1}|\sin x|)^2$, where $0 \leq x \leq 2\pi$, is

A. $\frac{1}{3} + \frac{\pi^2}{4}$ sq. units

B. $\frac{1}{6} + \frac{\pi^3}{8}$ sq. units

C. 2 sq. units

D. None of these

Answer: D



Watch Video Solution

36. Consider two curves $C_1: y^2 = 4[\sqrt{y}]x$ and $C_2: x^2 = 4[\sqrt{x}]y$, where $[\cdot]$ denotes the greatest integer function. Then the area of region enclosed by these two curves within the square formed by the lines $x = 1, y = 1, x = 4, y = 4$ is (a) $\frac{8}{3}$ sq units (b) $\frac{10}{3}$ sq units (c) $\frac{11}{3}$ sq units (d) $\frac{11}{4}$ sq units

A. $\frac{8}{3}$ sq. units

B. $\frac{10}{3}$ sq. units

C. $\frac{11}{3}$ sq. units

D. $\frac{11}{4}$ sq. units

Answer: C



Watch Video Solution

37. The area enclosed between the curve $y^2(2a - x) = x^3$ and the line $x=2$ above the x-axis is

A. πa^2 sq. units

B. $\frac{3\pi a^2}{2}$ sq. units

C. $2\pi a^2$ sq. units

D. $3\pi a^2$ sq. units

Answer: B



Watch Video Solution

38. The area of the region of the plane bounded by $\max(|x|, |y|) \leq 1$

and $xy \leq \frac{1}{2}$ is

(a) $\frac{1}{2} + \ln 2$ sq. units

(b) $3 + \ln 2$ sq. units

(c) $\frac{31}{4}$ sq. units

(d) $1 + 2\ln 2$ sq. units

A. $1/2 + \ln 2$ sq. units

B. $3 + \ln 2$ sq. units

C. $31/4$ sq. units

D. $1+2 \ln 2$ sq. units

Answer: B



[Watch Video Solution](#)

39. Find the area of the region containing the points (x, y) satisfying

$$4 \leq x^2 + y^2 \leq 2(|x| + |y|).$$

A. 8 sq. units

B. 2 sq. units

C. 4π sq. units

D. 2π sq. units

Answer: A



[Watch Video Solution](#)

40. Let $f(x)$ be a non-negative continuous function such that the area bounded by the curve $y=f(x)$, the x -axis, and the ordinates $x = \frac{\pi}{4}$ and $x = \beta > \frac{\pi}{4}$ is $\beta \sin \beta + \frac{\pi}{4} \cos \beta + \sqrt{2}\beta$. Then $f'\left(\frac{\pi}{2}\right)$ is

A. $\left(\frac{\pi}{2} - \sqrt{2} - 1\right)$

B. $\left(\frac{\pi}{4} + \sqrt{2} - 1\right)$

C. $-\frac{\pi}{2}$

D. $\left(1 - \frac{\pi}{2} - \sqrt{2}\right)$

Answer: C



Watch Video Solution

Multiple Correct Answers Type

1. Let $A(k)$ be the area bounded by the curves $y = x^2 - 3$ and $y = kx + 2$. The range of $A(k)$ is $\left(\frac{10\sqrt{5}}{3}, \infty\right)$. The range of $A(k)$ is

$\left(\frac{20\sqrt{5}}{3}, \infty\right)$ If function $k \vec{A}(k)$ is defined for $k \in [-2, \infty)$, then $A(k)$

is many-one function. The value of k for which area is minimum is 1.

A. The range of $A(k)$ is $\left[\frac{10\sqrt{5}}{3}, \infty\right)$

B. The range of $A(k)$ is $\left[\frac{20\sqrt{5}}{3}, \infty\right)$

C. If function $k \rightarrow A(k)$ is defined for $k \in [-2, \infty)$, then $A(k)$ is many-one function

D. The value of k for which area is minimum is 1

Answer: B::C



Watch Video Solution

2. The parabolas $y^2 = 4x$ and $x^2 = 4y$ divide the square region bounded by the lines $x = 4, y = 4$ and the coordinate axes. If S_1, S_2, S_3 are the areas of these parts numbered from top to bottom, respectively, then $S_1 : S_2 \equiv 1 : 1$ (b) $S_2 : S_3 \equiv 1 : 2$ $S_1 : S_3 \equiv 1 : 1$ (d) $S_1 : (S_1 + S_2) = 1 : 2$

A. $S_1 : S_2 \equiv 1 : 1$

B. $S_2 : S_3 \equiv 1 : 2$

C. $S_1 : S_3 \equiv 1 : 1$

D. $S_1 : (S_1 + S_2) = 1 : 2$

Answer: A::C::D

 [Watch Video Solution](#)

3. Which of the following have the same bounded area

$f(x) = s \in x, g(x) = \sin^2 x, \text{ where } 0 \leq x \leq 10\pi$

$f(x) = s \in x, g(x) = |s \in x|, \text{ where } 0 \leq x \leq 20\pi$

$f(x) = |s \in x|, g(x) = \sin^3 x, \text{ where } 0 \leq x \leq 10\pi$

$f(x) = s \in x, g(x) = \sin^4 x, \text{ where } 0 \leq x \leq 10\pi$

A. $f(x) = \sin x, g(x) = \sin^2 x, \text{ where } 0 \leq x \leq 10\pi$

B. $f(x) = \sin x, g(x) = |\sin|, \text{ where } 0 \leq x \leq 20\pi$

C. $f(x) = |\sin|, g(x) = \sin^3 x, \text{ where } 0 \leq x \leq 10\pi$

D. $f(x) = \sin x, g(x) = \sin^4 x$, where $0 \leq x \leq 10\pi$

Answer: A::C::D



Watch Video Solution

4. If the curve $y = ax^{\frac{1}{2}} + bx$ passes through the point $(1, 2)$ and lies above the x-axis for $0 \leq x \leq 9$ and the area enclosed by the curve, the x-axis, and the line $x = 4$ is 8 sq. units, then (a) $a = 1$ (b) $b = 1$ (c) $a = 3$ (d) $b = -1$

A. $a = 1$

B. $b = 1$

C. $a = 3$

D. $b = -1$

Answer: C::D



Watch Video Solution

5. The area bounded by the curve $x = a \cos^3 t$, $y = a \sin^3 t$, is :

A. $12a^2 \int_0^{\pi/2} \cos^4 t \sin^2 t dt$

B. $12a^2 \int_0^{\pi/2} \cos^2 t \sin^4 t dt$

C. $2 \int_{-a}^a (a^{2/3} - x^{2/3})^{3/2} dx$

D. $4 \int_0^a (a^{2/3} - x^{2/3}) dx$

Answer: A::C::D



Watch Video Solution

6. If A_1 is the area area bounded by $|x - a_i| + |y| = b_i$, $i \in N$, where

$a_{i+1} = a_i + \frac{3}{2}b_i$ and $b_{i+1} = \frac{b_i}{2}$, $a_i = 0$ and $b_i = 32$, then

A. $A_3 = 128$

B. $A_3 = 256$

$$C. \lim_{n \rightarrow \infty} \sum_{i=1}^n A_i = \frac{8}{3}(32)^2$$

$$D. \lim_{n \rightarrow \infty} \sum_{i=1}^n A_i = \frac{4}{3}(16)^2$$

Answer: A::C



Watch Video Solution

7. Find the area bounded by the curve $y = 2x - x^2$, and the line $y = x$

$$A. 2 \int_1^e \sqrt{\log_e y} dy$$

$$B. 2e - \int_{-1}^1 e^{x^2} dx$$

$$C. \int_{-1}^1 (e - e^{x^2}) dx$$

$$D. 2 \int_0^1 \sqrt{x} e^x dx$$

Answer: A::B::C::D



Watch Video Solution

8. The area bounded by the curves $y = |x| - 1$ and $y = -|x| + 1$ is 1 sq. units (b) 2 sq. units $2\sqrt{2}$ sq. units (d) 4 sq. units

A. $\alpha = e^2 + 1$

B. $\alpha = e^2 - 2$

C. $\beta = 1 + e^{-1}$

D. $\beta = 1 + e^{-2}$

Answer: A:D



Watch Video Solution

9. Consider curves

$S_1: \sqrt{|x|} + \sqrt{|y|} = \sqrt{a}$, $S_2: x^2 + y^2 = a^2$ and $S_3: |x| + |y| = a$. If α is

the area bounded by S_2 and S_3 , then

A. $\alpha = a^2 \left(\pi - \frac{2}{3} \right)$

B. $\beta = \frac{4a^2}{3}$

C. $\gamma = 2a^2(\pi - 1)$

D. the ratio in which S_3 divides area between

S_1 and S_2 is $4:3(\pi - 2)$

Answer: A::B::D



Watch Video Solution

10. Let $A(k)$ be the area bounded by the curves $y = x^2 + 2x - 3$ and $y = kx + 1$. Then

A. the value of k for which $A(k)$ is least is 2

B. the value of k for which $A(k)$ is least is $3/2$

C. least value of $A(k)$ is $32/3$

D. least value of $A(k)$ is $64/3$

Answer: A::C



Watch Video Solution

11. The area of the region bounded by the curve $y = e^x$ and lines $x=0$ and $y=e$ is

A. $e - 1$

B. $\int_1^e \ln(e + 1 - y) dy$

C. $e - \int_0^1 e^x dx$

D. $\int_1^e \ln y dy$

Answer: B::C::D



Watch Video Solution

12. The area of the region $R = \{(x, y) : |x| \leq |y| \text{ and } x^2 + y^2 \leq 1\}$ is

A. $\frac{1}{2} < \alpha < 1$

B. $\alpha^4 + 4\alpha^2 - 1 = 0$

C. $0 < \alpha \leq \frac{1}{2}$

D. $2\alpha^4 - 4\alpha^2 + 1 = 0$

Answer: A:D



Watch Video Solution

13. Let $f: R \rightarrow R$ be a differentiable function such that

$$f(x) = x^2 + \int_0^x e^{-t} f(x-t) dt.$$

$y = f(x)$ is

A. The curve $y=f(x)$ passes through the point (1,2)

B. The curve $y=f(x)$ passes through the point (2,-1)

C. The area of the region

$$\left\{ (x, y) \in [0, 1] \times R: f(x) \leq y \leq \sqrt{1-x^2} \right\} \text{ is } \frac{\pi-2}{4}$$

D. The area of the region

$$\left\{ (x, y) \in [0, 1] \times R: f(x) \leq y \leq \sqrt{1-x^2} \right\} \text{ is } \frac{\pi-1}{4}$$

Answer: B::C



Watch Video Solution

Linked Comprehension Type

1. Let A_r be the area of the region bounded between the curves $y^2 = (e^{-kr})x$ (where $k > 0, r \in N$) and the line $y = mx$ (where $m \neq 0$), k and m are some constants

A_1, A_2, A_3, \dots are in G.P. with common ratio

A. e^{-k}

B. e^{-2k}

C. e^{-4k}

D. None of these

Answer: B



Watch Video Solution

2. Let A_r be the area of the region bounded between the curves $y^2 = (e^{-kr})x$ (where $k > 0, r \in \mathbb{N}$) and the line $y = mx$ (where $m \neq 0$), k and m are some constants

$\lim_{n \rightarrow \infty} \sum_{i=1}^n A_i = \frac{1}{48(e^{2k} - 1)}$ then the value of m is

A. 3

B. 1

C. 2

D. 4

Answer: C




[Watch Video Solution](#)

3. If $y=f(x)$ is a monotonic function in (a,b) , then the area bounded by the ordinates _____ at $x = a, x = b, y = f(x)$ and $y = f(c)$ (where $c \in (a, b)$) is minimum when

Proof:
$$A = \int_a^c (f(c) - f(x))dx + \int_c^b (f(c))dx$$

$$= f(c)(c - a) - \int_a^c (f(x))dx + \int_a^b (f(x))dx - f(c)(b - c)$$

$$\Rightarrow A = [2c - (a + b)]f(c) + \int_c^b (f(x))dx - \int_a^c (f(x))dx$$



Differentiating w.r.t. c , we get

$$\frac{dA}{dc} = [2c - (a + b)]f'(c) + 2f(c) + 0 - f(c) - (f(c) - 0)$$

For maxima and minima, $\frac{dA}{dc} = 0$

$$\Rightarrow f'(c)[2c - (a + b)] = 0 \text{ (as } f'(c) \neq 0 \text{)}$$

Hence, $c = \frac{a + b}{2}$

Also for $c < \frac{a + b}{2}$, $\frac{dA}{dc} < 0$ and for $c > \frac{a + b}{2}$, $\frac{dA}{dc} > 0$

Hence, A is minimum when $c = \frac{a + b}{2}$.

If the area bounded by $f(x) = \frac{x^3}{3} - x^2 + a$ and the straight lines $x=0$, $x=2$, and the x-axis is minimum, then the value of a is

A. $1/2$

B. 2

C. 1

D. $2/3$

Answer: D



Watch Video Solution

4. If $y=f(x)$ is a monotonic function in (a,b) , then the area bounded by the ordinates _____ at


$x = a, x = b, y = f(x)$ and $y = f(c)$ (where $c \in (a, b)$) is minimum when

$$c = \frac{a + b}{2}.$$

Proof: $A = \int_a^c (f(c) - f(x))dx + \int_c^b (f(c))dx$

$$= f(c)(c - a) - \int_a^c (f(x))dx + \int_a^b (f(x))dx - f(c)(b - c)$$

$$\Rightarrow A = [2c - (a + b)]f(c) + \int_c^b (f(x))dx - \int_a^c (f(x))dx$$



Differentiating w.r.t. c , we get

$$\frac{dA}{dc} = [2c - (a + b)]f'(c) + 2f(c) + 0 - f(c) - (f(c) - 0)$$

For maxima and minima, $\frac{dA}{dc} = 0$

$$\Rightarrow f'(c)[2c - (a + b)] = 0 \text{ (as } f'(c) \neq 0)$$

Hence, $c = \frac{a + b}{2}$

Also for $c < \frac{a + b}{2}$, $\frac{dA}{dc} < 0$ and for $c > \frac{a + b}{2}$, $\frac{dA}{dc} > 0$

Hence, A is minimum when $c = \frac{a + b}{2}$.

The value of the parameter a for which the area of the figure bounded by

the abscissa axis, the graph of the function $y = x^3 + 3x^2 + x + a$, and the straight lines, which are parallel to the axis of ordinates and cut the abscissa axis at the point of extremum of the function, which is the least, is

A. 2

B. 0

C. -1

D. 1

Answer: C



[Watch Video Solution](#)


5. If $y=f(x)$ is a monotonic function in (a,b) , then the area bounded by the ordinates _____ at

$x = a, x = b, y = f(x)$ and $y = f(c)$ (where $c \in (a, b)$) is minimum when

Proof:
$$A = \int_a^c (f(c) - f(x))dx + \int_c^b (f(c))dx$$

$$= f(c)(c - a) - \int_a^c (f(x))dx + \int_a^b (f(x))dx - f(c)(b - c)$$

$$\Rightarrow A = [2c - (a + b)]f(c) + \int_c^b (f(x))dx - \int_a^c (f(x))dx$$



Differentiating w.r.t. c , we get

$$\frac{dA}{dc} = [2c - (a + b)]f'(c) + 2f(c) + 0 - f(c) - (f(c) - 0)$$

For maxima and minima, $\frac{dA}{dc} = 0$

$$\Rightarrow f'(c)[2c - (a + b)] = 0 \text{ (as } f'(c) \neq 0 \text{)}$$

$$\text{Hence, } c = \frac{a + b}{2}$$

Also for $c < \frac{a + b}{2}$, $\frac{dA}{dc} < 0$ and for $c > \frac{a + b}{2}$, $\frac{dA}{dc} > 0$

Hence, A is minimum when $c = \frac{a + b}{2}$.

If the area enclosed by $f(x) = \sin x + \cos x$, $y = a$ between two consecutive points of extremum is minimum, then the value of a is

- A. 0
- B. -1
- C. 1
- D. 2

Answer: A



[Watch Video Solution](#)

6. Consider the area S_0, S_1, S_2, \dots bounded by the x-axis and half-waves of the curve $y = e^{-x} \sin x$, where $x \geq 0$.

The value of S_0 is

- A. $\frac{1}{2}(1 + e^\pi)$ sq. units
- B. $\frac{1}{2}(1 + e^{-\pi})$ sq. units
- C. $\frac{1}{2}(1 - e^{-\pi})$ sq. units

D. $\frac{1}{2}(e^\pi - 1)$ sq. units

Answer: A



Watch Video Solution

7. Consider the sequence of natural numbers s_0, s_1, s_2, \dots such that $s_0 = 3, s_1 = 3$ and $s_n = 3 + s_{n-1}s_{n-2}$, then

A. $\frac{e^\pi}{2}$

B. $e^{-\pi}$

C. e^π

D. $\frac{e^{-\pi}}{2}$

Answer: C



Watch Video Solution

8. Consider the area S_0, S_1, S_2, \dots bounded by the x-axis and half-waves of the curve $y = e^{-x} \sin x$, where $x \geq 0$.

The value of S_0 is

A. $\frac{1 + e^\pi}{1 - e^{-\pi}}$

B. $\frac{\frac{1}{2}(1 + e^\pi)}{1 - e^\pi}$

C. $\frac{1}{2(1 - e^{-\pi})}$

D. None of these

Answer: B



Watch Video Solution

9. Two curves

$$C_1 \equiv [f(y)]^{2/3} + [f(x)]^{1/3} = 0 \text{ and } C_2 \equiv [f(y)]^{2/3} + [f(x)]^{2/3} = 12,$$

satisfying the relation

$$(x - y)f(x + y) - (x + y)f(x - y) = 4xy(x^2 - y^2)$$

The area bounded by the curve C_1 and C_2 is

A. (a) $2\pi - \sqrt{3}$ sq. units

B. (b) $2\pi + \sqrt{3}$ sq. units

C. (c) $\pi + \sqrt{6}$ sq. units

D. (d) $2\sqrt{3} - \pi$ sq. units

Answer: B



Watch Video Solution

10. Two curves

$C_1 \equiv [f(y)]^{2/3} + [f(x)]^{1/3} = 0$ and $C_2 \equiv [f(y)]^{2/3} + [f(x)]^{2/3} = 12$,

satisfying the relation

$$(x - y)f(x + y) - (x + y)f(x - y) = 4xy(x^2 - y^2)$$

The area bounded by the curve C_2 and $|x| + |y| = \sqrt{12}$ is

A. (a) $12\pi - 24$ sq. units

B. (b) $6 - \sqrt{12}$ sq. units

C. (c) $2\sqrt{12} - 6$ sq. units

D. (d) None of these

Answer: A

 [Watch Video Solution](#)

11. Two curves $C_1 \equiv [f(y)]^{2/3} + [f(x)]^{1/3} = 0$ and $C_2 \equiv [f(y)]^{2/3} + [f(x)]^{2/3} = 12$, satisfying the relation

$$(x - y)f(x + y) - (x + y)f(x - y) = 4xy(x^2 - y^2)$$

The area bounded by C_1 and $x + y + 2 = 0$ is

A. (a) $5/2$ sq. units

B. (b) $7/2$ sq. units

C. (c) $9/2$ sq. units

D. (d) None of these

Answer: C

 [Watch Video Solution](#)

12. Consider the two curves

$C_1: y = 1 + \cos x$ and $C_2: y = 1 + \cos(x - \alpha)$ for $\alpha \in \left(0, \frac{\pi}{2}\right)$, where

Also the area of the figure bounded by the curves C_1, C_2 , and $x = 0$ is same as that of the figure bounded by $C_2, y = 1$, and $x = \pi$.

The value of α is

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{6}$

D. $\frac{\pi}{8}$

Answer: C



Watch Video Solution

13. Consider two curves $C_1: y = \frac{1}{x}$ and $C_2: y = 1nx$ on the xy plane. Let

D_1 denotes the region surrounded by C_1, C_2 , and the line $x = 1$ and D_2

denotes the region surrounded by C_1, C_2 and the line $x = a$. If

$D_1 = D_2$, then the sum of logarithm of possible value of a is _____

- A. 1 sq. units
- B. 2 sq. units
- C. $2 + \sqrt{3}$ sq. units
- D. None of these

Answer: B



Watch Video Solution

14. Consider the function defined implicitly by the equation $y^2 - 2ye^{\sin^{-1}x} + x^2 - 1 + [x] + e^{2\sin^{-1}x} = 0$ (where $[x]$ denotes the greatest integer function).

The area of the region bounded by the curve and the line $x = -1$ is

- A. $\pi + 1$ sq. units
- B. $\pi - 1$ sq. units
- C. $\frac{\pi}{2} + 1$ sq. units

D. $\frac{\pi}{2} - 1$ sq. units

Answer: A



Watch Video Solution

15. Consider the function defined implicitly by the equation $y^2 - 2ye^{\sin^{-1}x} + x^2 - 1 + [x] + e^{2\sin^{-1}x} = 0$ (where $[x]$ denotes the greatest integer function). Line $x=0$ divides the region mentioned above in two parts. The ratio of area of left-hand side of line to that of right-hand side of line is

A. $1 + \pi : \pi$

B. $2 - \pi : \pi$

C. $1 : 1$

D. $\pi + 2 : \pi$

Answer: D



Watch Video Solution

16. Consider two functions

$$f(x) = \begin{cases} [x], & -2 \leq x \leq -1 \\ |x| + 1, & -1 < x \leq 2 \end{cases} \quad \text{and} \quad g(x) = \begin{cases} [x], & -\pi \leq x < 0 \\ \sin x, & 0 \leq x \leq \pi \end{cases}$$

where $[.]$ denotes the greatest integer function.

The exhaustive domain of $g(f(x))$ is

- A. $\frac{\sqrt{3}}{4} + \frac{\pi}{6}$ sq. units
- B. $\frac{\sqrt{3}}{2} + \frac{\pi}{6}$ sq. units
- C. $\frac{\sqrt{3}}{4} - \frac{\pi}{6}$ sq. units
- D. $\frac{\sqrt{3}}{2} - \frac{\pi}{6}$ sq. units

Answer: A

 [Watch Video Solution](#)

17. Computing area with parametrically represented boundaries : If the boundary of a figure is represented by parametric equation, i.e., $x = x(t), y = (t)$, then the area of the figure is evaluated by one of the three formulas :

$$S = - \int_{\alpha}^{\beta} y(t)x'(t)dt,$$

$$S = \int_{\alpha}^{\beta} x(t)y'(t)dt,$$

$$S = \frac{1}{2} \int_{\alpha}^{\beta} (xy' - yx')dt,$$

Where α and β are the values of the parameter t corresponding respectively to the beginning and the end of the traversal of the curve corresponding to increasing t .

The area of the region bounded by an arc of the cycloid $x = a(t - \sin t)$, $y = a(1 - \cos t)$ and the x-axis is

A. $6\pi a^2$ sq. units

B. $3\pi a^2$ sq. units

C. $4\pi a^2$ sq. units

D. None of these

Answer: B



Watch Video Solution

18. Computing area with parametrically represented boundaries : If the boundary of a figure is represented by parametric equation, i.e., $x = x(t), y = y(t)$, then the area of the figure is evaluated by one of the three formulas :

$$S = - \int_{\alpha}^{\beta} y(t)x'(t)dt,$$

$$S = \int_{\alpha}^{\beta} x(t)y'(t)dt,$$

$$S = \frac{1}{2} \int_{\alpha}^{\beta} (xy' - yx')dt,$$

Where α and β are the values of the parameter t corresponding respectively to the beginning and the end of the traversal of the curve corresponding to increasing t .

The area of the loop described as

$$x = \frac{t}{3}(6 - t), y = \frac{t^2}{8}(6 - t) \text{ is}$$

A. $\frac{27}{5}$ sq. units

B. $\frac{24}{5}$ sq. units

C. $\frac{27}{6}$ sq. units

D. $\frac{21}{5}$ sq. units

Answer: A



Watch Video Solution

19. Computing area with parametrically represented boundaries : If the boundary of a figure is represented by parametric equation, i.e., $x = x(t), y = y(t)$, then the area of the figure is evaluated by one of the three formulas :

$$S = - \int_{\alpha}^{\beta} y(t)x'(t)dt,$$

$$S = \int_{\alpha}^{\beta} x(t)y'(t)dt,$$

$$S = \frac{1}{2} \int_{\alpha}^{\beta} (xy' - yx')dt,$$

Where α and β are the values of the parameter t corresponding respectively to the beginning and the end of the traversal of the curve corresponding to increasing t .

If the curve given by parametric equation $x = t - t^3, y = 1 - t^4$ forms a loop for all values of $t \in [-1, 1]$ then the area of the loop is

A. $\frac{1}{7}$ sq. units

B. $\frac{3}{5}$ sq. units

C. $\frac{16}{35}$ sq. units

D. $\frac{8}{35}$ sq. units

Answer: C

 [Watch Video Solution](#)

20. Let $f(x)$ be a continuous function given by

$$f(x) = \begin{cases} 2x, & |x| \leq 1 \\ x^2 + ax + b, & |x| > 1 \end{cases}, \text{ then}$$

If $f(x)$ is continuous for all real x then the value of $a^2 + b^2$ is

A. 3

B. 4

C. 5

D. 11

Answer: C



Watch Video Solution

21. Let $f(x)$ be continuous function given by $f(x) = \{2x, |x| \leq 1$ and $x^2 + ax + b, |x| > 1\}$.

Find the area of the region in the third quadrant bounded by the curves $x = -2y^2$ and $y = f(x)$ lying on the left of the line $8x + 1 = 0$.

A. sq. units

B. $\frac{257}{192}$ sq. units

C. $\frac{257}{96}$ sq. units

D. $\frac{289}{192}$ sq. units

Answer: B



Watch Video Solution

Matrix Match Type

1. Match the following lists :

List I	List II
a. The area bounded by the curve $y = x x $, x-axis and the ordinates $x = 1, x = -1$	p. $10/3$ sq. units
b. The area of the region lying between the lines $x - y + 2 = 0, x = 0$, and the curve $x = \sqrt{y}$	q. $64/3$ sq. units
c. The area enclosed between the curves $y^2 = x$ and $y = x $	r. $2/3$ sq. units
d. The area bounded by parabola $y^2 = x$, straight line $y = 4$, and the y-axis	s. $1/6$ sq. units

 [Watch Video Solution](#)

2. show that the tangents to the curve $y = 2x^3 - 3$ at the point where $x=2$ and $x=-2$ are parallel

 [Watch Video Solution](#)

3. find the point on the curve $y = 2x^2 - 6x - 4$ at which the tangent is parallel to the x-axis



Watch Video Solution

4. Find the point on the curve $y = x^3 - 11x + 5$ at which the tangent has the equation $y = x - 11$



Watch Video Solution

5. find the equation of the tangent to the curve $y = -5x^2 + 6x + 7$ at the point $(1/2, 35/4)$



Watch Video Solution

Numerical Value Type

1. The area enclosed by the curve $c: y = x\sqrt{9 - x^2}$ ($x \geq 0$) and the x-axis is _____



Watch Video Solution

2. Let S be the area bounded by the curve $y = \sin x (0 \leq x \leq \pi)$ and the x-axis and T be the area bounded by the curves $y = \sin x (0 \leq x \leq \frac{\pi}{2})$, $y = a \cos x (0 \leq x \leq \frac{\pi}{2})$, and the x-axis (where $a \in R^+$). The value of $(3a)$ such that $S:T = 1:\frac{1}{3}$ is _____

 [Watch Video Solution](#)

3. Let C be a curve passing through $M(2,2)$ such that the slope of the tangent at any point to the curve is reciprocal of the ordinate of the point. If the area bounded by curve C and line $x=2$ is A , then the value of $\frac{3A}{2}$ is __.

 [Watch Video Solution](#)

4. The area enclosed by $f(x) = 12 + ax \pm x^2$ coordinates axes and the ordinates at $x = 3 (f(3) > 0)$ is 45 sq. units. If m and n are the x-axis intercepts of the graph of $y=f(x)$, then the value of $(m+n+a)$ is __.



Watch Video Solution

5. If the area bounded by the curve $f(x) = x^{1/3}(x - 1)$ and the x-axis is A , then the value of $28A$ is __.



Watch Video Solution

6. If the area bounded by the curve $y = x^2 + 1$ and the tangents to it drawn from the origin is A , then the value of $3A$ is __.



Watch Video Solution

7. If the area enclosed by the curve $y = \sqrt{x}$ and $x = -\sqrt{y}$, the circle $x^2 + y^2 = 2$ above the x-axis is A , then the value of $\frac{16}{\pi}A$ is __.



Watch Video Solution

8. The value of $a(a > 0)$ for which the area bounded by the curves $y = \frac{x}{6} + \frac{1}{x^2}$, $y = 0$, $x = a$, and $x = 2a$ has the least value is ___.

 [Watch Video Solution](#)

9. Area bounded by the curve $[|x|] + [|y|] = 3$, where $[.]$ denotes the greatest integer function

 [Watch Video Solution](#)

10. The area bounded by the curves $y = x(x - 3)^2$ and $y = x$ is _____ (in sq. units)

 [Watch Video Solution](#)

11. If the area of the region $\{(x, y) : 0 \leq y \leq x^2 + 1, 0 \leq y \leq x + 1, 0 \leq x \leq 2\}$ is A , then the

value of $3A - 17$ is ____

 [Watch Video Solution](#)

12. If S is the sum of possible values of c for which the area of the figure bounded by the curves $y = \sin 2x$, the straight lines $x = \frac{\pi}{6}$, $x = c$, and the abscissa axis is equal to $\frac{1}{2}$, then the value of π/S is ____

 [Watch Video Solution](#)

13. If A is the area bounded by the curves $y = \sqrt{1 - x^2}$ and $y = x^3 - x$, then of $\frac{\pi}{A}$.

 [Watch Video Solution](#)

14. Consider two curves $C_1: y = \frac{1}{x}$ and $C_2: y = 1nx$ on the xy plane. Let D_1 denotes the region surrounded by C_1 , C_2 , and the line $x = 1$ and D_2

denotes the region surrounded by C_1, C_2 and the line $x = a$. If

$D_1 = D_2$, then the sum of logarithm of possible value of a is _____



[Watch Video Solution](#)

15. If ' a ' ($a > 0$) is the value of parameter for each of which the area of

the figure bounded by the straight line $y = \frac{a^2 - ax}{1 + a^4}$ and the parabola

$y = \frac{x^2 + 2ax + 3a^2}{1 + a^4}$ is the greatest, then the value of a^4 is ___



[Watch Video Solution](#)

16. If S is the sum of cubes of possible value of c for which the area of the

figure bounded by the curve $y = 8x^2 - x^5$, then straight lines

$x = 1$ and $x = c$ and the abscissa axis is equal to $\frac{16}{3}$, then the value of

$[S]$, where $[.]$ denotest the greatest integer function, is ___



[Watch Video Solution](#)

17. For a point P in the plane, let $d_1(P)$ and $d_2(P)$ be the distances of the point P from the lines $x - y = 0$ and $x + y = 0$ respectively. The area of the region R consisting of all points P lying in the first quadrant of the plane and satisfying $2 \leq d_1(P) + d_2(P) \leq 4$, is

 [Watch Video Solution](#)

18. A farmer F_1 has a land in the shape of a triangle with vertices at $P(0, 0)$, $Q(1, 1)$ and $R(2, 0)$. From this land, a neighbouring farmer F_2 takes away the region which lies between the side PQ and a curve of the form $y = x^n$ ($n > 1$). If the area of the region taken away by the farmer F_2 is exactly 30% of the area of PQR , then the value of n is _____.

 [Watch Video Solution](#)

1. The area of the region bounded by the parabola $(y - 2)^2 = x - 1$, the tangent to the parabola at the point (2,3) and the X-axis is

- A. 3
- B. 6
- C. 9
- D. 12

Answer: C



[Watch Video Solution](#)

2. The area bounded by the curves $y = \cos x$ and $y = \sin x$ between the ordinates $x = 0$ and $x = \frac{3\pi}{2}$ is

- A. $4\sqrt{2} + 1$
- B. $4\sqrt{2} - 1$
- C. $4\sqrt{2} + 2$

D. $4\sqrt{2} - 2$

Answer: D



Watch Video Solution

3. The area of the region enclosed by the curve $y = x$, $x = e$, $y = \frac{1}{x}$ and the positive X-axis is

A. $\frac{5}{2}$ square units

B. $\frac{1}{2}$ square units

C. 1 square units

D. $\frac{3}{2}$ square units

Answer: D



Watch Video Solution

4. The area bounded between the parabolas $x^2 = \frac{y}{4}$ and $x^2 = 9y$ and the straight line $y=2$ is

A. $20\sqrt{2}$

B. $\frac{10\sqrt{2}}{3}$

C. $\frac{20\sqrt{2}}{3}$

D. $10\sqrt{2}$

Answer: C



[Watch Video Solution](#)

5. The area (in square units) bounded by the curves $y = \sqrt{x}$, $2y - x + 3 = 0$, x-axis, and lying in the first quadrant is

A. 9

B. 36

C. 18

D. $\frac{27}{4}$

Answer: A



Watch Video Solution

6. The area of the region described by

$$A = \{(x, y) : x^2 + y^2 \leq 1 \text{ and } y^2 \leq 1 - x\}$$
 is

A. $\frac{\pi}{2} + \frac{4}{3}$

B. $\frac{\pi}{2} - \frac{4}{3}$

C. $\frac{\pi}{2} - \frac{2}{3}$

D. $\frac{\pi}{2} + \frac{2}{3}$

Answer: D



Watch Video Solution

7. The area (in sq. units) of the region described by $\{x, y : y^2 \leq 2x \text{ and } y \geq 4x - 1\}$ is

A. $\frac{7}{32}$

B. $\frac{5}{64}$

C. $\frac{15}{64}$

D. $\frac{9}{32}$

Answer: D



[Watch Video Solution](#)

8. The area (in sq units) of the region $\{(x, y) : y^2 \geq 2x \text{ and } x^2 + y^2 \leq 4x, x \geq 0, y \geq 0\}$ is

A. $\pi - \frac{8}{3}$

B. $\pi - \frac{4\sqrt{2}}{3}$

C. $\frac{\pi}{2} - \frac{2\sqrt{2}}{3}$

D. $\pi - \frac{4}{3}$

Answer: A

 [Watch Video Solution](#)

9. The area (in sq units) of the region bounded by the curve $y = \sqrt{x}$ and the lines $y = 0$, $y = x - 2$, is

A. $\frac{5}{2}$

B. $\frac{59}{12}$

C. $\frac{3}{2}$

D. $\frac{7}{3}$

Answer: A

 [Watch Video Solution](#)

10. Let $g(x) = \cos^2 x$, $f(x) = \sqrt{x}$ and α, β (alpha

A. $\frac{1}{2}(\sqrt{2} - 1)$

B. $\frac{1}{2}(\sqrt{3} - 1)$

C. $\frac{1}{2}(\sqrt{3} + 1)$

D. $\frac{1}{2}(\sqrt{3} - \sqrt{2})$

Answer: B



Watch Video Solution

Single Correct Answer Type

1. Let the straight line $x = b$ divide the area enclosed by $y = (1 - x)^2$, $y = 0$, and $x = 0$ into two parts $R_1(0 \leq x \leq b)$ and $R_2(b \leq x \leq 1)$ such that $R_1 - R_2 = \frac{1}{4}$. Then b equals

A. $\frac{3}{4}$

B. $\frac{1}{2}$

C. $\frac{1}{3}$

D. $\frac{1}{4}$

Answer: B

 [Watch Video Solution](#)

2. The area enclosed by the curves

$y = \sin x + \cos x$ and $y = |\cos x - \sin x|$ over the interval $\left[0, \frac{\pi}{2}\right]$ is (a)

(a) $4(\sqrt{2} - 1)$ (b) $2\sqrt{2}(\sqrt{2} - 1)$ (c) $2(\sqrt{2} + 1)$ (d) $2\sqrt{2}(\sqrt{2} + 1)$

A. $4(\sqrt{2} - 1)$

B. $2\sqrt{2}(\sqrt{2} - 1)$

C. $2(\sqrt{2} + 1)$

D. $2\sqrt{2}(\sqrt{2} + 1)$

Answer: B



Watch Video Solution

3. Area of the region $\{(x, y) \in \mathbb{R}^2 : y \geq \sqrt{|x + 3|}, 5y \leq x + 9 \leq 15\}$ is equal to

A. $\frac{1}{6}$

B. $\frac{4}{3}$

C. $\frac{3}{2}$

D. $\frac{5}{3}$

Answer: C



Watch Video Solution

4. The area enclosed between the curve $y = \sin^2 x$ and $y = \cos^2 x$ in the interval $0 \leq x \leq \pi$ is _____ sq. units.

A. 2 sq unit

B. $\frac{1}{2}$ sq unit

C. 1 sq unit

D. None of these

Answer: C



Watch Video Solution

5. The area of the region enclosed by $y = x^2$ and $y = \sqrt{|x|}$ is

A. $1/3$

B. $2/3$

C. $1/6$

D. 1

Answer: B



Watch Video Solution

6. The area of the region bounded by $y = e^x$, $y = e^{-x}$, $x = 0$ and $x = 1$ is

(a) $e + \frac{1}{e}$ (b) $\log\left(\frac{4}{e}\right)$ (c) $4\log\left(\frac{4}{e}\right)$ (d) $e + \frac{1}{e} - 2$

A. $e + \frac{1}{e}$

B. $\log(4/e)$

C. $4\log(4/e)$

D. $e + \frac{1}{e} - 2$

Answer: D



Watch Video Solution

7. The area bounded by the curve $y = |\cos^{-1}(\sin x)| - |\sin^{-1}(\cos x)|$

and axis from $\frac{3\pi}{2} \leq x \leq 2\pi$

A. π^2 sq. units

B. $\pi^2 / 4$ sq. units

C. $\pi^2 / 2$ sq. units

D. none of these

Answer: B



[Watch Video Solution](#)

8. If $(a, 0)$, $a > 0$, is the point where the curve $y = \sin 2x - \sqrt{3} \sin x$ cuts the x-axis first, A is the area bounded by this part of the curve, the origin and the positive x-axis. Then

A. $4A + 8 \cos a = 7$

B. $4 + 8 \sin a = 7$

C. $4A - 8 \sin a = 7$

D. $4A - 8 \cos a = 7$

Answer: A



[Watch Video Solution](#)

9. The area in the first quadrant between $x^2 + y^2 = \pi^2$ and $y = \sin x$ is

(a) $\frac{(\pi^3 - 8)}{4}$ (b) $\frac{\pi^3}{4}$ (c) $\frac{(\pi^3 - 16)}{4}$ (d) $\frac{(\pi^3 - 8)}{2}$

A. $\frac{(\pi^3 - 8)}{4}$

B. $\frac{\pi^3}{4}$

C. $\frac{(\pi^3 - 16)}{4}$

D. $\frac{(\pi^3 - 8)}{2}$

Answer: A



Watch Video Solution

10. The area bounded by the curves

$y = \cos^{-1} x$, $y = \sin^{-1} x$ and $y = -\pi x^3$, where $-1 \leq x \leq 1$, is

A. $\frac{3\pi}{2} + 1 - \sqrt{2}$ sq. units

B. $\frac{3\pi}{4} + 1 + \sqrt{2}$ sq. units

C. $\frac{3\pi}{4} + 2 - \sqrt{2}$ sq. units

D. $\frac{3\pi}{4} + 1 - \sqrt{2}$ sq. units

Answer: D



[View Text Solution](#)

11. The area bounded by the curve $y = \sin^2 x - 2\sin x$ and the x-axis, where $x \in [0, 2\pi]$, is

A. 4 sq. units

B. 8 sq. units

C. 16 sq. units

D. 20 sq. units

Answer: B



[Watch Video Solution](#)

12. Consider the functions $f(x)$ and $g(x)$, both defined from $R \rightarrow R$ and are defined as $f(x) = 2x - x^2$ and $g(x) = x^n$ where $n \in N$. If the area between $f(x)$ and $g(x)$ is $1/2$, then the value of n is

- A. 5
- B. 6
- C. 7
- D. 8

Answer: A



[Watch Video Solution](#)

13. Let a function $f(x)$ be defined in $[-2, 2]$ as

$$f(x) = \begin{cases} \{x\}, & -2 \leq x < -1 \\ |\operatorname{sgn} x|, & -1 \leq x \leq 1 \\ \{-x\}, & 1 < x \leq 2 \end{cases} \quad \text{where } \{x\} \text{ and } \operatorname{sgn} x \text{ denote}$$

fractional part and signum functions, respectively. Then find the area bounded by the graph of $f(x)$ and the x-axis.

A. 2 sq. units

B. 3 sq. units

C. 4 sq. units

D. 5 sq. units

Answer: B



[Watch Video Solution](#)

14. The area bounded by $y = x^2 + 2$ and $y = 2|x| - \cos \pi x$ is equal to

A. $2/3$

B. $8/3$

C. $4/3$

D. $1/3$

Answer: B



[Watch Video Solution](#)

15. Area bounded by $f(x) = \frac{x^2 - 1}{x^2 + 1}$ and the line $y = 1$ is

- A. π sq. units
- B. 2π sq. units
- C. $\pi/2$ sq. units
- D. none of these

Answer: B



[Watch Video Solution](#)

16. The area bounded by the curve $y = xe^{-x}$; $xy = 0$ and $x = c$ where c is the x-coordinate of the curve's inflection point, is

- A. $1 - 3e^{-2}$
- B. $1 - 2e^{-2}$
- C. $1 - e^{-2}$

D. none of these

Answer: A

 [Watch Video Solution](#)

17. Area of region bounded by the curve $y = \frac{16 - x^2}{4}$ and $y = \sec^{-1}[-\sin^2 x]$ (where $[x]$ denotes the greatest integer function) is

A. $\frac{1}{3}(4 - \pi)^{3/2}$

B. $8(4 - \pi)^{3/2}$

C. $\frac{8}{3}(4 - \pi)^{3/2}$

D. $\frac{8}{3}(4 - \pi)^{1/2}$

Answer: C

 [Watch Video Solution](#)

18. Suppose $y = f(x)$ and $y = g(x)$ are two continuous functions whose graphs intersect at the three points $(0, 4)$, $(2, 2)$ and $(4, 0)$ with $f(x) > g(x)$ for $0 < x < 2$ and $f(x) < g(x)$ for $2 < x < 4$. If $\int_0^4 [f(x) - g(x)] dx = 10$ and $\int_2^4 [g(x) - f(x)] dx = 5$ the area between two curves for $0 < x < 4$ is (A) 5 (B) 10 (C) 15 (D) 20

A. 5

B. 10

C. 15

D. 20

Answer: C



Watch Video Solution

19. The ratio of the areas of two regions of the curve

$C_1 \equiv 4x^2 + \pi^2 y^2 = 4\pi^2$ divided by the curve

$C_2 \equiv y = -\left(\operatorname{sgn}\left(x - \frac{\pi}{2}\right)\right) \cos x$ (where $\operatorname{sgn}(x) = \operatorname{signum}(x)$) is

A. $\frac{\pi^2 - 2}{\pi^2 - 2\sqrt{2}}$

B. $\frac{\pi^2 + 2}{\pi^2 - 2\sqrt{2}}$

C. $\frac{\pi^2 + 6}{\pi^2 + 3\sqrt{2}}$

D. $\frac{\pi^2 - 1}{\pi^2 - \sqrt{2}}$

Answer: A



Watch Video Solution

20. The area bounded by the curves

$$x\sqrt{3} + y = 2\log_e(x - y\sqrt{3}) - 2\log_e 2, y = \sqrt{3}x,$$

$$y = -\frac{1}{\sqrt{3}}x + 2, \text{ is}$$

A. $2\log_e 2$ sq. units

B. $2\log_e 2 + 1$ sq. units

C. $2\log_e 2 - 1$ sq. units

D. $4\log_e 2 - 1$ sq. units

Answer: C

 [Watch Video Solution](#)

21. Area of region bounded by the curve $y = \frac{4 - x^2}{4 + x^2}$, $25y^2 = 9x$ and $y = \frac{3}{5}|x| - \frac{6}{5}$ which contains (1, 0) point in its interior is

A. $\left\{ \pi - 4 \tan^{-1} \cdot \frac{1}{2} + \frac{1}{10} \right\}$ sq. units

B. $\left\{ \pi - 2 \tan^{-1} \cdot \frac{1}{2} - \frac{1}{5} \right\}$ sq. units

C. $\left\{ \pi + 4 \tan^{-1} \cdot \frac{1}{2} - \frac{1}{5} \right\}$ sq. units

D. none of these

Answer: A

 [Watch Video Solution](#)

22. Area bounded by the min. $\{|x|, |y|\} = 1$ and the max. $\{|x|, |y|\} = 2$ is

A. 4

B. 8

C. 16

D. 9

Answer: A



Watch Video Solution

23. Consider $f(x) = \begin{cases} \cos x & 0 \leq x < \frac{\pi}{2} \\ \left(\frac{\pi}{2} - x\right)^2 & \frac{\pi}{2} \leq x < \pi \end{cases}$ such that f is periodic

with period π . Then which of the following is not true?

A. The range of f is $\left[0, \frac{\pi^2}{4}\right)$.

B. f is discontinuous for infinite values of x .

C. The area bounded by $y = f(x)$ and the X-axis from $x = 0$ to $x = n\pi$ is

$n\left(1 + \frac{\pi^3}{24}\right)$ for a given $n \in \mathbb{N}$.

D. none of these

Answer: D



Watch Video Solution

24. The area made by curve $f(x) = [x] + \sqrt{x - [x]}$ and x-axis when $0 \leq x \leq n$ ($n \in N$) is equal to { where $[x]$ is greatest integer function}

A. $\frac{2n}{3} + \frac{n(n+1)}{2}$

B. $\frac{n}{3} + \frac{n(n+1)}{2}$

C. $\frac{2n}{3} + \frac{n(n-1)}{2}$

D. $\frac{n}{3} + \frac{n(n-1)}{2}$

Answer: C



View Text Solution

25. Consider the regions

$A = \{(x, y) \mid x^2 + y^2 \leq 100\}$ and $B = \{x \mid \sin(x + y) > 0\}$ in the

plane. Then the area of the region $A \cap B$ is

A. 10π

B. 100

C. 100π

D. 50π

Answer: D



[Watch Video Solution](#)

26. Let R be the region containing the point (x, y) on the X - Y plane, satisfying $2 \leq |x + 3y| + |x - y| \leq 4$. Then the area of this region is

A. 5 sq. units

B. 6 sq. units

C. 7 sq. units

D. 8 sq. units

Answer: B

 [Watch Video Solution](#)

27. If $f(x) = \begin{cases} \sqrt{\{x\}} & \text{for } x \notin Z \\ 1 & \text{for } x \in Z \end{cases}$ and $g(x) = \{x\}^2$ where $\{.\}$ denotes

fractional part of x then area bounded by $f(x)$ and $g(x)$ for $x \in 0, 6$ is

A. $\frac{2}{3}$

B. 2

C. $\frac{10}{3}$

D. 6

Answer: B

 [Watch Video Solution](#)

28. Let S is the region of points which satisfies $y^2 < 16x$, $x < 4$ and $\frac{xy(x^2 - 3x + 2)}{x^2 - 7x + 12} > 0$. Its area is

A. $\frac{8}{3}$

B. $\frac{64}{3}$

C. $\frac{32}{3}$

D. none of these

Answer: B



Watch Video Solution

29. The area of the region $\{(x, y) : x^2 + y^2 \leq 5, ||x| - |y| | \geq 1$ is

A. $4\left(\pi - \tan^{-1}\left(\frac{24}{7}\right)\right) - 4$

B. $5\left(\pi - \tan^{-1}\left(\frac{24}{7}\right)\right) - 4$

C. $3\left(\pi - \tan^{-1}\left(\frac{24}{7}\right)\right) - 4$

$$D. 2\left(\pi - \tan^{-1}\left(\frac{24}{7}\right)\right) - 1$$

Answer: B



Watch Video Solution

30. The following figure shows the graph of a continuous function $y = f(x)$ on the interval $[1, 3]$. The points A, B, C have coordinates $(1,1)$, $(3,2)$, $(2,3)$, respectively, and the lines L_1 and L_2 are parallel, with L_1 being tangent to the curve at C. If the area under the graph of $y = f(x)$ from $x = 1$ to $x = 3$ is 4 square units, then the area of the shaded region is



- A. 1
- B. 2
- C. 3
- D. 4

Answer: B



Watch Video Solution

Multiple Correct Answer Type

1. If the area of bounded between the x-axis and the graph of $y = 6x - 3x^2$ between the ordinates $x = 1$ and $x = a$ is 19 units, then a can take the value 4 or -2 two value are in (2,3) and one in $(-1, 0)$ two value are in (3,4) and one in $(-2, -1)$ none of these

A. one value in (2, 3)

B. one value in $(-2, -1)$

C. one value in $(-1, 0)$

D. one value in (3, 4)

Answer: B::D



Watch Video Solution

2. Which of the following is the possible value/values of c for which the area of the figure bounded by the curves $y = \sin 2x$, the straight lines $x = \pi/6$, $x = c$ and the abscissa axis is equal to $1/2$?

A. $-\frac{\pi}{6}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{6}$

D. none of these

Answer: B



Watch Video Solution

3. Area of the region bounded by the curve $y = \tan x$ and lines $y = 0$ and $x = 1$ is

A. $\int_0^1 (1 - x) dx$

B. $\tan 1 - \int_0^{\tan 1} \tan^{-1} y dy$

$$C. \int_0^{\tan^{-1} 1} \tan^{-1} y dy$$

$$D. \int_0^1 \tan^{-1} x dx$$

Answer: A:B



Watch Video Solution

Comprehension Type

1. In the following figure, the graphs of two functions $y = f(x)$ and $y = \sin x$ are given. They intersect at origin, $A(a, f(a))$, $B(\pi, 0)$ and $C(2\pi, 0)$. $A_i (i = 1, 2, 3)$ is the area bounded by the curves as shown in the figure, respectively, for $x \in (0, a)$, $x \in (a, \pi)$, $x \in (\pi, 2\pi)$.

If $A_1 = 1 + (a - 1)\cos a - \sin a$, then



The function $f(x)$ is

A. $x^2 \sin x$

B. $x \sin x$

C. $2x \sin x$

D. $x^3 \sin x$

Answer: B



Watch Video Solution

2. In the following figure, the graphs of two functions $y = f(x)$ and $y = \sin x$ are given. They intersect at origin, $A(a, f(a))$, $B(\pi, 0)$ and $C(2\pi, 0)$. $A_i (i = 1, 2, 3)$ is the area bounded by the curves as shown in the figure, respectively, for $x \in (0, a)$, $x \in (a, \pi)$, $x \in (\pi, 2\pi)$.

If $A_1 = 1 + (a - 1)\cos a - \sin a$, then



The function $f(x)$ is

A. $(\pi - 1)\text{units}^2$

B. $(\pi/2 - 1)\text{units}^2$

C. $(\pi - \sin 1 - 1)\text{units}^2$

D. $\pi/2\text{units}^2$

Answer: C



Watch Video Solution