



## MATHS

### BOOKS - CENGAGE MATHS (ENGLISH)

#### CROSS PRODUCTS

#### Dpp 2 2

1. Let  $\vec{a}$  and  $\vec{b}$  be two vectors of equal magnitude 5 units. Let  $\vec{p}$ ,  $\vec{q}$  be vectors such that  $\vec{p} = \vec{a} - \vec{b}$  and  $\vec{q} = \vec{a} + \vec{b}$ . If

$$|\vec{p} \times \vec{q}| = 2 \left\{ \lambda - (\vec{a} \cdot \vec{b})^2 \right\}^{\frac{1}{2}}, \text{ then value of } \lambda \text{ is}$$

A. 25

B. 125

C. 625

D. none of these

**Answer: C**



**Watch Video Solution**

2. Let  $\vec{u} = 2\hat{i} - \hat{j} + \hat{k}$ ,  $\vec{v} = -3\hat{j} + 2\hat{k}$  be vectors and  $\vec{w}$  be a unit vector in the xy-plane. Then the maximum possible value of  $\left|(\vec{u} \times \vec{v}) \cdot \vec{w}\right|$  is

- A.  $\sqrt{5}$
- B.  $\sqrt{12}$
- C.  $\sqrt{13}$
- D.  $\sqrt{17}$

**Answer: D**



**Watch Video Solution**

3. Let  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are three unit vectors in a plane such that they are equally inclined to each other, then the value of  $(\vec{a} \times \vec{b}) \cdot (\vec{b} \times \vec{c}) + (\vec{b} \times \vec{c}) \cdot (\vec{c} \times \vec{a}) + (\vec{c} \times \vec{a}) \cdot (\vec{a} \times \vec{b})$  can be

A.  $\frac{9}{4}$

B.  $-\frac{9}{4}$

C.  $\frac{3}{4}$

D.  $-\frac{3}{4}$

**Answer: A**



[Watch Video Solution](#)

4. The coordinates of the mid-points of the sides of  $\Delta PQR$ , are  $(3a, 0, 0)$ ,  $(0, 3b, 0)$  and  $(0, 0, 3c)$  respectively, then the area of  $\Delta PQR$  is

A.  $18\sqrt{b^2c^2 + c^2a^2 + a^2b^2}$

B.  $9\sqrt{b^2c^2 + c^2a^2 + a^2b^2}$

C.  $\frac{9}{12}\sqrt{b^2c^2 + c^2a^2 + a^2b^2}$

D.  $\frac{9}{2}\sqrt{b^2c^2 + c^2a^2 + a^2b^2}$

**Answer: A**



**Watch Video Solution**

5. If  $\vec{a} = (\hat{i} + \hat{j} + \hat{k})$ ,  $\vec{a} \cdot \vec{b} = 1$  and  $\vec{a} \times \vec{b} = \hat{j} - \hat{k}$ , then  $\vec{b}$  is

(a)  $\hat{i} - \hat{j} + \hat{k}$  (b)  $2\hat{i} - \hat{k}$  (c)  $\hat{i}$  (d)  $2\hat{i}$

A. 3

B. 9

C. 10

D. 12

**Answer: A**



**Watch Video Solution**

6. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors such that  $\vec{a} \cdot \vec{b} = 0$ ,  $(\vec{a} - \vec{c}) \cdot (\vec{b} + \vec{c}) = 0$  and  $\vec{c} = \lambda \vec{a} + \mu \vec{b} + \omega (\vec{a} \times \vec{b})$ , where  $\lambda, \mu, \omega$  are scalars, then

A.  $\mu^2 + \omega^2 = 1$

B.  $\lambda + \mu = 1$

C.  $(\mu + 1)^2 + \mu^2 + \omega^2 = 1$

D.  $\lambda^2 + \mu^2 = 1$

**Answer: C**



**Watch Video Solution**

7. Let  $\triangle ABC$  be a given triangle. If  $\left| \overrightarrow{BA} - t\overrightarrow{BC} \right| \geq \left| \overrightarrow{AC} \right|$  for any  $t \in R$ , then  $\triangle ABC$  is

A. Equilateral

B. Right angled

C. Isosceles

D. None of these

**Answer: B**



[Watch Video Solution](#)

8. If  $\vec{a}, \vec{b}$  are vectors perpendicular to each other and  $|\vec{a}| = 2, |\vec{b}| = 3, \vec{c} \times \vec{a} = \vec{b}$ , then the least value of  $2|\vec{c} - \vec{a}|$  is

A. 1

B. 2

C. 3

D. 4

**Answer: C**



[Watch Video Solution](#)

9.  $\vec{a}$  and  $\vec{b}$  are two vectors such that  $|\vec{a}| = 1$ ,  $|\vec{b}| = 4$  and  $\vec{a} \cdot \vec{b} = 2$ . If  $\vec{c} = (2\vec{a} \times \vec{b}) - 3\vec{b}$  then find angle between  $\vec{b}$  and  $\vec{c}$ .

A.  $\frac{\pi}{3}$

B.  $\frac{\pi}{6}$

C.  $\frac{3\pi}{4}$

D.  $\frac{5\pi}{6}$

**Answer: D**



**Watch Video Solution**

10. If  $\vec{a}$  and  $\vec{b}$  are non-zero, non parallel vectors, then the value of

$$\left| \vec{a} + \vec{b} + \vec{a} \times \vec{b} \right|^2 + \left| \vec{a} + \vec{b} - \vec{a} \times \vec{b} \right|^2 \text{ equals}$$

A.  $(1 + \vec{a} \cdot \vec{a})(1 + \vec{b} \cdot \vec{b})$

B.  $2(1 + \vec{a} \cdot \vec{a})(1 + \vec{b} \cdot \vec{b})$

$$C. 2 \left\{ \left( 1 + \vec{a} \cdot \vec{a} \right) \left( 1 + \vec{b} \cdot \vec{b} \right) - \left( 1 - \vec{a} \cdot \vec{b} \right)^2 \right\}$$

$$D. 2 \left\{ \left( 1 - \vec{a} \cdot \vec{a} \right) \left( 1 - \vec{b} \cdot \vec{b} \right) + \left( 1 - \vec{a} \cdot \vec{b} \right)^2 \right\}$$

**Answer: C**



**Watch Video Solution**

11. If  $a^2 + b^2 + c^2 = 1$  where,  $a, b, c \in R$ , then the maximum value of  $(4a - 3b)^2 + (5b - 4c)^2 + (3c - 5a)^2$  is

A. 25

B. 50

C. 144

D. none of these

**Answer: B**



**Watch Video Solution**



12. Three vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are such that  $\vec{a} \times \vec{b} = 4(\vec{a} \times \vec{c})$  and  $|\vec{a}| = |\vec{b}| = 1$  and  $|\vec{c}| = \frac{1}{4}$ . If the angle between  $\vec{b}$  and  $\vec{c}$  is  $\frac{\pi}{3}$  then  $\vec{b}$  is

A.  $\vec{a} + 4\vec{c}$

B.  $\vec{a} - 4\vec{c}$

C.  $4\vec{c} - \vec{a}$

D.  $2\vec{c} - \vec{a}$

Answer: A:C

 Watch Video Solution

13. If  $2\vec{a}$ ,  $3\vec{b}$ ,  $2(\vec{a} \times \vec{b})$  are position vectors of the vertices A, B, C, of  $\triangle ABC$  and  $|\vec{a}| = |\vec{b}| = 1$ ,  $\vec{OA} \cdot \vec{OB} = -3$  (where O is the origin), then

A. Triangle ABC is right-angled triangle

B. Angle B is  $90^\circ$

C.  $A = \cos^{-1}\left(\sqrt{\frac{7}{19}}\right)$

D. The position vector of orthocenter is  $2\left(\vec{a} \times \vec{b}\right)$

**Answer: A::C::D**



**Watch Video Solution**