



## MATHS

### BOOKS - CENGAGE MATHS (ENGLISH)

#### HYPERBOLA

#### Illustration

1. If the base of a triangle and the ratio of tangent of half of base angles are given, then identify the locus of the opposite vertex.

 [Watch Video Solution](#)

2. Prove that the locus of centre of the circle which touches two given disjoint circles externally is hyperbola.

 [Watch Video Solution](#)

3. The equation of one of the directrices of a hyperbola is  $2x + y = 1$ , the corresponding focus is  $(1, 2)$  and  $e = \sqrt{3}$ . Find the equation of the hyperbola and the coordinates of the center and the second focus.

 [Watch Video Solution](#)

4. The eccentricity of the conic represented by  $2x^2 + 5xy + 2y^2 + 11x - 7y - 4 = 0$  is

 [Watch Video Solution](#)

5. If equation

$$\left| \sqrt{(x - \tan \theta)^2 + (y - \sqrt{3} \tan \theta)^2} - \sqrt{(x - 2 \tan \theta)^2 + y^2} \right| = 2, \theta \in [0, \pi]$$

represents hyperbola, then find the value of  $\theta$ .

 [Watch Video Solution](#)

6. The distance between the foci of a hyperbola is 16 and its eccentricity is  $\sqrt{2}$  then equation of the hyperbola is

 [Watch Video Solution](#)

7. If the foci of the ellipse  $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$  and the hyperbola  $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$  coincide, then find the value

 [Watch Video Solution](#)

8. If hyperbola  $\frac{x^2}{b^2} - \frac{y^2}{a^2} = 1$  passes through the focus of ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , then find the eccentricity of hyperbola.

 [Watch Video Solution](#)

9. Find the eccentricity of the hyperbola given by equations  $x = \frac{e^t + e^{-1}}{2}$  and  $y = \frac{e^t - e^{-1}}{3}$ ,  $t \in R$ .

 [Watch Video Solution](#)

10. An ellipse and a hyperbola have their principal axes along the coordinate axes and have a common foci separated by distance  $2\sqrt{3}$ . The difference of their focal semi-axes is equal to 4. If the ratio of their eccentricities is  $3/7$ , find the equation of these curves.

 [Watch Video Solution](#)

11. If the latus rectum subtends a right angle at the center of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then find its eccentricity.

 [Watch Video Solution](#)

12. Find the equation of hyperbola if centre is  $(1, 0)$ , one focus is  $(6, 0)$  and transverse axis 6 .

 [Watch Video Solution](#)

13. Two rods are rotating about two fixed points in opposite directions. If they start from their position of coincidence and one rotates at the rate double that of the other, then find the locus of point of the intersection of the two rods.

 [Watch Video Solution](#)

14. Find the coordinates of the foci, the eccentricity, the latus rectum, and the equations of directrices for the hyperbola  $9x^2 - 16y^2 - 72x + 96y - 144 = 0$

 [Watch Video Solution](#)

15. Find the coordinates of the foci and the center of the hyperbola,  $x^2 - 3y^2 - 4x = 8$

 [Watch Video Solution](#)

16. about to only mathematics

A.  $x^2 + 2y^2 < 1$

B.  $\max\{|x|, |y|\} < 1$

C.  $x^2 - y^2 < 1$

D.  $y^2 - x < 0$

Answer:



Watch Video Solution

17. Find the locus of the midpoints of chords of hyperbola  $3x^2 - 2y^2 + 4x - 6y = 0$  parallel to  $y = 2x$ .



Watch Video Solution

18. If  $PQ$  is a double ordinate of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  such that  $OPQ$  is an equilateral triangle,  $O$  being the center of the hyperbola, then

find the range of the eccentricity  $e$  of the hyperbola.

 [Watch Video Solution](#)

19. If  $(a \sec \theta, b \tan \theta)$  and  $(a \sec \phi, b \tan \phi)$  are the ends of the focal chord of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  then prove that  $\tan\left(\frac{x}{a}\right)\tan\left(\frac{\phi}{2}\right) = \frac{1 - e}{1 + e}$

 [Watch Video Solution](#)

20. Find the point on the hyperbola  $x^2 - 9y^2 = 9$  where the line  $5x + 12y = 9$  touches it.

 [Watch Video Solution](#)

21. Find the value of  $m$  for which  $y = mx + 6$  is a tangent to the hyperbola  $\frac{x^2}{100} - \frac{y^2}{49} = 1$

 [Watch Video Solution](#)

22. Find the equation of tangents to the curve  $4x^2 - 9y^2 = 1$  which are parallel to  $4y = 5x + 7$ .

 [Watch Video Solution](#)

23. If it is possible to draw the tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  having slope 2, then find the range of eccentricity

 [Watch Video Solution](#)

24. Find the equation of tangents to hyperbola  $x^2 - y^2 - 4x - 2y = 0$  having slope 2.

 [Watch Video Solution](#)

25. Find the minimum value of  $(2 - a - 4 \sec \theta)^2 + (a - 3 \tan \theta)^2, a \in \mathbb{R}$ .



 [Watch Video Solution](#)

26. Find the locus of the mid points of the chords of the circle  $x^2 + y^2 = 16$ , which are tangent to the hyperbola  $9x^2 - 16y^2 = 144$

 [Watch Video Solution](#)

27. Find the equation of tangent to the conic  $x^2 - y^2 - 8x + 2y + 11 = 0$  at  $(2, 1)$ .

 [Watch Video Solution](#)

28. A tangent to the hyperbola  $x^2 - 2y^2 = 4$  meets x-axis at P and y-axis at Q. Lines PR and QR are drawn such that OPRQ is a rectangle (where O is origin). Find the locus of R.

 [Watch Video Solution](#)

29. Find the equations of the tangents to the hyperbola  $x^2 - 9y^2 = 9$  that are drawn from (3, 2).

 [Watch Video Solution](#)

30. Find the equation of pair of tangents drawn from point (4, 3) to the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ . Also, find the angle between the tangents.

 [Watch Video Solution](#)

31. Tangents drawn from the point (c, d) to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  make angles  $\alpha$  and  $\beta$  with the x-axis.

If  $\tan \alpha \tan \beta = 1$ , then find the value of  $c^2 - d^2$ .

 [Watch Video Solution](#)

32. On which curve does the perpendicular tangents drawn to the hyperbola  $\frac{x^2}{25} - \frac{y^2}{16} = 1$  intersect?



[Watch Video Solution](#)

**33.** Find the equation of hyperbola having foci  $S(2, 1)$  and  $S'(10, 1)$  and a straight line  $x + y - 9 = 0$  as its tangent.



[Watch Video Solution](#)

**34.** Find the eccentricity of the hyperbola with asymptotes  $3x + 4y = 2$  and  $4x - 3y = 2$ .



[Watch Video Solution](#)

**35.** Find the equation of the hyperbola which has  $3x - 4y + 7 = 0$  and  $4x + 3y + 1 = 0$  as its asymptotes and which passes through the origin.



[Watch Video Solution](#)

36. Find the equation of the asymptotes of the hyperbola

$$3x^2 + 10xy + 9y^2 + 14x + 22y + 7 = 0$$



Watch Video Solution

37. If a hyperbola passing through the origin has  $3x - 4y - 1 = 0$  and

$4x - 3y - 6 = 0$  as its asymptotes, then find the equation of its

transvers and conjugate axes.



Watch Video Solution

38. Show that the locus represented by

$$x = \frac{1}{2}a\left(t + \frac{1}{t}\right), y = \frac{1}{2}a\left(t - \frac{1}{t}\right) \text{ is a rectangular hyperbola.}$$



Watch Video Solution

39. If two distinct tangents can be drawn from the point  $(\alpha, \alpha + 1)$  on different branches of the hyperbola  $\frac{x^2}{9} - \frac{y^2}{16} = 1$ , then find the values of  $\alpha$ .

 [Watch Video Solution](#)

40. From a point  $P(1, 2)$ , two tangents are drawn to a hyperbola  $H$  in which one tangent is drawn to each arm of the hyperbola. If the equations of the asymptotes of hyperbola  $H$  are  $\sqrt{3}x - y + 5 = 0$  and  $\sqrt{3}x + y - 1 = 0$ , then the eccentricity of  $H$  is 2 (b)  $\frac{2}{\sqrt{3}}$  (c)  $\sqrt{2}$  (d)  $\sqrt{3}$

 [Watch Video Solution](#)

41. Find the equation of normal to the hyperbola  $x^2 - 9y^2 = 7$  at point  $(4, 1)$ .

 [Watch Video Solution](#)

42. Find the equation of normal to the hyperbola  $3x^2 - y^2 = 1$  having slope  $\frac{1}{3}$ .

 [Watch Video Solution](#)

43. If the normal at  $P(\theta)$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{2a^2} = 1$  meets the transverse axis at  $G$ , then prove that  $AG \cdot A'G = a^2(e^4 \sec^2 \theta - 1)$ , where  $A$  and  $A'$  are the vertices of the hyperbola.

 [Watch Video Solution](#)

44. Normals are drawn to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at point  $\theta_1$  and  $\theta_2$  meeting the conjugate axis at  $G_1$  and  $G_2$ , respectively. If  $\theta_1 + \theta_2 = \frac{\pi}{2}$ , prove that  $CG_1 \cdot CG_2 = \frac{a^2 e^4}{e^2 - 1}$ , where  $C$  is the center of the hyperbola and  $e$  is the eccentricity.

 [Watch Video Solution](#)

45. Let  $P(6,3)$  be a point on the hyperbola parabola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  If the normal at the point intersects the x-axis at  $(9,0)$ , then the eccentricity of the hyperbola is

 [Watch Video Solution](#)

46. Prove that any hyperbola and its conjugate hyperbola cannot have common normal.

 [Watch Video Solution](#)

47. A ray emerging from the point  $(5, 0)$  is incident on the hyperbola  $9x^2 - 16y^2 = 144$  at the point  $P$  with abscissa 8. Find the equation of the reflected ray after the first reflection if point  $P$  lies in the first quadrant.

 [Watch Video Solution](#)

48. Normal to a rectangular hyperbola at P meets the transverse axis at N.

If foci of hyperbola are S and S', then find the value of  $\frac{SN}{SP}$ .

 [Watch Video Solution](#)

49. Consider hyperbola  $xy = 16$  to find the equation of tangent at point (2, 8).

 [Watch Video Solution](#)

50. A triangle has its vertices on a rectangular hyperbola. Prove that the orthocentre of the triangle also lies on the same hyperbola.

 [Watch Video Solution](#)

51. If  $A$ ,  $B$ , and  $C$  are three points on the hyperbola  $xy = c^2$  such that  $AB$  subtends a right angle at  $C$ , then prove that  $AB$  is parallel to the



normal to the hyperbola at point  $C$ .

 [Watch Video Solution](#)

52. Prove that product of parameters of four concyclic points on the hyperbola  $xy = c^2$  is 1. Also, prove that the mean of these four concyclic points bisects the distance between the centres of the hyperbola and the circle.

 [Watch Video Solution](#)

## Solved Examples

1. A variable line  $y = mx - 1$  cuts the lines  $x = 2y$  and  $y = -2x$  at points  $A$  and  $B$ . Prove that the locus of the centroid of triangle  $OAB$  ( $O$  being the origin) is a hyperbola passing through the origin.

 [Watch Video Solution](#)

2. Let  $P$  be a point on the hyperbola  $x^2 - y^2 = a^2$ , where  $a$  is a parameter, such that  $P$  is nearest to the line  $y = 2x$ . Find the locus of  $P$ .

 [Watch Video Solution](#)

3. Show that the midpoints of focal chords of a hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  lie on another similar hyperbola.

 [Watch Video Solution](#)

4. From the center  $C$  of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , perpendicular  $CN$  is drawn on any tangent to it at the point  $P(a \sec \theta, b \tan \theta)$  in the first quadrant. Find the value of  $\theta$  so that the area of  $CPN$  is maximum.

 [Watch Video Solution](#)

5. Semi transverse axis of hyperbola is 5. Tangent at point P and normal to this tangent meet conjugate axis at A and B, respectively. The circle on AB as diameter passes through two fixed points, the distance between which is 20. Find the eccentricity of hyperbola.

 [Watch Video Solution](#)

6. The exhaustive set of values of  $\alpha^2$  such that there exists a tangent to the ellipse  $x^2 + \alpha^2 y^2 = \alpha^2$  and the portion of the tangent intercepted by the hyperbola  $\alpha^2 x^2 - y^2 = 1$  subtends a right angle at the center of the curves is:

 [Watch Video Solution](#)

7. Prove that the part of the tangent at any point of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  intercepted between the point of contact and the transverse axis is a harmonic mean between the lengths of the perpendiculars drawn from the foci on the normal at the same point.



[Watch Video Solution](#)

8. If one of varying central conic (hyperbola) is fixed in magnitude and position, prove that the locus of the point of contact of a tangent drawn to it from a fixed point on the other axis is a parabole.



[Watch Video Solution](#)

9. If normal at  $P$  to a hyperbola of eccentricity  $e$  intersects its transverse and conjugate axes at  $L$  and  $M$ , respectively, then prove that the locus of midpoint of  $LM$  is a hyperbola. Find the eccentricity of this hyperbola



[Watch Video Solution](#)

10. If the normal at a point  $P$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets the  $x$ -axis at  $G$ , show that the  $SG = eSP$ .  $S$  being the focus of the hyperbola.



[Watch Video Solution](#)

11.  $(x - 1)(y - 2) = 5$  and  $(x - 1)^2 + (y + 2)^2 = r^2$  intersect at four points A, B, C, D and if centroid of  $\triangle ABC$  lies on line  $y = 3x - 4$ , then locus of D is

 [Watch Video Solution](#)

### Concept Application Exercise 7 1

1. The equation  $\sqrt{(x - 4)^2 + (y - 2)^2} + \sqrt{(x + 4)^2 + (y - 2)^2} = 8$  represents a

 [Watch Video Solution](#)

2. If  $\cot \theta = \frac{3}{4}$  find the value of :  
$$\frac{\sin \theta - \cos \theta}{\sin \theta + \cos \theta}$$

 [Watch Video Solution](#)

3. The equation of the transvers axis of the hyperbola  $(x - 3)^2 + (y - 1)^2 + (4x + 3y)^2$  is  $x + 3y = 0$  (b)  $4x + 3y = 9$   
 $3x - 4y = 13$  (d)  $4x + 3y = 0$

 [Watch Video Solution](#)

### Concept Application Exercise 7 2

1. Write the length of the latus rectum of the hyperbola  $16x^2 - 9y^2 = 144$ .

 [Watch Video Solution](#)

2. If the latus rectum of a hyperbola forms an equilateral triangle with the vertex at the center of the hyperbola, then find the eccentricity of the hyperbola.

 [Watch Video Solution](#)

3. The distance between two directrices of a rectangular hyperbola is 10 units. Find the distance between its foci.



[Watch Video Solution](#)

4. An ellipse and a hyperbola are confocal (have the same focus) and the conjugate axis of the hyperbola is equal to the minor axis of the ellipse. If  $e_1$  and  $e_2$  are the eccentricities of the ellipse and the hyperbola, respectively, then prove that  $\frac{1}{e_1^2} + \frac{1}{e_2^2} = 2$ .



[Watch Video Solution](#)

5. If  $S$  and  $S'$  are the foci,  $C$  is the center, and  $P$  is a point on the rectangular hyperbola, show that  $SP \times S'P = (CP)^2$



[Watch Video Solution](#)

6. Find the equation of the hyperbola whose foci are  $(8, 3)$  and  $(0, 3)$  and eccentricity  $= \frac{4}{3}$ .

 Watch Video Solution

7. Find all the aspects of hyperbola  $16x^2 - 3y^2 - 32x + 12y - 44 = 0$ .

 Watch Video Solution

8. Show that the locus represented by  $x = \frac{1}{2}a\left(t + \frac{1}{t}\right)$ ,  $y = \frac{1}{2}a\left(t - \frac{1}{t}\right)$  is a rectangular hyperbola.

 Watch Video Solution

9. Two straight lines pass through the fixed points  $(\pm a, 0)$  and have slopes whose products is  $p > 0$ . Show that the locus of the points of intersection of the lines is a hyperbola.



 [Watch Video Solution](#)

10. If  $AOB$  and  $COD$  are two straight lines which bisect one another at right angles, show that the locus of a point  $P$  which moves so that  $PA \cdot PB = PC \cdot PD$  is a hyperbola. Find its eccentricity.

 [Watch Video Solution](#)

11. Find the equation of the chord of the hyperbola  $25x^2 - 16y^2 = 400$  which is bisected at the point  $(5, 3)$ .

 [Watch Video Solution](#)

12.  $PN$  is the ordinate of any point  $P$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\forall'$  is its transverse axis. If  $Q$  divides  $AP$  in the ratio  $a^2 : b^2$ , then prove that  $NQ$  is perpendicular to  $A'P$ .

 [Watch Video Solution](#)

## Concept Application Exercise 7.3

1. The tangents from  $(1, 2\sqrt{2})$  to the hyperbola  $16x^2 - 25y^2 = 400$  include between them an angle equal to:

 [Watch Video Solution](#)

2. Tangents are drawn to the hyperbola  $3x^2 - 2y^2 = 25$  from the point  $(0, \frac{5}{2})$ . Find their equations.

 [Watch Video Solution](#)

3. A common tangent to  $9x^2 - 16y^2 = 144$  and  $x^2 + y^2 = 9$ , is

 [Watch Video Solution](#)

4. The locus a point  $P(\alpha, \beta)$  moving under the condition that the line  $y = \alpha x + \beta$  is a tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is (A) a parabola (B) an ellipse (C) a hyperbola (D) a circle

 [Watch Video Solution](#)

5. A normal to the hyperbola,  $4x^2 - 9y^2 = 36$  meets the co-ordinate axes  $x$  and  $y$  at  $A$  and  $B$ . respectively. If the parallelogram  $OABP$  ( $O$  being the origin) is formed, then the locus of  $P$  is :-

 [Watch Video Solution](#)

6. A point  $P$  moves such that the chord of contact of the pair of tangents from  $P$  on the parabola  $y^2 = 4ax$  touches the rectangular hyperbola  $x^2 - y^2 = c^2$ . Show that the locus of  $P$  is the ellipse  $\frac{x^2}{c^2} + \frac{y^2}{(2a)^2} = 1$ .

 [Watch Video Solution](#)

7. If a tangent to the parabola  $y^2 = 4ax$  intersects the  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  at  $A$  and  $B$ , then the locus of the point of intersection of tangents at  $A$  and  $B$  to the ellipse is

 [Watch Video Solution](#)

8. If the chords of contact of tangents from two points  $(-4, 2)$  and  $(2, 1)$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are at right angle, then find the eccentricity of the hyperbola.

 [Watch Video Solution](#)

9. Statement 1 : If from any point  $P(x_1, y_1)$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$ , tangents are drawn to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then the corresponding chord of contact lies on an other branch of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$  Statement 2 : From any point outside the hyperbola, two tangents can be drawn to the hyperbola.

 [Watch Video Solution](#)

10. Let 'p' be the perpendicular distance from the centre C of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  to the tangent drawn at a point R on the hyperbola. If S & S' are the two foci of the hyperbola, then show that  $(RS + RS')^2 = 4a^2 \left(1 + \frac{b^2}{p^2}\right)$ .

 Watch Video Solution

### Concept Application Exercise 7 4

1. Find the angle between the asymptotes of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ .

 Watch Video Solution

2. Find the asymptotes of the curve  $xy - 3y - 2x = 0$ .

 Watch Video Solution

3. If asymptotes of hyperbola bisect the angles between the transverse axis and conjugate axis of hyperbola, then what is eccentricity of hyperbola?

 Watch Video Solution

4. The asymptote of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  form with an tangent to the hyperbola triangle whose area is  $a^2 \tan \lambda$  in magnitude then its eccentricity is: (a)  $\sec \lambda$  (b)  $\cos ec \lambda$  (c)  $\sec^2 \lambda$  (d)  $\cos ec^2 \lambda$

 Watch Video Solution

5. If the foci of a hyperbola lie on  $y = x$  and one of the asymptotes is  $y = 2x$ , then the equation of the hyperbola, given that it passes through (3, 4), is (a)  $x^2 - y^2 - \frac{5}{2}xy + 5 = 0$  (b)  $2x^2 - 2y^2 + 5xy + 5 = 0$  (c)  $2x^2 + 2y^2 - 5xy + 10 = 0$  (d) none of these

[Watch Video Solution](#)

### Concept Application Exercise 7 5

1. If any line perpendicular to the transverse axis cuts the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and the conjugate hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$  at points  $P$  and  $Q$ , respectively, then prove that normal at  $P$  and  $Q$  meet on the  $x$ -axis.

[Watch Video Solution](#)

2. A normal to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets the axes at  $M$  and  $N$  and lines  $MP$  and  $NP$  are drawn perpendicular to the axes meeting at  $P$ . Prove that the locus of  $P$  is the hyperbola  $a^2x^2 - b^2y^2 = (a^2 + b^2)$ .

[Watch Video Solution](#)

3. Prove that the locus of the point of intersection of the tangents at the ends of the normal chords of the hyperbola  $x^2 - y^2 = a^2$  is  $a^2(y^2 - x^2) = 4x^2y^2$ .



Watch Video Solution

4. Find the value of  $m$ , for which the line  $y = mx + 25\frac{\sqrt{3}}{3}$  is a normal to the conic  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ .



Watch Video Solution

5. Normal is drawn at one of the extremities of the latus rectum of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  which meets the axes at points  $A$  and  $B$ . Then find the area of triangle  $OAB$  ( $O$  being the origin).



Watch Video Solution



1. Find the asymptotes and axes of hyperbola having equation  $xy - 3y - 4x + 7 = 0$ .

 [Watch Video Solution](#)

2. The chord  $PQ$  of the rectangular hyperbola  $xy = a^2$  meets the axis of  $x$  at  $A$ ;  $C$  is the midpoint of  $PQ$ ; and  $O$  is the origin. Then  $ACO$  is equilateral (b) isosceles right-angled (d) right isosceles

 [Watch Video Solution](#)

3. If  $P(x_1, y_1)$ ,  $Q(x_2, y_2)$ ,  $R(x_3, y_3)$  and  $S(x_4, y_4)$  are four concyclic points on the rectangular hyperbola ) and  $xy = c^2$ , then coordinates of the orthocentre of the triangle  $PQR$  is

 [Watch Video Solution](#)

4. If the sum of the slopes of the normal from a point  $P$  to the hyperbola  $xy = c^2$  is equal to  $\lambda (\lambda \in R^+)$ , then the locus of point  $P$  is (a)  $x^2 = \lambda c^2$  (b)  $y^2 = \lambda c^2$  (c)  $xy = \lambda c^2$  (d) none of these



Watch Video Solution

## Exercises

1. If the distance between the foci and the distance between the two directrices of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are in the ratio 3:2, then  $b : a$  is (a)  $1 : \sqrt{2}$  (b)  $\sqrt{3} : \sqrt{2}$  (c)  $1 : 2$  (d)  $2 : 1$

A.  $1 : \sqrt{2}$

B.  $\sqrt{3} : \sqrt{2}$

C.  $1 : 2$

D.  $2 : 1$

**Answer: A**



Watch Video Solution

2. There is a point P on the hyperbola  $\frac{x^2}{16} - \frac{y^2}{6} = 1$  such that its distance from the right directrix is the average of its distance from the two foci. Then the x-coordinate of P is (a)  $-\frac{64}{5}$  (b)  $-\frac{32}{9}$  (c)  $-\frac{64}{9}$  (d) none of these

A.  $-\frac{64}{5}$

B.  $-\frac{32}{9}$

C.  $-\frac{64}{9}$

D. none of these

**Answer: A**



Watch Video Solution

3. The equation,  $2x^2 + 3y^2 - 8x - 18y + 35 = K$  represents (a) no locus if  $k > 0$  (b) an ellipse if  $k < 0$  (c) a point if  $k = 0$  (d) a hyperbola if  $k > 0$

A. no locus if  $k > 0$

B. an ellipse if  $k < 0$

C. a point if  $k = 0$

D. a hyperbola if  $k > 0$

**Answer: C**

 [Watch Video Solution](#)

4. Let 'a' and 'b' be non-zero real numbers. Then, the equation

$(ax^2 + by^2 + c)(x^2 - 5xy + 6y^2)$  represents :

A. four straight lines, when  $c = 0$  and a, b are of the same sign

B. two straight lines and a circle, when  $a = b$  and c is of sign opposite to that of a

C. two straight lines and a hyperbola, when a and b are of the same sign and c is of sign opposite to that of a

D. a circle and an ellipse, when a and b are of the same sign and c is of sign opposite to that of a

**Answer: B**



[Watch Video Solution](#)

5. For the hyperbola  $\frac{x^2}{\cos^2 \alpha} - \frac{y^2}{\sin^2 \alpha} = 1; \left(0 < \alpha < \frac{\pi}{4}\right)$ . Which of the following remains constant when alpha varies?

- A. Eccentricity
- B. Abscissa of foci
- C. Directrix
- D. Vertex

**Answer: B**



[Watch Video Solution](#)

6. Which of the following pairs may represent the eccentricities of two conjugate hyperbolas, for  $\alpha \in (0, \pi/2)$ ?

a.  $\sin \theta, \cos \theta$  b.  $\tan \theta, \cot \theta$  c.  $\sec \theta, \operatorname{cosec} \theta$  d.  $1 + \sin \theta, 1 + \cos \theta$

A.  $\sin \theta, \cos \theta$

B.  $\tan \theta, \cot \theta$

C.  $\sec \theta, \operatorname{cosec} \theta$

D.  $1 + \sin \theta, 1 + \cos \theta$

**Answer: C**



**Watch Video Solution**

7. If a variable line has its intercepts on the coordinate axes  $e$  and  $e'$ , where  $\frac{e}{2}$  and  $\frac{e'}{2}$  are the eccentricities of a hyperbola and its conjugate hyperbola, then the line always touches the circle  $x^2 + y^2 = r^2$ , where  $r =$  (a) 1 (b) 2 (c) 3 (d) cannot be decided

A. 1

B. 2

C. 3

D. cannot be decided

**Answer: C**



**Watch Video Solution**

8. A hyperbola, having the transverse axis of length  $2\sin\theta$ , is confocal with the ellipse  $3x^2 + 4y^2 = 12$ . Then its equation is

A. (a)  $x^2 \operatorname{cosec}^2 \theta - y^2 \sec^2 \theta = 1$

B. (b)  $x^2 \sec^2 \theta - y^2 \operatorname{cosec}^2 \theta = 1$

C. (c)  $x^2 \sin^2 \theta - y^2 \cos^2 \theta = 1$

D. (d)  $x^2 \cos^2 \theta - y^2 \cos^2 \theta = 1$

**Answer: A**



**Watch Video Solution**

9. If the distances of one focus of hyperbola from its directrices are 5 and 3, then its eccentricity is

A.  $\sqrt{2}$

B. 2

C. 4

D. 8

**Answer: B**



**Watch Video Solution**

10. Let  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{A^2} - \frac{y^2}{B^2} = 1$  be confocal ( $a > A$  and  $a > b$ ) having the foci at  $s_1$  and  $S_2$ , respectively. If P is their point of intersection, then  $S_1P$  and  $S_2P$  are the roots of quadratic equation



A. (a)  $x^2 + 2Ax + (a^2 - A^2) = 0$

B. (b)  $x^2 + 2ax + (a^2 - A^2) = 0$

C. (c)  $x^2 - 2Ax + (a^2 + A^2) = 0$

D. (d)  $x^2 - 2ax + (a^2 - A^2) = 0$

**Answer: D**

 [Watch Video Solution](#)

11. Two tangents are drawn from a point on hyperbola  $x^2 - y^2 = 5$  to the ellipse  $\frac{x^2}{9} + \frac{y^2}{4} = 1$ . If they make angle  $\alpha$  and  $\beta$  with x-axis, then

A.  $\alpha - \beta = \pm \frac{\pi}{2}$

B.  $\alpha + \beta = \frac{\pi}{2}$

C.  $\alpha + \beta = \pi$

D.  $\alpha + \beta = 0$

**Answer: B**



Watch Video Solution

12. Equation of the rectangular hyperbola whose focus is  $(1, -1)$  and the corresponding directrix is  $x - y + 1 = 0$ .

a.  $x^2 - y^2 = 1$       b.  $xy = 1$       c.  $2xy - 4x + 4y + 1 = 0$       d.

$2xy + 4x - 4y - 1 = 0$

A.  $x^2 - y^2 = 1$

B.  $xy = 1$

C.  $2xy - 4x + 4y + 1 = 0$

D.  $2xy + 4x - 4y - 1 = 0$

Answer: C



Watch Video Solution

13. If two circles  $(x + 4)^2 + y^2 = 1$  and  $(x - 4)^2 + y^2 = 9$  are touched externally by a circle, then locus of centre of variable circle is

A. (a)  $\frac{x^2}{15} - \frac{y^2}{1} = 1$

B. (b)  $\frac{x^2}{4} - \frac{y^2}{12} = 1$

C. (c)  $\frac{x^2}{1} - \frac{y^2}{15} = 1$

D. (d)  $\frac{x^2}{12} - \frac{y^2}{4} = 1$

**Answer: C**



**Watch Video Solution**

14. If the vertex of a hyperbola bisects the distance between its center and the corresponding focus, then the ratio of the square of its conjugate axis to the square of its transverse axis is (a) 2 (b) 4 (c) 6 (d) 3

A. 2

B. 4

C. 6

D. 3

**Answer: C**



**Watch Video Solution**

15. The eccentricity of the hyperbola whose length of the latus rectum is equal to 8 and the length of its conjugate axis is equal to half of the distance between its foci, is : (1)  $\frac{4}{3}$  (2)  $\frac{4}{\sqrt{3}}$  (3)  $\frac{2}{\sqrt{3}}$  (4)  $\sqrt{3}$

A.  $3/4$

B.  $4/\sqrt{3}$

C.  $2/\sqrt{3}$

D. none of these

**Answer: C**



**Watch Video Solution**

16. Let  $LL'$  be the latus rectum through the focus of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $A'$  be the farther vertex. If  $A'LL'$  is equilateral, then the eccentricity of the hyperbola is (axes are coordinate axes). (a)  $\sqrt{3}$  (b)

$\sqrt{3} + 1$  (c)  $\left(\frac{\sqrt{3} + 1}{\sqrt{2}}\right)$  (d)  $\frac{(\sqrt{3} + 1)}{\sqrt{3}}$

A.  $\sqrt{3}$

B.  $\sqrt{3} + 1$

C.  $(\sqrt{3} + 1) / \sqrt{2}$

D.  $(\sqrt{3} + 1) / \sqrt{3}$

**Answer: D**



**Watch Video Solution**

17. The eccentricity of the conjugate hyperbola of the hyperbola  $x^2 - 3y^2 = 1$  is (a) 2 (b)  $2\sqrt{3}$  (c) 4 (d)  $\frac{4}{5}$

A. 2

B.  $2/\sqrt{3}$

C. 4

D.  $4/5$

**Answer: A**



**Watch Video Solution**

18. The locus of the point of intersection of the lines  $\sqrt{3}x - y - 4\sqrt{3}t = 0$  &  $\sqrt{3}tx + ty - 4\sqrt{3} = 0$  (where  $t$  is a parameter) is a hyperbola whose eccentricity is: (a)  $\sqrt{3}$  (b) 2 (c)  $\frac{2}{\sqrt{3}}$  (d)  $\frac{4}{3}$

A.  $\sqrt{3}$

B. 2

C.  $2/\sqrt{3}$

D.  $4/3$

**Answer: B**

 [Watch Video Solution](#)

19. If the eccentricity of the hyperbola  $x^2 - y^2 \sec^2 \alpha = 5$  is  $\sqrt{3}$  times the eccentricity of the ellipse  $x^2 (\sec^2 \alpha) + y^2 = 25$ , then a value of  $\alpha$  is : (a)

$\frac{\pi}{6}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{2}$

A.  $\pi / 6$

B.  $\pi / 4$

C.  $\pi / 3$

D.  $\pi / 2$

**Answer: B**

 [Watch Video Solution](#)

20. The equation of the transvers and conjugate axes of a hyperbola are, respectively,  $x + 2y - 3 = 0$  and  $2x - y + 4 = 0$ , and their respective lengths are  $\sqrt{2}$  and  $\frac{2}{\sqrt{3}}$ . The equation of the hyperbola is

$$\text{a) } \frac{2}{5}(x + 2y - 3)^2 - \frac{3}{5}(2x - y + 4)^2 = 1$$

$$\text{b) } \frac{2}{5}(x - y - 4)^2 - \frac{3}{5}(x + 2y - 3)^2 = 1$$

$$\text{c) } \frac{2}{5}(2x - y + 4)^2 - \frac{3}{5}(x + 2y - 3)^2 = 1$$

$$\text{d) } 2(x + 2y - 3)^2 - 3(2x - y + 4)^2 = 1$$

$$\text{A. } \frac{2}{5}(x + 2y - 3)^2 - \frac{3}{5}(2x - y + 4)^2 = 1$$

$$\text{B. } \frac{2}{5}(2x - y + 4)^2 - \frac{3}{5}(x + 2y - 3)^2 = 1$$

$$\text{C. } 2(2x - y + 4)^2 - 3(x + 2y - 3)^2 = 1$$

$$\text{D. } 2(x + 2y - 3)^2 - 3(2x - y + 4)^2 = 1$$

**Answer: B**



**Watch Video Solution**

**21.** about to only mathematics

$$\text{A. } 1 - \sqrt{2/3}$$

$$\text{B. } \sqrt{3/2} - 1$$

$$\text{C. } 1 + \sqrt{2/3}$$



D.  $\sqrt{3/2} + 1$

**Answer: B**



**Watch Video Solution**

22. If two points  $P$  &  $Q$  on the hyperbola,  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  whose centre is  $C$  be such that  $CP$  is perpendicular to  $CQ$  and  $a < b$ , then prove that

$$\frac{1}{CP^2} + \frac{1}{CQ^2} = \frac{1}{a^2} - \frac{1}{b^2}.$$

A.  $\frac{b^2 - a^2}{2ab}$

B.  $\frac{1}{a^2} + \frac{1}{b^2}$

C.  $\frac{2ab}{b^2 - a^2}$

D.  $\frac{1}{a^2} - \frac{1}{b^2}$

**Answer: D**



**Watch Video Solution**

23. The angle between the lines joining the origin to the points of intersection of the line  $\sqrt{3}x + y = 2$  and the curve  $y^2 - x^2 = 4$  is

A.  $\tan^{-1}(2/\sqrt{3})$

B.  $\pi/6$

C.  $\tan^{-1}(\sqrt{3}/2)$

D.  $\pi/2$

**Answer: C**



**Watch Video Solution**

24. A variable chord of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, (b > a)$ , subtends a right angle at the center of the hyperbola if this chord touches a fixed circle concentric with the hyperbola a fixed ellipse concentric with the hyperbola a fixed hyperbola concentric with the hyperbola a fixed parabola having vertex at  $(0, 0)$ .

A. a fixed circle concentric with the hyperbola

B. a fixed ellipse concentric with the hyperbola

C. a fixed hyperbola concentric with the hyperbola

D. a fixed parabola having vertex at (0, 0)

**Answer: A**



[Watch Video Solution](#)

25. If the distance between two parallel tangents having slope  $m$  drawn to the hyperbola  $\frac{x^2}{9} - \frac{y^2}{49} = 1$  is 2, then the value of  $2|m|$  is \_\_\_\_\_

A.  $\pm 5/2$

B.  $\pm 4/5$

C.  $\pm 7/2$

D. none of these

**Answer: A**



[Watch Video Solution](#)

26. If  $ax + by = 1$  is tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $a^2 - b^2$  is equal to (a)  $\frac{1}{a^2e^2}$  (b)  $a^2e^2$  (c)  $b^2e^2$  (d) none of these

A.  $1/a^2e^2$

B.  $a^2e^2$

C.  $b^2e^2$  none of these

D. none of these

**Answer: A**



**Watch Video Solution**

27. A tangent drawn to hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at  $P\left(\frac{\pi}{6}\right)$  forms a triangle of area  $3a^2$  square units, with the coordinate axes, then the square of its eccentricity is (A) 15 (B) 24 (C) 17 (D) 14

A. 15

B. 24

C. 17

D. 14

**Answer: C**



**Watch Video Solution**

**28.** about to only mathematics

A. 2

B. 4

C. zero

D. none of these

**Answer: C**



**Watch Video Solution**

29. The locus of a point whose chord of contact with respect to the circle  $x^2 + y^2 = 4$  is a tangent to the hyperbola  $xy = 1$  is a/an ellipse (b) circle hyperbola (d) parabola

- A. ellipse
- B. circle
- C. hyperbola
- D. parabola

**Answer: C**



[Watch Video Solution](#)

30. The sides  $AC$  and  $AB$  of a  $ABC$  touch the conjugate hyperbola of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . If the vertex  $A$  lies on the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , then the side  $BC$  must touch parabola (b) circle hyperbola (d) ellipse

A. parabola

B. circle

C. hyperbola

D. ellipse

**Answer: D**



[Watch Video Solution](#)

**31.** The number of possible tangents which can be drawn to the curve  $4x^2 - 9y^2 = 36$ , which are perpendicular to the straight line  $5x + 2y - 10 = 0$ , is zero (b) 1 (c) 2 (d) 4

A. a. zero

B. b. 1

C. c. 2

D. d. 4

**Answer: A**



**Watch Video Solution**

32. The tangent at a point  $P$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  passes through the point  $(0, -b)$  and the normal at  $P$  passes through the point  $(2a\sqrt{2}, 0)$ . Then the eccentricity of the hyperbola is

A. a. 2

B. b.  $\sqrt{2}$

C. c. 3

D. d.  $\sqrt{3}$

**Answer: B**



**Watch Video Solution**



33. Locus of the feet of the perpendiculars drawn from either foci on a variable tangent to the hyperbola  $16y^2 - 9x^2 = 1$  is

A.  $x^2 + y^2 = 9$

B.  $x^2 + y^2 = 1/9$

C.  $x^2 + y^2 = 7/144$

D.  $x^2 + y^2 = 1/16$

**Answer: D**



[Watch Video Solution](#)

34. P is a point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , and N is the foot of the perpendicular from P on the transverse axis. The tangent to the hyperbola at P meets the transverse axis at T. If O is the centre of the hyperbola, then  $OT \cdot ON$  is equal to

A.  $e^2$

B.  $a^2$

C.  $b^2$

D.  $b^2/a^2$

**Answer: B**



**Watch Video Solution**

35. The coordinates of a point on the hyperbola  $\frac{x^2}{24} - \frac{y^2}{18} = 1$  which is nearest to the line  $3x + 2y + 1 = 0$  are (a) (6, 3) (b) ( - 6, - 3) (c) 6, - 3) (d) ( - 6, 3)

A. (6, 3)

B. ( - 6, - 3)

C. ( - 6, 3)

D. (6, - 3)

**Answer: C**

 [Watch Video Solution](#)

36. The tangent at a point  $P$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets one of the directrix at  $F$ . If  $PF$  subtends an angle  $\theta$  at the corresponding focus, then  $\theta = \frac{\pi}{4}$  (b)  $\frac{\pi}{2}$  (c)  $\frac{3\pi}{4}$  (d)  $\pi$

A.  $\pi/4$

B.  $\pi/2$

C.  $3\pi/4$

D.  $\pi$

**Answer: B**

 [Watch Video Solution](#)

37. The locus of a point, from where the tangents to the rectangular hyperbola  $x^2 - y^2 = a^2$  contain an angle of  $45^\circ$ , is (a)  $(x^2 + y^2)^2 + a^2(x^2 - y^2) = 4a^2$  (b)

$$2(x^2 + y^2)^2 + 4a^2(x^2 - y^2) = 4a^2 \quad (c)$$

$$(x^2 + y^2)^2 + 4a^2(x^2 - y^2) = 4a^2 \quad (d) \quad (x^2 + y^2) + a^2(x^2 - y^2) = a^4$$

A.  $(x^2 + y^2)^2 + a^2(x^2 - y^2) = 4a^2$

B.  $2(x^2 + y^2)^2 + 4a^2(x^2 - y^2) = 4a^2$

C.  $(x^2 + y^2)^2 + 4a^2(x^2 - y^2) = 4a^4$

D.  $(x^2 + y^2)^2 + a^2(x^2 - y^2) = a^4$

**Answer: C**



**Watch Video Solution**

**38.** If tangents  $PQ$  and  $PR$  are drawn from a variable point  $P$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , ( $a > b$ ), so that the fourth vertex  $S$  of parallelogram  $PQSR$  lies on the circumcircle of triangle  $PQR$ , then the locus of  $P$  is (a)  $x^2 + y^2 = b^2$  (b)  $x^2 + y^2 = a^2$  (c)  $x^2 + y^2 = a^2 - b^2$  (d) none of these

A.  $x^2 + y^2 = b^2$

B.  $x^2 + y^2 = a^2$

C.  $x^2 + y^2 = a^2 - b^2$

D. none of these

**Answer: C**



**Watch Video Solution**

39. The number of points on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 3$  from which mutually perpendicular tangents can be drawn to the circle  $x^2 + y^2 = a^2$  is/are (a) 0 (b) 2 (c) 3 (d) 4

A. 0

B. 2

C. 3

D. 4

**Answer: A**

 Watch Video Solution

40. If a ray of light incident along the line  $3x + (5 - 4\sqrt{2})y = 15$  gets reflected from the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ , then its reflected ray goes along the line.  $x\sqrt{2} - y + 5 = 0$  (b)  $\sqrt{2}y - x + 5 = 0$   $\sqrt{2}y - x - 5 = 0$   
(d) none of these

A.  $x\sqrt{2} - y + 5 = 0$

B.  $\sqrt{2}y - x + 5 = 0$

C.  $\sqrt{2}y - x - 5 = 0$

D. none of these

**Answer: D**

 Watch Video Solution

41. The chord of contact of a point  $P$  w.r.t a hyperbola and its auxiliary circle are at right angle. Then the point  $P$  lies (a) on conjugate hyperbola

(b) one of the directrix (c) one of the asymptotes (d) none of these

A. conjugate hyperbola

B. one of the directrix

C. asymptotes

D. none of these

**Answer: C**



**Watch Video Solution**

**42.** The ellipse  $4x^2 + 9y^2 = 36$  and the hyperbola  $a^2x^2 - y^2 = 4$  intersect at right angles. Then the equation of the circle through the points of intersection of two conics is (a)  $x^2 + y^2 = 5$  (b)  $\sqrt{5}(x^2 + y^2) - 3x - 4y = 0$  (c)  $\sqrt{5}(x^2 + y^2) + 3x + 4y = 0$  (d)  $x^2 + y^2 = 25$

A.  $x^2 + y^2 = 5$

B.  $\sqrt{5}(x^2 + y^2) - 3x - 4y = 0$

C.  $\sqrt{5}(x^2 + y^2) + 3x + 4y = 0$

D.  $x^2 + y^2 = 25$

**Answer: A**



**Watch Video Solution**

**43.** The locus of the point which is such that the chord of contact of tangents drawn from it to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  forms a triangle of constant area with the coordinate axes is (a) straight line (b) a hyperbola (c) an ellipse (d) a circle

A. a straight line

B. a hyperbola

C. an ellipse

D. a circle

**Answer: B**





[Watch Video Solution](#)

44. If  $x = 9$  is the chord of contact of the hyperbola  $x^2 - y^2 = 9$  then the equation of the corresponding pair of tangents is (A)  $9x^2 - 8y^2 + 18x - 9 = 0$  (B)  $9x^2 - 8y^2 - 18x + 9 = 0$  (C)  $9x^2 - 8y^2 - 18x - 9 = 0$  (D)  $9x^2 - 8y^2 + 18x + 9 = 0$

A.  $9x^2 - 8y^2 + 18x - 9 = 0$

B.  $9x^2 - 8y^2 - 18x = 0$

C.  $9x^2 - 8y^2 - 9 = 0$

D.  $9x^2 - 8y^2 + 18x + 9 = 0$

**Answer: B**

[Watch Video Solution](#)

45. If the tangent at point  $P(h, k)$  on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  cuts the circle  $x^2 + y^2 = a^2$  at points  $Q(x_1, y_1)$  and  $R(x_2, y_2)$ , then the value

of  $\frac{1}{y_1} + \frac{1}{y_2}$  is

A.  $\frac{1}{k}$

B.  $\frac{2}{k}$

C.  $\frac{ab}{k}$

D.  $\frac{a+b}{k}$

**Answer: B**



**Watch Video Solution**

**46.** Let  $P(a \sec \theta, b \tan \theta)$  and  $Q(a \sec \phi, b \tan \phi)$  (where  $\theta + \phi = \frac{\pi}{2}$ ) be two points on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  If  $(h, k)$  is the point of intersection of the normals at  $P$  and  $Q$  then  $k$  is equal to

(A)  $\frac{a^2 + b^2}{a}$  (B)  $-\left(\frac{a^2 + b^2}{a}\right)$  (C)  $\frac{a^2 + b^2}{b}$  (D)  $-\left(\frac{a^2 + b^2}{b}\right)$

A.  $\frac{a^2 + b^2}{a}$

B.  $-\left(\frac{a^2 + b^2}{a}\right)$

C.  $\frac{a^2 + b^2}{b}$

D.  $-\left(\frac{a^2 + b^2}{b}\right)$

**Answer: D**



**Watch Video Solution**

47. A normal to the hyperbola  $\frac{x^2}{4} - \frac{y^2}{1} = 1$  has equal intercepts on the positive x- and y-axis. If this normal touches the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , then  $a^2 + b^2$  is equal to 5 (b) 25 (c) 16 (d) none of these

A. 5

B. 25

C. 16

D. none of these

**Answer: D**



**Watch Video Solution**

48. Portion of asymptote of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  (between centre and the tangent at vertex) in the first quadrant is cut by the line  $y + \lambda(x - a) = 0$  ( $\lambda$  is a parameter) then (A)  $\lambda \in R$  (B)  $\lambda \in (0, \infty)$  (C)  $\lambda \in (-\infty, 0)$  (D)  $\lambda \in R - \{0\}$

A.  $\lambda \in R$

B.  $\lambda \in (0, \infty)$

C.  $\lambda \in (-\infty, 0)$

D. none of these

**Answer: B**



**Watch Video Solution**

49. If the angle between the asymptotes of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is  $120^\circ$  and the product of perpendiculars drawn from the foci upon its any tangent is 9, then the locus of the point of intersection of perpendicular

tangents of the hyperbola can be  $x^2 + y^2 = 6$  (b)  $x^2 + y^2 = 9$

$x^2 + y^2 = 3$  (d)  $x^2 + y^2 = 18$

A.  $x^2 + y^2 = 6$

B.  $x^2 + y^2 = 9$

C.  $x^2 + y^2 = 3$

D.  $x^2 + y^2 = 18$

**Answer: D**



**Watch Video Solution**

50. Let any double ordinate  $PNP'$  of the hyperbola  $\frac{x^2}{25} - \frac{y^2}{16} = 1$  be produced on both sides to meet the asymptotes in  $Q$  and  $Q'$ . Then  $PQ \dot{P}' Q'$  is equal to 25 (b) 16 (c) 41 (d) none of these

A. 25

B. 16

C. 41

D. none of these

**Answer: B**



**Watch Video Solution**

51. For hyperbola whose center is at  $(1, 2)$  and the asymptotes are parallel to lines  $2x + 3y = 0$  and  $x + 2y = 1$ , the equation of the hyperbola passing through  $(2, 4)$  is (a)  $(2x + 3y - 5)(x + 2y - 8) = 40$  (b)  $(2x + 3y - 8)(x + 2y - 8) = 40$  (c)  $(2x + 3y - 8)(x + 2y - 5) = 30$  (d) none of these

A.  $(2x + 3y - 5)(x + 2y - 8) = 40$

B.  $(2x + 3y - 8)(x + 2y - 5) = 40$

C.  $(2x + 3y - 8)(x + 2y - 8) = 30$

D. none of these

**Answer: B**



**Watch Video Solution**

52. The asymptotes of the hyperbola  $\frac{x^2}{a_1^2} - \frac{y^2}{b_1^2} = 1$  and  $\frac{x^2}{a_2^2} - \frac{y^2}{b_2^2} = 1$  are perpendicular to each other. Then, (a)  $\frac{a_1}{a_2} = \frac{b_1}{b_2}$  (b)  $a_1a_2 = b_1b_2$  (c)  $a_1a_2 + b_1b_2 = 0$  (d)  $a_1 - a_2 = b_1 - b_2$

A.  $a_1/a_2 = b_1/b_2$

B.  $a_1a_2 = b_1b_2$

C.  $a_1a_2 + b_1b_2 = 0$

D.  $a_1 - a_2 = b_1 - b_2$

**Answer: C**



**Watch Video Solution**

53. If  $S = 0$  is the equation of the hyperbola  $x^2 + 4xy + 3y^2 - 4x + 2y + 1 = 0$ , then the value of  $k$  for which  $S + K = 0$  represents its asymptotes is (a) 20 (b) -16 (c) -22 (d) 18

A. 20

B. -16

C. -22

D. 18

**Answer: C**



**Watch Video Solution**

54. If two distinct tangents can be drawn from the Point  $(\alpha, 2)$  on different branches of the hyperbola  $\frac{x^2}{9} - \frac{y^2}{16} = 1$  then (1)  $|\alpha| < \frac{3}{2}$  (2)  $|\alpha| > \frac{2}{3}$  (3)  $|\alpha| > 3$  (4)  $\alpha = 1$

A.  $|\alpha| < 3/2$

B.  $|\alpha| > 2/3$

C.  $|\alpha| > 3$

D. none of these



**Answer: A**



**Watch Video Solution**

55. A hyperbola passes through (2,3) and has asymptotes  $3x - 4y + 5 = 0$  and  $12x + 5y - 40 = 0$ . Then, the equation of its transverse axis is  $77x - 21y - 265 = 0$   $21x - 77y + 265 = 0$   $21x - 77y - 265 = 0$   $21x + 77y - 265 = 0$

A.  $77x - 21y - 265 = 0$

B.  $21x - 77y + 265 = 0$

C.  $21x - 77y - 265 = 0$

D.  $21x + 77y - 265 = 0$

**Answer: B**



**Watch Video Solution**

56. From any point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , tangents are drawn to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$ . The area cut-off by the chord of contact on the asymptotes is equal to  $\frac{a}{2}$  (b)  $ab$  (c)  $2ab$  (d)  $4ab$

A.  $a/2$

B.  $ab$

C.  $2ab$

D.  $4ab$

**Answer: D**



**Watch Video Solution**

57. The combined equation of the asymptotes of the hyperbola  $2x^2 + 5xy + 2y^2 + 4x + 5y = 0$  is -

A.  $2x^2 + 5xy + 2y^2 + 4x + 5y + 2 = 0$

B.  $2x^2 + 5xy + 2y^2 + 4x + 5y - 2 = 0$

C.  $2x^2 + 5xy + 2y^2 = 0$

D. none of these

**Answer: A**



[Watch Video Solution](#)

58. The asymptotes of the hyperbola  $xy = hx + ky$  are (1)  $x - k = 0$  and  $y - h = 0$  (2)  $x + h = 0$  and  $y + k = 0$  (3)  $x - k = 0$  and  $y + h = 0$  (4)  $x + k = 0$  and  $y - h = 0$

A.  $x - k = 0$  and  $y - h = 0$

B.  $x + h = 0$  and  $y + k = 0$

C.  $x - k = 0$  and  $y + h = 0$

D.  $x + k = 0$  and  $y - h = 0$

**Answer: A**



[Watch Video Solution](#)

59. The center of a rectangular hyperbola lies on the line  $y = 2x$ . If one of the asymptotes is  $x + y + c = 0$ , then the other asymptote is  $6x + 3y - 4c = 0$  (b)  $3x + 6y - 5c = 0$   $3x - 6y - c = 0$  (d) none of these

A.  $6x + 3y - 4c = 0$

B.  $3x + 6y - 5c = 0$

C.  $3x - 6y - c = 0$

D. none of these

**Answer: D**



[Watch Video Solution](#)

60. The equation of a rectangular hyperbola whose asymptotes are  $x = 3$  and  $y = 5$  and passing through (7,8) is (a)  $xy - 3y + 5x + 3 = 0$  (b)

$$xy + 3y + 4x + 3 = 0$$

(c)

$$xy - 3y + 5x - 3 = 0$$

(d)

$$xy - 3y + 5x + 3 = 0$$

A.  $xy - 3y + 5x + 3 = 0$

B.  $xy + 3y + 4x + 3 = 0$

C.  $xy - 3y + 5x - 3 = 0$

D.  $xy - 3y - 5x + 3 = 0$

**Answer: D**



**Watch Video Solution**

61. If tangents  $OQ$  and  $OR$  are drawn to variable circles having radius  $r$  and the center lying on the rectangular hyperbola  $xy = 1$ , then the locus of the circumcenter of triangle  $OQR$  is ( $O$  being the origin)

(a).  $xy = 4$  (b)  $xy = \frac{1}{4}$  (c)  $xy = 1$  (d) none of these

A.  $xy = 4$

B.  $xy = 1/4$

C.  $xy = 1$

D. none of these

**Answer: B**



[Watch Video Solution](#)

62. Four points are such that the line joining any two points is perpendicular to the line joining other two points. If three point out of these lie on a rectangular hyperbola, then the fourth point will lie on

A. the same hyperbola

B. the conjugate hyperbola

C. one of the directrix

D. one of the asymptotes

**Answer: A**



[Watch Video Solution](#)

63. If  $S_1$  and  $S_2$  are the foci of the hyperbola whose length of the transverse axis is 4 and that of the conjugate axis is 6, and  $S_3$  and  $S_4$  are the foci of the conjugate hyperbola, then the area of quadrilateral  $S_1S_3S_2S_4$  is 24 (b) 26 (c) 22 (d) none of these

A. 24

B. 26

C. 22

D. none of these

**Answer: B**



[Watch Video Solution](#)

64. Suppose the circle having equation  $x^2 + y^2 = 3$  intersects the rectangular hyperbola  $xy = 1$  at points  $A, B, C,$  and  $D$ . The equation  $x^2 + y^2 - 3 + \lambda(xy - 1) = 0, \lambda \in R,$  represents. a pair of lines

through the origin for  $\lambda = -3$  an ellipse through  $A, B, C,$  and  $D$  for  $\lambda = -3$  a parabola through  $A, B, C,$  and  $D$  for  $\lambda = -3$  a circle for any  $\lambda \in R$

A. a pair of lines through the origin for  $\lambda = -3$

B. an ellipse through  $A, B, C$  and  $D$  for  $\lambda = -3$

C. a parabola through  $A, B, C$  and  $D$  for  $\lambda = -3$

D. a circle for any  $\lambda \in R$

**Answer: A**



**Watch Video Solution**

**65.** The equation to the chord joining two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on

the rectangular hyperbola  $xy = c^2$  is:  $\frac{x}{x_1 + x_2} + \frac{y}{y_1 + y_2} = 1$

$$\frac{x}{x_1 - x_2} + \frac{y}{y_1 - y_2} = 1$$

$$\frac{x}{y_1 + y_2} + \frac{y}{x_1 + x_2} = 1 \quad \text{(d)}$$

$$\frac{x}{y_1 - y_2} + \frac{y}{x_1 - x_2} = 1$$

A.  $\frac{x}{x_1 + x_2} + \frac{y}{y_1 + y_2} = 1$



$$\text{B. } \frac{x}{x_1 - x_2} + \frac{y}{y_1 - y_2} = 1$$

$$\text{C. } \frac{x}{y_1 + y_2} + \frac{y}{x_1 + x_2} = 1$$

$$\text{D. } \frac{x}{y_1 - y_2} + \frac{y}{x_1 - x_2} = 1$$

**Answer: A**



**Watch Video Solution**

**66.** The locus of the foot of the perpendicular from the center of the hyperbola  $xy = 1$  on a variable tangent is (a)  $(x^2 + y^2)^2 = 4xy$  (b)  $(x^2 - y^2) = \frac{1}{9}$  (c)  $(x^2 - y^2) = \frac{7}{144}$  (d)  $(x^2 - y^2) = \frac{1}{16}$

$$\text{A. } (x^2 - y^2)^2 = 4xy$$

$$\text{B. } (x^2 + y^2)^2 = 2xy$$

$$\text{C. } (x^2 + y^2) = 4xy$$

$$\text{D. } (x^2 + y^2)^2 = 4xy$$

**Answer: D**

[Watch Video Solution](#)

67. The curve  $xy = c (c > 0)$  and the circle  $x^2 + y^2 = 1$  touch at two points, then distance between the points of contact is

A. (a) 1

B. (b) 2

C. (c)  $2\sqrt{2}$

D. (d) none of these

**Answer: B**

[Watch Video Solution](#)

68. Let  $C$  be a curve which is the locus of the point of intersection of lines  $x = 2 + m$  and  $my = 4 - m$ . A circle  $s \equiv (x - 2)^2 + (y + 1)^2 = 25$  intersects the curve  $C$  at four points:  $P, Q, R,$  and  $S$ . If  $O$  is center of

the curve  $C$ , then  $OP^2 + OP^2 + OR^2 + OS^2$  is (a) 50 (b) 100 (c) 25 (d)

$$\frac{25}{2}$$

A. 50

B. 100

C. 25

D. 25/5

**Answer: B**



**Watch Video Solution**

### Multiple Correct Answers Type

1. If the circle  $x^2 + y^2 = a^2$  intersects the hyperbola  $xy = c^2$  at four points  $P(x_1, y_1)$ ,  $Q(x_2, y_2)$ ,  $R(x_3, y_3)$ , and  $S(x_4, y_4)$ , then

$$x_1 + x_2 + x_3 + x_4 = 0 \quad y_1 + y_2 + y_3 + y_4 = 0 \quad x_1x_2x_3x_4 = C^4$$

$$y_1y_2y_3y_4 = C^4$$

A.  $x_1 + x_2 + x_3 + x_4 = 0$

B.  $y_1 + y_2 + y_3 + y_4 = 0$

C.  $x_1x_2x_3x_4 = c^4$

D.  $y_1y_2y_3y_4 = c^4$

**Answer: A::B::C::D**



**Watch Video Solution**

2. The equation  $(x - \alpha)^2 + (y - \beta)^2 = k(lx + my + n)^2$  represents (a) a parabola for  $k = (l^2 + m^2)^{-1}$  (b) an ellipse for  $0 < k < (l^2 + m^2)^{-1}$  (c) a hyperbola for  $k > (l^2 + m^2)^{-1}$  (d) a point circle for  $k = 0$

A. a parabola for  $k < (l^2 + m^2)^{-1}$

B. an ellipse for  $0 < k < (l^2 + m^2)^{-1}$

C. a hyperbola for  $k > (l^2 + m^2)^{-1}$

D. a point circle for  $k = 0$

Answer: B::C::D

 [Watch Video Solution](#)

3. If  $(5, 12)$  and  $(24, 7)$  are the foci of a hyperbola passing through the origin, then  $e = \frac{\sqrt{386}}{12}$  (b)  $e = \frac{\sqrt{386}}{13}$   $LR = \frac{121}{6}$  (d)  $LR = \frac{121}{3}$

A.  $e = \frac{\sqrt{386}}{12}$

B.  $e = \frac{\sqrt{386}}{13}$

C.  $LR = 121/6$

D.  $LR = 121/3$

Answer: A::C

 [Watch Video Solution](#)

4. Show that the equation  $9x^2 - 16y^2 - 18x + 32y - 151 = 0$  represents a hyperbola.

A. one of the directrix is  $x = 21/5$

B. the length of latus rectum is  $9/2$

C. foci are  $(6, 1)$  and  $(-4, 1)$

D. the eccentricity is  $5/4$

**Answer: A::B::C::D**

 [Watch Video Solution](#)

5. If a hyperbola passes through the foci of the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$ . Its transverse and conjugate axes coincide respectively with the major and minor axes of the ellipse and if the product of eccentricities of hyperbola and ellipse is 1 then the equation of hyperbola is  $\frac{x^2}{9} - \frac{y^2}{16} = 1$  b. the equation of hyperbola is  $\frac{x^2}{9} - \frac{y^2}{25} = 1$  c. focus of hyperbola is  $(5, 0)$  d. focus of hyperbola is  $(5\sqrt{3}, 0)$

A. the equation of hyperbola is  $\frac{x^2}{9} - \frac{y^2}{16} = 1$

B. the equation of the hyperbola is  $\frac{x^2}{9} - \frac{y^2}{25} = 1$

C. the vertex of the hyperbola is (5, 0)

D. the vertex of the hyperbola is  $(5\sqrt{3}, 0)$

**Answer: A::C**



**Watch Video Solution**

6. If the foci of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  coincide with the foci of  $\frac{x^2}{25} + \frac{y^2}{9} = 1$  and the eccentricity of the hyperbola is 2, then  $a^2 + b^2 = 16$  there is no director circle to the hyperbola the center of the director circle is (0, 0). the length of latus rectum of the hyperbola is 12

A.  $a^2 + b^2 = 16$

B. there is no director circle to the hyperbola

C. the centre of the director circle is (0, 0)

D. the length of latus rectum of the hyperbola is 12

**Answer: A::B::D**



[Watch Video Solution](#)

7. The differential equation  $\frac{dy}{dx} = \frac{2x}{3y}$  represents a family of hyperbolas (except when it represents a pair of lines) with eccentricity. (a)  $\sqrt{\frac{3}{5}}$  (b)

$\sqrt{\frac{5}{3}}$  (c)  $\sqrt{\frac{2}{5}}$  (d)  $\sqrt{\frac{5}{2}}$

A.  $\sqrt{3/5}$

B.  $\sqrt{5/3}$

C.  $\sqrt{2/5}$

D.  $\sqrt{5/2}$

**Answer: B::D**

[Watch Video Solution](#)

8. find the equation of parabola which is symmetric about y-axis, and passes through point (2,-3).

[Watch Video Solution](#)



9. If the ellipse  $x^2 + 2y^2 = 4$  and the hyperbola  $S = 0$  have same end points of the latus rectum, then the eccentricity of the hyperbola can be

A.  $\operatorname{cosec} \frac{\pi}{4}$

B.  $\operatorname{cosec} \frac{\pi}{3}$

C.  $2 \sin. \frac{\pi}{3} + \sin. \frac{\pi}{4}$

D.  $\sqrt{2} \sin. \frac{\pi}{3} + \sin. \frac{\pi}{4}$

Answer: A:D



Watch Video Solution

10. For which of the hyperbolas, can we have more than one pair of perpendicular tangents?  $\frac{x^2}{4} - \frac{y^2}{9} = 1$  (b)  $\frac{x^2}{4} - \frac{y^2}{9} = -1$   
 $x^2 - y^2 = 4$  (d)  $xy = 44$

A.  $\frac{x^2}{4} - \frac{y^2}{9} = 1$

$$B. \frac{x^2}{4} - \frac{y^2}{9} = -1$$

$$C. x^2 - y^2 = 4$$

$$D. xy = 44$$

**Answer: B**



**Watch Video Solution**

11. The lines parallel to the normal to the curve  $xy = 1$  is/are (a)

$$3x + 4y + 5 = 0 \quad (b) \quad 3x - 4y + 5 = 0 \quad (c) 4x + 3y + 5 = 0 \quad (d)$$

$$3y - 4x + 5 = 0$$

$$A. 3x + 4y + 5 = 0$$

$$B. 3x - 4y + 5 = 0$$

$$C. 4x + 3y + 5 = 0$$

$$D. 3y - 4x + 5 = 0$$

**Answer: B::D**



[Watch Video Solution](#)

12. From the point (2, 2) tangent are drawn to the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ . Then the point of contact lies in the (a) first quadrant (b) second quadrant (c) third quadrant (d) fourth quadrant

- A. first quadrant
- B. second quadrant
- C. third quadrant
- D. forth quadrant

**Answer: C::D**



[Watch Video Solution](#)

13. For hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , let n be the number of points on the plane through which perpendicular tangents are drawn.

A. If  $n = 1$ , then  $e = \sqrt{2}$

B. If  $n > 1$ , then  $0 < e < \sqrt{2}$ .

C. If  $n = 0$ , then  $e > \sqrt{2}$ .

D. none of these

**Answer: A::B::C**



**Watch Video Solution**

14. If the normal at P to the rectangular hyperbola  $x^2 - y^2 = 4$  meets the axes in G and g and C is the centre of the hyperbola, then

A.  $PG = PC$

B.  $Pg = PC$

C.  $PG = Pg$

D.  $Gg = 2PC$

**Answer: A::B::C::D**

 [Watch Video Solution](#)

15. Find the equation of tangent to the hyperbola  $y = \frac{x + 9}{x + 5}$  which passes through  $(0, 0)$  origin

A.  $x + 25y = 0$

B.  $x + y = 0$

C.  $5x - y = 0$

D.  $x - 25y = 0$

**Answer: A::B**

 [Watch Video Solution](#)

16. Tangents which are parallel to the line  $2x + y + 8 = 0$  are drawn to hyperbola  $x^2 - y^2 = 3$ . The points of contact of these tangents is/are

A.  $(2,1)$

B.  $(2, -1)$

C.  $(-2, -1)$

D.  $(-2, 1)$

**Answer: B::D**



**Watch Video Solution**

17. Find the equations of the tangents to the hyperbola  $x^2 - 9y^2 = 9$  that are drawn from  $(3, 2)$ .

A. equation of one of the tangents is  $x = 3$

B. equation of one of the tangents is  $5x - 12y + 9 = 0$

C. the area of triangle that these tangents form with their chord of contact is 12 sq. units

D. the area of triangle that these tangents form with their chord of contact is 8 sq. units

Answer: A::B::D

 Watch Video Solution

18. Find the equation of the circle which passes through the points  $(2, -2)$ , and  $(3,4)$  and whose centre lies on the line  $x + y = 2$ .

 Watch Video Solution

19. If  $P(\alpha, \beta)$ , the point of intersection of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{a^2}(1 - e^2) = 1$  and hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{a^2(E^2 - 1)} = \frac{1}{4}$  is equidistant from the foci of the curves all lying in the right of y-axis then

A.  $2\alpha = a(2e + E)$

B.  $a - e\alpha = E\alpha - \alpha/2$

C.  $E = \frac{\sqrt{e^2 + 24} - 3e}{2}$

D.  $E = \frac{\sqrt{e^2 + 12} - 3e}{2}$

Answer: B::C



Watch Video Solution

20. A hyperbola having the transverse axis of length  $\frac{1}{2}$  unit is confocal with the ellipse  $3x^2 + 4y^2 = 12$ , then

A. Equation of the hyperbola is  $\frac{x^2}{15} - \frac{y^2}{1} = \frac{1}{16}$

B. Eccentricity of the hyperbola is 4

C. Distance between the directrices of the hyperbola is  $\frac{1}{8}$  units

D. Length of latus rectum of the hyperbola is  $\frac{15}{2}$  units

Answer: B::C::D



Watch Video Solution

21. In X-Y plane, the path defined by the equation

$\frac{1}{x^m} + \frac{1}{y^m} + \frac{k}{(x+y)^n} = 0$ , is (a) a parabola if



$m = \frac{1}{2}, k = -1, n = 0$  (b) a hyperbola if  $m = 1, k = -1, n = 0$  (c) a pair of lines if  $m = k = n = 1$  (d) a pair of lines if  $m = k = -1, n = 1$

A. a parabola if  $m = \frac{1}{2}, k = -1, n = 0$

B. a hyperbola if  $m = 1, k = -1, n = 0$

C. a pair of lines if  $m = k = n = 1$

D. a pair of lines if  $m = k = -1, n = 1$

**Answer: A::B::C::D**



**Watch Video Solution**

**22.** A point moves such that the sum of the squares of its distances from the two sides of length 'a' of a rectangle is twice the sum of the squares of its distances from the other two sides of length b. The locus of the point can be:

A. a circle

B. an ellipse

C. a hyperbola

D. a pair of lines

**Answer: C::D**



[Watch Video Solution](#)

**23.** The equation of a hyperbola with co-ordinate axes as principal axes, and the distances of one of its vertices from the foci are 3 and 1 can be

A.  $3x^2 - y^2 = 3$

B.  $x^2 - 3y^2 + 3 = 0$

C.  $x^2 - 3y^2 - 3 = 0$

D. none of these

**Answer: A::B**



[Watch Video Solution](#)

24. Three points A, B and C taken on rectangular hyperbola  $xy = 4$  where  $B(-2, -2)$  and  $C(6, 2/3)$ . The normal at A is parallel to BC, then

A. circumcentre of  $\triangle ABC$  is  $(2, -2/3)$

B. equation of circumcircle of  $\triangle ABC$  is

$$3x^2 + 3y^2 - 12x + 4y - 40 = 0$$

C. orthocenter of  $\triangle ABC$  is  $\left(\frac{2}{\sqrt{3}}, 2\sqrt{3}\right)$

D. none of these

**Answer: A::B::C**



**Watch Video Solution**

25. A tangent is drawn at any point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . If this tangent is intersected by the tangents at the vertices at points P and Q, then which of the following is/are true

A. S, S', P and Q are concyclic

B. PQ is diameter of the circle

C. S,S', P and Q forms rhombus

D. PQ is diagonal of acute angle of the rhombus formed by S,S',P and Q

**Answer: A::B**



**Watch Video Solution**

26. If two tangents can be drawn the different branches of hyperbola

$$\frac{x^2}{1} - \frac{y^2}{4} = 1 \text{ from } (\alpha, \alpha^2), \text{ then}$$

A.  $\alpha \in (-2, 0)$

B.  $\alpha \in (0, 2)$

C.  $\alpha \in (-\infty, -2)$

D.  $\alpha \in (2, \infty)$

**Answer: C::D**



**Watch Video Solution**

27. The director circle of a hyperbola is  $x^2 + y^2 - 4y = 0$ . One end of the major axis is (2,0) then a focus is (a)  $(\sqrt{3}, 2 - \sqrt{3})$  (b)  $(-\sqrt{3}, 2 + \sqrt{3})$  (c)  $(\sqrt{6}, 2 - \sqrt{6})$  (d)  $(-\sqrt{6}, 2 + \sqrt{6})$

A.  $(\sqrt{3}, 2 - \sqrt{3})$

B.  $(-\sqrt{3}, 2 + \sqrt{3})$

C.  $(\sqrt{6}, 2 - \sqrt{6})$

D.  $(-\sqrt{6}, 2 + \sqrt{6})$

Answer: C::D



Watch Video Solution

28. The points on the ellipse  $\frac{x^2}{2} + \frac{y^2}{10} = 1$  from which perpendicular tangents can be drawn to the hyperbola  $\frac{x^2}{5} - \frac{y^2}{1} = 1$  is/are

A.  $\left(\sqrt{\frac{3}{2}}, \sqrt{\frac{5}{2}}\right)$

B.  $\left(\sqrt{\frac{3}{2}}, -\sqrt{\frac{5}{2}}\right)$

C.  $\left(-\sqrt{\frac{3}{2}}, \sqrt{\frac{5}{2}}\right)$

D.  $\left(\sqrt{\frac{5}{2}}, \sqrt{\frac{3}{2}}\right)$

**Answer: A::B::C**



**Watch Video Solution**

## Comorehension Type

1. Consider an ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  Let a hyperbola is having its vertices at the extremities of minor axis of an ellipse and length of major axis of an ellipse is equal to the distance between the foci of hyperbola. Let  $e_1$  and  $e_2$  be the eccentricities of an ellipse and hyperbola respectively. Again let A be the area of the quadrilateral formed by joining all the foci and A, be the area of the quadrilateral formed by all the directrices. The relation between  $e_1$  and  $e_2$  is given by

A.  $e_1 e_2 = 1$

B.  $e_2^2(1 - e_1^2) = 1$

C.  $e_1^2(e_2^2 - 1) = 1$

D.  $e_1 e_2(1 - e_1^2) = 1$

**Answer: B**

 [Watch Video Solution](#)

2. Consider an ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 (a > b)$ . A hyperbola has its vertices at the extremities of minor axis of the ellipse and the length of major axis of the ellipse is equal to the distance between the foci of hyperbola. Let  $e_1$  and  $e_2$  be the eccentricities of ellipse and hyperbola, respectively. Also, let  $A_1$  be the area of the quadrilateral formed by joining all the foci and  $A_2$  be the area of the quadrilateral formed by all the directrices.

The relation between  $e_1$  and  $e_2$  is given by

A.  $\tan^{-1} \left( \frac{1}{\sqrt{1 - e_1^2}} \right)$

B.  $\tan^{-1} \left( \frac{e_1}{\sqrt{1 - e_1^2}} \right)$

C.  $\tan^{-1} \left( \frac{1}{\sqrt{e_2^2 - 1}} \right)$

D.  $\tan^{-1} \sqrt{1 - e_1^2}$

**Answer: C**



**Watch Video Solution**

3. Consider the ellipse  $E_1$ ,  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , ( $a > b$ ). An ellipse  $E_2$  passes through the extremities of the major axis of  $E_1$  and has its foci at the ends of its minor axis. Consider the following property: Sum of focal distances of any point on an ellipse is equal to its major axis. Equation of  $E_2$  is

A. 2 : 1

B. 3 : 2

C.  $\sqrt{2} : 1$



D. 5 : 2

**Answer: D**

 [Watch Video Solution](#)

4. Consider the hyperbola  $\frac{X^2}{9} - \frac{y^2}{a^2} = 1$  and the circle  $x^2 + (y - 3) = 9$ .

Also, the given hyperbola and the ellipse  $\frac{x^2}{41} + \frac{y^2}{16} = 1$  are orthogonal to each other.

The number of points on the hyperbola and the circle from which tangents drawn to the circle and the hyperbola, respectively, are perpendicular to each other is

A.  $x^2 - y^2 = 0$

B.  $x^2 - 9 = 0$

C.  $9y^2 - 19x^2 = 0$

D. No common tangent.

**Answer: B**



**Watch Video Solution**

5. Consider the hyperbola  $\frac{X^2}{9} - \frac{y^2}{a^2} = 1$  and the circle  $x^2 + (y - 3) = 9$ .

Also, the given hyperbola and the ellipse  $\frac{x^2}{41} + \frac{y^2}{16} = 1$  are orthogonal to each other.

The number of points on the hyperbola and the circle from which tangents drawn to the circle and the hyperbola, respectively, are perpendicular to each other is

- A. 0
- B. 2
- C. 4
- D. 6

**Answer: C**



Watch Video Solution

6. Consider the hyperbola  $\frac{x^2}{9} - \frac{y^2}{a^2} = 1$  and the circle  $x^2 + (y - 3) = 9$ .

Also, the given hyperbola and the ellipse  $\frac{x^2}{41} + \frac{y^2}{16} = 1$  are orthogonal to each other.

Combined equation of pair of common tangents between the hyperbola and the circle is given be: (1). $x^2 - y^2 = 0$  (2). $x^2 - 9 = 0$  (3). $9y^2 - 19x^2 = 0$  (4).No common tangent.

A.  $x^2 + y^2 \pm 5x - 3y = 0$

B.  $x^2 + y^2 + 5x \pm 3y = 0$

C.  $x^2 - y^2 \pm 5x - 3y = 0$

D.  $x^2 - y^2 + 3x \pm 3y = 0$

**Answer: A**



Watch Video Solution

7. The locus of the foot of perpendicular from my focus of a hyperbola upon any tangent to the hyperbola is the auxiliary circle of the hyperbola. Consider the foci of a hyperbola as  $(-3, -2)$  and  $(5,6)$  and the foot of perpendicular from the focus  $(5, 6)$  upon a tangent to the hyperbola as  $(2, 5)$ .

The conjugate axis of the hyperbola is

A.  $4\sqrt{11}$

B.  $2\sqrt{11}$

C.  $4\sqrt{22}$

D.  $2\sqrt{22}$

**Answer: D**



[Watch Video Solution](#)

8. The locus of the foot of perpendicular from my focus of a hyperbola upon any tangent to the hyperbola is the auxiliary circle of the hyperbola.

Consider the foci of a hyperbola as  $(-3, -2)$  and  $(5,6)$  and the foot of perpendicular from the focus  $(5,6)$  upon a tangent to the hyperbola as  $(2, 5)$ .

The directrix of the hyperbola corresponding to the focus  $(5, 6)$  is

A.  $2x + 2y - 1 = 0$

B.  $2x + 2y - 11 = 0$

C.  $2x + 2y - 7 = 0$

D.  $2x + 2y - 9 = 0$

**Answer: B**



[Watch Video Solution](#)

9. The locus of the foot of perpendicular from any focus of a hyperbola upon any tangent to the hyperbola is the auxiliary circle of the hyperbola.

Consider the foci of a hyperbola as  $(-3, -2)$  and  $(5,6)$  and the foot of perpendicular from the focus  $(5,6)$  upon a tangent to the hyperbola as  $(2,$

5).

The point of contact of the tangent with the hyperbola is

A.  $(2/9, 31/3)$

B.  $(7/4, 23/4)$

C.  $(2/3, 9)$

D.  $(7/9, 7)$

**Answer: C**



**Watch Video Solution**

10. Let  $P(x, y)$  is a variable point such that

$$\left| \sqrt{(x-1)^2 + (y-2)^2} - \sqrt{(x-5)^2 + (y-5)^2} \right| = 3, \quad \text{which}$$

represents hyperbola. The eccentricity  $e'$  of the corresponding conjugate

hyperbola is (A)  $\frac{5}{3}$  (B)  $\frac{4}{3}$  (C)  $\frac{5}{4}$  (D)  $\frac{3}{\sqrt{7}}$

A.  $5/3$

B.  $4/3$

C.  $5/4$

D.  $3/\sqrt{7}$

**Answer: C**



**Watch Video Solution**

11. Let  $P(x, y)$  is a variable point such that  $\left| \sqrt{(x-1)^2 + (y-2)^2} - \sqrt{(x-5)^2 + (y-5)^2} \right| = 3$ , which represents hyperbola. The eccentricity  $e$  of the corresponding conjugate hyperbola is (A)  $\frac{5}{3}$  (B)  $\frac{4}{3}$  (C)  $\frac{5}{4}$  (D)  $\frac{3}{\sqrt{7}}$

A.  $(x-3)^2 + \left(y - \frac{7}{2}\right)^2 = \frac{55}{4}$

B.  $(x-3)^2 + \left(y - \frac{7}{2}\right)^2 = \frac{25}{4}$

C.  $(x-3)^2 + \left(y - \frac{7}{2}\right)^2 = \frac{7}{4}$

D. none of these

**Answer: D**



[Watch Video Solution](#)

12. Let  $P(x, y)$  is a variable point such that  $\left| \sqrt{(x-1)^2 + (y-2)^2} - \sqrt{(x-5)^2 + (y-5)^2} \right| = 3$ , which represents hyperbola. The eccentricity  $e'$  of the corresponding conjugate hyperbola is (A)  $\frac{5}{3}$  (B)  $\frac{4}{3}$  (C)  $\frac{5}{4}$  (D)  $\frac{3}{\sqrt{7}}$

A.  $\tan^{-1}(4/3)$

B.  $\tan^{-1}(3/4)$

C.  $\tan^{-1}(5/3)$

D.  $\tan^{-1}(3/5)$

**Answer: B**

[Watch Video Solution](#)

13. In a hyperbola, the portion of the tangent intercepted between the asymptotes is bisected at the point of contact.



Consider a hyperbola whose center is at the origin. A line  $x + y = 2$  touches this hyperbola at  $P(1,1)$  and intersects the asymptotes at A and B such that  $AB = 6\sqrt{2}$  units.

The equation of the pair of asymptotes is

A.  $5xy + 2x^2 + 2y^2 = 0$

B.  $3x^2 + 4y^2 + 6xy = 0$

C.  $2x^2 + 2y^2 - 5xy = 0$

D. none of these

**Answer: A**



[Watch Video Solution](#)

**14.** In a hyperbola, the portion of the tangent intercepted between the asymptotes is bisected at the point of contact.

Consider a hyperbola whose center is at the origin. A line  $x + y = 2$  touches this hyperbola at  $P(1,1)$  and intersects the asymptotes at A and B

such that  $AB = 6\sqrt{2}$  units.

The equation of the pair of asymptotes is

A.  $\sin^{-1} \cdot \frac{4}{5}$

B.  $\sin^{-1} \cdot \frac{2}{5}$

C.  $\sin^{-1} \cdot \frac{3}{5}$

D. none of these

**Answer: C**



**Watch Video Solution**

**15.** In a hyperbola, the portion of the tangent intercepted between the asymptotes is bisected at the point of contact.

Consider a hyperbola whose center is at the origin. A line  $x + y = 2$  touches this hyperbola at  $P(1,1)$  and intersects the asymptotes at  $A$  and  $B$  such that  $AB = 6\sqrt{2}$  units.

The equation of the tangent to the hyperbola at  $(-1, 7/2)$  is

A.  $5x + 2y = 2$

B.  $3x + 2y = 4$

C.  $3x + 4y = 11$

D. none of these

**Answer: B**



[Watch Video Solution](#)

**16.** A point P moves such that sum of the slopes of the normals drawn from it to the hyperbola  $xy=16$  is equal to the sum of ordinates of feet of normals. The locus of P is a curve C

A. (A)  $x^2 = 4y$

B. (B)  $x^2 = 16y$

C. (C)  $x^2 = 12y$

D. (D)  $y^2 = 8x$

**Answer: B**



[Watch Video Solution](#)

17. A point P moves such that the sum of the slopes of the normals drawn from it to the hyperbola  $xy = 16$  is equal to the sum of ordinates of feet of normals. The locus of P is a curve C.

If the tangent to the curve C cuts the coordinate axes at A and B, then the locus of the middle point of AB is

A.  $x^2 = 16y$

B.  $x^2 = 2y$

C.  $x^2 + 2y = 0$

D.  $x^2 + 4y = 0$

**Answer: C**



[Watch Video Solution](#)

18. A point  $P$  moves such that the sum of the slopes of the normals drawn from it to the hyperbola  $xy = 16$  is equal to the sum of ordinates of feet of normals. The locus of  $P$  is a curve  $C$ .

the equation of the curve  $C$  is

 [Watch Video Solution](#)

19. A triangle has its vertices on a rectangular hyperbola. Prove that the orthocentre of the triangle also lies on the same hyperbola.

A.  $xy - 1 = x - y$

B.  $xy + 1 = x + y$

C.  $2xy = x + y$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

20. A triangle has its vertices on a rectangular hyperbola. Prove that the orthocentre of the triangle also lies on the same hyperbola.

A.  $xy = 2x + y - 2$

B.  $2xy = x + 2y + 5$

C.  $xy = x + y + 1$

D. none of these

**Answer: C**



[Watch Video Solution](#)

21. A triangle has its vertices on a rectangular hyperbola. Prove that the orthocentre of the triangle also lies on the same hyperbola.

A. 4

B. 0

C. 3

Answer: D



Watch Video Solution

## Matrix Mathc Type

1. Let the foci of the hyperbola  $\frac{X^2}{A^2} - \frac{y^2}{B^2} = 1$  be the vertices of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and the foci of the ellipse be the vertices of the hyperbola. Let the eccentricities of the ellipse and hyperbola be  $e_E$  and  $e_H$ , respectively. Then match the following lists.



View Text Solution

2. Match the following lists:



### 3. Match the following lists:

List I	List II
<b>a.</b> Four lines $x + 3y - 10 = 0$ , $x + 3y - 20 = 0$ , $3x - y + 5 = 0$ , and $3x - y - 5 = 0$ form a figure which is	<b>p.</b> a quadrilateral which is neither a parallelogram nor a trapezium
<b>b.</b> The points $A(1, 2)$ , $B(2, -3)$ , $C(-1, -5)$ , and $D(-2, 4)$ in order are the vertices of	<b>q.</b> a parallelogram
<b>c.</b> The lines $7x + 3y - 33 = 0$ , $3x - 7y + 19 = 0$ , $3x - 7y - 10$ , and $7x + 3y - 4 = 0$ form a figure which is	<b>r.</b> a rectangle of area 10 sq. units
<b>d.</b> Four lines $4y - 3x - 7 = 0$ , $3y - 4x + 7 = 0$ , $4y - 3x - 21 = 0$ , $3y - 4x + 14 = 0$ form a figure which is	<b>s.</b> a square

6 Match the following lists:

### 4. Match the following lists:





5. Match the following lists:

List I	List II
a. Four lines $x + 3y - 10 = 0$ , $x + 3y - 20 = 0$ , $3x - y + 5 = 0$ , and $3x - y - 5 = 0$ form a figure which is	p. a quadrilateral which is neither a parallelogram nor a trapezium
b. The points $A(1, 2)$ , $B(2, -3)$ , $C(-1, -5)$ , and $D(-2, 4)$ in order are the vertices of	q. a parallelogram
c. The lines $7x + 3y - 33 = 0$ , $3x - 7y + 19 = 0$ , $3x - 7y - 10 = 0$ , and $7x + 3y - 4 = 0$ form a figure which is	r. a rectangle of area 10 sq. units
d. Four lines $4y - 3x - 7 = 0$ , $3y - 4x + 7 = 0$ , $4y - 3x - 21 = 0$ , $3y - 4x + 14 = 0$ form a figure which is	s. a square

6. Match the following lists:



Watch Video Solution

6. Consider all possible permutations of the letters of the word ENDEANOEL. Match the statements/ expressions in column I with the

statement/expressions in

	Column I		Column II
(A)	The number of permutations containing the word ENDEA is	(p)	$5!$
(B)	The number of permutations in which the letters E occurs in the first and the last positions, is	(q)	$2 \times 5!$
(C)	The number of permutations in which none of the letters D, L, N occurs in the last five positions, is	(r)	$7 \times 5!$
(D)	The number of permutations in which the letters A, E, O occur only in odd positions, is	(s)	$21 \times 5!$

 [Watch Video Solution](#)

7. Match the following and mark the correct options

Animal	Respiratory Organ
A. Earthworm	i. Moist cuticle
B. Aquatic Arthropods	ii. Gills
C. Fishes	iii. Lungs
D. Birds/Reptiles	iv. Trachea

A. (II) (iii) (R)

B. (IV) (iv) (S)

C. (IV) (iii) (S)

D. (II) (iv) (R)

**Answer: D**

 **Watch Video Solution**

**8. Match the items of columns I and II.**

Column-I		Column-II	
(P)	Z for ideal gas behaviour	(1)	$3/8$
(Q)	Z for real gas at low pressure	(2)	$\left(1 + \frac{Pb}{RT}\right)$
(R)	Z for real gas at high pressure	(3)	1
(S)	Z for critical state	(4)	$\left(1 - \frac{a}{RTV}\right)$

A. (II) (i) (P)

B. (III) (iO) (Q)

C. (II) (iv) (R)

D. (I) (ii) (Q)

**Answer: A**



[Watch Video Solution](#)

9. Lists I, II and III contains conics, equation of tangents to the conics and points of contact, respectively.



For  $a = \sqrt{2}$  if a tangent is drawn to a suitable conic (List I) at the point of contact  $(-1, 1)$ , which of the following options is the only CORRECT combination for obtaining its equation?

A. (II) (ii) (Q)

B. (III) (i) (P)

C. (I) (i) (P)

D. (I) (ii) (Q)

**Answer: D**



[View Text Solution](#)

10. The equivalent capacitance between points A and B will be



A.  $P \rightarrow IV, Q \rightarrow II, R \rightarrow I, S \rightarrow III$

B.  $P \rightarrow IV, Q \rightarrow III, R \rightarrow I, S \rightarrow II$

C.  $P \rightarrow IV, Q \rightarrow I, R \rightarrow III, S \rightarrow II$

D.  $P \rightarrow III, Q \rightarrow IV, R \rightarrow II, S \rightarrow I$

**Answer: B**



**Watch Video Solution**

**Numerical Value Type**

1. The eccentricity of the hyperbola

$$\left| \sqrt{(x-3)^2 + (y-2)^2} - \sqrt{(x+1)^2 + (y+1)^2} \right| = 1 \text{ is } \underline{\hspace{2cm}}$$



**Watch Video Solution**

2. If  $y = mx + c$  is tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , having eccentricity 5, then the least positive integral value of  $m$  is \_\_\_\_

 [Watch Video Solution](#)

3. Consider the graphs of  $y = Ax^2$  and  $y^2 + 3 = x^2 + 4y$ , where  $A$  is a positive constant and  $x, y \in R$ . Number of points in which the two graphs intersect, is

 [Watch Video Solution](#)

4. If  
 $4(x - \sqrt{2})^2 + \lambda(y - \sqrt{3})^2 = 45$  and  $(x - \sqrt{2})^2 - 4(y - \sqrt{3})^2 = 5$   
cut orthogonally, then integral value of  $\lambda$  is \_\_\_\_\_.

 [Watch Video Solution](#)

5. If the hyperbola  $x^2 - y^2 = 4$  is rotated by  $45^\circ$  in the anticlockwise direction about its center keeping the axis intact, then the equation of the hyperbola is  $xy = a^2$ , where  $a^2$  is equal to \_\_\_\_\_



[Watch Video Solution](#)

6. Tangents are drawn from the point  $(\alpha, \beta)$  to the hyperbola  $3x^2 - 2y^2 = 6$  and are inclined at an angle  $\theta$  and  $\phi$  to the x-axis. If  $\tan \theta \cdot \tan \phi = 2$ , prove that  $\beta^2 = 2\alpha^2 - 7$ .



[Watch Video Solution](#)

7. The area of triangle formed by the tangents from the point  $(3, 2)$  to the hyperbola  $x^2 - 9y^2 = 9$  and the chord of contact w.r.t. the point  $(3, 2)$  is \_\_\_\_\_



[Watch Video Solution](#)

8. The values of 'm' for which a line with slope m is common tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and parabola  $y^2 = 4ax$  can lie in interval:

 [Watch Video Solution](#)

9. If tangents drawn from the point  $(a, 2)$  to the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  are perpendicular, then the value of  $a^2$  is \_\_\_\_\_

 [Watch Video Solution](#)

10. If radii of director circles of  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are  $2r$  and  $r$  respectively, let  $e_E$  and  $e_H$  are the eccentricities of ellipse and hyperbola respectively, then

 [Watch Video Solution](#)

11. If  $L$  is the length of the latus rectum of the hyperbola for which  $x = 3$  and  $y = 2$  are the equations of asymptotes and which passes



through the point  $(4, 6)$ , then the value of  $\frac{L}{\sqrt{2}}$  is \_\_\_\_\_

 [Watch Video Solution](#)

12. If the angle between the asymptotes of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is  $\frac{\pi}{3}$ , then the eccentricity of conjugate hyperbola is \_\_\_\_\_.

 [Watch Video Solution](#)

13. If the chord  $x \cos \alpha + y \sin \alpha = p$  of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{18} = 1$  subtends a right angle at the center, and the diameter of the circle, concentric with the hyperbola, to which the given chord is a tangent is  $d$ , then the value of  $\frac{d}{4}$  is \_\_\_\_\_

 [Watch Video Solution](#)

14. The line  $2x + y = 1$  is tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . If this line passes through the point of intersection of the nearest directrix and

the x-axis, then the eccentricity of the hyperbola is

 [Watch Video Solution](#)

Jee Main

1. The eccentricity of the hyperbola whose length of the latus rectum is equal to 8 and the length of its conjugate axis is equal to half of the distance between its foci, is :

A. (a)  $4/\sqrt{3}$

B. (b)  $2/\sqrt{3}$

C. (c)  $\sqrt{3}$

D. (d)  $4/3$

**Answer: B**

 [Watch Video Solution](#)

2. A hyperbola passes through the point  $(\sqrt{2}, \sqrt{3})$  and has foci at  $(\pm 2, 0)$ . Then the tangent to this hyperbola at P also passes through the point:

A. (a)  $(-\sqrt{2}, -\sqrt{3})$

B. (b)  $(3\sqrt{2}, 2\sqrt{3})$

C. (c)  $(2\sqrt{2}, 3\sqrt{3})$

D. (d)  $(\sqrt{3}, \sqrt{2})$

**Answer: C**



**Watch Video Solution**

3. Tangents are drawn to the hyperbola  $4x^2 - y^2 = 36$  at the points P and Q. If these tangents intersect at the point T(0,3) then the area (in sq units) of  $\triangle PTQ$  is

A.  $36\sqrt{5}$

B.  $45\sqrt{5}$

C.  $54\sqrt{3}$

D.  $60\sqrt{3}$

**Answer: B**



**Watch Video Solution**

**Jee Advanced**

1. Let P(6,3) be a point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  If the normal at the point intersects the x-axis at (9,0), then the eccentricity of the hyperbola is

A.  $\sqrt{5/2}$

B.  $\sqrt{3/2}$

C.  $\sqrt{2}$

D.  $\sqrt{3}$

**Answer: B**



**Watch Video Solution**

2. An ellipse intersects the hyperbola  $2x^2 - 2y = 1$  orthogonally. The eccentricity of the ellipse is reciprocal to that of the hyperbola. If the axes of the ellipse are along the coordinate axes, then (a) the foci of ellipse are  $(\pm 1, 0)$  (b) equation of ellipse is  $x^2 + 2y^2 = 2$  (c) the foci of ellipse are  $(t2, 0)$  (d) equation of ellipse is  $(x^2 2y)$

A. the equation of the ellipse is  $x^2 + 2y^2 = 1$

B. the foci of the ellipse are  $(\pm 1, 0)$

C. the equation of the ellipse is  $x^2 + 2y^2 = 4$

D. the foci of the ellipse are  $(\pm \sqrt{2}, 0)$

**Answer: A::B**



**Watch Video Solution**

### 3. about to only mathematics

A. the equation of the hyperbola is  $\frac{x^2}{3} - \frac{y^2}{2} = 1$

B. a focus of the hyperbola is (2,0)

C. the eccentricity of the hyperbola is  $\frac{2}{\sqrt{3}}$

D. the equation of the hyperbola is  $x^2 - 3y^2 = 3$

**Answer: B::D**



**Watch Video Solution**

4. Tangents are drawn to the hyperbola  $\frac{x^2}{9} - \frac{y^2}{4} = 1$  parallel to the straight line  $2x - y = 1$ . The points of contact of the tangents on the hyperbola are (A)  $\left(\frac{9}{2\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  (B)  $\left(-\frac{9}{2\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$  (C)  $(3\sqrt{3}, -2\sqrt{2})$  (D)  $(-3\sqrt{3}, 2\sqrt{2})$

A.  $\left(\frac{9}{2\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

B.  $\left(-\frac{9}{2\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$

C.  $(3\sqrt{3}, -2\sqrt{2})$

D.  $(3\sqrt{3}, -2\sqrt{2})$

**Answer: A::B**



**Watch Video Solution**

5. Find the equations of the tangent and the normal to  $16x^2 + 9y^2 = 144$  at  $(x_1, y_1)$  where  $x_1 = 2$  and  $y_1 > 0$ .

A.  $\frac{dl}{dx_1} = 1 - \frac{1}{3x_1^2}$  for  $x_1 > 1$

B.  $\frac{dm}{dx_1} = \frac{x_1}{3\sqrt{x_1^2 - 1}}$  for  $x_1 > 1$

C.  $\frac{dl}{dx_1} = 1 + \frac{1}{3x_1^2}$  for  $x_1 > 1$

D.  $\frac{dm}{dy_1} = \frac{1}{3}$  for  $x_1 > 0$

**Answer: A::B::D**



**Watch Video Solution**

6. If  $2x - y + 1 = 0$  is a tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{16} = 1$  then which of the following CANNOT be sides of a right angled triangle?

A. (a)  $2a, 4, 1$

B. (b)  $2a, 8, 1$

C. (c)  $a, 4, 1$

D. (d)  $a, 4, 2$

**Answer: B::C::D**



**Watch Video Solution**

## Comprehension Type

1. The circle  $x^2 + y^2 - 8x = 0$  and hyperbola  $\frac{x^2}{9} - \frac{y^2}{4} = 1$  intersect at the points A and B. Equation of a common tangent with positive slope to the circle as well as to the hyperbola is



A.  $2x - \sqrt{5}y - 20 = 0$

B.  $2x - \sqrt{5}y + 4 = 0$

C.  $3x - 4y + 8 = 0$

D.  $4x - 3y + 4 = 0$

**Answer: B**



**Watch Video Solution**

2. The circle  $x^2 + y^2 - 8x = 0$  and hyperbola  $\frac{x^2}{9} - \frac{y^2}{4} = 1$  intersect at the points A and B. Then the equation of the circle with AB as its diameter is

A. a.  $x^2 + y^2 - 12x + 24 = 0$

B. b.  $x^2 + y^2 + 12x + 24 = 0$

C. c.  $x^2 + y^2 + 24x - 12 = 0$

D. d.  $x^2 + y^2 - 24x - 12 = 0$

**Answer: A**



**Watch Video Solution**

3. Consider a hyperbola  $xy = 4$  and a line  $y = 2x = 4$ . O is the centre of hyperbola. Tangent at any point P of hyperbola intersect the coordinate axes at A and B.

Locus of circumcentre of triangle OAB is

- A. an ellipse with eccentricity  $\frac{1}{\sqrt{2}}$
- B. a parabola
- C. a hyperbola with eccentricity  $\sqrt{2}$
- D. a circle

**Answer: C**



**Watch Video Solution**

4. Consider a hyperbola  $xy = 4$  and a line  $y = 2x = 4$ . O is the centre of hyperbola. Tangent at any point P of hyperbola intersect the coordinate axes at A and B.

Shortest distance between the line and hyperbola is

A.  $\frac{8\sqrt{2}}{\sqrt{5}}$

B.  $\frac{4(\sqrt{2} - 1)}{\sqrt{5}}$

C.  $\frac{2\sqrt{2}}{\sqrt{5}}$

D.  $\frac{4(\sqrt{2} - 1)}{\sqrt{5}}$

**Answer: B**



[Watch Video Solution](#)

5. Consider a hyperbola  $xy = 4$  and a line  $y = 2x = 4$ . O is the centre of hyperbola. Tangent at any point P of hyperbola intersect the coordinate axes at A and B.

Let the given line intersects the x-axis at R. if a line through R. intersect the hyperbolas at S and T, then minimum value of  $RS \times RT$  is

A. 2

B. 4

C. 6

D. 8

**Answer: D**

 [Watch Video Solution](#)

6. Consider a hyperbola:  $\frac{(x - 7)^2}{a} - \frac{(y + 3)^2}{b^2} = 1$ . The line  $3x - 2y - 25 = 0$ , which is not a tangent, intersect the hyperbola at  $H\left(\frac{11}{3}, -7\right)$  only. A variable point  $P(\alpha + 7, \alpha^2 - 4) \forall \alpha \in \mathbb{R}$  exists in the plane of the given hyperbola.

The eccentricity of the hyperbola is

A.  $\sqrt{\frac{7}{5}}$

B.  $\sqrt{2}$

C.  $\frac{\sqrt{13}}{2}$

D.  $\frac{3}{2}$

**Answer: C**



**Watch Video Solution**

7. Consider a hyperbola:  $\frac{(x - 7)^2}{a} - \frac{(y + 3)^2}{b^2} = 1$ . The line  $3x - 2y - 25 = 0$ , which is not a tangent, intersect the hyperbola at  $H\left(\frac{11}{3}, -7\right)$  only. A variable point  $P(\alpha + 7, \alpha^2 - 4) \forall \alpha \in R$  exists in the plane of the given hyperbola.

The eccentricity of the hyperbola is

A.  $(2, \infty)$

B.  $(-\infty, -2)$

C.  $\left(-\frac{1}{2}, \frac{1}{2}\right)$

D. None of these

**Answer: D**



[Watch Video Solution](#)

### Single Correct Answer Type

1. The locus of  $P(x, y)$  such that

$$\sqrt{x^2 + y^2 + 8y + 16} - \sqrt{x^2 + y^2 - 6x + 9} = 5, \text{ is}$$

- A. hyperbola
- B. circle
- C. finite line segment
- D. infinite ray

**Answer: D**



[Watch Video Solution](#)

2. The distance of the focus of  $x^2 - y^2 = 4$ , from the directrix, which is nearer to it, is

A.  $2\sqrt{2}$

B.  $\sqrt{2}$

C.  $4\sqrt{2}$

D.  $8\sqrt{2}$

**Answer: B**



[Watch Video Solution](#)

3. If  $\frac{x^2}{36} - \frac{y^2}{k^2} = 1$  is a hyperbola, then which of the following points lie on hyperbola?

A. (3, 1)

B. (-3, 1)

C. (5, 2)

D. (10, 4)

**Answer: D**



[Watch Video Solution](#)

4. The ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  and the hyperbola  $\frac{x^2}{25} - \frac{y^2}{16} = 1$  have in common

- A. centre and vertices only
- B. centre, foci and vertices
- C. centre, foci and directrices
- D. centre only

**Answer: A**



[Watch Video Solution](#)



5. The equation to the hyperbola having its eccentricity 2 and the distance between its foci is 8 is

A.  $\frac{x^2}{12} - \frac{y^2}{4} = 1$

B.  $\frac{x^2}{4} - \frac{y^2}{12} = 1$

C.  $\frac{x^2}{8} - \frac{y^2}{2} = 1$

D.  $\frac{x^2}{16} - \frac{y^2}{9} = 1$

**Answer: B**



[Watch Video Solution](#)

6. If the centre, vertex and focus of a hyperbola be (0,0), (4,0) and (6,0) respectively, then the equation of the hyperbola is

A.  $4x^2 - 5y^2 = 8$

B.  $4x^2 - 5y^2 = 80$

C.  $5x^2 - 4y^2 = 80$

D.  $5x^2 - 4y^2 = 8$

Answer: C



Watch Video Solution

7. The equation  $\frac{x^2}{9 - \lambda} + \frac{y^2}{4 - \lambda} = 1$  represents a hyperbola when  $a < \lambda < b$  then  $(b - a) =$

A. 3

B. 4

C. 5

D. 6

Answer: C



Watch Video Solution

8. If  $e$  and  $e'$  are the eccentricities of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$ , then the point  $\left(\frac{1}{e}, \frac{1}{e'}\right)$  lies on the circle (A)  $x^2 + y^2 = 1$  (B)  $x^2 + y^2 = 2$  (C)  $x^2 + y^2 = 3$  (D)  $x^2 + y^2 = 4$

A.  $x^2 + y^2 = 1$

B.  $x^2 + y^2 = 2$

C.  $x^2 + y^2 = 3$

D.  $x^2 + y^2 = 4$

**Answer: A**



[Watch Video Solution](#)

9. the eccentricity of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{25} = 1$  is



[Watch Video Solution](#)

10. The length of the transverse axis of the hyperbola  $9x^2 - 16y^2 - 18x - 32y - 151 = 0$  is

A. 8

B. 2

C. 6

D. 2

**Answer: A**



[Watch Video Solution](#)

11. A hyperbola has centre 'C' and one focus at  $P(6, 8)$ . If its two directrices are  $3x + 4y + 10 = 0$  and  $3x + 4y - 10 = 0$  then  $CP =$

A. 14

B. 8

C. 10

D. 6

**Answer: C**



**Watch Video Solution**

12. If the foci of  $\frac{x^2}{16} + \frac{y^2}{4} = 1$  and  $\frac{x^2}{a^2} - \frac{y^2}{3} = 1$  coincide, the value of  $a$  is

A. 3

B. 2

C.  $\frac{1}{\sqrt{3}}$

D.  $\sqrt{3}$

**Answer: A**



**Watch Video Solution**

13. A rectangular hyperbola of latus rectum 4 units passes through (0,0) and has (2,0) as its one focus. The equation of locus of the other focus is

A.  $x^2 + y^2 = 36$

B.  $x^2 + y^2 = 4$

C.  $x^2 - y^2 = 4$

D.  $x^2 + y^2 = 9$

**Answer: A**



[Watch Video Solution](#)

14. If the curves  $x^2 - y^2 = 4$  and  $xy = \sqrt{5}$  intersect at points A and B, then the possible number of points (s) C on the curve  $x^2 - y^2 = 4$  such that triangle ABC is equilateral is

A. 0

B. 1

C. 2

D. 4

**Answer: A**



[Watch Video Solution](#)

15. The point  $(3 \tan(\theta + 60^\circ), 2 \tan(\theta + 30^\circ))$  lies on the conic, then its centre is ( $\theta$  is the parameter)

A.  $(-3\sqrt{3}, 2\sqrt{3})$

B.  $(3\sqrt{3}, -2\sqrt{3})$

C.  $(-3\sqrt{3}, -2\sqrt{3})$

D. (0,0)

**Answer: A**



[Watch Video Solution](#)

16. The equation of a tangent to the hyperbola  $3x^2 - y^2 = 3$ , parallel to the line  $y = 2x + 4$  is

A.  $y = 2x + 3$

B.  $y = 2x + 1$

C.  $y = 2x + 4$

D.  $y = 2x + 2$

**Answer: B**



**Watch Video Solution**

17. A tangent to the hyperbola  $y = \frac{x + 9}{x + 5}$  passing through the origin is

A.  $x + 25y = 0$

B.  $5x + y = 0$

C.  $5x - y = 0$

D.  $x - 25y = 0$



**Answer: A**



[Watch Video Solution](#)

**18.** The absolute value of slope of common tangents to parabola  $y^2 = 8x$  and hyperbola  $3x^2 - y^2 = 3$  is

- A. 1
- B. 2
- C. 3
- D. 4

**Answer: B**



[Watch Video Solution](#)

**19.** For the hyperbola  $xy = 8$  any tangent of it at P meets co-ordinates at Q and R then area of triangle CQR where 'C' is centre of the hyperbola is

A. 16 sq. units

B. 12 sq. units

C. 24 sq. units

D. 18 sq. units

**Answer: A**



**Watch Video Solution**

20. The tangents and normal at a point on  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  cut the y-axis A and B. Then the circle on AB as diameter passes through the foci of the hyperbola

A. one of the vertex of the hyperbola

B. one of the foot of directrix on x-axis of the hyperbola

C. foci of the hyperbola

D. none of these

**Answer: C**



**Watch Video Solution**

21. If  $4x^2 + py^2 = 45$  and  $x^2 - 4y^2 = 5$  cut orthogonally, then the value of p is

A.  $1/9$

B.  $1/3$

C. 3

D. 9

**Answer: D**



**Watch Video Solution**

22. A tangent drawn to hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at  $P\left(\frac{\pi}{6}\right)$  forms a triangle of area  $3a^2$  square units, with coordinate axes, then the square of

its eccentricity is equal to

A. 15

B. 16

C. 17

D. 18

**Answer: C**



**Watch Video Solution**

**23.** If  $m$  is the slope of a tangent to the hyperbola

$$\frac{x^2}{a^2 - b^2} - \frac{y^2}{a^3 - b^3} = 1 \text{ where } a > b > 1 \text{ when}$$

A.  $(a + b)m^2 + ab \geq (a + b)^2$

B.  $(a + b)^2 m + ab \geq (a + b)$

C.  $abm^2 + (a + b) \geq (a + b)^2$

D.  $(a + b)m^2 + a^2 b^2 \geq (a + b)^2$

**Answer: A**



**Watch Video Solution**

**24.** Two tangents to the hyperbola  $\frac{x^2}{25} - \frac{y^2}{9} = 1$ , having slopes 2 and  $m$  where  $(m \neq 2)$  cuts the axes at four concyclic points then the slope  $m$  is/are

A.  $-\frac{1}{2}$

B.  $-2$

C.  $\frac{1}{2}$

D.  $2$

**Answer: C**



**Watch Video Solution**

25. The equation of that chord of hyperbola  $25x^2 - 16y = 400$ , whose mid point is (5,3) is

A.  $115x - 117y = 17$

B.  $125x - 48y = 481$

C.  $127x + 33y = 341$

D.  $15x - 121y = 105$

**Answer: B**



[Watch Video Solution](#)

26. If a chord joining  $P(a \sec \theta, a \tan \theta)$ ,  $Q(a \sec \alpha, a \tan \alpha)$  on the hyperbola  $x^2 - y^2 = a^2$  is the normal at P, then  $\tan \alpha$  is (a)  $\tan \theta(4 \sec^2 \theta + 1)$  (b)  $\tan \theta(4 \sec^2 \theta - 1)$  (c)  $\tan \theta(2 \sec^2 \theta - 1)$  (d)  $\tan \theta(1 - 2 \sec^2 \theta)$

A.  $\tan \theta(4 \sec^2 \theta + 1)$

B.  $\tan \theta (4 \sec^2 \theta - 1)$

C.  $\tan \theta (2 \sec^2 \theta - 1)$

D.  $\tan \theta (1 - 2 \sec^2 \theta)$

**Answer: B**

 [Watch Video Solution](#)

27. The number of normal (s) of a rectangular hyperbola which can touch its conjugate is equal to

A. 0

B. 2

C. 4

D. 8

**Answer: C**

 [Watch Video Solution](#)

28. If the normal at a point P to the hyperbola meets the transverse axis at G, and the value of SG/SP is 6, then the eccentricity of the hyperbola is (where S is focus of the hyperbola)

A. 2

B. 4

C. 6

D. 8

**Answer: C**



[Watch Video Solution](#)

29. If the normal at  $P(a \sec \theta, b \tan \theta)$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets the transverse axis in G then minimum length of PG is

A.  $\frac{b^2}{a}$



B.  $\left| \frac{a}{b}(a + b) \right|$

C.  $\left| \frac{a}{b}(a - b) \right|$

D.  $\left| \frac{a}{b}(a - b) \right|$

**Answer: A**



**Watch Video Solution**

30. If normal to hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  drawn at an extremity of its latus-rectum has slope equal to the slope of line which meets hyperbola only once, then the eccentricity of hyperbola is

A.  $e = \sqrt{\frac{1 + \sqrt{5}}{2}}$

B.  $e = \sqrt{\frac{\sqrt{5} + 3}{2}}$

C.  $e = \sqrt{\frac{2}{\sqrt{5} - 1}}$

D. None of these

**Answer: A**



Watch Video Solution

31. At the point of intersection of the rectangular hyperbola  $xy = c^2$  and the parabola  $y^2 = 4ax$  tangents to the rectangular hyperbola and the parabola make angles  $\theta$  and  $\phi$ , respectively with x-axis, then

A.  $\theta = \tan^{-1}(-2 \tan \phi)$

B.  $\theta = \frac{1}{2} \tan^{-1}(-\tan \phi)$

C.  $\phi = \tan^{-1}(-2 \tan \theta)$

D.  $\phi = \frac{1}{2} \tan^{-1}(-\tan \theta)$

Answer: A



Watch Video Solution

32. The number of points from where a pair of perpendicular tangents can be drawn to the hyperbola,  $x^2 \sec^2 \alpha - y^2 \cos^2 \alpha = 1$ ,  $\alpha \in \left(0, \frac{\pi}{4}\right)$ , is

A. 0

B. 1

C. 2

D. infinite

**Answer: D**



**Watch Video Solution**

33. If  $e$  is the eccentricity of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\theta$  is the angle between the asymptotes, then  $\cos. \frac{\theta}{2}$  is equal to

A.  $\frac{1 - e}{e}$

B.  $\frac{2}{e} - e$

C.  $\frac{1}{e}$

D.  $\frac{2}{e}$

**Answer: C**



Watch Video Solution

34. The equation of a hyperbola whose asymptotes are  $3x \pm 5y = 0$  and vertices are  $(\pm 5, 0)$  is

A.  $9x^2 - 25y^2 = 225$

B.  $25x^2 - 9y^2 = 225$

C.  $5x^2 - 3y^2 = 225$

D.  $3x^2 - 5y^2 = 25$

Answer: A



Watch Video Solution

35. The tangent at P on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets one of the asymptote in Q. Then the locus of the mid-point of PQ is

A.  $3\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) = 4$

$$\text{B. } \frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$$

$$\text{C. } \frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{1}{2}$$

$$\text{D. } 4\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) = 3$$

**Answer: D**



**Watch Video Solution**

**36.** Find the locus of the foot of perpendicular from the centre upon any normal to line hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ .

$$\text{A. } (x^2 - + y^2)^2 \left(\frac{a^2}{x^2} + \frac{b^2}{y^2}\right) = (a^2 - b^2)^2$$

$$\text{B. } (x^2 + y^2)^2 \left(\frac{a^2}{x^2} - \frac{b^2}{y^2}\right) = (a^2 + b^2)^2$$

$$\text{C. } (x^2 + y^2)^2 \left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) = (a^2 + b^2)^2$$

**D.** None of these

**Answer: B**



**Watch Video Solution**

37. Let the transverse axis of a varying hyperbola be fixed with length of transverse axis being  $2a$ . Then the locus of the point of contact of any tangent drawn to it from a fixed point on conjugate axis is

- A. a parabola
- B. a circle
- C. an ellipse
- D. a hyperbola

**Answer: A**

 [Watch Video Solution](#)

38. The locus of the foot of the perpendicular from the centre of the hyperbola  $xy = c^2$  on a variable tangent is (A)  $(x^2 - y^2) = 4c^2xy$  (B)  $(x^2 + y^2)^2 = 2c^2xy$  (C)  $(x^2 + y^2) = 4c^2xy$  (D)  $(x^2 + y^2)^2 = 4c^2xy$

A.  $(x^2 - y^2)^2 = 4c^2xy$

B.  $(x^2 + y^2)^2 = 2c^2xy$

C.  $(x^2 - y^2)^2 = 2c^2xy$

D.  $(x^2 + y^2)^2 = 4c^2xy$

**Answer: D**



**Watch Video Solution**