



## MATHS

### BOOKS - CENGAGE MATHS (ENGLISH)

#### MATRICES

##### Example

1. If  $e^A$  is defined as  $e^A = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots = \frac{1}{2} \begin{bmatrix} f(x) & g(x) \\ g(x) & f(x) \end{bmatrix}$ , where  $A = \begin{bmatrix} x & x \\ x & x \end{bmatrix}$ ,  $0 < x < 1$  and  $I$  is identity matrix, then find the functions  $f(x)$  and  $g(x)$ .



Watch Video Solution

2. Prove that matrix  $\begin{bmatrix} \frac{b^2 - a^2}{a^2 + b^2} & \frac{-2ab}{a^2 + b^2} \\ \frac{-2ab}{a^2 + b^2} & \frac{a^2 - b^2}{a^2 + b^2} \end{bmatrix}$  is orthogonal.



 [Watch Video Solution](#)

3. If  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , where  $a, b, c$  and  $d$  are real numbers, then prove that  $A^2 - (a + d)A + (ad - bc)I = O$ . Hence or otherwise, prove that if  $A^3 = O$  then  $A^2 = O$

 [Watch Video Solution](#)

4. If  $A = ([a_{ij}])_{n \times n}$  is such that  $(a)_{ij} = \overline{a_{ji}}, \forall i, j$  and  $A^2 = O$ , then  
Statement 1: Matrix  $A$  null matrix.

Statement 2:  $|A| = 0$ .

 [Watch Video Solution](#)

5. Find the possible square roots of the two rowed unit matrix  $I$ .

 [Watch Video Solution](#)

6. Prove the orthogonal matrices of order two are of the form

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \text{ or } \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix}$$

 [Watch Video Solution](#)

7. Let  $A = \begin{bmatrix} \tan \frac{\pi}{3} & \sec \frac{2\pi}{3} \\ \cot \left( 2013 \frac{\pi}{3} \right) & \cos(2012\pi) \end{bmatrix}$  and  $P$  be a  $2 \times 2$  matrix such

that  $PP^T = I$ , where  $I$  is an identity matrix of order 2. If  $Q = PAP^T$

and  $R = [r_{ij}]_{2 \times 2} = P^T Q^8 P$ , then find  $r_{11}$ .

 [Watch Video Solution](#)

8. Consider,  $A = \begin{bmatrix} a & 2 & 1 \\ 0 & b & 0 \\ 0 & -3 & c \end{bmatrix}$ , where  $a$ ,  $b$  and  $c$  are the roots of the

equation  $x^3 - 3x^2 + 2x - 1 = 0$ . If matrix  $B$  is such that

$AB = BA$ ,  $A + B - 2I \neq O$  and  $A^2 - B^2 = 4I - 4B$ , then find the

value of  $\det. (B)$

 [Watch Video Solution](#)

9. If  $A$  and  $B$  are square matrices of order 3 such that  $|A| = 3$  and  $|B| = 2$ , then the value of  $|A^{-1}adj(B^{-1})adj(3A^{-1})|$  is equal to



[Watch Video Solution](#)

## Illustration

1. If a matrix has 28 elements, what are the possible orders it can have ?



[Watch Video Solution](#)

2. Construct a  $2 \times 2$  matrix, where

$$(i) a_{ij} = \frac{(i - 2j)^2}{2} \quad (ii) a_{ij} = | -2i + 3j |$$



[Watch Video Solution](#)

3. What is the maximum number of different elements required to form a symmetric matrix of order 12 ?

 [Watch Video Solution](#)

4. If a square matrix  $A$  of order three is defined  $A = [a_{ij}]$  where  $a_{ij} = \text{sgn}(i - j)$ , then prove that  $A$  is skew-symmetric matrix.

 [Watch Video Solution](#)

5. For what values of  $x$  and  $y$  are the following matrices equal ?

$$A = \begin{bmatrix} 2x + 1 & 3y \\ 0 & y^2 - 5y \end{bmatrix}, B = \begin{bmatrix} x + 3 & y^2 + 2 \\ 0 & -6 \end{bmatrix}$$

 [Watch Video Solution](#)

6. For  $\alpha, \beta, \gamma \in R$ , let

$$A = \begin{bmatrix} \alpha^2 & 6 & 8 \\ 3 & \beta^2 & 9 \\ 4 & 5 & \gamma^2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2\alpha & 3 & 5 \\ 2 & 2\beta & 6 \\ 1 & 4 & 2\gamma - 3 \end{bmatrix}$$

 [Watch Video Solution](#)

7. Find the values of  $x$  for which matrix  $\begin{bmatrix} 3 & -1+x & 2 \\ 3 & -1 & x+2 \\ x+3 & -1 & 2 \end{bmatrix}$  is singular.

 [Watch Video Solution](#)

8. If  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$  and  $B = \begin{bmatrix} -3 & -2 \\ 1 & -5 \\ 4 & 3 \end{bmatrix}$ , then find  $D = \begin{bmatrix} p & q \\ r & s \\ t & u \end{bmatrix}$  such that  $A + B - D = O$ .

 [Watch Video Solution](#)

9.  $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$  and  $A + A^T = I$ , find the value of  $\alpha$ .

 [Watch Video Solution](#)

10. Let  $A$  be a square matrix. Then prove that (i)  $A + A^T$  is a symmetric matrix, (ii)  $A - A^T$  is a skew-symmetric matrix and (iii)  $\forall^T$  and  $A^T A$  are symmetric matrices.

 [Watch Video Solution](#)

11. If  $A = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 4 \\ 7 & 2 \end{bmatrix}$ , find  $3A - 2B$ .

 [Watch Video Solution](#)

12. Find non-zero values of  $x$  satisfying the matrix equation:

$$x \begin{bmatrix} 2x & 2 \\ 3 & x \end{bmatrix} + 2 \begin{bmatrix} 8 & 5x \\ 4 & 4x \end{bmatrix} = 2 \begin{bmatrix} x^2 + 8 & 24 \\ 10 & 6x \end{bmatrix}$$

 [Watch Video Solution](#)

13. Let  $A + 2B = \begin{bmatrix} 1 & 2 & 0 \\ 6 & -3 & 3 \\ -5 & 3 & 1 \end{bmatrix}$  and  $2A - B = \begin{bmatrix} 2 & -1 & 5 \\ 2 & -1 & 6 \\ 0 & 1 & 2 \end{bmatrix}$ , then

find  $tr(A) - tr(B)$ .



[Watch Video Solution](#)

14. If  $\begin{bmatrix} \lambda^2 - 2\lambda + 1 & \lambda - 2 \\ 1 - \lambda^2 + 3\lambda & 1 - \lambda^2 \end{bmatrix} = A\lambda^2 + B\lambda + C$ , where A, B and C are

matrices then find matrices B and C.



[Watch Video Solution](#)

15. Prove that every square matrix can be uniquely expressed as the sum of a symmetric matrix and a skew-symmetric matrix.



[Watch Video Solution](#)



16. Matrix A has  $m$  rows and  $n + 5$  columns; matrix B has  $m$  rows and  $11 - n$  columns. If both  $AB$  and  $BA$  exist, then (A)  $AB$  and  $BA$  are square matrix (B)  $AB$  and  $BA$  are of order  $8 \times 8$  and  $3 \times 13$ , respectively (C)  $AB = BA$  (D) None of these

 [Watch Video Solution](#)

17. If  $A = \begin{bmatrix} 2 & 3 & -1 \\ 1 & 4 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} 2 & 3 \\ 4 & 5 \\ 2 & 1 \end{bmatrix}$  then  $AB$  and  $BA$  are defined and equal.

 [Watch Video Solution](#)

18. Find the value of  $x$  and  $y$  that satisfy the equations

$$\begin{bmatrix} 3 & -2 \\ 3 & 0 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} y & y \\ x & x \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 3y & 3y \\ 10 & 10 \end{bmatrix}$$

 [Watch Video Solution](#)

19. Find the values of  $x$ ,  $y$ ,  $z$  if the matrix  $A = [0 \ 2yz \ xy - zx - yz]$  satisfy the equation  $A^T A = I_3$ .

 [Watch Video Solution](#)

20. If  $A = [\cos \theta \ \sin \theta - \sin \theta \ \cos \theta]$ , then prove that  $A^n = [\cos n\theta \ \sin n\theta - \sin n\theta \ \cos n\theta]$ ,  $n \in N$ .

 [Watch Video Solution](#)

21. If  $A = \begin{pmatrix} p & q \\ 0 & 1 \end{pmatrix}$ , then show that  $A^8 = \begin{pmatrix} p^8 & q\left(\frac{p^8-1}{p-1}\right) \\ 0 & 1 \end{pmatrix}$

 [Watch Video Solution](#)

22. Let  $A = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix}$  be a matrix. If  $A^{10} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  then prove that  $a + d$  is divisible by 13.

 [Watch Video Solution](#)

23. Show that the solutions of the equation

$$\begin{bmatrix} x & y \\ z & t \end{bmatrix}^2 = 0 \text{ are } \begin{bmatrix} x & y \\ z & t \end{bmatrix} = \begin{bmatrix} \pm\sqrt{\alpha\beta} & -\beta \\ \alpha & \pm\sqrt{\alpha\beta} \end{bmatrix}, \text{ where } \alpha, \beta \text{ are}$$

arbitrary.



[Watch Video Solution](#)

24. Let  $A$  be square matrix. Then prove that  $AA^T$  and  $A^T A$  are symmetric matrices.



[Watch Video Solution](#)

25. If  $A, B$  are square matrices of same order and  $B$  is a skewsymmetric matrix, show that  $A^T B A$  is skew-symmetric.



[Watch Video Solution](#)

26. If  $A$  and  $B$  are square matrices of same order such that  $AB + BA = O$ , then prove that  $A^3 - B^3 = (A + B)(A^2 - AB - B^2)$ .

 [Watch Video Solution](#)

27. Let  $A = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$ . If  $A^6 = kA - 205I$  then the numerical quantity of  $k - 40$  should be

 [Watch Video Solution](#)

28. Let  $A, B, C, D$  be (not necessarily square) real matrices such that  $A^T = BCD; B^T = CDA; C^T = DAB$  and  $D^T = ABC$ . For the matrix  $S = ABCD$ , consider the two statements. I.  $S^3 = S$  II.  $S^2 = S^4$   
(A) II is true but not I (B) I is true but not II (C) both I and II are true (D) both I and II are false

 [Watch Video Solution](#)

29. If  $A$  and  $B$  are square matrices of the same order such that  $AB = BA$ , then prove by induction that  $AB^n = B^n A$ . Further, prove that  $(AB)^n = A^n B^n$  for all  $n \in \mathbb{N}$ .

 [Watch Video Solution](#)

30. If  $A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -2 & 0 \end{bmatrix}$ , then prove that  $A^2 + 3A + 2I = O$ . Hence, find  $B$  and  $C$  matrices of order 2 with integer elements, if  $A = B^3 + C^3$ .

 [Watch Video Solution](#)

31. If  $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$  then find  $\text{tr.}(A^{2012})$ .

 [Watch Video Solution](#)

32. If  $A$  is a nonsingular matrix satisfying  $AB - BA = A$ , then prove that  $\det.(B + I) = \det.(B - I)$ .

 [Watch Video Solution](#)

[Watch Video Solution](#)

33. If  $\det(A - B) \neq 0$ ,  $A^4 = B^4$ ,  $C^3A = C^3B$  and  $B^3A = A^3B$ , then find the value of  $\det(A^3 + B^3 + C^3)$ .

 [Watch Video Solution](#)

34. Given a matrix  $A = [abc|bcacab]$ , where  $a, b, c$  are real positive numbers  $abc = 1$  and  $A^T A = I$ , then find the value of  $a^3 + b^3 + c^3$ .

 [Watch Video Solution](#)

35. If  $M$  is a  $3 \times 3$  matrix, where  $\det M = 1$  and  $MM^T = I$ , where  $I$  is an identity matrix, prove that  $\det(M - I) = 0$ .

 [Watch Video Solution](#)

36. Consider point  $P(x, y)$  in first quadrant. Its reflection about x-axis is

$Q(x_1, y_1)$ . So,  $x_1 = x$  and  $y_1 = -y$ .

This may be written as : 
$$\begin{cases} x_1 = 1 \cdot x + 0 \cdot y \\ y_1 = 0 \cdot x + (-1)y \end{cases}$$

This system of equations can be put in the matrix as :

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Here, matrix  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$  is the matrix of reflection about x-axis. Then find the matrix of reflection about the line  $y = x$ .



[Watch Video Solution](#)

37. If  $A = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$  then A is 1) an idempotent matrix 2)

nilpotent matrix 3) involutory 4) orthogonal matrix



[Watch Video Solution](#)

38. If  $A = \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}$  then find  $A^{14} + 3A - 2I$



Watch Video Solution

39. The matrix  $A = \begin{bmatrix} -5 & -8 & 3 & 5 & 0 & 1 & 2 & - \end{bmatrix}$  is a. idempotent matrix b. involutory matrix c. nilpotent matrix d. none of these

A. idempotent matrix

B. involutory matrix

C. nilpotent matrix

D. none of these

**Answer: involutory matrix**



Watch Video Solution

40. If  $abc = p$  and  $A = \begin{bmatrix} a & b & c \\ c & a & b \\ b & c & a \end{bmatrix}$ , prove that A is orthogonal if and

only if a, b, c are the roots of the equation  $x^3 \pm x^2 - p = 0$ .



Watch Video Solution



41. Let  $A$  be an orthogonal matrix, and  $B$  is a matrix such that  $AB = BA$ , then show that  $AB^T = B^T A$ .



Watch Video Solution

42. Find the adjoint of the matrix  $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 1 & -3 \\ -1 & 2 & 3 \end{bmatrix}$ .



Watch Video Solution

43. If  $S = \begin{bmatrix} \frac{\sqrt{3}-1}{2\sqrt{2}} & \frac{\sqrt{3}+1}{2\sqrt{2}} \\ -\left(\frac{\sqrt{3}+1}{2\sqrt{2}}\right) & \frac{\sqrt{3}-1}{2\sqrt{2}} \end{bmatrix}$ ,  $A = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$  and  $P = S(\text{adj.}A)S^T$ ,

then find matrix  $S^T P^{10} S$ .



Watch Video Solution

44. If  $A$  is a square matrix such that  $A(\text{adj}A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$ , then

$$= \frac{|\text{adj}(\text{adj}A)|}{2|\text{adj}A|}$$
 is equal to

 [Watch Video Solution](#)

45. Let  $A$  be a square matrix of order 3 such that

$$\text{adj.}(\text{adj.}(\text{adj.}A)) = \begin{bmatrix} 16 & 0 & -24 \\ 0 & 4 & 0 \\ 0 & 12 & 4 \end{bmatrix}. \text{ Then find}$$

(i)  $|A|$  (ii)  $\text{adj.}A$

 [Watch Video Solution](#)

46. Let  $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}$  and  $10B = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix}$ . If  $B$  is the

inverse of  $A$ , then  $\alpha$  is :

 [Watch Video Solution](#)

47. Matrices  $A$  and  $B$  satisfy  $AB = B^{-1}$ , where  $B = \begin{bmatrix} 2 & -1 \\ 2 & 0 \end{bmatrix}$ . Find

(i) without finding  $B^{-1}$ , the value of  $K$  for which

$$KA - 2B^{-1} + I = O.$$

(ii) without finding  $A^{-1}$ , the matrix  $X$  satisfying  $A^{-1}XA = B$ .

 [Watch Video Solution](#)

48. Given the matrices  $A$  and  $B$  as  $A = \begin{bmatrix} 1 & -1 \\ 4 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix}$ .

The two matrices  $X$  and  $Y$  are such that  $XA = B$  and  $AY = B$ , then find the matrix  $3(X + Y)$

 [Watch Video Solution](#)

49. If  $M$  is the matrix  $\begin{bmatrix} 1 & -3 \\ -1 & 1 \end{bmatrix}$  then find matrix  $\sum_{r=0}^{\infty} \left(\frac{-1}{3}\right)^r M^{r+1}$

 [Watch Video Solution](#)

50. Let  $p$  be a non singular matrix, and  $I + P + p^2 + \dots + p^n = 0$ , then find  $p^{-1}$ .



Watch Video Solution

51. If  $A$  and  $B$  are square matrices of same order such that  $AB = O$  and  $B \neq O$ , then prove that  $|A| = 0$ .



Watch Video Solution

52. If  $A$  is a symmetric matrix,  $B$  is a skew-symmetric matrix,  $A + B$  is nonsingular and  $C = (A + B)^{-1}(A - B)$ , then prove that

(i)  $C^T(A + B)C = A + B$  (ii)  $C^T(A - B)C = A - B$

(iii)  $C^T AC = A$



Watch Video Solution

53. If the matrices, A, B and  $(A + B)$  are non-singular, then prove that

$$\left[ A(A + B)^{-1}B \right]^{-1} = B^{-1} + A^{-1}.$$

 [Watch Video Solution](#)

54. If matrix  $A$  satisfies the equation  $A^2 = A^{-1}$ , then prove that

$$A^{2^n} = A^{2^{(n-2)}}, n \in \mathbb{N}.$$

 [Watch Video Solution](#)

55. If A and B are non-singular symmetric matrices such that  $AB = BA$ , then prove that  $A^{-1}B^{-1}$  is symmetric matrix.

 [Watch Video Solution](#)

56. If A is a matrix of order n such that  $A^T A = I$  and X is any matrix such that  $X = (A + I)^{-1}(A - I)$ , then show that X is skew symmetric

matrix.

 [Watch Video Solution](#)

57. Show that two matrices

$$A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 1 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 & 0 & 1 \\ 0 & 3 & 1 \end{bmatrix} \text{ are row equivalent.}$$

 [Watch Video Solution](#)

58. Using elementary transformations, find the inverse of the matrix :

$$(20 - 1510013)$$

 [Watch Video Solution](#)

59. Let  $a$  be a  $3 \times 3$  matrix such that

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 3 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \text{ then find } A^{-1}.$$

 [Watch Video Solution](#)

60. Solve the following system of equations, using matrix method.

$$x + 2y + z = 7, x + 3z = 11, 2x - 3y = 1$$

 [Watch Video Solution](#)

61. Using matrix method, show that following system of equation is

inconsistent :  $2x + 3y - z + 4 = 0$        $x - y + 2z - 7 = 0$

$$x + 4y - 3z + 5 = 0$$

 [Watch Video Solution](#)

62. about to only mathematics

 [Watch Video Solution](#)

63. Find the characteristic roots of the two-rowed orthogonal matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \text{ and verify that they are of unit modulus.}$$

 [Watch Video Solution](#)

64. Show that if  $\lambda_1, \lambda_2, \dots, \lambda_n$  are  $n$  eigenvalues of a square matrix  $A$  of order  $n$ , then the eigenvalues of the matrix  $A^2$  are  $\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2$ .

 [Watch Video Solution](#)

65. If  $A$  is nonsingular, prove that the eigenvalues of  $A^{-1}$  are the reciprocals of the eigenvalues of  $A$ .

 [Watch Video Solution](#)



66. If one of the eigenvalues of a square matrix a order  $3 \times 3$  is zero, then prove that  $\det A = 0$ .

 [Watch Video Solution](#)

Case 13 1

1. Construct a  $3 \times 4$  matrix, whose elements are given by:

$$a_{ij} = \frac{1}{2} | -3i + j |$$

 [Watch Video Solution](#)

2. Find the value of  $a$  if  $[a - b2a + c2a - b3c + d] = [-15013]$

 [Watch Video Solution](#)

3. Find the number of all possible matrices of order  $3 \times 3$  with each entry 0 or 1. How many of these are symmetric ?

 [Watch Video Solution](#)

4. Find the value of  $x$  for which the matrix  $A = \begin{bmatrix} 2/x & -1 & 2 \\ 1 & x & 2x^2 \\ 1 & 1/x & 2 \end{bmatrix}$  is singular.

 [Watch Video Solution](#)

5. If matrix  $A$  is skew-symmetric matrix of odd order, then show that  $\text{tr. } A = \det. A$ .

 [Watch Video Solution](#)

1. Solve for  $x$  and  $y$ ,  $x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} 3 \\ 5 \end{bmatrix} + \begin{bmatrix} -8 \\ -11 \end{bmatrix} = 0$ .

 [Watch Video Solution](#)

2. If  $A = \begin{bmatrix} 1 & 5 \\ 7 & 12 \end{bmatrix}$  and  $B = \begin{bmatrix} 9 & 1 \\ 7 & 8 \end{bmatrix}$  then find a matrix  $C$  such that  $3A + 5B + 2C$  is a null matrix.

 [Watch Video Solution](#)

3. Solve the following equations for  $X$  and  $Y$  :

$$2X - Y = \begin{bmatrix} 3 & -3 & 0 \\ 3 & 3 & 2 \end{bmatrix}, 2Y + X = \begin{bmatrix} 4 & 1 & 5 \\ -1 & 4 & -4 \end{bmatrix}$$

 [Watch Video Solution](#)

4. If  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 2 & 2 & 3 \end{bmatrix}$   $B = \begin{bmatrix} 1 & 2 & 2 \\ -2 & -1 & -2 \\ 2 & 2 & 3 \end{bmatrix}$  and  $C = \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & 2 \\ 2 & 2 & 3 \end{bmatrix}$  then find the value of  $\text{tr.}(A + B^T + 3C)$ .

 [Watch Video Solution](#)

5. If  $A = \begin{bmatrix} 3 & -2 \\ 4 & -1 \end{bmatrix}$ , then find all the possible values of  $\lambda$  such that the matrix  $(A - \lambda I)$  is singular.

 [Watch Video Solution](#)

6. If matrix  $A = \begin{bmatrix} 0 & 1 & -1 \\ 4 & -3 & 4 \\ 3 & -3 & 4 \end{bmatrix} = B + C$ , where  $B$  is symmetric matrix and  $C$  is skew-symmetric matrix, then find matrices  $B$  and  $C$ .

 [Watch Video Solution](#)

### Case 13.3

1. Consider the matrices

$$A = \begin{bmatrix} 4 & 6 & -1 \\ 3 & 0 & 2 \\ 1 & -2 & 5 \end{bmatrix}, B = \begin{bmatrix} 2 & 4 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}, C = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

Out of the given matrix products, which one is not defined ?

A.  $(AB)^T C$

B.  $C^T C (AB)^T$

C.  $C^T AB$

D.  $A^T A B B^T C$

**Answer: B**

 [Watch Video Solution](#)

2. Let  $A = BB^T + CC^T$ , where  $B = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ ,  $C = \begin{bmatrix} \sin \theta \\ -\cos \theta \end{bmatrix}$ ,  $\theta \in R$ .

Then prove that  $A$  is unit matrix.

 [Watch Video Solution](#)

3. The matrix  $R(t)$  is defined by  $R(t) = \begin{bmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{bmatrix}$ . Show that

$$R(s)R(t) = R(s + t).$$

 [Watch Video Solution](#)

4. if  $A = \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix}$  where  $i = \sqrt{-1}$  and  $x \in \mathbb{N}$  then  $A^{4x}$  equals to:

 [Watch Video Solution](#)

5. if  $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$  prove that  $A^k = \begin{bmatrix} 1 + 2k & -4k \\ k & 1 - 2k \end{bmatrix}$  where  $k$  is any positive integer.

 [Watch Video Solution](#)

6. if  $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$  and  $X$  is a matrix such that  $A = BX$ , then  $X =$

 [Watch Video Solution](#)

7. for what values of x:

$$\begin{bmatrix} 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 \\ 2 & 0 & 1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \\ x \end{bmatrix} = 0?$$



Watch Video Solution

8. Find the matrix X so that  $X[123456] = [-7 - 8 - 9246]$



Watch Video Solution

9. If  $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ , then  $\lim_{x \rightarrow \infty} \frac{1}{n} A^n$  is

A. (A) an identity matrix

B. (B)  $[0 \ 10 \ -1 \ 0]$

C. (C) a null matrix

D. (D) none of these

**Answer: Zero Matrix**



Watch Video Solution

10.  $A = \begin{bmatrix} 3 & a & -1 \\ 2 & 5 & c \\ b & 8 & 2 \end{bmatrix}$  is symmetric and  $B = \begin{bmatrix} d & 3 & a \\ b - a & e & -2b - c \\ -2 & 6 & -f \end{bmatrix}$  is

skew-symmetric, then find  $AB$ .



Watch Video Solution

Cae 13 4

1. If  $A$  and  $B$  are matrices of the same order, then  $AB^T - B^T A$  is a (a) skew-symmetric matrix (b) null matrix (c) unit matrix (d) symmetric matrix



Watch Video Solution

2. If  $A$  and  $B$  are square matrices such that  $AB = BA$  then prove that  $A^3 - B^3 = (A - B)(A^2 + AB + B^2)$ .



Watch Video Solution



3. If  $A$  is a square matrix such that  $A^2 = I$ , then

$(A - I)^3 + (A + I)^3 - 7A$  is equal to

 [Watch Video Solution](#)

4. If  $B, C$  are square matrices of order  $n$  and if  $A = B + C, BC = CB, C^2 = O$ , then without using mathematical induction, show that for any positive integer  $p$ ,  $A^{p+1} = B^p[B + (p + 1)C]$ .

 [Watch Video Solution](#)

5. Let  $A$  be any  $3 \times 2$  matrix. Then prove that  $\det. (AA^T) = 0$ .

 [Watch Video Solution](#)

6. Let  $A$  be a matrix of order 3, such that  $A^T A = I$ . Then find the value of  $\det. (A^2 - I)$ .

 [Watch Video Solution](#)

7.  $A$  and  $B$  are different matrices of order  $n$  satisfying  $A^3 = B^3$  and  $A^2 B = B^2 A$ . If  $\det. (A - B) \neq 0$ , then find the value of  $\det. (A^2 + B^2)$ .

 [Watch Video Solution](#)

8. If  $D = \text{diag}[d_1, d_2, d_n]$ , then prove that  $f(D) = \text{diag}[f(d_1), f(d_2), , f(d_n)]$ , where  $f(x)$  is a polynomial with scalar coefficient.

 [Watch Video Solution](#)

9. Point  $P(x, y)$  is rotated by an angle  $\theta$  in anticlockwise direction. The new position of point  $P$  is  $Q(x_1, y_1)$ . If  $\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = A \begin{bmatrix} x \\ y \end{bmatrix}$ , then find matrix  $A$ .

 [Watch Video Solution](#)

10. How many different diagonal matrices of order  $n$  can be formed which are involutory ?

 [Watch Video Solution](#)

11. How many different diagonal matrices of order  $n$  can be formed which are involutory ?

 [Watch Video Solution](#)

12. If  $A$  and  $B$  are  $n$ -rowed unitary matrices, then  $AB$  and  $BA$  are also unitary matrices.

 Watch Video Solution

Cae 13 5

1. By the method of matrix inversion, solve the system.

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 5 & 7 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \\ x_3 & y_3 \end{bmatrix} = \begin{bmatrix} 9 & 2 \\ 52 & 15 \\ 0 & -1 \end{bmatrix}$$

 Watch Video Solution

2. Let  $A = \begin{bmatrix} 2 & 0 & 7 \\ 0 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} -x & 14x & 7x \\ 0 & 1 & 0 \\ x & -4x & -2x \end{bmatrix}$  are two

matrices such that  $AB = (AB)^{-1}$  and  $AB \neq I$  then

$$\text{Tr}\left((AB) + (AB)^2 + (AB)^3 + (AB)^4 + (AB)^5 + (AB)^6\right) =$$

 Watch Video Solution

3. Find  $A^{-1}$  if  $A = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{vmatrix}$  and show that  $A^{-1} = \frac{A^2 - 3I}{2}$

 [Watch Video Solution](#)

4. For the matrix  $A = [3175]$ , find  $x$  and  $y$  so that  $A^2 + xI = yA$ .

 [Watch Video Solution](#)

5. If  $A^3 = O$ , then prove that  $(I - A)^{-1} = I + A + A^2$ .

 [Watch Video Solution](#)

6. If  $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ ,  $B = \begin{bmatrix} \cos 2\beta & \sin 2\beta \\ \sin \beta & -\cos \beta \end{bmatrix}$  where

$0 < \beta < \frac{\pi}{2}$ , then prove that  $BAB = A^{-1}$ . Also, find the least

value of  $\alpha$  of which  $BA^4B = A^{-1}$

 [Watch Video Solution](#)

7. If  $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 2 & 3 \\ 1 & -1 & 3 \end{bmatrix}$ ,  $C = \begin{bmatrix} 2 & 1 & 1 \\ 2 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ ,  $D = \begin{bmatrix} 10 \\ 13 \\ 9 \end{bmatrix}$ , and  $CB = D$ .

Solve the equation  $AX = B$ .

 [Watch Video Solution](#)

8. If  $A$  is a  $2 \times 2$  matrix such that  $A^2 - 4A + 3I = O$ , then prove that

$$(A + 3I)^{-1} = \frac{7}{24}I - \frac{1}{24}A.$$

 [Watch Video Solution](#)

9. For two unimobular complex numbers  $z_1$  and  $z_2$ , find

$$\begin{bmatrix} \bar{z}_1 & -z_2 \\ \bar{z}_2 & z_1 \end{bmatrix}^{-1} \begin{bmatrix} z_1 & z_2 \\ -\bar{z}_2 & \bar{z}_1 \end{bmatrix}^{-1}$$

 [Watch Video Solution](#)

10. Prove that inverse of a skew-symmetric matrix (if it exists) is skew-symmetric.

 [Watch Video Solution](#)

11. If square matrix  $A$  is orthogonal, then prove that its inverse is also orthogonal.

 [Watch Video Solution](#)

12. If  $A$  is a skew symmetric matrix, then  $B = (I - A)(I + A)^{-1}$  is (where  $I$  is an identity matrix of same order as of  $A$ )

 [Watch Video Solution](#)

13. Prove that  $(\text{adj. } A)^{-1} = (\text{adj. } A^{-1})$ .

 [Watch Video Solution](#)

14. Using elementary transformation, find the inverse of the matrix

$$A = \begin{bmatrix} a & b \\ c & \left(\frac{1+bc}{a}\right) \end{bmatrix}.$$

 [Watch Video Solution](#)

15. If  $A$  and  $P$  are the square matrices of the same order and if  $P$  be invertible, show that the matrices  $A$  and  $P^{-1}AP$  have the same characteristic roots.

 [Watch Video Solution](#)

16. Show that the characteristics roots of an idempotent matrix are either 0 or 1

 [Watch Video Solution](#)



17. If  $\alpha$  is a characteristic root of a nonsingular matrix, then prove that  $\frac{|A|}{\alpha}$  is a characteristic root of  $\text{adj } A$ .



[Watch Video Solution](#)

## Exercises

1. If  $A$  is symmetric as well as skew-symmetric matrix, then  $A$  is

- A. diagonal matrix
- B. null matrix
- C. triangular matrix
- D. none of these

**Answer: B**



[Watch Video Solution](#)

2. Elements of a matrix A of order  $10 \times 10$  are defined as  $a_{ij} = \omega^{i+j}$

(where  $\omega$  is cube root unity), then  $\text{tr}(A)$  of matrix is

A. 0

B. 1

C. 3

D. none of these

**Answer: D**



[Watch Video Solution](#)

3. If  $A_1, A_2, \dots, A_{2n-1}$  are skew-symmetric matrices of same order, then

$B = \sum_{r=1}^n (2r - 1) (A^{2r-1})^{2r-1}$  will be i) symmetric ii) skew-symmetric iii)

neither symmetric nor skew-symmetric iv) data not adequate

A. symmetric

B. skew-symmetric

C. neither symmetric nor skew-symmetric

D. data not adequate

**Answer: B**



**Watch Video Solution**

4. The equation  $[1xy] \begin{bmatrix} 1 & 3 & 1 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ x \\ y \end{bmatrix} = [0]$  has

i)  $y=0$  (p) rational roots

ii)  $y=-1$  (q) irrational roots

(r) integral roots

- A. (i) (ii)  
(p) (r)
- B. (i) (ii)  
(q) (p)
- C. (i) (ii)  
(p) (q)
- D. (i) (ii)  
(r) (p)

**Answer: C**



Watch Video Solution

5. Let  $A$  and  $B$  be two  $2 \times 2$  matrices. Consider the statements (i)  $AB = O \Rightarrow A = O$  or  $B = O$  (ii)  $AB = I_2 \Rightarrow A = B^{-1}$  (iii)  $(A + B)^2 = A^2 + 2AB + B^2$  (a)(i) and (ii) are false, (iii) is true (b)(ii) and (iii) are false, (i) is true (c)(i) is false (ii) and, (iii) are true (d)(i) and (iii) are false, (ii) is true

- A. (i) and (ii) are false, (iii) is true
- B. (ii) and (iii) are false, (i) is true
- C. (i) is false, (ii) and (iii) are true
- D. (i) and (iii) are false, (ii) is true

**Answer: D**



Watch Video Solution

6. The number of diagonal matrix,  $A$  of order  $n$  which  $A^3 = A$  is a. 1  
b. 0 c.  $2^n$  d.  $3^n$

A. 1

B. 0

C.  $2^n$

D.  $3^n$

**Answer: D**



[Watch Video Solution](#)

7.  $A$  is a  $2 \times 2$  matrix such that  $A \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$  and  $A^2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . The sum of the elements of  $A$  is a.  $-1$  b. 0 c. 2 d. 5

A.  $-1$

B. 0

C. 2

D. 5

**Answer: D**



**Watch Video Solution**

8. If  $\theta - \phi = \frac{\pi}{2}$ , prove that,

$$\begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix} \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix} = 0$$

A.  $2n\pi, n \in \mathbb{Z}$

B.  $n\frac{\pi}{2}, n \in \mathbb{Z}$

C.  $(2n + 1)\frac{\pi}{2}, n \in \mathbb{X}$

D.  $n\pi, n \in \mathbb{Z}$

**Answer: C**



**Watch Video Solution**

9. If  $A = \begin{bmatrix} a & b & 0 & a \end{bmatrix}$  is  $n$ th root of  $I_2$ , then choose the correct statements: If  $n$  is odd,  $a = 1, b = 0$  If  $n$  is odd,  $a = -1, b = 0$  If  $n$  is even,  $a = 1, b = 0$  If  $n$  is even,  $a = -1, b = 0$  a. i, ii, iii, iv b. ii, iii, iv c. i, ii, iii, iv d. i, iii, iv

A. i, ii, iii

B. ii, iii, iv

C. i, ii, iii, iv

D. i, iii, iv

**Answer: D**



**Watch Video Solution**

10. If  $\begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$  is to be square root of two-rowed unit matrix, then  $\alpha, \beta$  and  $\gamma$  should satisfy the relation. 1.  $1 - \alpha^2 + \beta\gamma = 0$  b.  $\alpha^2 + \beta\gamma = 0$  c.  $1 + \alpha^2 + \beta\gamma = 0$  d.  $1 - \alpha^2 - \beta\gamma = 0$

A.  $1 - \alpha^2 + \beta\gamma = 0$

B.  $\alpha^2 + \beta\gamma - 1 = 0$

C.  $1 + \alpha^2 + \beta\gamma = 0$

D.  $1 - \alpha^2 - \beta\gamma = 0$

**Answer: B**



**Watch Video Solution**

11. If  $A = \begin{bmatrix} i & -i \\ -i & i \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$ , then,  $A^8$  equals a.  $4B$  b.

$128B$  c.  $-128B$  d.  $-64B$

A.  $4B$

B.  $128B$

C.  $-128B$

D.  $-64B$

**Answer: B**



**Watch Video Solution**



12. If  $\begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 4 \end{bmatrix} A = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}$ , then sum of all the elements of matrix  $A$  is 0 b. 1 c. 2 d.  $-3$

A. 0

B. 1

C. 2

D.  $-3$

**Answer: B**



**Watch Video Solution**

13. For each real  $x$ ,  $-1 < x < 1$ . Let  $A(x)$  be the matrix  $(1-x)^{-1} \begin{bmatrix} 1 & -x \\ -x & 1 \end{bmatrix}$  and  $z = \frac{x+y}{1+xy}$ . Then

A.  $A(z) = A(x)A(y)$

B.  $A(z) = A(x) - A(y)$

C.  $A(z) = A(x) + A(y)$

D.  $A(z) = A(x)[A(y)]^{-1}$

**Answer: A**



**Watch Video Solution**

**14.** Let  $A = [0 - \tan(\alpha/2) \tan(\alpha/2) 0]$  and  $I$  be the identity matrix of order 2. Show that  $I + A = (I - A)[\cos \alpha - \sin \alpha \sin \alpha \cos \alpha]$ .

A.  $-I + A$

B.  $I - A$

C.  $-I - A$

D. none of these

**Answer: B**



**Watch Video Solution**

15. The number of solutions of the matrix equation  $X^2 = [1123]$  is a. more than 2 b. 2 c. 0 d. 1

A. more than 2

B. 2

C. 0

D. 1

**Answer: A**



**Watch Video Solution**

16. If  $A = [abcd]$  (where  $bc \neq 0$ ) satisfies the equations  $x^2 + k = 0$ , then  $a + d = 0$  b.  $k = -|A|$  c.  $k = |A|$  d. none of these

A.  $a + d = 0$

B.  $k = -|A|$

C.  $k = |A|$

D. none of these

**Answer: C**



**Watch Video Solution**

17.  $A = \begin{bmatrix} 2 & 1 \\ 4 & 1 \end{bmatrix}; B = \begin{bmatrix} 3 & 4 \\ 2 & 3 \end{bmatrix}$  &  $c = \begin{bmatrix} 3 & -4 \\ -2 & 3 \end{bmatrix}$ ,

$tr(A) + tr\left[\frac{ABC}{2}\right] + tr\left[\frac{A(BC)^2}{4}\right] + tr\left[\frac{A(BC)^2}{8}\right] + \dots \dots \infty$  is:

A. 6

B. 9

C. 12

D. none of these

**Answer: A**



**Watch Video Solution**

18. If  $\left[ \frac{\cos(2\pi)}{7} - \frac{\sin(2\pi)}{7} \frac{\sin(2\pi)}{7} \frac{\cos(2\pi)}{7} \right] = [1001]$ , then the least positive integral value of  $k$  is (a) 3 (b) 4 (c) 6 (d) 7

A. 3

B. 6

C. 7

D. 14

**Answer: C**



[Watch Video Solution](#)

19. If  $A$  and  $B$  are square matrices of order  $n$ , then prove that  $A$  and  $B$  will commute iff  $A - \lambda I$  and  $B - \lambda I$  commute for every scalar  $\lambda$ .

A.  $AB = BA$

B.  $AB + BA = O$

C.  $A = -B$

D. none of these

**Answer: A**

 [Watch Video Solution](#)

20. Matrix  $A$  such that  $A^2 = 2A - I$ , where  $I$  is the identity matrix, the for  $n \geq 2$ .  $A^n$  is equal to  $2^{n-1}A - (n-1)I$  b.  $2^{n-1}A - I$  c.  $nA - (n-1)I$  d.  $nA - I$

A.  $2^{n-1}A - (n-1)I$

B.  $2^{n-1}A - I$

C.  $nA - (n-1)I$

D.  $nA - I$

**Answer: C**

 [Watch Video Solution](#)

21. Let  $A = \begin{bmatrix} 0 & \alpha \\ 0 & 0 \end{bmatrix}$  and  $(A + I)^{50} = 50A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  Then the value of  $a + b + c + d$  is (A) 2 (B) 1 (C) 4 (D) none of these

A. 2

B. 1

C. 4

D. none of these

**Answer: A**



**Watch Video Solution**

22. If  $Z$  is an idempotent matrix, then  $(I + Z)^n$

A.  $I + 2^n Z$

B.  $I + (2^n - 1)Z$

C.  $I - (2^n - 1)Z$

D. none of these

**Answer: B**



[Watch Video Solution](#)

23. if  $A$  and  $B$  are square matrices such that  $A^{2006} = O$  and  $AB = A + B$ , then  $\det(B)$  equals 0 b. 1 c.  $-1$  d. none of these

A. 0

B. 1

C.  $-1$

D. none of these

**Answer: A**



[Watch Video Solution](#)



24. If matrix  $A$  is given by  $A = \begin{bmatrix} 6 & 11 \\ 2 & 4 \end{bmatrix}$  then determinant of  $A^{2005} - 6A^{2004}$  is

A.  $2^{2006}$

B.  $(-11)2^{2005}$

C.  $-2^{2005} \cdot 7$

D.  $(-9)2^{2004}$

**Answer: B**



[Watch Video Solution](#)

25. If  $A$  is a non-diagonal involutory matrix, then

A.  $A - I = O$

B.  $A + I = O$

C.  $A - I$  is nonzero singular

D. none of these

**Answer: C**



[Watch Video Solution](#)

26. If  $A$  and  $B$  are two nonzero square matrices of the same order such that the product  $AB = O$ , then

- A. both  $A$  and  $B$  must be singular
- B. exactly one of them must be singular
- C. both of them are nonsingular
- D. none of these

**Answer: A**



[Watch Video Solution](#)

27. If  $A$  and  $B$  are symmetric matrices of the same order and  $X = AB + BA$  and  $Y = AB - BA$ , then  $(XY)^T$  is equal to : (A)  $XY$   
(B)  $YX$  (C)  $-YX$  (D) non of these

A.  $XY$

B.  $YX$

C.  $-YX$

D. none of these

**Answer: C**



**Watch Video Solution**

28. If  $A, B, A + I, A + B$  are idempotent matrices, then  $AB$  is equal to  
 $BA$  b.  $-BA$  c.  $I$  d.  $O$

A.  $BA$

B.  $-BA$

C.  $I$

D.  $O$

**Answer: B**

 [Watch Video Solution](#)

29. If  $A = \begin{bmatrix} 0 & x \\ y & 0 \end{bmatrix}$  and  $A^3 + A = O$  then sum of possible values of  $xy$  is

A. 0

B.  $-1$

C. 1

D. 2

**Answer: B**

 [Watch Video Solution](#)

30. Which of the following is an orthogonal matrix? (a)  $\begin{bmatrix} 6/7 & 2/7 & -3/7 \\ 2/7 & 3/7 & 6/7 \\ 3/7 & -6/7 & 2/7 \end{bmatrix}$  (b)  $\begin{bmatrix} 6/7 & 2/7 & 3/7 \\ 2/7 & -3/7 & 6/7 \\ 3/7 & 6/7 & -2/7 \end{bmatrix}$  (c)  $\begin{bmatrix} -6/7 & -2/7 & -3/7 \\ 2/7 & 3/7 & 6/7 \\ 3/7 & 6/7 & -2/7 \end{bmatrix}$  (d)  $\begin{bmatrix} 6/7 & -2/7 & 3/7 \\ 2/7 & 2/7 & -3/7 \\ -6/7 & 2/7 & 3/7 \end{bmatrix}$

- A.  $\begin{bmatrix} 6/7 & 2/7 & -3/7 \\ 2/7 & 3/7 & 6/7 \\ 3/7 & -6/7 & 2/7 \end{bmatrix}$
- B.  $\begin{bmatrix} 6/7 & 2/7 & 3/7 \\ 2/7 & -3/7 & 6/7 \\ 3/7 & 6/7 & -2/7 \end{bmatrix}$
- C.  $\begin{bmatrix} -6/7 & -2/7 & -3/7 \\ 2/7 & 3/7 & 6/7 \\ -3/7 & 6/7 & 2/7 \end{bmatrix}$
- D.  $\begin{bmatrix} 6/7 & -2/7 & 3/7 \\ 2/7 & 2/7 & -3/7 \\ -6/7 & 2/7 & 3/7 \end{bmatrix}$

Answer: A



Watch Video Solution

31. Let A and B be two square matrices of the same size such that  $AB^T + BA^T = O$ . If A is a skew-symmetric matrix then BA is

- A. a symmetric matrix
- B. a skew-symmetric matrix
- C. an orthogonal matrix
- D. an invertible matrix

**Answer: B**



[Watch Video Solution](#)

32. In which of the following type of matrix inverse does not exist always?

a. idempotent b. orthogonal c. involuntary d. none of these

- A. idempotent
- B. orthogonal
- C. involuntary

D. none of these

**Answer: A**



[Watch Video Solution](#)

**33.** Let  $A$  be an  $n$ th-order square matrix and  $B$  be its adjoint, then  $|AB + KI_n|$  is (where  $K$  is a scalar quantity)  $(|A| + K)^{n-2}$  b.  $(|A| + K)^n$  c.  $(|A| + K)^{n-1}$  d. none of these

A.  $(|A| + K)^{n-2}$

B.  $(|A| + K)^n$

C.  $(|A| + K)^{n-1}$

D. none of these

**Answer: B**



[Watch Video Solution](#)

34. If  $A = \begin{bmatrix} a & b & c \\ x & y & z \\ p & q & r \end{bmatrix}$ ,  $B = \begin{bmatrix} q & -b & y \\ -p & a & -x \\ r & -c & z \end{bmatrix}$  and If A is invertible, then

which of the following is not true ?

A.  $|A| = |B|$

B.  $|A| = -|B|$

C.  $|\text{adj } A| = |\text{adj } B|$

D. A is invertible if and only if B is invertible

**Answer: A**



**Watch Video Solution**

35. If  $A(\alpha, \beta) = \begin{bmatrix} \cos \alpha & s \in \alpha & 0 \\ -s \in \alpha & \cos \alpha & 0 \\ 0 & 0 & e^\beta \end{bmatrix}$ , then  $A(\alpha, \beta)^{-1}$  is equal to  $A(-\alpha, -\beta)$  b.  $A(-\alpha, \beta)$  c.  $A(\alpha, -\beta)$  d.  $A(\alpha, \beta)$

A.  $A(-\alpha, -\beta)$

B.  $A(-\alpha, \beta)$

C.  $A(\alpha, -\beta)$



D.  $A(\alpha, \beta)$

**Answer: A**



**Watch Video Solution**

36. If  $A = \begin{bmatrix} a + ib & c + id \\ -c + id & a - ib \end{bmatrix}$  and  $a^2 + b^2 + c^2 + d^2 = 1$ , then  $A^{-1}$  is equal to

A.  $\begin{bmatrix} a - ib & -c - id \\ c - id & a + ib \end{bmatrix}$

B.  $\begin{bmatrix} a + ib & -c + id \\ -c + id & a - ib \end{bmatrix}$

C.  $\begin{bmatrix} a - ib & -c - id \\ -c - id & a + ib \end{bmatrix}$

D. none of these

**Answer: A**



**Watch Video Solution**

37. If  $[1/250x^{1/25}] = [50 - a5]^{-2}$ , then the value of  $x$  is  $a/125$  b.

$2a/125$  c.  $2a/25$  d. none of these

A.  $a/125$

B.  $2a/125$

C.  $2a/25$

D. none of these

**Answer: B**



**Watch Video Solution**

38. If  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$  and  $f(x) = \frac{1+x}{1-x}$ , then  $f(A)$  is

A.  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$

C.  $\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$

D. none of these

**Answer: C**



**Watch Video Solution**

**39.** There are two possible values of A in the solution of the matrix equation

$$\begin{bmatrix} 2A + 1 & -5 \\ -4 & A \end{bmatrix}^{-1} \begin{bmatrix} A - 5 & B \\ 2A - 2 & C \end{bmatrix} = \begin{bmatrix} 14 & D \\ E & F \end{bmatrix}$$

where A, B, C, D, E and F are real numbers. The absolute value of the difference of these two solutions, is

A.  $\frac{8}{3}$

B.  $\frac{19}{3}$

C.  $\frac{1}{3}$

D.  $\frac{11}{3}$

**Answer: B**



**Watch Video Solution**

40. If  $A$  and  $B$  are two square matrices such that  $B = -A^{-1}BA$ , then  $(A + B)^2$  is equal to

A.  $A^2 + B^2$

B.  $O$

C.  $A^2 + 2AB + B^2$

D.  $A + B$

Answer: A



Watch Video Solution

41. If  $A = [1 \tan x - \tan x 1]$ , show that

$$A^T A^{-1} = [\cos 2x - \sin 2x \sin 2x \cos 2x]$$

A.  $\begin{bmatrix} -\cos 2x & \sin 2x \\ -\sin 2x & \cos 2x \end{bmatrix}$

B.  $\begin{bmatrix} \cos 2x & -\sin 2x \\ \sin 2x & \cos 2x \end{bmatrix}$

C.  $\begin{bmatrix} \cos 2x & \cos 2x \\ \cos 2x & \sin 2x \end{bmatrix}$

D. none of these

**Answer: B**



[Watch Video Solution](#)

42. If  $A$  is order 3 square matrix such that  $|A| = 2$ , then  $|\text{adj}(\text{adj}(\text{adj } A))|$  is

A. 512

B. 256

C. 64

D. none of these

**Answer: B**



[Watch Video Solution](#)

43. If  $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & a & 1 \end{bmatrix}$  and  $A^{-1} = \begin{bmatrix} 1/2 & -1/2 & 1/2 \\ -4 & 3 & b \\ 5/2 & -3/2 & 1/2 \end{bmatrix}$  then

A. 1, 1

B. 1, -1

C. 1, 2

D. -1, 1

**Answer: B**



[Watch Video Solution](#)

44. If  $n$ th-order square matrix  $A$  is a orthogonal, then  $|\text{adj}(\text{adj} A)|$  is

A. always -1 if  $n$  is even

B. always 1 if  $n$  is odd

C. always 1

D. none of these

**Answer: B**



**Watch Video Solution**

45. Let  $a$  and  $b$  be two real numbers such that  $a > 1, b > 1$ . If  $A = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ , then  $(\lim)_{n \rightarrow \infty} A^{-n}$  is a. unit matrix b. null matrix c.  $2I$  d. none of these

A. unit matrix

B. null matrix

C.  $2I$

D. none of these

**Answer: B**



**Watch Video Solution**

46. If  $A = [a_{ij}]_{4 \times 4}$ , such that  $a_{ij} = \begin{cases} 2, & \text{when } i = j \\ 0, & \text{when } i \neq j \end{cases}$ , then  $\left\{ \frac{\det(\text{adj}(\text{adj} A))}{7} \right\}$  is (where  $\{ \cdot \}$  represents fractional part function)

A.  $1/7$

B.  $2/7$

C.  $3/7$

D. none of these

**Answer: A**



**Watch Video Solution**

47.  $A$  is an involutory matrix given by  $A = [01 - 14 - 343 - 34]$ , then

the inverse of  $A/2$  will be  $2A$  b.  $\frac{A^{-1}}{2}$  c.  $\frac{A}{2}$  d.  $A^2$

A.  $2A$

B.  $\frac{A^{-1}}{2}$



C.  $\frac{A}{2}$

D.  $A^2$

**Answer: A**



[Watch Video Solution](#)

48. If  $A$  is a nonsingular matrix such that  $AA^T = A^T A$  and  $B = A^{-1}A^T$ ,

then matrix  $B$  is

A. involuntary

B. orthogonal

C. idempotent

D. none of these

**Answer: B**



[Watch Video Solution](#)

49. If  $P$  is an orthogonal matrix and  $Q = PAP^T$  and  $x = P^T A b$ .  $I$  c.  $A^{1000}$  d. none of these

A.  $A$

B.  $I$

C.  $A^{1000}$

D. none of these

**Answer: B**



[Watch Video Solution](#)

50. If  $A$  and  $B$  are two non-singular matrices of the same order such that  $B^r = I$ , for some positive integer  $r > 1$ , then  $(A^{-1}B^{r-1}A) - (A^{-1}B^{-1}A) =$  a.  $I$  b.  $2I$  c.  $O$  d.  $-I$

A.  $I$

B.  $2I$

C.  $O$

D.  $-I$

**Answer: C**



[Watch Video Solution](#)

51. If  $\text{adj}B = A$ ,  $|P| = |Q| = 1$ , then  $\text{adj}(Q^{-1}BP^{-1})$  is  $PQ$  b.  $QAP$  c.  $PAQ$  d.  $PA^1Q$

A.  $PQ$

B.  $QAP$

C.  $PAQ$

D.  $PA^{-1}Q$

**Answer: C**



[Watch Video Solution](#)

52. If  $A$  is non-singular and  $(A - 2I)(A - 4I) = O$ , then  $\frac{1}{6}A + \frac{4}{3}A^{-1}$  is equal to  $O$  **I** **b.  $2I$**  **c.  $6I$**  **d.  $I$**

A.  $O$

B.  $I$

C.  $2I$

D.  $6I$

**Answer: B**



**Watch Video Solution**

53. Let  $f(x) = \frac{1+x}{1-x}$ . If  $A$  is matrix for which  $A^3 = 0$ , then  $f(A)$  is (a)  $I + A + A^2$  (b)  $I + 2A + 2A^2$  (c)  $I - A - A^2$  (d) none of these

A.  $I + A + A^2$

B.  $I + 2A + 2A^2$

C.  $I - A - A^2$

D. none of these

**Answer: B**



[Watch Video Solution](#)

**54.** Find the matrix A satisfying the matrix equation

$$\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

A.  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

B.  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$

C.  $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$

D.  $-\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

**Answer: A**



[Watch Video Solution](#)

55. If  $A^2 - A + I = 0$ , then the inverse of  $A$  is: (A)  $A + I$  (B)  $A$  (C)  $A - I$   
(D)  $I - A$

A.  $A^{-2}$

B.  $A + I$

C.  $I - A$

D.  $A - I$

**Answer: C**



**Watch Video Solution**

56. If  $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$  and  $G(y) = \begin{bmatrix} \cos y & 0 & \sin y \\ 0 & 1 & 0 \\ -\sin y & 0 & \cos y \end{bmatrix}$ ,

then  $[F(x)G(y)]^{-1}$  is equal to

A.  $F(-x)G(-y)$

B.  $G(-y)F(-x)$

C.  $F(x^{-1})G(y^{-1})$

D.  $G(y^{-1})F(x^{-1})$

**Answer: B**



**Watch Video Solution**

57. about to only mathematics

A.  $A^{-n}B^nA^n$

B.  $A^nB^nA^{-n}$

C.  $A^{-1}B^nA$

D.  $n(A^{-1}BA)$

**Answer: C**



**Watch Video Solution**

58. If  $k \in R$ , then  $\det\{\text{adj}(kI_n)\}$  is equal to  $K^{n-1}$  b.  $K^{n(n-1)}$  c.  $K^n$  d.  $k$

A.  $k^{n-1}$

B.  $k^{n(n-1)}$

C.  $k^n$

D.  $k$

**Answer: B**



**Watch Video Solution**

59. Given that matrix  $A = \begin{bmatrix} x & 3 & 2 \\ 1 & y & 4 \\ 2 & 2 & z \end{bmatrix}$ . If  $xyz = 60$  and  $8x + 4y + 3z = 20$ ,

then  $A(\text{adj } A)$  is equal to

A.  $\begin{bmatrix} 64 & 0 & 0 \\ 0 & 64 & 0 \\ 0 & 0 & 64 \end{bmatrix}$

B.  $\begin{bmatrix} 88 & 0 & 0 \\ 0 & 88 & 0 \\ 0 & 0 & 88 \end{bmatrix}$



- C.  $\begin{bmatrix} 68 & 0 & 0 \\ 0 & 68 & 0 \\ 0 & 0 & 68 \end{bmatrix}$
- D.  $\begin{bmatrix} 34 & 0 & 0 \\ 0 & 34 & 0 \\ 0 & 0 & 34 \end{bmatrix}$

**Answer: C**



**Watch Video Solution**

60. Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 0 & 5 \\ 0 & 2 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 \\ -3 \\ 1 \end{bmatrix}$ . Which of the following is a true? a.  $AX = B$  has a unique solution b.  $AX = B$  has exactly three solutions c.  $Ax = B$  has infinitely many solutions d.  $AX = B$  is inconsistent

- A.  $AX = B$  has a unique solution
- B.  $AX = B$  has exactly three solutions
- C.  $AX = B$  has infinitely many solutions
- D.  $AX = B$  is inconsistent

**Answer: A**



**Watch Video Solution**

61. If  $A$  is a square matrix of order less than 4 such that  $|A - A^T| \neq 0$  and  $B = \text{adj.}(A)$ , then  $\text{adj.}(B^2 A^{-1} B^{-1} A)$  is

A.  $A$

B.  $B$

C.  $|A|A$

D.  $|B|B$

**Answer: A**



**Watch Video Solution**

62. Let  $A$  be a square matrix of order 3 such that  $\det.(A) = \frac{1}{3}$ , then the value of  $\det.(\text{adj. } A^{-1})$  is

A.  $1/9$

B.  $1/3$

C. 3

D. 9

**Answer: D**



**Watch Video Solution**

**63.** If A and B are two non-singular matrices of order 3 such that

$AA^T = 2I$  and  $A^{-1} = A^T - A \cdot \text{Adj.}(2B^{-1})$ , then  $\det.(B)$  is equal to

A. 4

B.  $4\sqrt{2}$

C. 16

D.  $16\sqrt{2}$

**Answer: D**

 [Watch Video Solution](#)

64. If  $A$  is a square matrix of order 5 and  $2A^{-1} = A^T$ , then the remainder when  $|\text{adj.}(\text{adj.}(\text{adj.} A))|$  is divided by 7 is

A. 2

B. 3

C. 4

D. 5

**Answer: A**

 [Watch Video Solution](#)

65. Let  $P = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 3 & 1 & 1 \end{bmatrix}$ . If the product  $PQ$  has inverse  $R = \begin{bmatrix} -1 & 0 & 1 \\ 1 & 1 & 3 \\ 2 & 0 & 2 \end{bmatrix}$  then  $Q^{-1}$  equals

- A.  $\begin{bmatrix} 3 & 2 & 9 \\ -1 & 1 & 1 \\ 0 & 1 & 8 \end{bmatrix}$
- B.  $\begin{bmatrix} 5 & 2 & 9 \\ -1 & 1 & 1 \\ 0 & 1 & 7 \end{bmatrix}$
- C.  $\begin{bmatrix} 2 & -1 & 0 \\ 10 & 6 & 3 \\ 8 & 6 & 4 \end{bmatrix}$

D. none of these

**Answer: C**



[Watch Video Solution](#)

### Multiple Correct Answer

1. If  $A$  is unimodular, then which of the following is unimodular? –  $A$  b.  $A^{-1}$  c.  $adjA$  d.  $\omega A$ , where  $\omega$  is cube root of unity

A.  $-A$

B.  $A^{-1}$

C.  $\text{adj } A$

D.  $\omega A$ , where  $\omega$  is cube root of unity

**Answer: B::C**



**Watch Video Solution**

2. Let  $A = (a_{ij})$  be a matrix of order 3, where  $a_{ij} = x$ ; if  $i = j$ ,  $x \in \mathbb{R}$ , 1 if  $|i - j| = 1$ , 0; otherwise then when of the following Hold (s) good: for  $x = 2$ , (a)  $A$  is a diagonal matrix (b)  $A$  is a symmetric matrix for  $x = 2$ , (c)  $\det A$  has the value equal to 6 (d) Let  $f(x) = \det A$ , then the function  $f(x)$  has both the maxima and minima.

A. for  $x = 2$ ,  $A$  is a diagonal matrix

B.  $A$  is a symmetric matrix

C. for  $x = 2$ ,  $\det A$  has the value equal to 6

D. Let  $f(x) = \det A$ , then the function  $f(x)$  has both the maxima and minima

**Answer: B::D**

 [Watch Video Solution](#)

3. If

$A = [1 \ -1 \ 2]$ ,  $B = [a \ 1 \ b]$  and  $(A + B)^2 = A^2 + B^2 + 2AB$ , then

$a = -1$  b.  $a = 1$  c.  $b = 2$  d.  $b = -2$

A.  $a = -1$

B.  $a = 1$

C.  $b = 2$

D.  $b = -2$

**Answer: A::D**

 [Watch Video Solution](#)

4. If  $AB=A$  and  $BA=B$  then which of the following is/are true ?

- A. (a)  $A$  is idempotent
- B. (b)  $B$  is idempotent
- C. (c)  $A^T$  is idempotent
- D. (d) none of these

**Answer: A::B::C**



**Watch Video Solution**

5. If  $A(\theta) = \begin{bmatrix} \sin \theta & i \cos \theta \\ i \cos \theta & \sin \theta \end{bmatrix}$ , then which of the following is not true ?

- A.  $A(\theta)^{-1} = A(\pi - \theta)$
- B.  $A(\theta) + A(\pi + \theta)$  is a null matrix
- C.  $A(\theta)$  is invertible for all  $\theta \in R$
- D.  $A(\theta)^{-1} = A(-\theta)$



**Answer: A::B::C**



**Watch Video Solution**

6. Let  $A$  and  $B$  be two nonsingular square matrices,  $A^T$  and  $B^T$  are the tranpose matrices of  $A$  and  $B$ , respectively, then which of the following are correct ?

- A.  $B^T AB$  is symmetric matrix if  $A$  is symmetric
- B.  $B^T AB$  is symmetric matrix if  $B$  is symmetric
- C.  $B^T AB$  is skew-symmetric matrix for every matrix  $A$
- D.  $B^T AB$  is skew-symmetric matrix if  $A$  is skew-symmetric

**Answer: A::D**



**Watch Video Solution**

7. If  $B$  is an idempotent matrix, and  $A = I - B$ , then

A.  $A^2 = A$

B.  $A^2 = I$

C.  $AB = O$

D.  $BA = O$

Answer: A::C::D



Watch Video Solution

8. If  $A_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$ ,  $A_2 = \begin{bmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{bmatrix}$ , then  $A_i A_k + A_k A_i$

is equal to  $2I$  if  $i = k$  b.  $O$  if  $i \neq k$  c.  $2I$  if  $i \neq k$  d.  $O$  always

A.  $2I$  if  $i = k$

B.  $O$  if  $i \neq k$

C.  $2I$  if  $i \neq k$

D.  $O$  always

Answer: A::B



Watch Video Solution

9. Suppose  $a_1, a_2, \dots$  Are real numbers, with  $a_1 \neq 0$ . If  $a_1, a_2, a_3, \dots$  Are in A.P., then

A.  $A = \begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_5 & a_6 & a_7 \end{bmatrix}$  is singular (where  $i = \sqrt{-1}$ )

B. the system of equations

$$a_1x + a_2y + a_3z = 0, a_4x + a_5y + a_6z = 0, a_7x + a_8y + a_9z = 0$$

has infinite number of solutions

C.  $B = \begin{bmatrix} a_1 & ia_2 \\ ia_2 & a_1 \end{bmatrix}$  is nonsingular

D. none of these

Answer: A::B::C



Watch Video Solution

10. If  $\alpha, \beta, \gamma$  are three real numbers and

$$A = \begin{bmatrix} 1 & \cos(\alpha - \beta) & \cos(\alpha - \gamma) \\ \cos(\beta - \alpha) & 1 & \cos(\beta - \gamma) \\ \cos(\gamma - \alpha) & \cos(\gamma - \beta) & 1 \end{bmatrix}, \text{ then which of following}$$

is/are true? a.  $A$  is singular b.  $A$  is symmetric c.  $A$  is orthogonal d.  $A$  is not invertible

A.  $A$  is singular

B.  $A$  is symmetric

C.  $A$  is orthogonal

D.  $A$  is not invertible

**Answer: A::B::D**



**Watch Video Solution**

11. If  $D_1$  and  $D_2$  are two  $3 \times 3$  diagonal matrices, then which of the following is/are true ?

A.  $D_1 D_2$  is a diagonal matrix

B.  $D_1 D_2 = D_2 D_1$

C.  $D_1^2 + D_2^2$  is a diagonal matrix

D. none of these

**Answer:**



**Watch Video Solution**

**12.** Let  $A$  be the  $2 \times 2$  matrix given by  $A = [a_{ij}]$  where  $a_{ij} \in \{0, 1, 2, 3, 4\}$  such that  $a_{11} + a_{12} + a_{21} + a_{22} = 4$  then which of the following statement(s) is/are true ?

A. Number of matrices  $A$  such that the trace of  $A$  equal to 4, is 5

B. Number of matrices  $A$ , such that  $A$  is invertible is 18

C. Absolute difference between maximum value and minimum value of  $\det(A)$  is 8

D. Number of matrices  $A$  such that  $A$  is either symmetric (or) skew symmetric and  $\det(A)$  is divisible by 2, is 5.

**Answer:**



[Watch Video Solution](#)

13.

If

$$S = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \text{ and } A = \begin{bmatrix} b+c & c+a & b-c \\ c-b & c+b & a-b \\ b-c & a-c & a+b \end{bmatrix} \quad (a, b, c \neq 0), \text{ then } SAS^{-1}$$

is a. symmetric matrix b. diagonal matrix c. invertible matrix d. singular matrix

A. symmetric matrix

B. diagonal matrix

C. invertible matrix

D. singular matrix

**Answer:**



[Watch Video Solution](#)

14.  $P$  is a non-singular matrix and  $A, B$  are two matrices such that

$B = P^{-1}AP$ . The true statements among the following are

A.  $A$  is invertible iff  $B$  is invertible

B.  $B^n = P^{-1}A^nP \forall n \in \mathbb{N}$

C.  $\forall \lambda \in \mathbb{R}, B - \lambda I = P^{-1}(A - \lambda I)P$

D.  $A$  and  $B$  are both singular matrices

**Answer:**



[Watch Video Solution](#)

15. Let  $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$ . Then  $A^2 - 4A - 5I_3 = O$  b.

$A^{-1} = \frac{1}{5}(A - 4I_3)$  c.  $A^3$  is not invertible d.  $A^2$  is invertible

A.  $A^2 - 4A - 5I_3 = O$

B.  $A^{-1} = \frac{1}{5}(A - 4I_3)$

C.  $A^3$  is not invertible

D.  $A^2$  is invertible

**Answer:**

 [Watch Video Solution](#)

16. If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , then which is true a.  $A^3 - A^2 = A - I$  b.  $\det.$

$(A^{100} - I) = 0$  c.  $A^{200} = \begin{bmatrix} 1 & 0 & 0 \\ 100 & 1 & 0 \\ 100 & 0 & 1 \end{bmatrix}$  d.  $A^{100} = \begin{bmatrix} 1 & 1 & 0 \\ 50 & 1 & 0 \\ 50 & 0 & 1 \end{bmatrix}$

A.  $A^3 - A^2 = A - I$

B.  $\det. (A^{100} - I) = 0$

C.  $A^{200} = \begin{bmatrix} 1 & 0 & 0 \\ 100 & 1 & 0 \\ 100 & 0 & 1 \end{bmatrix}$

D.  $A^{100} = \begin{bmatrix} 1 & 1 & 0 \\ 50 & 1 & 0 \\ 50 & 0 & 1 \end{bmatrix}$

**Answer:**

 [Watch Video Solution](#)



17. If  $A$  is symmetric and  $B$  is skew-symmetric matrix, then which of the following is/are CORRECT ?

- A.  $ABA^T$  is skew-symmetric matrix
- B.  $AB^T + BA^T$  is symmetric matrix
- C.  $(A + B)(A - B)$  is skew-symmetric
- D.  $(A + I)(B - I)$  is symmetric

**Answer:**

 [Watch Video Solution](#)

18. If  $A = ((a_{ij}))_{n \times n}$  and  $f$  is a function, we define

$f(A) = ((f(a_{ij})))_{n \times n}$ , Let  $A = \begin{bmatrix} \pi/2 - \theta & \theta \\ -\theta & \pi/2 - \theta \end{bmatrix}$ . Then

a.  $\sin A$  is invertible b.  $\sin A = \cos A$  c.  $\sin A$  is orthogonal

d.  $\sin(2A) = 2A \sin A \cos A$

A.  $\sin A$  is invertible

B.  $\sin A = \cos A$

C.  $\sin A$  is orthogonal

D.  $\sin(2A) = 2 \sin A \cos A$

**Answer:**



**Watch Video Solution**

19. If  $A$  is a matrix such that  $A^2 + A + 2I = O$ ; the which of the following is/are true? (a)  $A$  is non-singular (b)  $A$  is symmetric (c)  $A$  cannot be skew-symmetric (d)  $A^{-1} = -\frac{1}{2}(A + I)$

A.  $A$  is nonsingular

B.  $A$  is symmetric

C.  $A$  cannot be skew-symmetric

D.  $A^{-1} = -\frac{1}{2}(A + I)$

**Answer:**



Watch Video Solution

20. If  $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ , then the trace of the matrix  $Adj(AdjA)$  is

A.  $adj(adjA) = A$

B.  $|adj(adjA)|=1$

C.  $|adjA|=1$

D. none of these

Answer: B



Watch Video Solution

21. If  $\begin{bmatrix} 1 & -\tan\theta \\ \tan\theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan\theta \\ -\tan\theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ , then

A.  $a = \cos 2\theta$

B.  $a = 1$

$$C. b = \sin 2\theta$$

$$D. b = -1$$

**Answer:**

 [Watch Video Solution](#)

22. If  $A^{-1} = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & -1/3 \end{bmatrix}$ , then :

A.  $|A| = -1$

B.  $\text{adj } A = \begin{bmatrix} -1 & 1 & -2 \\ 0 & -3 & -1 \\ 0 & 0 & 1/3 \end{bmatrix}$

C.  $A = \begin{bmatrix} 1 & 1/3 & 7 \\ 0 & 1/3 & 1 \\ 0 & 0 & -3 \end{bmatrix}$

D.  $A = \begin{bmatrix} 1 & -1/3 & -7 \\ 0 & -3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

**Answer:**

 [Watch Video Solution](#)

23. If  $A$  is an invertible matrix, then  $(\text{adj} A)^{-1}$  is equal to a.  $\text{adj} A^{-1}$  b.

$\frac{A}{\det A}$  c.  $A$  d.  $(\det A)A$

A.  $\text{adj.} (A^{-1})$

B.  $\frac{A}{\det. A}$

C.  $A$

D.  $(\det. A) A$

**Answer:**



**Watch Video Solution**

24. If  $A$  and  $B$  are two invertible matrices of the same order, then  $\text{adj} (AB)$

is equal to

A.  $\text{adj} (B) \text{adj} (A)$

B.  $|B||A|B^{-1}A^{-1}$

C.  $|B||A|A^{-1}B^{-1}$

D.  $|A||B|(AB)^{-1}$

**Answer:**

 [Watch Video Solution](#)

25. If  $A$ ,  $B$ , and  $C$  are three square matrices of the same order, then

$AB = AC \Rightarrow B = C$ . Then

A.  $|A| \neq 0$

B.  $A$  is invertible

C.  $A$  may be orthogonal

D.  $A$  is symmetric

**Answer:**

 [Watch Video Solution](#)

26. If  $A$  and  $B$  are two non singular matrices and both are symmetric and commute each other, then

A.  $A^{-1}B$

B.  $AB^{-1}$

C.  $A^{-1}B^{-1}$

D. none of these

**Answer:**



[Watch Video Solution](#)

27. If  $A$  and  $B$  are square matrices of order 3 such that  $A^3 = 8B^3 = 8I$  and  $\det. (AB - A - 2B + 2I) \neq 0$ , then identify the correct statement(s), where  $I$  is identity matrix of order 3.

A.  $A^2 + 2A + 4I = O$

B.  $A^2 + 2A + 4I \neq O$

$$C. B^2 + B + I = O$$

$$D. B^2 + B + I \neq O$$

**Answer:**  $A^2 + 2A + 4I = O$  and  $B^2 + B + I = O$



[Watch Video Solution](#)

28. Let  $A, B$  be two matrices different from identity matrix such that  $AB = BA$  and  $A^n - B^n$  is invertible for some positive integer  $n$ . If  $A^n - B^n = A^{n+1} - B^{n+1} = A^{n+1} - B^{n+2}$ , then

A.  $I - A$  is non-singular

B.  $I - B$  is non-singular

C.  $I - A$  is singular

D.  $I - B$  is singular

**Answer:**



[Watch Video Solution](#)



29. Let  $A$  and  $B$  be square matrices of the same order such that  $A^2 = I$  and  $B^2 = I$ , then which of the following is CORRECT ?

A. If  $A$  and  $B$  are inverse to each other, then  $A = B$ .

B. If  $AB = BA$ , then there exists matrix  $C = \frac{AB + BA}{2}$  such that  $C^2 = C$ .

C. If  $AB = BA$ , then there exists matrix  $D = AB - BA$  such that  $D^n = O$  for some  $n \in \mathbb{N}$ .

D. If  $AB = BA$  then  $(A + B)^5 = 16(A + B)$ .

**Answer:**



[Watch Video Solution](#)

30. Let  $B$  is an invertible square matrix and  $B$  is the adjoint of matrix  $A$  such that  $AB = B^T$ . Then

A. A is an identity matrix

B. B is symmetric matrix

C. A is a skew-symmetric matrix

D. B is skew symmetric matrix

**Answer: A**



**Watch Video Solution**

**31.** First row of a matrix A is  $[1, 3, 2]$ . If

$$\text{adj } A = \begin{bmatrix} -2 & 4 & \alpha \\ -1 & 2 & 1 \\ 3\alpha & -5 & -2 \end{bmatrix}, \text{ then a det (A) is}$$

A.  $-2$

B.  $-1$

C.  $0$

D.  $1$

**Answer:**



**Watch Video Solution**

**32.** Let  $A$  be a square matrix of order 3 satisfies the relation

$A^3 - 6A^2 + 7A - 8I = O$  and  $B = A - 2I$ . Also,  $\det. A = 8$ , then

A.  $\det. \left( \text{adj.} (I - 2A^{-1}) \right) = \frac{25}{16}$

B.  $\text{adj.} \left( \left( \frac{B}{2} \right)^{-1} \right) = \frac{B}{10}$

C.  $\det. \left( \text{adj.} (I - 2A^{-1}) \right) = \frac{75}{32}$

D.  $\text{adj.} \left( \left( \frac{B}{2} \right)^{-1} \right) = \frac{2B}{5}$

**Answer:**



**Watch Video Solution**

**33.** Which of the following matrices have eigen values as 1 and -1 ? (a)

$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  (b)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  (c)  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$  (d)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

A.  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

B.  $\begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$

C.  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

D.  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

**Answer:**



[Watch Video Solution](#)

**34.** Let  $M$  and  $N$  be two  $3 \times 3$  non singular skew-symmetric matrices such that  $MN = NM$ . If  $P^T$  denote the transpose of  $P$ , then  $M^2 N^2 (M^T N^{-1})^T$  is equal to  $M^2$  b.  $-N^2$  c.  $-M^2$  d.  $MN$

A.  $M^2$

B.  $-N^2$

C.  $-M^2$

D.  $MN$

**Answer: C**



**Watch Video Solution**

**35.** Let  $\omega$  be a complex cube root of unity with  $\omega \neq 1$  and  $P = [p_{ij}]$  be a  $n \times n$  matrix with  $p_{ij} = \omega^{i+j}$ . Then  $P^2 \neq O$ , when = a.57 b. 55 c. 58 d.

56

A. 57

B. 55

C. 58

D. 56

**Answer: B::C::D**



**Watch Video Solution**

36. For  $3 \times 3$  matrices  $M$  and  $N$ , which of the following statement (s) is (are) NOT correct ?

Statement - I :  $N^T MN$  is symmetric or skew-symmetric, according as  $M$  is symmetric or skew-symmetric.

Statement - II :  $MN - NM$  is skew-symmetric for all symmetric matrices  $M$  and  $N$ .

Statement - III :  $MN$  is symmetric for all symmetric matrices  $M$  and  $N$ .

Statement - IV :  $(adj M)(adj N) = adj(MN)$  for all invertible matrices  $M$  and  $N$ .

A.  $N^T MN$  is symmetric or skew-symmetric, according as  $M$  is symmetric or skew-symmetric

B.  $MN - NM$  is skew-symmetric for all symmetric matrices  $M$  and  $N$

C.  $MN$  is symmetric for all symmetric matrices  $M$  and  $N$

D.  $(adj M)(adj N) = adj(MN)$  for all invertible matrices  $M$  and  $N$ .

Answer: C::D



Watch Video Solution

37. Let  $M$  be a  $2 \times 2$  symmetric matrix with integer entries. Then  $M$  is invertible if

- a. The first column of  $M$  is the transpose of the second row of  $M$
- b. The second row of  $M$  is the transpose of the first column of  $M$
- c.  $M$  is a diagonal matrix with non-zero entries in the main diagonal
- d. The product of entries in the main diagonal of  $M$  is not the square of an integer

A. the first column of  $M$  is the transpose of the second row of  $M$

B. the second row of  $M$  is the transpose of the column of  $M$

C.  $M$  is a diagonal matrix with non-zero entries in the main diagonal

D. the product of entries in the main diagonal of  $M$  is not the square of an integer

**Answer: C::D**



**Watch Video Solution**

38. Let  $m$  and  $N$  be two  $3 \times 3$  matrices such that  $MN=NM$ . Further if  $M \neq N^2$  and  $M^2 = N^4$  then which of the following are correct.

A. determinant of  $(M^2 + Mn^2)$  is 0

B. there is a  $3 \times 3$  non-zero matrix  $U$  such that  $(M^2 + MN^2)U$  is the zero matrix

C. determinant of  $(M^2 + MN^2) \geq 1$

D. for a  $3 \times 3$  matrix  $U$ , is the zero matrix

**Answer: A:B**



[Watch Video Solution](#)

39. Let  $X$  and  $Y$  be two arbitrary,  $3 \times 3$ , non-zero, skew-symmetric matrices and  $Z$  be an arbitrary  $3 \times 3$ , non-zero, symmetric matrix. Then which of the following matrices is (are) skew symmetric?

A.  $Y^3Z^4 - Z^4Y^3$



B.  $X^{44} + Y^{44}$

C.  $X^4 Z^3 - Z^3 X^4$

D.  $X^{23} + Y^{23}$

Answer: C::D



Watch Video Solution

40. Let  $p = \begin{bmatrix} 3 & -1 & -2 \\ 2 & 0 & \alpha \\ 3 & -5 & 0 \end{bmatrix}$ , where  $\alpha \in \mathbb{R}$ . Suppose  $Q = [q_{ij}]$  is a matrix such that  $PQ = kI$ , where  $k \in \mathbb{R}, k \neq 0$  and  $I$  is the identity matrix of order 3. If  $q_{23} = -\frac{k}{8}$  and  $\det(Q) = \frac{k^2}{2}$ , then

A.  $\alpha = 0, k = 8$

B.  $4\alpha - k + 8 = 0$

C.  $\det(P \operatorname{adj}(Q)) = 2^9$

D.  $\det(Q \operatorname{adj}(P)) = 2^{13}$

Answer: B::C



Watch Video Solution

41. Which of the following is(are) NOT of the square of a  $3 \times 3$  matrix with real entries?

A. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

B. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

C. 
$$\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

D. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: A::C



Watch Video Solution

42. Let  $S$  be the set of all column matrices  $\begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$  such that  $b_1, b_2, b_3 \in \mathbb{R}$

and the system of equations (in real variables)

$$-x + 2y + 5z = b_1$$

$$2x - 4y + 3z = b_2$$

$$x - 2y + 2z = b_3$$

has at least one solution. The, which of the following system (s) (in real

variables) has (have) at least one solution for each  $\begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} \in S$ ?

A.  $x + 2y + 3z = b_1$ ,  $4y + 5z = b_2$  and  $x + 2y + 6z = b_3$

B.  $x + y + 3z = b_1$ ,  $5x + 2y + 6z = b_2$  and  $-2x - y - 3z = b_3$

C.  $x + 2y - 5z = b_1$ ,  $2x - 4y + 10z = b_2$  and  $x - 2y + 5z = b_3$

D.  $x + 2y + 5z = b_1$ ,  $2x + 3z = b_2$  and  $x + 4y - 5z = b_3$

**Answer: A:C:D**



**Watch Video Solution**

43. If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ , then

A.  $A^3 - A^2 = A - I$

B.  $\text{Det}(A^{2010} - I) = 0$

C.  $A^{50} = \begin{bmatrix} 1 & 0 & 0 \\ 25 & 1 & 0 \\ 25 & 0 & 1 \end{bmatrix}$

D.  $A^{50} = \begin{bmatrix} 1 & 1 & 0 \\ 25 & 1 & 0 \\ 25 & 0 & 1 \end{bmatrix}$

**Answer: A::B::C**



**Watch Video Solution**

44. If the elements of a matrix  $A$  are real positive and distinct such that

$\det(A + A^T)^T = 0$  then

A.  $\det A > 0$

B.  $\det A \geq 0$

C.  $\det(A - A^T) > 0$

D.  $\det(A \cdot A^T) > 0$

**Answer: A::C::D**



**Watch Video Solution**

45. If  $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$  and  $X$  is a non zero column matrix such

that  $AX = \lambda X$ , where  $\lambda$  is a scalar, then values of  $\lambda$  can be

A. 3

B. 6

C. 12

D. 15

**Answer: A::D**



**Watch Video Solution**

46. If  $A, B$  are two square matrices of same order such that  $A + B = AB$  and  $I$  is identity matrix of order same as that of  $A, B$ , then

A.  $AB = BA$

B.  $|A - I| = 0$

C.  $|B - I| \neq 0$

D.  $|A - B| = 0$

Answer: A::C



Watch Video Solution

47. If  $A$  is a non-singular matrix of order  $n \times n$  such that  $3ABA^{-1} + A = 2A^{-1}BA$ , then

A.  $A$  and  $B$  both are identity matrices

B.  $|A + B| = 0$

C.  $|ABA^{-1} - A^{-1}BA| = 0$

D.  $A + B$  is not a singular matrix

**Answer: B::C**



**Watch Video Solution**

**48.** If the matrix  $A$  and  $B$  are of  $3 \times 3$  and  $(I - AB)$  is invertible, then which of the following statement is/are correct ?

A.  $I - BA$  is not invertible

B.  $I - BA$  is invertible

C.  $I - BA$  has for its inverse  $I + B(I - AB)^{-1}A$

D.  $I - BA$  has for its inverse  $I + A(I - BA)^{-1}B$

**Answer: B::C**



**Watch Video Solution**

49. If  $A$  is a square matrix such that  $A \cdot (\text{Adj}A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$ , then

A.  $|A| = 4$

B.  $|\text{adj}A| = 16$

C.  $\frac{|\text{adj}(\text{adj}A)|}{|\text{adj}A|} = 16$

D.  $|\text{adj}^2A| = 128$

Answer: A::B::C



Watch Video Solution

### Linked Comprehension Type

1. Let  $a$  be a matrix of order  $2 \times 2$  such that  $A^2 = O$ .

$A^2 - (a + d)A + (ad - bc)I$  is equal to

A.  $I$



B.  $O$

C.  $-I$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

2. Let  $A$  be a matrix of order  $2 \times 2$  such that  $A^2 = O$ .

$\text{tr}(A)$  is equal to

A. 1

B. 0

C.  $-1$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

3. Let  $A$  be a matrix of order  $2 \times 2$  such that  $A^2 = O$ .

$$(I + A)^{100} =$$

A.  $100A$

B.  $100(I + A)$

C.  $100I + A$

D.  $I + 100A$

**Answer: D**



[Watch Video Solution](#)

4. If  $A$  and  $B$  are two square matrices of order  $3 \times 3$  which satisfy  $AB = A$

and  $BA = B$ , then

Which of the following is true ?

A. If matrix  $A$  is singular, then matrix  $B$  is nonsingular.

B. If matrix A is nonsingular, then matrix B is singular.

C. If matrix A is singular, then matrix B is also singular.

D. Cannot say anything.

**Answer: C**



[Watch Video Solution](#)

5. if  $A$  and  $B$  are two matrices of order  $3 \times 3$  so that  $AB = A$  and  $BA = B$  then  $(A + B)^7 =$

A.  $7(A + B)$

B.  $7 \cdot I_{3 \times 3}$

C.  $64(A + B)$

D.  $128I$

**Answer: C**



[Watch Video Solution](#)

6. If  $A$  and  $B$  are two square matrices of order  $3 \times 3$  which satisfy  $AB = A$  and  $BA = B$ , then

$(A + I)^5$  is equal to (where  $I$  is identity matrix)

A.  $I + 60I$

B.  $I + 16A$

C.  $I + 31A$

D. none of these

**Answer: C**



[Watch Video Solution](#)

7. Consider an arbitrary  $3 \times 3$  non-singular matrix  $A [a_{ij}]$ . A matrix  $B = [b_{ij}]$  is formed such that  $b_{ij}$  is the sum of all the elements except  $a_{ij}$  in the  $i$ th row of  $A$ . Answer the following questions :

If there exists a matrix  $X$  with constant elements such that  $AX=B$ , then  $X$  is

A. skew-symmetric

B. null matrix

C. diagonal matrix

D. none of these

**Answer: D**



**Watch Video Solution**

8. Let  $A = [a_{ij}]$  be  $3 \times 3$  matrix and  $B = [b_{ij}]$  be  $3 \times 3$  matrix such that  $b_{ij}$  is the sum of the elements of  $i^{th}$  row of A except  $a_{ij}$ . If  $\det(A) = 19$ , then the value of  $\det(B)$  is \_\_\_\_\_ .

A.  $|A|$

B.  $|A|/2$

C.  $2|A|$

D. none of these

Answer: C



Watch Video Solution

9. Let  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  satisfies  $A^n = A^{n-2} + A^2 - I$  for  $n \geq 3$ . And

trace of a square matrix X is equal to the sum of elements in its principal diagonal.

Further consider a matrix  $U_{3 \times 3}$  with its column as  $U_1, U_2, U_3$  such that

$$A^{50} U_1 = \begin{bmatrix} 1 \\ 25 \\ 25 \end{bmatrix}, A^{50} U_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, A^{50} U_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Then answer the following question :

Trace of  $A^{50}$  equals

A. 0

B. 1

C. -1

D. 25

Answer: B



Watch Video Solution

10. Let  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  satisfies  $A^n = A^{n-2} + A^2 - I$  for  $n \geq 3$ . And

trace of a square matrix X is equal to the sum of elements in its principal diagonal.

Further consider a matrix  $U_{3 \times 3}$  with its column as  $U_1, U_2, U_3$  such

that

$$A^{50} U_1 = \begin{bmatrix} 1 \\ 25 \\ 25 \end{bmatrix}, A^{50} U_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, A^{50} U_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Then answer the following question :

Trace of  $A^{50}$  equals

A. 0

B. 1

C. 2

D. 3

Answer: D

 Watch Video Solution

11. Let  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  satisfies  $A^n = A^{n-2} + A^2 - I$  for  $n \geq 3$ . And

trace of a square matrix X is equal to the sum of elements in its principal diagonal.

Further consider a matrix  $U_{3 \times 3}$  with its column as  $U_1, U_2, U_3$  such that

$$A^{50} U_1 = \begin{bmatrix} 1 \\ 25 \\ 25 \end{bmatrix}, A^{50} U_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, A^{50} U_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Then answer the following question :

The value of  $|U|$  equals

A. 0

B. 1

C. 2

D. -1



**Answer: B**



**Watch Video Solution**

12. Let for  $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix}$ , there be three row matrices  $R_1, R_2$  and  $R_3$ ,

satisfying the relations,  $R_1A = [1 \ 0 \ 0]$ ,  $R_2A = [2 \ 3 \ 0]$  and  $R_3A = [2 \ 3 \ 1]$ . If B is square matrix of order 3 with rows  $R_1, R_2$  and  $R_3$  in order, then

The value of  $\det. (2A^{100}B^3 - A^{99}B^4)$  is

A.  $-2$

B.  $-1$

C.  $2$

D.  $-27$

**Answer: D**



**Watch Video Solution**

13. Let for  $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix}$ , there be three row matrices  $R_1, R_2$  and  $R_3$ ,

satisfying the relations,  $R_1A = [1 \ 0 \ 0]$ ,  $R_2A = [2 \ 3 \ 0]$  and

$R_3A = [2 \ 3 \ 1]$ . If B is square matrix of order 3 with rows  $R_1, R_2$  and

$R_3$  in order, then

The value of  $\det. (B)$  is

A.  $-27$

B.  $-9$

C.  $-3$

D.  $9$

**Answer: A**



[Watch Video Solution](#)

14. A and B are square matrices such that  $\det. (A) = 1$ ,  $BB^T = I$ ,  $\det (B) > 0$ , and  $A(\text{adj. } A + \text{adj. } B) = B$ .

The value of  $\det(A + B)$  is

A.  $-2$

B.  $-1$

C.  $0$

D.  $1$

**Answer: D**



[Watch Video Solution](#)

15. A and B are square matrices such that  $\det(A) = 1$ ,  $BB^T = I$ ,  $\det(B) > 0$ ,  $B + A = B^2$  and  $A(\text{adj. } A + \text{adj. } B) = B$ .

$AB^{-1} =$

A.  $B^{-1}A$

B.  $AB^{-1}$

C.  $A^T B^{-1}$

$$D. B^T A^{-1}$$

**Answer: A**



**Watch Video Solution**

**16.** Let  $A$  be an  $m \times n$  matrix. If there exists a matrix  $L$  of type  $n \times m$  such that  $LA = I_n$ , then  $L$  is called left inverse of  $A$ . Which of the following

matrices is NOT left inverse of matrix  $\begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 3 \end{bmatrix}$  ?

- A.  $\begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}$
- B.  $\begin{bmatrix} 2 & -7 & 3 \\ -\frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}$
- C.  $\begin{bmatrix} -\frac{1}{2} & \frac{1}{2} & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}$
- D.  $\begin{bmatrix} 0 & 3 & -1 \\ -\frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}$

**Answer: C**



**Watch Video Solution**

17. Let  $A$  be an  $m \times n$  matrix. If there exists a matrix  $L$  of type  $n \times m$  such that  $LA = I_n$ , then  $L$  is called left inverse of  $A$ . Similarly, if there exists a matrix  $R$  of type  $n \times m$  such that  $AR = I_m$ , then  $R$  is called right inverse of  $A$ .

For example, to find right inverse of matrix

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 3 \end{bmatrix}, \text{ we take } R = \begin{bmatrix} x & y & z \\ u & v & w \end{bmatrix}$$

and solve  $AR = I_3$ , i.e.,

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x & y & z \\ u & v & w \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\implies \begin{array}{lll} x - u = 1 & y - v = 0 & z - w = 0 \\ x + u = 0 & y + v = 1 & z + w = 0 \\ 2x + 3u = 0 & 2y + 3v = 0 & 2z + 3w = 1 \end{array}$$

As this system of equations is inconsistent, we say there is no right inverse for matrix  $A$ .

The number of right inverses for the matrix  $\begin{bmatrix} 1 & -1 & 2 \\ 2 & -1 & 1 \end{bmatrix}$  is

A. 0

B. 1

C. 2

D. infinite

Answer: D



Watch Video Solution

**18.** Let  $A$  be an  $m \times n$  matrix. If there exists a matrix  $L$  of type  $n \times m$  such that  $LA = I_n$ , then  $L$  is called left inverse of  $A$ . Similarly, if there exists a matrix  $R$  of type  $n \times m$  such that  $AR = I_m$ , then  $R$  is called right inverse of  $A$ .

For example, to find right inverse of matrix

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 3 \end{bmatrix}, \text{ we take } R = \begin{bmatrix} x & y & z \\ u & v & w \end{bmatrix}$$

and solve  $AR = I_3$ , i.e.,

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x & y & z \\ u & v & w \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{aligned} \implies x - u &= 1 & y - v &= 0 & z - w &= 0 \\ x + u &= 0 & y + v &= 1 & z + w &= 0 \\ 2x + 3u &= 0 & 2y + 3v &= 0 & 2z + 3w &= 1 \end{aligned}$$

As this system of equations is inconsistent, we say there is no right inverse for matrix  $A$ .

For which of the following matrices, the number of left inverses is greater than the number of right inverses ?

A.  $\begin{bmatrix} 1 & 2 & 4 \\ -3 & 2 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} 3 & 2 & 1 \\ 3 & 2 & 1 \end{bmatrix}$

C.  $\begin{bmatrix} 1 & 4 \\ 2 & -3 \\ 2 & -3 \end{bmatrix}$

D.  $\begin{bmatrix} 3 & 3 \\ 1 & 1 \\ 4 & 4 \end{bmatrix}$

**Answer: C**



[Watch Video Solution](#)

**19.** Let A be the set of all  $3 \times 3$  symmetric matrices all of whose entries are either 0 or 1. Five of these entries are 1 and four of them are 0. The number of matrices in A is

A. 12

B. 6

C. 9

D. 3

**Answer: A**



**Watch Video Solution**

**20.** Let  $A$  be the set of all  $3 \times 3$  symmetric matrices all of whose either 0 or 1. Five of these entries are 1 and four of them are 0.

The number of matrices  $A$  in  $A$  for which the system of linear equations

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

is inconsistent is

A. less than 4

B. at least 4 but less than 7

C. at least 7 but less than 10

D. at least 10



**Answer: B**



[Watch Video Solution](#)

21. Let  $A$  be the set of all  $3 \times 3$  symmetric matrices all of whose either 0 or 1. Five of these entries are 1 and four of them are 0.

The number of matrices  $A$  in  $A$  for which the system of linear equations

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

is inconsistent is

- A. 0
- B. more than 2
- C. 2
- D. 1

**Answer: B**



[Watch Video Solution](#)

22. Let  $P$  be an odd prime number and  $T_p$  be the following set of  $2 \times 2$  matrices :

$$T_p = \left\{ A = \begin{bmatrix} a & b \\ c & a \end{bmatrix} : a, b, c \in \{0, 1, \dots, p-1\} \right\}$$

The number of  $A$  in  $T_p$  such that  $\det(A)$  is not divisible by  $p$  is

A.  $(p-1)^2$

B.  $2(p-1)$

C.  $(p-1)^2 + 1$

D.  $2p-1$

**Answer: D**



[Watch Video Solution](#)

23. Let  $P$  be an odd prime number and  $T_p$  be the following set of  $2 \times 2$  matrices :

$$T_p = \left\{ A = \begin{bmatrix} a & b \\ c & a \end{bmatrix} : a, b, c \in \{0, 1, \dots, p-1\} \right\}$$

The number of  $A$  in  $T_p$  such that  $\det(A)$  is not divisible by  $p$  is

A.  $(p - 1)(p^2 - p + 1)$

B.  $p^3 - (p - 1)^2$

C.  $(p - 1)^2$

D.  $(p - 1)(p^2 - 2)$

**Answer: C**



**Watch Video Solution**

**24.** Let  $p$  be an odd prime number and  $T_p$ , be the following set of  $2 \times 2$

matrices  $T_p = \left\{ A = \begin{bmatrix} a & b \\ c & a \end{bmatrix} : a, b, c \in \{0, 1, 2, \dots, p - 1\} \right\}$  The

number of  $A$  in  $T_p$ , such that  $A$  is either symmetric or skew-symmetric or

both, and  $\det(A)$  divisible by  $p$  is

A.  $2p-1$

B.  $p^3 - 5p$

C.  $3p-4$

D.  $p^3 - p^2$

**Answer: D**



**Watch Video Solution**

25. Let  $a, b,$  and  $c$  be three real numbers satisfying

$$[a, b, c] \begin{bmatrix} 1 & 9 & 7 \\ 8 & 2 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0, 0, 0]$$

If the point  $P(a, b, c)$  with reference to (E),

lies on the plane  $2x + y + z = 1$ , the the value of  $7a + b + c$  is (A) 0 (B)

12 (C) 7 (D) 6

A. 0

B. 12

C. 7

D. 6

**Answer: D**



**Watch Video Solution**

26. Let  $a, b,$  and  $c$  be three real numbers satisfying

$$[a, b, c] \begin{bmatrix} 1 & 9 & 7 \\ 8 & 2 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0, 0, 0] \text{ Let } \omega \text{ be a solution of } x^3 - 1 = 0 \text{ with}$$

$\text{Im}(\omega) > 0$ . If  $a = 2$  with  $b$  and  $c$  satisfying (E) then the value of

$$\frac{3}{\omega^a} + \frac{1}{\omega^b} + \frac{3}{\omega^c} \text{ is equal to (A) } -2 \text{ (B) } 2 \text{ (C) } 3 \text{ (D) } -3$$

A.  $-2$

B.  $2$

C.  $3$

D.  $-3$

**Answer: A**



**Watch Video Solution**

27. Let  $a, b,$  and  $c$  be three real numbers satisfying

$$[a, b, c] \begin{bmatrix} 1 & 9 & 7 \\ 8 & 2 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0, 0, 0] \text{ Let } b=6, \text{ with } a \text{ and } c \text{ satisfying (E). If alpha}$$

and  $\beta$  are the roots of the quadratic equation

$ax^2 + bx + c = 0$  then  $\sum_{n=0}^{\infty} \left( \frac{1}{\alpha} + \frac{1}{\beta} \right)^n$  is (A) 6 (B) 7 (C)  $\frac{6}{7}$  (D)  $\infty$

A. 6

B. 7

C.  $\frac{6}{7}$

D.  $\infty$

**Answer: B**



[Watch Video Solution](#)

Matrix Type

1. Match the following lists :

List I	List II
a. $(I - A)^n$ is if $A$ is idempotent	p. $2^{n-1}(I - A)$
b. $(I - A)^n$ is if $A$ is involutory	q. $I - nA$
c. $(I - A)^n$ is if $A$ is nilpotent of index 2	r. $A$
d. If $A$ is orthogonal, then $(A^T)^{-1}$	s. $I - A$



Watch Video Solution

2. Match the following lists :

List I	List II
a. If $A$ is an idempotent matrix and $I$ is an identity matrix of the same order, then the value of $n$ , such that $(A + I)^n = I + 127I$ is	p. 9
b. If $(I - A)^{-1} = I + A + A^2 + \dots + A^7$ , then $A^n = O$ , where $n$ is	q. 10
c. If $A$ is matrix such that $a_{ij} = (i + j)(i - j)$ , then $A$ is singular if order of matrix is	r. 7
d. If a nonsingular matrix $A$ is symmetric, show that $A^{-1}$ is also symmetric, then order of $A$ can be	s. 8



Watch Video Solution

### 3. Match the following lists :

List I ( $A, B, C$ are matrices)	List II
a. If $ A  = 2$ , then $ 2A^{-1}  =$ (where $A$ is of order 3)	p. 1
b. If $ A  = 1/8$ , then $ \text{adj}(\text{adj}(2A))  =$ (where $A$ is of order 3)	q. 4
c. If $(A + B)^2 = A^2 + B^2$ , and $ A  = 2$ , then $ B  =$ (where $A$ and $B$ are of odd order)	r. 24
d. $ A_{2 \times 2}  = 2$ , $ B_{3 \times 3}  = 3$ and $ C_{4 \times 4}  = 4$ , then $ ABC $ is equal to	s. 0
	t. does not exist



[Watch Video Solution](#)

4. Consider a matrix  $A = [a_{ij}]$  of order  $3 \times 3$  such that  $a_{ij} = (k)^{i+j}$  where  $k \in I$ .

Match List I with List II and select the correct answer using the codes



given below the lists.

List I	List II
a. $A$ is singular if	p. $k \in \{0\}$
b. $A$ is null matrix if	q. $k \in \phi$
c. $A$ is skew-symmetric which is not null matrix if	r. $k \in I$
d. $A^2 = 3A$ if	s. $k \in \{-1, 0, 1\}$

A.  $a \ b \ c \ d$   
 $r \ p \ s \ q$

B.  $a \ b \ c \ d$   
 $s \ p \ q \ r$

C.  $a \ b \ c \ d$   
 $r \ p \ q \ s$

D.  $a \ b \ c \ d$   
 $q \ p \ r \ s$

**Answer: C**



**Watch Video Solution**

5. Match the following lists :

List I	List II
<p>a. If <math>M_r = \begin{bmatrix} r-1 &amp; \frac{1}{r} \\ 1 &amp; \frac{1}{(r-1)^2} \end{bmatrix}</math> and <math> M_r </math> is the corresponding determinant, then <math>\lim_{n \rightarrow \infty} ( M_2  +  M_3  + \dots +  M_n ) =</math></p>	p. 0
<p>b. If <math>(A + B)^2 = A^2 + B^2</math> and <math> A  = 2</math> then <math> B  =</math> (where A and B are matrices of odd order)</p>	q. 1
<p>c. If <math>A = \begin{bmatrix} 3 &amp; 1 \\ -1 &amp; 1 \end{bmatrix}</math> and a matrix C is defined as <math>C = (BAB^{-1})(B^{-1}A^T B)</math>, where <math> C  = K^2</math> (<math>K \in N</math>) then <math>K =</math></p>	r. 2
<p>d. If <math>A = \begin{bmatrix} 1 &amp; 1 \\ -1 &amp; 1 \end{bmatrix}</math> and <math>A^4 = -\lambda I</math> then <math>\lambda - 2</math> is equal to</p>	s. 4

- A.  $a \ b \ c \ d$   
 $s \ r \ q \ p$
- B.  $a \ b \ c \ d$   
 $s \ p \ q \ r$
- C.  $a \ b \ c \ d$   
 $q \ p \ s \ r$
- D.  $a \ b \ c \ d$   
 $s \ q \ r \ p$

Answer: C



Watch Video Solution

## Numerical Value Type

1.  $A = \begin{bmatrix} 0 & 1 \\ 3 & 0 \end{bmatrix}$  and  $(A^8 + A^6 + A^4 + A^2 + I)V = \begin{bmatrix} 0 \\ 11 \end{bmatrix}$  (where  $I$  is the  $2 \times 2$  identity matrix), then the product of all elements of matrix  $V$  is \_\_\_\_\_.



Watch Video Solution

2. If  $[abc \ 1 - a]$  is an idempotent matrix and  $f(x) = x^{-2} = bc = 1/4$ , then the value of  $1/f(a)$  is \_\_\_\_\_.



Watch Video Solution

3. Let  $x$  be the solution set of equation  $A^x = I$ ; where  $A = \begin{bmatrix} 0 & 1 & -14 & -34 \\ 1 & -14 & -34 & -34 \end{bmatrix}$  and  $I$  is the corresponding unit matrix and  $x \subseteq N$ , then the minimum value of  $\sum (\cos^x \theta + \sin^x \theta)$ ,  $\theta \in R$ .



Watch Video Solution

4.  $A = [1 \tan x \ - \tan x \ 1]$  and  $f(x)$  is defined as  $f(x) = \det A^T A^{-1}$  then the value of  $(f(f(f(f(f(x)))))$  is ( $n \geq 2$ ) \_\_\_\_\_.



Watch Video Solution

5. The equation  $\begin{bmatrix} 1 & 2 & 2 \\ 1 & 3 & 4 \\ 2 & 4 & k \end{bmatrix} \begin{bmatrix} x & y & z \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$  has a solution for  $(x, y, z)$  besides  $(0, 0, 0)$ . Then the value of  $k$  is \_\_\_\_\_.



Watch Video Solution

6. If  $A$  is an idempotent matrix satisfying,  $(I - 0.4A)^{-1} = I - \alpha A$ , where  $I$  is the unit matrix of the same order as that of  $A$ , then the value of  $|9\alpha|$  is equal to \_\_\_\_\_.



Watch Video Solution

7.

Let

$$A = \begin{bmatrix} 3x^2 \\ 1 \\ 6x \end{bmatrix}, B = [a \ b \ c], \text{ and } C = \begin{bmatrix} (x+2)^2 & 5x^2 & 2x \\ 5x^2 & 2x & (x+2)^2 \\ 2x & (x+2)^2 & 5x^2 \end{bmatrix}$$

be three given matrices, where  $a, b, c,$  and  $x \in R$ . Given that

$f(x) = ax^2 + bx + c$ , then the value of  $f(I)$  is \_\_\_\_\_.



[Watch Video Solution](#)

8. Let  $A$  be the set of all  $3 \times 3$  skew-symmetric matrices whose entries are either  $-1, 0,$  or  $1$ . If there are exactly three 0s three 1s, and there  $(-1)$ 's, then the number of such matrices is \_\_\_\_\_.



[Watch Video Solution](#)

9. Let  $A = [a_{ij}]_{3 \times 3}$  be a matrix such that  $AA^T = 4I$  and  $a_{ij} + 2c_{ij} = 0$ , where  $C_{ij}$  is the cofactor of  $a_{ij}$  and  $I$  is the unit matrix of order 3.

$$\begin{vmatrix} a_{11} + 4 & a_{12} & a_{13} \\ a_{21} & a_{22} + 4 & a_{23} \\ a_{31} & a_{32} & a_{33} + 4 \end{vmatrix} + 5\lambda \begin{vmatrix} a_{11} + 1 & a_{12} & a_{13} \\ a_{21} & a_{22} + 1 & a_{23} \\ a_{31} & a_{32} & a_{33} + 1 \end{vmatrix} = 0$$

then the value of  $\lambda$  is

 [Watch Video Solution](#)

10. Let  $S$  be the set which contains all possible values for  $I, m, n, p, q, r$  for which  $A = [I^2 - 3p00m^2 - 8qr0n^2 - 15]$  be non-singular idempotent matrix. Then the sum of all the elements of the set  $S$  is \_\_\_\_\_.

 [Watch Video Solution](#)

11. If  $A$  is a diagonal matrix of order  $3 \times 3$  is commutative with every square matrix of order  $3 \times 3$  under multiplication and trace  $(A)=12$ , then find  $|A|$

 [Watch Video Solution](#)

12. If  $A$  is a square matrix of order 3 such that  $|A| = 2$ , then  $\left|(\text{adj}A^{-1})^{-1}\right|$  is \_\_\_\_\_.

 [Watch Video Solution](#)

13. If  $A$  and  $B$  are two matrices of order 3 such that  $AB = O$  and  $A^2 + B = I$ , then  $\text{tr.}(A^2 + B^2)$  is equal to \_\_\_\_\_.

 [Watch Video Solution](#)

14. If  $a, b$ , and  $c$  are integers, then number of matrices  $A = \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix}$  which are possible such that  $AA^T = I$  is \_\_\_\_\_.

 [Watch Video Solution](#)

15. Let  $A = [a_{ij}]$  be  $3 \times 3$  matrix and  $B = [b_{ij}]$  be  $3 \times 3$  matrix such that  $b_{ij}$  is the sum of the elements of  $i^{\text{th}}$  row of  $A$  except  $a_{ij}$ . If  $\det(A) = 19$ ,

then the value of  $\det. (B)$  is \_\_\_\_\_ .

 [Watch Video Solution](#)

16. A square matrix  $M$  of order 3 satisfies  $M^2 = I - M$ , where  $I$  is an identity matrix of order 3. If  $M^n = 5I - 8M$ , then  $n$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)

17. Let  $A = [a_{ij}]_{3 \times 3}$ ,  $B = [b_{ij}]_{3 \times 3}$  and  $C = [c_{ij}]_{3 \times 3}$  be any three matrices, where  $b_{ij} = 3^{i-j}a_{ij}$  and  $c_{ij} = 4^{i-j}b_{ij}$ . If  $\det. A = 2$ , then  $\det. (BC)$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)

18. If  $A$  is a square matrix of order  $2 \times 2$  such that  $|A| = 27$ , then sum of the infinite series  $|A| + \left| \frac{1}{2}A \right| + \left| \frac{1}{4}A \right| + \left| \frac{1}{8}A \right| + \dots$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)



19. If  $A$  is a square matrix of order 2 and  $\det. A = 10$ , then  $\left( (tr. A)^2 - tr. (A^2) \right)$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)

20. Let  $A$  and  $B$  are two square matrices of order 3 such that  $\det. (A) = 3$  and  $\det. (B) = 2$ , then the value of  $\det. \left( (adj. (B^{-1}A^{-1}))^{-1} \right)$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)

21. Let  $P$ ,  $Q$  and  $R$  be invertible matrices of order 3 such  $A = PQ^{-1}$ ,  $B = QR^{-1}$  and  $C = RP^{-1}$ . Then the value of  $\det. (ABC + BCA + CAB)$  is equal to \_\_\_\_\_ .

 [Watch Video Solution](#)

22. If  $A = \begin{bmatrix} 1 & x & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 4 \end{bmatrix}$  is the adjoint of a  $3 \times 3$  matrix B and  $\det(B) = 4$ , then the value of x is \_\_\_\_\_.

 [Watch Video Solution](#)

23. A, B and C are three square matrices of order 3 such that  $A = \text{diag}(x, y, z)$ ,  $\det(B) = 4$  and  $\det(C) = 2$ , where  $x, y, z \in I^+$ . If  $\det(\text{adj}(\text{adj}(ABC))) = 2^{16} \times 3^8 \times 7^4$ , then the number of distinct possible matrices A is \_\_\_\_\_.

 [Watch Video Solution](#)

24. Let  $A = [a_{ij}]$  be a matrix of order 2 where  $a_{ij} \in \{-1, 0, 1\}$  and  $\text{adj. } A = -A$ . If  $\det(A) = -1$ , then the number of such matrices is \_\_\_\_\_.

 [Watch Video Solution](#)

25. Let  $K$  be a positive real number and  $A = [2k - 12\sqrt{k} \ 2\sqrt{k} \ 2\sqrt{k} \ 1 - 2k - 2\sqrt{k} \ 2k - 1]$  and  $B = [0 \ 2k - 1 \ \sqrt{k} \ 1 - 2k]$ . If  $\det(\text{adj}A) + \det(\text{adj}B) = 10^6$ , then  $[k]$  is equal to. [Note:  $\text{adj}M$  denotes the adjoint of a square matrix  $M$  and  $[k]$  denotes the largest integer less than or equal to  $K$ ].

 [Watch Video Solution](#)

26. Let  $M$  be a  $3 \times 3$  matrix satisfying  $M[010] = [-123]$ ,  $M[1-10] = [11-1]$ , and  $M[111] = [0012]$ . Then the sum of the diagonal entries of  $M$  is \_\_\_\_\_.

 [Watch Video Solution](#)

27. let  $z = \frac{-1 + \sqrt{3}i}{2}$ , where  $i = \sqrt{-1}$  and  $r, s \in P1, 2, 3$ . Let  $P = \begin{bmatrix} (-z)^r \\ z^{2s} \end{bmatrix}$

and  $I$  be the identity matrix of order 2. Then the total number of ordered pairs  $(r,s)$  for which  $P^2 = -I$  is

 [Watch Video Solution](#)

## Archives Single Correct Answer Type

1. Let  $A$  be a  $2 \times 2$  matrix

Statement-1  $\text{adj}(\text{adj}A) = A$

Statement-2  $|\text{adj}A| = |A|$

- A. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.
- B. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.
- C. Statement 1 is true, statement 2 is false.
- D. Statement 1 is false, statement 2 is true.

**Answer: B**



[Watch Video Solution](#)

2. The number of  $3 \times 3$  non-singular matrices, with four entries as 1 and all other entries as 0, is:- (1) 5 (2) 6 (3) at least 7 (4) less than 4

A. 5

B. 6

C. at least 7

D. less than 4

**Answer: C**



[Watch Video Solution](#)

3. Let  $A$  be a  $2 \times 2$  matrix with non-zero entries and let  $A^2 = I$ , where  $I$  is a  $2 \times 2$  identity matrix,  $\text{Tr}(A)$  = sum of diagonal elements of  $A$  and  $|A| =$

determinant of matrix A. Statement 1:  $\text{Tr}(A)=0$  Statement 2:  $|A|=1$

- A. Statement 1 is false, statement 2 is true.
- B. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.
- C. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.
- D. Statement 1 is true, statement 2 is false.

**Answer: D**



[Watch Video Solution](#)

4. Let A and B two symmetric matrices of order 3.

Statement 1 :  $A(BA)$  and  $(AB)A$  are symmetric matrices.

Statement 2 :  $AB$  is symmetric matrix if matrix multiplication of A with B is commutative.

- A. Statement 1 is false, statement 2 is true.

B. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.

C. Statement 1 is true, statement 2 is true, statement 2 is not a correct explanation for statement 1.

D. Statement 1 is true, statement 2 is false.

**Answer: C**

 [Watch Video Solution](#)

5. Let  $A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{pmatrix}$ . If  $u_1$  and  $u_2$  are column matrices such that

$Au_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$  and  $Au_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ , then  $u_1 + u_2$  is equal to :

A.  $\begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}$

B.  $\begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}$

- C.  $\begin{pmatrix} -1 \\ -1 \\ 0 \end{pmatrix}$
- D.  $\begin{pmatrix} 1 \\ -1 \\ -1 \end{pmatrix}$

**Answer: D**

 [Watch Video Solution](#)

6. Let  $P$  and  $Q$  be  $3 \times 3$  matrices with  $P \neq Q$ . If  $P^3 = Q^3$  and  $P^2Q = Q^2P$ , then determinant of  $(P^2 + Q^2)$  is equal to  
(1) 2 (2) 1 (3) 0 (4) 1

- A.  $-2$
- B. 1
- C. 0
- D.  $-1$

**Answer: C**

 [Watch Video Solution](#)



7. If  $P = \begin{bmatrix} 1 & \alpha & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 4 \end{bmatrix}$  is the adjoint of a  $3 \times 3$  matrix  $A$  and  $|A| = 4$ , then

$\alpha$  is equal to

A. 4

B. 11

C. 5

D. 0

**Answer: B**



Watch Video Solution

8. If  $A$  is an  $3 \times 3$  non-singular matrix such that  $AA' = A'A$  and  $B = A^{-1}A'$ , then  $BB'$  equals (1)  $I + B$  (2)  $I$  (3)  $B^{-1}$  (4)  $(B^{-1})'$

A.  $I + B$

B.  $I$

C.  $B^{-1}$

D.  $(B^{-1})'$

**Answer: B**



**Watch Video Solution**

9. If  $A = [12221 - 2a2b]$  is a matrix satisfying the equation  $\forall^T = 9I$ , where  $I$  is  $3 \times 3$  identity matrix, then the ordered pair (a, b) is equal to :

(1) (2, -1) (2) (-2, 1) (3) (2, 1) (4) (-2, -1)

A. (2, -1)

B. (-2, 1)

C. (2, 1)

D. (-2, -1)

**Answer: D**



Watch Video Solution

10. If  $A = [5a - b \ 32]$  and  $A \operatorname{adj} A = \sqrt[3]{I}$ , then  $5a + b$  is equal to:

A. 5

B. 4

C. 13

D. -1

Answer: A



Watch Video Solution

11. if  $A = \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix}$  then  $(3A^2 + 12A) = ?$

A.  $\begin{bmatrix} 72 & -63 \\ -84 & 51 \end{bmatrix}$

B.  $\begin{bmatrix} 72 & -84 \\ -63 & 51 \end{bmatrix}$

C.  $\begin{bmatrix} 51 & 63 \\ 84 & 72 \end{bmatrix}$

D.  $\begin{bmatrix} 51 & 84 \\ 63 & 72 \end{bmatrix}$

**Answer: C**



**Watch Video Solution**

### Jee Advanced Single Correct Answer Type

1. The number of  $3 \times 3$  matrices  $A$  whose entries are either 0 or 1 and

for which the system  $A \begin{vmatrix} x \\ y \\ z \end{vmatrix} = \begin{vmatrix} 1 \\ 0 \\ 0 \end{vmatrix}$  has exactly two distinct solution is a. 0

b.  $2^9 - 1$  c. 168 d. 2

A. 0

B.  $2^9 - 1$

C. 168

D. 2

**Answer: A**



Watch Video Solution

2. Let  $\omega \neq 1$  be cube root of unity and  $S$  be the set of all non-singular matrices of the form  $\begin{bmatrix} 1 & a & b \\ \omega & 1 & c \\ \omega^2 & \omega & 1 \end{bmatrix}$ , where each of  $a, b,$  and  $c$  is either  $\omega$  or  $\omega^2$ . Then the number of distinct matrices in the set  $S$  is a. 2 b. 6 c. 4 d. 8

A. 2

B. 6

C. 4

D. 8

Answer: A



Watch Video Solution

3. Let  $P = [a_{ij}]$  be a  $3 \times 3$  matrix and let  $Q = [b_{ij}]$ , where  $b_{ij} = 2^{i+j} a_{ij}$  or  $1 \leq i, j \leq 3$ . If the determinant of

$P$  is 2, then the determinant of the matrix  $Q$  is  $2^{10}$  b.  $2^{11}$  c.  $2^{12}$  d.  $2^{13}$

A.  $2^{10}$

B.  $2^{11}$

C.  $2^{12}$

D.  $2^{13}$

**Answer: D**



**Watch Video Solution**

4. Let  $P = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{bmatrix}$  and  $Q = [q_{ij}]$  be two  $3 \times 3$  matrices such that  $Q - P^5 = I_3$ . Then  $\frac{q_{21} + q_{31}}{q_{32}}$  is equal to

A. 52

B. 103

C. 201

D. 205

**Answer: B**



[Watch Video Solution](#)

5. How many  $3 \times 3$  matrices  $M$  with entries from  $\{0, 1, 2\}$  are there, for which the sum of the diagonal entries of  $M^T M$  is 5?

A. 126

B. 198

C. 135

D. 162

**Answer: B**



[Watch Video Solution](#)

Single Correct Answer

1. If  $A = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$ , then  $(A + B)^2 =$

A.  $A$

B.  $B$

C.  $I$

D.  $A^2 + B^2$

**Answer: D**



[Watch Video Solution](#)

2. If the value of  $\prod_{k=1}^{50} \begin{bmatrix} 1 & 2k - 1 \\ 0 & 1 \end{bmatrix}$  is equal to  $\begin{bmatrix} 1 & r \\ 0 & 1 \end{bmatrix}$  then  $r$  is equal to

A. 62500

B. 2500

C. 1250

D. 12500



**Answer: B**



[Watch Video Solution](#)

3. A square matrix  $P$  satisfies  $P^2 = I - P$ , where  $I$  is identity matrix. If

$P^n = 5I - 8P$ , then  $n$  is :

A. 4

B. 5

C. 6

D. 7

**Answer: C**



[Watch Video Solution](#)

4.  $A$  and  $B$  are two square matrices such that  $A^2B = BA$  and if

$(AB)^{10} = A^k B^{10}$ , then  $k$  is

A. 1001

B. 1023

C. 1042

D. none of these

**Answer: B**



**Watch Video Solution**

5. If matrix  $A = [a_{ij}]_{3 \times 3}$ , matrix  $B = [b_{ij}]_{3 \times 3}$ , where  $a_{ij} + a_{ji} = 0$  and  $b_{ij} - b_{ji} = 0 \forall i, j$ , then  $A^4 \cdot B^3$  is

A. Singular

B. Zero matrix

C. Symmetric

D. Skew-Symmetric matrix

**Answer: A**



Watch Video Solution

6. If  $A \begin{pmatrix} 1 & 3 & 4 \\ 3 & -1 & 5 \\ -2 & 4 & -3 \end{pmatrix} = \begin{pmatrix} 3 & -1 & 5 \\ 1 & 3 & 4 \\ +4 & -8 & 6 \end{pmatrix}$ , then  $A =$

A.  $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}$

B.  $\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

C.  $\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -2 \end{pmatrix}$

D.  $\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -2 \end{pmatrix}$

Answer: D



Watch Video Solution

7. Let  $A = \begin{bmatrix} -5 & -8 & -7 \\ 3 & 5 & 4 \\ 2 & 3 & 3 \end{bmatrix}$  and  $B = \begin{bmatrix} x \\ y \\ 2 \end{bmatrix}$ . If  $AB$  is a scalar multiple of  $B$ , then the value of  $x + y$  is

A.  $-1$

B.  $-2$

C.  $1$

D.  $2$

**Answer: B**

 [Watch Video Solution](#)

8.  $A = \begin{bmatrix} a & b \\ b & -a \end{bmatrix}$  and  $MA = A^{2m}$ ,  $m \in N$  for some matrix  $M$ , then

which one of the following is correct ?

A.  $M = \begin{bmatrix} a^{2m} & b^{2m} \\ b^{2m} & -a^{2m} \end{bmatrix}$

B.  $M = (a^2 + b^2)^m \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

C.  $M = (a^m + b^m) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

D.  $M = (a^2 + b^2)^{m-1} \begin{bmatrix} a & b \\ b & -a \end{bmatrix}$

**Answer: D**



Watch Video Solution

9. If  $A = [a_{ij}]_{m \times n}$  and  $a_{ij} = (i^2 + j^2 - ij)(j - i)$ ,  $n$  odd, then which of the following is not the value of  $Tr(A)$

A. 0

B.  $|A|$

C.  $2|A|$

D. none of these

Answer: D



Watch Video Solution

10.  $|A - B| \neq 0$ ,  $A^4 = B^4$ ,  $C^3A = C^3B$ ,  $B^3A = A^3B$ , then  $|A^3 + B^3 + C^3| =$

A. 0

B. 1

C.  $3|A|^3$

D. 6

**Answer: A**



[Watch Video Solution](#)

11. If  $AB + BA = 0$ , then which of the following is equivalent to  $A^3 - B^3$

A.  $(A - B)(A^2 + AB + B^2)$

B.  $(A - B)(A^2 - AB - B^2)$

C.  $(A + B)(A^2 - AB - B^2)$

D.  $(A + B)(A^2 + AB - B^2)$

**Answer: C**



[Watch Video Solution](#)

12.  $A, B, C$  are three matrices of the same order such that any two are symmetric and the 3<sup>rd</sup> one is skew symmetric. If  $X = ABC + CBA$  and  $Y = ABC - CBA$ , then  $(XY)^T$  is

A. symmetric

B. skew symmetric

C.  $I - XY$

D.  $-YX$

**Answer: D**



**Watch Video Solution**

13. If  $A$  and  $P$  are different matrices of order  $n$  satisfying  $A^3 = P^3$  and  $A^2P = P^2A$  (where  $|A - P| \neq 0$ ) then  $|A^2 + P^2|$  is equal to

A.  $n$

B. 0

C.  $|A||P|$

D.  $|A + P|$

**Answer: B**



[Watch Video Solution](#)

14. Let  $A, B$  are square matrices of same order satisfying  $AB = A$  and  $BA = B$  then  $(A^{2010} + B^{2010})^{2011}$  equals.

A.  $A + B$

B.  $2010(A + B)$

C.  $2011(A + B)$

D.  $2^{2011}(A + B)$

**Answer: D**



[Watch Video Solution](#)



15. The number of  $2 \times 2$  matrices  $A$ , that are there with the elements as real numbers satisfying  $A + A^T = I$  and  $AA^T = I$  is

- A. zero
- B. one
- C. two
- D. infinite

**Answer: C**

 [Watch Video Solution](#)

16. If the orthogonal square matrices  $A$  and  $B$  of same size satisfy  $\det A + \det B = 0$  then the value of  $\det(A + B)$

- A.  $-1$
- B.  $1$

C. 0

D. none of these

**Answer: C**

 [Watch Video Solution](#)

17. If  $A = \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$ ,  $C = ABA^T$ , then

$A^T C^n A$ ,  $n \in I^+$  equals to

A.  $\begin{bmatrix} -n & 1 \\ 1 & 0 \end{bmatrix}$

B.  $\begin{bmatrix} 1 & -n \\ 0 & 1 \end{bmatrix}$

C.  $\begin{bmatrix} 0 & 1 \\ 1 & -n \end{bmatrix}$

D.  $\begin{bmatrix} 1 & 0 \\ -n & 1 \end{bmatrix}$

**Answer: D**

 [Watch Video Solution](#)

18. Let  $A$  be a  $3 \times 3$  matrix given by  $A = (a_{ij})_{3 \times 3}$ . If for every column vector  $X$  satisfies  $X'AX = 0$  and  $a_{12} = 2008$ ,  $a_{13} = 2010$  and  $a_{23} = -2012$ . Then the value of  $a_{21} + a_{31} + a_{32} =$

- A.  $-6$
- B.  $2006$
- C.  $-2006$
- D.  $0$

**Answer: C**



[Watch Video Solution](#)

19. Let  $A$  and  $B$  be two non-singular matrices such that  $A \neq I$ ,  $B^2 = I$  and  $AB = BA^2$ , where  $I$  is the identity matrix, the least value of  $k$  such that  $A^{(k)} = I$  is

- A.  $31$

B. 32

C. 64

D. 63

**Answer: D**



[Watch Video Solution](#)

20. Let  $A$  be a  $2 \times 3$  matrix, whereas  $B$  be a  $3 \times 2$  matrix. If  $\det. (AB) = 4$ , then the value of  $\det. (BA)$  is

A.  $-4$

B. 2

C.  $-2$

D. 0

**Answer: D**



[Watch Video Solution](#)

21. Let  $A$  be a square matrix of order 3 so that sum of elements of each row is 1. Then the sum elements of matrix  $A^2$  is

A. 1

B. 3

C. 0

D. 6

**Answer: B**



[Watch Video Solution](#)

22.  $A$  and  $B$  be  $3 \times 3$  matrices such that  $AB + A + B = 0$ , then

A.  $(A + B)^2 = A^2 + 2AB + B^2$

B.  $|A| = |B|$

C.  $A^2 = B^2$

D. none of these

**Answer: A**



[Watch Video Solution](#)

23. If  $(A + B)^2 = A^2 + B^2$  and  $|A| \neq 0$ , then  $|B| =$  (where  $A$  and  $B$  are matrices of odd order)

A. 2

B.  $-2$

C. 1

D. 0

**Answer: D**



[Watch Video Solution](#)

24. If  $A$  is a square matrix of order 3 such that  $|A| = 5$ , then  $|Adj(4A)| =$

A.  $5^3 \times 4^2$

B.  $5^2 \times 4^3$

C.  $5^2 \times 16^3$

D.  $5^3 \times 16^2$

**Answer: C**



[Watch Video Solution](#)

25. If  $A$  and  $B$  are two non singular matrices and both are symmetric and commute each other, then

A. Both  $A^{-1}B$  and  $A^{-1}B^{-1}$  are symmetric.

B.  $A^{-1}B$  is symmetric but  $A^{-1}B^{-1}$  is not symmetric.

C.  $A^{-1}B^{-1}$  is symmetric but  $A^{-1}B$  is not symmetric.

D. Neither  $A^{-1}B$  nor  $A^{-1}B^{-1}$  are symmetric

**Answer: A**



[Watch Video Solution](#)

26. If  $A$  is a square matrix of order 3 such that  $|A| = 2$ , then

$\left|(\text{adj}A^{-1})^{-1}\right|$  is

A. 1

B. 2

C. 4

D. 8

**Answer: C**



[Watch Video Solution](#)



27. Let matrix  $A = \begin{bmatrix} x & y & -z \\ 1 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$ , where  $x, y, z \in N$ . If

$|adj(adj(adj(adjA)))| = 4^8 \cdot 5^{16}$ , then the number of such  $(x, y, z)$  are

A. 28

B. 36

C. 45

D. 55

**Answer: B**



**Watch Video Solution**

28. A be a square matrix of order 2 with  $|A| \neq 0$  such that  $|A + |A|adj(A)| = 0$ , where  $adj(A)$  is a adjoint of matrix  $A$ , then the value of  $|A - |A|adj(A)|$  is

A. 1

B. 2

C. 3

D. 4

**Answer: D**



[Watch Video Solution](#)

29. If  $A$  is a skew symmetric matrix, then  $B = (I - A)(I + A)^{-1}$  is (where  $I$  is an identity matrix of same order as of  $A$ )

A. idempotent matrix

B. symmetric matrix

C. orthogonal matrix

D. none of these

**Answer: C**



[Watch Video Solution](#)

30. If  $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ , then the trace of the matrix  $Adj(AdjA)$  is

- A. 1
- B. 2
- C. 3
- D. 4

**Answer: A**

 [Watch Video Solution](#)

31. If  $A = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 1 & 0 \end{bmatrix}$  and  $B = (adjA)$  and  $C = 5A$ , then find the value of  $\frac{|adjB|}{|C|}$

- A. 25
- B. 2

C. 1

D. 5

**Answer: C**



[Watch Video Solution](#)

**32.** Let  $A$  and  $B$  be two non-singular square matrices such that  $B \neq I$  and  $AB^2 = BA$ . If  $A^3 = B^{-1}A^3B^n$ , then value of  $n$  is

A. 4

B. 5

C. 8

D. 7

**Answer: C**



[Watch Video Solution](#)

33. If  $A$  is an idempotent matrix satisfying  $(I - 0.4A)^{-1} = I - \alpha A$  where  $I$  is the unit matrix of the same order as that of  $A$  then the value of  $\alpha$  is

A.  $-1/3$

B.  $1/3$

C.  $-2/3$

D.  $2/3$

**Answer: C**



[Watch Video Solution](#)

34. If  $A$  and  $B$  are two non-singular matrices which commute, then

$$\left( A(A + B)^{-1}B \right)^{-1} (AB) =$$

A.  $A + B$

B.  $A^{-1} + B^{-1}$

C.  $A^{-1} + B$

D. none of these

**Answer: A**



**Watch Video Solution**