



MATHS

BOOKS - CENGAGE MATHS (ENGLISH)

SOLUTIONS AND PROPERTIES OF TRIANGLE

Single Correct Answer Type

1. In a triangle $\angle A = 55^\circ$, $\angle B = 15^\circ$, $\angle C = 110^\circ$. Then $c^2 - a^2$ is equal to

A. ab

B. $2ab$

C. $-ab$

D. none of these

Answer: A



Watch Video Solution

2. In a triangle ABC if $2a = \sqrt{3}b + c$, then possible relation is

A. $c^2 = a^2 + b^2 - ab$

B. $a^2 = b^2 + c^2$

C. $b^2 = a^2 + c^2 - ac\sqrt{3}$

D. $a = b = c$

Answer: B



Watch Video Solution

3. A circle of area 20 sq. units is centered at the point O. Suppose $\triangle ABC$ is inscribed in that circle and has area 8 sq. units. The central angles α , β and γ are as shown in the figure. The value of $(\sin \alpha + \sin \beta + \sin \gamma)$ is equal to



A. $\frac{4\pi}{5}$

B. $\frac{3\pi}{4}$

C. $\frac{2\pi}{5}$

D. $\frac{\pi}{4}$

Answer: A



Watch Video Solution

4.



 Watch Video Solution

5. Given a triangle $\triangle ABC$ such that $\sin^2 A + \sin^2 C = 1001 \cdot \sin^2 B$. Then the value of $\frac{2(\tan A + \tan C) \cdot \tan^2 B}{\tan A + \tan B + \tan C}$ is

A. $\frac{1}{2000}$

B. $\frac{1}{1000}$

C. $\frac{1}{500}$

D. $\frac{1}{250}$

Answer: D

 Watch Video Solution

6. $\triangle ABC$ has different side lengths a, b, c . If a^2, b^2, c^2 as sides form another $\triangle PQR$, then $\triangle ABC$ will always be

A. acute angled triangle only

B. obtuse angled triangle only

C. sometimes acute or sometimes obtuse depending on values
of a, b and c

D. none of these

Answer: A

 [Watch Video Solution](#)

7. In $\triangle ABC$, $\angle B = \frac{\pi}{4}$, $\angle C = \frac{\pi}{6}$. D is a point on BC which divides it in the ratio 1 : 3, $\angle DAB = \beta$, then

A. $\left(\sec. \frac{\pi}{6}\right) AB + \cot \beta = \cot\left(\frac{\pi}{6}\right) AC + (\sqrt{3} - 5)$

B. $\left(\sec. \frac{\pi}{4}\right) AB \cot \beta = \cot\left(\frac{\pi}{4}\right) AC(4\sqrt{3} - 5)$

C. $\left(\sec. \frac{\pi}{4}\right) AB + \cot \beta = \cot\left(\frac{\pi}{6}\right) AC + (4\sqrt{3} + 5)$

$$D. \left(\sec. \frac{\pi}{6}\right) AB \cot \beta = \cot\left(\frac{\pi}{4}\right) AC(\sqrt{3} + 5)$$

Answer: B

 [Watch Video Solution](#)

8. The acute angle of a rhombus whose side is a mean proportional between its diagonals is

A. 15°

B. 20°

C. 30°

D. 80°

Answer: C

 [Watch Video Solution](#)

9. Let side a, b and c of $\triangle ABC$ be related by the relation $a : b : c = 3 : 5 : 4$. Altitudes AD, BE and CF are dropped on BC, CA and AB , respectively. If $P_1D + P_2E + P_3F = 42$, then the value of $a + b + c$ is

A. 1200

B. 120

C. 12

D. none of these

Answer: B



[Watch Video Solution](#)

10. Triangle ABC is right angle at A . The points P and Q are on hypotenuse BC such that $BP = PQ = QC$. if $AP = 3$ and $AQ = 4$, then length BC is equal to

A. $3\sqrt{5}$

B. $5\sqrt{3}$

C. $4\sqrt{5}$

D. 7

Answer: A



[Watch Video Solution](#)

11. ABC is a right angled triangle of which A is the right angle, BD is drawn perpendicular to BC meets CA produced in D. If $AB = 12$, $AC = 16$, $BC = 20$, then $BD =$

A. 15

B. 25

C. 10

D. 225

Answer: A

 [Watch Video Solution](#)

12. In a $\triangle ABC$, the median AD is perpendicular to AC . If $b = 5$ and $c = 11$, then $a =$

A. 10

B. 12

C. 14

D. $\sqrt{221}$

Answer: C

 [Watch Video Solution](#)

13. ABC is an equilateral triangle where $AB = a$ and P is any point in its plane such that $PA = PB + PC$. Then $\frac{PA^2 + PB^2 + PC^2}{a^2}$ is

A. 3

B. $\frac{\sqrt{3}}{4}$

C. $\frac{3}{4}$

D. 2

Answer: D



Watch Video Solution

14. In ΔABC if $\tan\left(\frac{A}{2}\right)\tan\left(\frac{B}{2}\right) + \tan\left(\frac{B}{2}\right)\tan\left(\frac{C}{2}\right) = \frac{2}{3}$

then $a + c$

A. $3b$

B. $2b$

C. $\frac{3b}{2}$

D. $4b$

Answer: B



Watch Video Solution

15. In a triangle ABC if $\tan. \frac{A}{2} \tan. \frac{B}{2} = \frac{1}{3}$ and $ab = 4$, then the value of c can be

A. 1

B. 1.5

C. 2.5

D. none of these

Answer: C



Watch Video Solution

16. In a triangle ABC if

$$\cot\left(\frac{A}{2}\right)\cot\left(\frac{B}{2}\right) = c, \cot\left(\frac{B}{2}\right)\cot\left(\frac{C}{2}\right) = a \quad \text{and}$$

$$\cot\left(\frac{C}{2}\right)\cot\left(\frac{A}{2}\right) = b \text{ then } \frac{1}{s-a} + \frac{1}{s-b} + \frac{1}{s-c} =$$

A. -1

B. 0

C. 1

D. 2

Answer: D



Watch Video Solution

17. The perimeter of a triangle ABC right angled at C is 70 and the inradius is 6, then $|a - b| =$

A. 1

B. 2

C. 8

D. 9

Answer: A



[Watch Video Solution](#)

18. The diagonals of a parallelogram are inclined to each other at an angle of 45° , while its sides a and b ($a > 0$) are inclined to each other at an angle of 30° , then the value of $\frac{a}{b}$ is

A. $\frac{3}{2}$

B. $\frac{3 + \sqrt{5}}{2}$

C. $\frac{3 + \sqrt{5}}{4}$

D. $\frac{\sqrt{5} + 1}{2}$

Answer: D



Watch Video Solution

19. In a triangle ABC if $2\Delta^2 = \frac{a^2b^2c^2}{a^2 + b^2 + c^2}$, then it is

A. equilateral

B. isosceles but not right angled

C. isosceles right angled

D. right angled

Answer: D



Watch Video Solution

20. If in any triangle, the area $\Delta ABC \leq \frac{b^2 + c^2}{\lambda}$, then the largest possible numerical value of λ is

- A. 1
- B. 2
- C. 3
- D. 4

Answer: D



Watch Video Solution

21. In any triangle ABC, if $2\Delta a - b^2c = c^3$, (where Δ is the area of triangle), then which of the following is possible ?

A. B is obtuse

B. A is obtuse

C. C is obtuse

D. B is right angle

Answer: B



Watch Video Solution

22. In ΔABC , $a^2(s - a) + b^2(s - b) + c^2(s - c) =$

A. $4R\Delta(\cos A + \sin B + \cos C)$

B. $4R\Delta(\sin A + \sin B + \sin C)$

C. $4R\Delta\left(1 + 4\sin. \frac{A}{2}\sin. \frac{B}{2}\sin. \frac{C}{2}\right)$

D. none of these

Answer: C



Watch Video Solution

23. Let ABC be an equilateral triangle, let KLMN be a rectangle with K, L on BC, M on AC and N on AB. Suppose $AN/NB = 2$ and the area of triangle BKN is 6. The area of the triangle ABC is -

A. 54

B. 108

C. 48

D. none of these

Answer: B



Watch Video Solution

24. ABC is an acute angled triangle with circumcenter O and orthocentre H. If $AO=AH$, then find the angle A.

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: C



Watch Video Solution

25. In a triangle ABC if $\angle ABC = 60^\circ$, then

$$\left(\frac{AB - BC + CA}{r} \right)^2 =$$

A. 10

B. 11

C. 12

D. 14

Answer: C



Watch Video Solution

26. The area of an acute triangle ABC is Δ , the area of its pedal triangle is 'p', where $\cos B = \frac{2p}{\Delta}$ and $\sin B = \frac{2\sqrt{3}p}{\Delta}$. The value of $8(\cos^2 A \cos B + \cos^2 C)$ is

A. 1

B. 2

C. 3

D. none of these

Answer: C



Watch Video Solution

27. In a triangle ABC , D is a point on BC such that AD is the internal bisector of $\angle A$. Let $\angle B = 2\angle C$ and $CD = AB$. Then $\angle A$ is

A. 18°

B. 36°

C. 54°

D. 72°

Answer: D



Watch Video Solution

28. In $\triangle ABC$, circumradius is 3 and radius is 1.5 units. The value of $a \cot^2 A + b^2 \cot^3 B + c^3 \cot^4 C$ is

A. $13\sqrt{3}$

B. $11\sqrt{6}$

C. 21

D. none of these

Answer: A



Watch Video Solution

29. AD, BE, CF are internal angular bisectors of ΔABC and I is the incentre.

If
$$a(b+c)\sec\frac{A}{2}ID + b(a+c)\sec\frac{B}{2}IE + c(a+b)\sec\frac{C}{2}IF = kabc$$

, then the value of k is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

30. In $\triangle ABC$ it is given distance between the circumcentre (O) and orthocentre (H) is $R\sqrt{1 - 8 \cos A \cos B \cos C}$. If Q is the midpoint of OH, then AQ is

A. $\frac{R}{2} \sqrt{1 + 8 \cos A \sin B \sin C}$

B. $R\sqrt{1 + 8 \cos A \sin B \sin C}$

C. $2R\sqrt{1 + 8 \cos A \sin B \sin C}$

D. $\frac{R}{2} \sqrt{1 + 8 \sin A \cos B \cos C}$

Answer: A



Watch Video Solution

31. In any $\triangle ABC$ line joining circumcentre (O) and incentre (I) is parallel to AC, then OI is equal to

A. $R \left| \tan \left(\frac{A - C}{2} \right) \right|$

B. $R |\tan(A - C)|$

C. $R \left| \sec \left(\frac{A - C}{2} \right) \right|$

D. $R |\sec(A - C)|$

Answer: A



Watch Video Solution

32. Let the incircle of a $\triangle ABC$ touches sides BC, CA and AB at D, E and F, respectively. Let area of $\triangle ABC$ be Δ and that of DEF be Δ' .

If a , b and c are side of ΔABC , then the value of

$$abc(a + b + c) \frac{\Delta'}{\Delta^3} \text{ is}$$

A. 1

B. 2

C. 3

D. 4

Answer: D



[Watch Video Solution](#)

33. Let H be the orthocentre of triangle ABC . Then angle subtended

by side BC at the centre of incircle of ΔCHB is

A. $\frac{A}{2} + 90^\circ$

B. $\frac{B + C}{2} + 90^\circ$

C. $\frac{B - C}{2} + 90^\circ$

D. none of these

Answer: B



[Watch Video Solution](#)

34. If in a triangle ABC , $r_1 + r_2 + r_3 = 9r$, then the triangle is necessarily

A. right angled

B. equilateral

C. obtuse angled

D. none of these

Answer: B



[Watch Video Solution](#)

35. In triangle ABC , $r = \frac{R}{6}$ and $r_1 = 7r$. Then the measure of angle $A =$

A. $\frac{\pi}{12}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{3}$

Answer: D

 [Watch Video Solution](#)

36. $(r_2 + r_3) \sqrt{\frac{rr_1}{r_2r_3}} =$

A. a

B. b

C. c

D. bc

Answer: A



Watch Video Solution

37. In $\triangle ABC$, right angled at A, $\cos^{-1}\left(\frac{R}{r_2 + r_3}\right)$ is

A. 30°

B. 60°

C. 90°

D. 45°

Answer: B



Watch Video Solution

38. In $\triangle ABC$ if $r_1 = 2r_2 = 3r_3$ and D is the mid point of BC then

$\cos \angle ADC$ is (a) $\frac{7}{25}$ (b) $-\frac{7}{25}$ (c) $\frac{24}{25}$ (d) $-\frac{24}{25}$

A. $\frac{7}{25}$

B. $-\frac{7}{25}$

C. $\frac{24}{25}$

D. $-\frac{24}{25}$

Answer: D



Watch Video Solution

39. In the ambiguous case if the remaining angles of a triangle with

given a, b, A and B_1, B_2, C_1, C_2 then $\frac{\sin C_1}{\sin B_1} + \frac{\sin C_2}{\sin B_2} =$

A. $2 \cos A$

B. $2 \sin B$

C. $2 \tan A$

D. $2 \cot A$

Answer: A



[Watch Video Solution](#)

40. Two sides of a triangle are of lengths $\sqrt{6}$ and 4 and the angle opposite to smaller side is 30° . How many such triangles are possible? Find the length of their third side and area.

A. 0

B. 1

C. 2

D. infinite

Answer: C



Watch Video Solution

41. If circumradius of triangle is 2, then the maximum value of

$$\frac{abc}{a + b + c} \text{ is}$$

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

42. Let ABC and AB'C be two non-congruent triangles with sides $BC=B'C=5$, $AC=6$, and $\angle A$ is fixed. If A_1 and A_2 are the area of the two triangles ABC and AB'C, then the value of $\frac{A_1^2 + A_2^2 - 2A_1A_2 \cos 2A}{(A_1 + A_2)^2}$ is

- A. $9/36$
- B. $25/36$
- C. $25/16$
- D. $16/25$

Answer: B



[Watch Video Solution](#)

Multiple Correct Answers Type

1. Let a, b, c be the sides of a triangle ABC , $a=2c, \cos(A-C)+\cos B=1$.

then the value of C is

A. $\pi / 6$

B. $\pi / 3$

C. $2\pi / 3$

D. $5\pi / 6$

Answer: A:D



[Watch Video Solution](#)

2. If A, B, C are the angles of a triangle such that

$$\sin^2 A + \sin^2 B = \sin^2 C, \text{ then}$$

A. $\sin A + \sin B > 1$

B. $\tan A \tan B = 1$

C. $\sin A + \sin B = 1$

D. $\tan A \tan B < 1$

Answer: A::B



Watch Video Solution

3. In $\triangle ABC$, $\angle C = 2\angle A$ and $AC=2BC$. Then which of the following is/are True ?

A. Angles A,B,C are in arithmetic progression

B. Angles A,C,B are in arithmetic progression

C. $\triangle ABC$ is a right angled isosceles triangle

D. $BC^2 + CA^2 + AB^2 = 8R^2$, where R is the circum-radius of

$\triangle ABC$

Answer: B::D



Watch Video Solution

4. ΔABC is isosceles with $AB=AC=7\text{cm}$ and $BC=9\text{cm}$. The height AD from A to BC , is 6cm . Find the area of ΔABC . What will be the height from C to AB i.e., CE ?



Watch Video Solution

5. If in a triangle ABC , θ is the angle determined by $\cos \theta = (a - b) / c$, then

A.
$$\frac{(a + b)\sin \theta}{2\sqrt{ab}} = \frac{\cos(A - B)}{2}$$

B.
$$\frac{(a + b)\sin \theta}{2\sqrt{ab}} = \frac{\cos(A + B)}{2}$$

C.
$$\frac{c \sin \theta}{2\sqrt{ab}} = \frac{\cos(A - B)}{2}$$

$$D. \frac{c \sin \theta}{2\sqrt{ab}} = \frac{\cos(A + B)}{2}$$

Answer: A::D

 [Watch Video Solution](#)

6. Let a, b, c be the sides BC, CA, AB of $\triangle ABC$ on xy plane. If abscissa and ordinate of vertices of the triangle are integers and R is the circumradius, then $2R$ can be equal to

A. $\frac{8}{9}abc$

B. abc

C. $\frac{9}{8}abc$

D. $\frac{abc}{2}$

Answer: A::B::D

 [Watch Video Solution](#)

7. In a ΔABC , if $\tan. \frac{A}{2} = \frac{5}{6}$, $\tan. \frac{B}{2} = \frac{20}{37}$, then which of the following is/are correct ?

A. $\angle B > \angle C$

B. $\angle B < \angle C$

C. $a > b > c$

D. $a < b < c$

Answer: A:C



Watch Video Solution

8. If area of $\Delta ABC(\Delta)$ and angle C are given and if c opposite to given angle is minimum, then

A. $a = \sqrt{\frac{2\Delta}{\sin C}}$

$$B. b = \sqrt{\frac{2\Delta}{\sin C}}$$

$$C. a = \frac{4\Delta}{\sin C}$$

$$D. b = \frac{4\Delta}{\sin^2 C}$$

Answer: A:B



Watch Video Solution

9. Let 'P' be an interior point of $\triangle ABC$. If $\angle A = 45^\circ$, $\angle B = 60^\circ$ and $\angle C = 75^\circ$. If $X = \text{area of } \triangle PBC$, $Y = \text{area of } \triangle PAC$ and $Z = \text{area of } \triangle PAB$, then which of the following ratios is/are true ?

A. If P is the centroid, then $X : Y : Z$ is $1 : 1 : 1$

B. If P is the incentre, then $X : Y : Z$ is $2 : \sqrt{6} : (\sqrt{3} + 1)$

C. If P the orthocentre, then $X : Y : Z$ is $1 : \sqrt{3} : (2 + \sqrt{3})$

D. If P is the circumcentre, then $X : Y : Z$ is $2 : \sqrt{3} : 1$

Answer: A::B::C::D



View Text Solution

10. Let 'l' is the length of median from the vertex A to the side BC of a ΔABC . Then

A. $4l^2 = 2b^2 + 2c^2 - a^2$

B. $4l^2 = b^2 + c^2 + 2bc \cos A$

C. $4l^2 = a^2 + 4bc \cos A$

D. $4l^2 = (2s - a)^2 \sin^2 \frac{A}{2}$

Answer: A::B::C::D



Watch Video Solution

11. A circle having centre as O' and radius r' touches the incircle of $\triangle ABC$ externally at F , where F is on BC and also touches its circumcircle internally at G . If O is the circumcentre of $\triangle ABC$ and I is its incentre, then

A. $OO' = R - r'$

B. Perpendicular distance from O to line joining IO' is $\left| \frac{b - c}{2} \right|$

C. Projection of OO' on line joining $IO' = r' + R \cos A$

D. $r' = \frac{\Delta}{a} \tan^2 A$

Answer: A::B::C::D



[View Text Solution](#)

12. In triangle ABC , if $r_1 + r_2 = 3R$ and $r_2 + r_3 = 2R$, then

A. $\angle A = 90^\circ$

B. $\angle B = 45^\circ$

C. $\angle C = 60^\circ$

D. triangle ABC is right angled isosceles

Answer: A:C



Watch Video Solution

13. The radii r_1, r_2, r_3 of the escribed circles of the triangle ABC are in H.P. If the area of the triangle is 24cm^2 and its perimeter is 24cm, then the length of its largest side is 10 (b) 9 (c) 8 (d) none of these

A. The length of longest side of triangle ABC is equal to 10

B. The radius of circle inscribed in triangle ABC is equal to 4.

C. The circumradius of triangle ABC is equal to 5.

D. The sides of triangle ABC are in A.P.

Answer: A::C::D

 [Watch Video Solution](#)

Comprehension Type

1. Let ABC be a triangle in which the line joining the circumcentre and incentre is parallel to base BC of the triangle. Then answer the following questions :

Then range of $\angle A$ is

A. $\left[\frac{\pi}{6}, \frac{\pi}{3} \right]$

B. $\left[\frac{\pi}{3}, \frac{\pi}{2} \right)$

C. $\left[\frac{\pi}{3}, \frac{2\pi}{3} \right] = \left\{ \frac{\pi}{3} \right\}$

D. $\left[0, \frac{\pi}{2} \right]$

Answer: B



Watch Video Solution

2. Let ABC be a triangle in which the line joining the circumcentre and incentre is parallel to base BC of the triangle. Then answer the following questions :

If ODEI is a square where O and I stands for circumcentre and incentre, respectively and D and E are the point of perpendicular from O and I on the base BC, then

A. $\frac{r}{R} = \frac{3}{8}$

B. $\frac{r}{R} = 2 - \sqrt{3}$

C. $\frac{r}{R} = \sqrt{2} - 1$

D. $\frac{r}{R} = \frac{1}{4}$

Answer: C



[Watch Video Solution](#)

3. Let ABC be a triangle in which the line joining the circumcentre and incentre is parallel to base BC of the triangle. Then answer the following questions :

If $\angle A = 60^\circ$, then $\triangle ABC$ is

- A. isosceles
- B. right angled
- C. right angled isosceles
- D. equilateral

Answer: D



[Watch Video Solution](#)

4. Incircle of $\triangle ABC$ touches AB, BC, CA at R, P, Q, respectively. If

$$\frac{2}{AR} + \frac{5}{BP} + \frac{5}{CQ} = \frac{6}{r}$$
 and the perimeter of the triangle is the

smallest integer, then answer the following questions :

$\triangle ABC$ is

- A. scalene
- B. isosceles
- C. equilateral
- D. right angled

Answer: B

 [Watch Video Solution](#)

5. Incircle of $\triangle ABC$ touches AB, BC, CA at R, P, Q, respectively. If

$$\frac{2}{AR} + \frac{5}{BP} + \frac{5}{CQ} = \frac{6}{r}$$
 and the perimeter of the triangle is the

smallest integer, then answer the following questions :

The inradius of incircle of ΔABC is

A. 4

B. 3

C. 2

D. 1

Answer: C



[Watch Video Solution](#)

6. Incircle of ΔABC touches AB, BC, CA at R, P, Q, respectively. If

$\frac{2}{AR} + \frac{5}{BP} + \frac{5}{CQ} = \frac{6}{r}$ and the perimeter of the triangle is the

smallest integer, then answer the following questions :

The area of ΔABC is

A. 15 sq. units

B. 21 sq. units

C. 24 sq. units

D. 27 sq. units

Answer: D



Watch Video Solution