



## MATHS

### BOOKS - CENGAGE MATHS (ENGLISH)

#### STRAIGHT LINE

##### Illustration

1. Find the equation of line passing through point (2,3) which is

(i) parallel of the x-axis

(ii) parallel to the y-axis



[Watch Video Solution](#)

2. Find the equation of line passing through point (2,-5) which is

(i) parallel to the line  $3x + 2y - 4 = 0$

(ii) perpendicular to the line  $3x + 2y - 4 = 0$

 [Watch Video Solution](#)

3. Find the equation of the perpendicular bisector of the line segment joining the points  $A(2, 3)$  and  $B(6, -5)$ .

 [Watch Video Solution](#)

4. Find the locus of a point  $P$  which moves such that its distance from the line  $y = \sqrt{3}x - 7$  is the same as its distance from  $(2\sqrt{3}, -1)$

 [Watch Video Solution](#)

5. Consider a triangle with vertices  $A(1, 2)$ ,  $B(3, 1)$ , and  $C(-3, 0)$ . Find the equation of altitude through vertex  $A$ . the equation of median through vertex  $A$ . the equation of internal angle bisector of  $\angle A$ .

 [Watch Video Solution](#)

6. Find the coordinates of the foot of the perpendicular drawn from the point P(1,-2) on the line  $y = 2x + 1$ . Also, find the image of P in the line.

 [Watch Video Solution](#)

7. If the line  $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 1$  moves in such a way that  $\left(\frac{1}{a^2}\right) + \left(\frac{1}{b^2}\right) = \left(\frac{1}{c^2}\right)$ , where  $c$  is a constant, prove that the foot of the perpendicular from the origin on the straight line describes the circle  $x^2 + y^2 = c^2$ .

 [Watch Video Solution](#)

8. In what ratio does the line joining the points (2, 3) and (4, 1) divide the segment joining the points (1, 2) and (4, 3)?

 [Watch Video Solution](#)

9.  $ABCD$  is a square whose vertices are  $A(0, 0)$ ,  $B(2, 0)$ ,  $C(2, 2)$ , and  $D(0, 2)$ . The square is rotated in the  $XY$  - plane through an angle  $30^\circ$  in the anticlockwise sense about an axis passing through  $A$  perpendicular to the  $XY$  - plane. Find the equation of the diagonal  $BD$  of this rotated square.

 [Watch Video Solution](#)

10. In a triangle  $ABC$ , side  $AB$  has equation  $2x + 3y = 29$  and side  $AC$  has equation  $x + 2y = 16$ . If the midpoint of  $BC$  is  $(5, 6)$ , then find the equation of  $BC$ .

 [Watch Video Solution](#)

11. Two consecutive sides of a parallelogram are  $4x + 5y = 0$  and  $7x + 2y = 0$ . If the equation of one diagonal is  $11x - 7y = 9$ , find the equation of the other diagonal.

 [Watch Video Solution](#)



12. If one of the sides of a square is  $3x - 4y - 12 = 0$  and the center is  $(0, 0)$ , then find the equations of the diagonals of the square.

 [Watch Video Solution](#)

13. A vertex of an equilateral triangle is  $2, 3$  and the opposite side is  $x + y = 2$ . Find the equations of other sides.

 [Watch Video Solution](#)

14. A line  $4x + y = 1$  passes through the point  $A(2, -7)$  and meets line BC at B whose equation is  $3x - 4y + 1 = 0$ , the equation of line AC such that  $AB = AC$  is (a)  $52x + 89y + 519 = 0$  (b)  $52x + 89y - 519 = 0$  (c)  $82x + 52y + 519 = 0$  (d)  $89x + 52y - 519 = 0$

 [Watch Video Solution](#)

15. A ray of light is sent along the line  $x - 2y - 3 = 0$  upon reaching the line  $3x - 2y - 5 = 0$ , the ray is reflected from it. Find the equation of the line containing the reflected ray.

 [Watch Video Solution](#)

16. Find the equation of the line which intersects the y-axis at a distance of 2 units above the origin and makes an angle of  $30^\circ$  with the positive direction of the x-axis.

 [Watch Video Solution](#)

17. Find the equation of a straight line cutting off an intercept -1 from y-axis and being equally inclined to the axes.

 [Watch Video Solution](#)

18. Find the equation of a line that has  $y$ -intercept 4 and is a perpendicular to the line joining  $(2, -3)$  and  $(4, 2)$ .

 [Watch Video Solution](#)

19. Find equation of the line passing through the point  $(2, 2)$  and cutting off intercepts on the axes whose sum is 9.

 [Watch Video Solution](#)

20. Find the equation of the straight line that (i) makes equal intercepts on the axes and passes through the point  $(2;3)$  (ii) passes through the point  $(-5;4)$  and is such that the portion intercepted between the axes is divided by the point in the ratio 1: 2

 [Watch Video Solution](#)

21. Line segment AB of fixed length  $c$  slides between coordinate axes such that its ends A and B lie on the axes. If O is origin and rectangle OAPB is completed, then show that the locus of the foot of the perpendicular drawn from P to AB is  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = c^{\frac{2}{3}}$ .

 [Watch Video Solution](#)

22. Reduce the line  $2x - 3y + 5 = 0$  in slope-intercept, intercept, and normal forms.

 [Watch Video Solution](#)

23. Find the equation of the line which satisfy the given conditions : Perpendicular distance from the origin is 5 units and the angle made by the perpendicular with the positive xaxis is  $30^\circ$  .

 [Watch Video Solution](#)

24. A straight line is drawn through the point  $P(2;3)$  and is inclined at an angle of  $30^\circ$  with the x-axis . Find the coordinates of two points on it at a distance 4 from point P.

 [Watch Video Solution](#)

25. The line joining two points  $A(2,0)$  and  $B(3,1)$  is rotated about A in anticlockwise direction through an angle of  $15^\circ$  . find the equation of line in the new position. If b goes to c in the new position what will be the coordinates of C.

 [Watch Video Solution](#)

26. A line through point  $A(1,3)$  and parallel to the line  $x-y+1 = 0$  meets the line  $2x-3y + 9 = 0$  at point P. Find distance AP without finding point P.

 [Watch Video Solution](#)

27. Two adjacent vertices of a square are  $(1, 2)$  and  $(-2, 6)$  Find the other vertices.

 [Watch Video Solution](#)

28. A Line through the variable point  $A(1 + k; 2k)$  meets the lines  $7x + y - 16 = 0$ ;  $5x - y - 8 = 0$  and  $x - 5y + 8 = 0$  at B;C;D respectively.

Prove that AC;AB and AD are in HP.

 [Watch Video Solution](#)

29. if  $P$  is the length of perpendicular from origin to the line  $\frac{x}{a} + \frac{y}{b} = 1$  then prove that  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2}$

 [Watch Video Solution](#)

**30.** Find the coordinates of a point on  $x + y + 3 = 0$ , whose distance from  $x + 2y + 2 = 0$  is  $\sqrt{5}$ .

 [Watch Video Solution](#)

**31.** Find the least and greatest values of the distance of the point  $(\cos \theta, \sin \theta)$ ,  $\theta \in R$ , from the line  $3x - 4y + 10 = 0$ .

 [Watch Video Solution](#)

**32.** Prove that the product of the lengths of the perpendiculars drawn from the points  $(\sqrt{a^2 - b^2}, 0)$  and  $(-\sqrt{a^2 - b^2}, 0)$  to the line  $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$  is  $b^2$ .

 [Watch Video Solution](#)

33. Find the least value of  $(x - 1)^2 + (y - 2)^2$  under the condition  $3x + 4y - 2 = 0$ .

 [Watch Video Solution](#)

34.  $ABC$  is an equilateral triangle with  $A(0, 0)$  and  $B(a, 0)$ , ( $a > 0$ ).  $L$ ,  $M$  and  $N$  are the foot of the perpendiculars drawn from a point  $P$  to the side  $AB$ ,  $BC$ , and  $CA$ , respectively. If  $P$  lies inside the triangle and satisfies the condition  $PL^2 = PM^2 + PN^2$ , then find the locus of  $P$ .

 [Watch Video Solution](#)

35. Line  $L$  has intercepts  $a$  and  $b$  on the coordinate axes. When the axes are rotated through a given angle keeping the origin fixed, the same line  $L$  has intercepts  $p$  and  $q$ . Then  $a^2 + b^2 = p^2 + q^2$  (a)  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$  (b)  $a^2 + p^2 = b^2 + q^2$  (c)  $\frac{1}{a^2} + \frac{1}{p^2} = \frac{1}{b^2} + \frac{1}{q^2}$  (d)  $\frac{1}{a^2} + \frac{1}{q^2} = \frac{1}{b^2} + \frac{1}{p^2}$

A. (a)  $a^2 + b^2 = p^2 + q^2$



$$B. (b) \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$$

$$C. (c) a^2 + p^2 = b^2 + q^2$$

$$D. (d) \frac{1}{a^2} + \frac{1}{p^2} = \frac{1}{b^2} + \frac{1}{q^2}$$

**Answer:**

 [Watch Video Solution](#)

**36.** Two sides of a square lie on the lines  $x + y = 1$  and  $x + y + 2 = 0$ .

What is its area?

 [Watch Video Solution](#)

**37.** Find equation of the line which is equidistant from parallel lines

$$9x + 6y - 7 = 0 \text{ and } 3x + 2y + 6 = 0.$$

 [Watch Video Solution](#)

38. If one side of the square is  $2x - y + 6 = 0$ , then one of the vertices is  $(2, 1)$ . Find the other sides of the square.

 [Watch Video Solution](#)

39. Prove that the area of the parallelogram contained by the lines  $4y - 3x - a = 0$ ,  $3y - 4x + a = 0$ ,  $4y - 3x - 3a = 0$ , and  $3y - 4x + 2a = 0$  is  $\left(\frac{2}{7}\right)a^2$ .

 [Watch Video Solution](#)

40. The equation of straight line passing through  $(-2, -7)$  and having an intercept of length 3 between the straight lines :  $4x + 3y = 12$  ,  $4x + 3y = 3$  are : (A)  $7x + 24y + 182 = 0$  (B)  $7x + 24y + 18 = 0$  (C)  $x + 2 = 0$  (D)  $x - 2 = 0$

 [Watch Video Solution](#)

41. A line  $L$  is drawn from  $P(4, 3)$  to meet the lines  $L_1$  and  $L_2$  given by  $3x + 4y + 5 = 0$  and  $3x + 4y + 15 = 0$  at points  $A$  and  $B$ , respectively. From  $A$ , a line perpendicular to  $L$  is drawn meeting the line  $L_2$  at  $A_1$ . Similarly, from point  $B$ , a line perpendicular to  $L$  is drawn meeting the line  $L_1$  at  $B_1$ . Thus, a parallelogram  $AA_1BB_1$  is formed. Then the equation of  $L$  so that the area of the parallelogram  $AA_1BB_1$  is the least is  $x - 7y + 17 = 0$   $7x + y + 31 = 0$   
 $x - 7y - 17 = 0$   $x + 7y - 31 = 0$



Watch Video Solution

42. Are the points  $(3, 4)$  and  $(2, -6)$  on the same or opposite sides of the line  $3x - 4y = 8$ ?



Watch Video Solution

43. Find the set of positive values of  $b$  for which the origin and the point  $(1, 1)$  lie on the same side of the straight line,  $a^2x + aby + 1 = 0, \forall a \in R, b > 0$



Watch Video Solution

44. If the point  $(a^2, a + 1)$  lies in the angle between the lines  $3x - y + 1 = 0$  and  $x + 2y - 5 = 0$  containing the origin, then find the value of  $a$ .



Watch Video Solution

45. If the point  $(a, a)$  is placed in between the lines  $|x + y| = 4$ , then find the values of  $a$ .



Watch Video Solution

46. The complete set of real values of 'a' such that the point lies triangle  $p(a, \sin a)$  lies inside the triangle formed by the lines  $x - 2y + 2 = 0$ ;  $x + y = 0$  and  $x - y - \pi = 0$



Watch Video Solution

47. Determine all the values of  $\alpha$  for which the point  $(\alpha, \alpha^2)$  lies inside the triangle formed by the lines.  $2x + 3y - 1 = 0$   $x + 2y - 3 = 0$   
 $5x - 6y - 1 = 0$

 [Watch Video Solution](#)

48. Sketch the origin in which the points satisfying the following inequality lie.

(i)  $2x - 3y - 5 > 0$       (ii)  $-3x + 4y + 7 > 0$

(iii)  $x > 2$       (iv)  $y > -3$

 [Watch Video Solution](#)

49. Sketch the origin in which the points satisfying the following inequalities lie.

(i)  $|x + y| < 2$       (ii)  $|2x - y| > 3$       (iii)  $|x| > |y|$

 [Watch Video Solution](#)

50. Find the values of  $b$  for which the points  $(2b + 3, b^2)$  lies above of the line  $3x - 4y - a(a - 2) = 0 \quad \forall a \in R$ .

 [Watch Video Solution](#)

51. Plot the region of the points  $P(x, y)$  satisfying  $|x| + |y| < 1$ .

 [Watch Video Solution](#)

52. Plot the region of the points  $P(x, y)$  satisfying  $2 > \max\{|x|, |y|\}$ .

 [Watch Video Solution](#)

53. IF one of the vertices of a square is  $(3, 2)$  and one of the diagonals is along the line  $3x + 4y + 8 = 0$ , then find the centre of the square and other

vertices.



[Watch Video Solution](#)

54. In  $\triangle ABC$ , vertex A is (1, 2). If the internal angle bisector of  $\angle B$  is  $2x - y + 10 = 0$  and the perpendicular bisector of AC is  $y = x$ , then find the equation of BC



[Watch Video Solution](#)

55. Find the locus of image of the variable point  $(\lambda^2, 2\lambda)$  in the line mirror  $x-y+1=0$ , where  $\lambda$  is a parameter.



[Watch Video Solution](#)

56. Lines  $L_1 \equiv ax + by + c = 0$  and  $L_2 \equiv lx + my + n = 0$  intersect at the point  $P$  and make an angle  $\theta$  with each other. Find the equation of a

line different from  $L_2$  which passes through  $P$  and makes the same angle  $\theta$  with  $L_1$ .

 [Watch Video Solution](#)

57. For the straight lines  $4x + 3y - 6 = 0$  and  $5x + 12y + 9 = 0$ , find the equation of the bisector of the obtuse angle between them, bisector of the acute angle between them, and bisector of the angle which contains  $(1, 2)$

 [Watch Video Solution](#)

58. The equations of bisectors of two lines  $L_1$  &  $L_2$  are  $2x - 16y - 5 = 0$  and  $64x + 8y + 35 = 0$ . If the line  $L_1$  passes through  $(-11, 4)$ , the equation of acute angle bisector of  $L_1$  &  $L_2$  is:

 [Watch Video Solution](#)



59. If  $x + y = 0$  is the angle bisector of the angle containing the point  $(1,0)$ , for the line  $3x + 4y + b = 0$ ;  $4x + 3y + b = 0$ ,  $4x + 3y - , b = 0$  then



Watch Video Solution

60. Two equal sides of an isosceles triangle are given by  $7x - y + 3 = 0$  and  $x + y = 3$ , and its third side passes through the point  $(1, -10)$ . Find the equation of the third side.



Watch Video Solution

61. The vertices  $B$  and  $C$  of a triangle  $ABC$  lie on the lines  $3y = 4x$  and  $y = 0$ , respectively, and the side  $BC$  passes through the point  $\left(\frac{2}{3}, \frac{2}{3}\right)$ . If  $ABOC$  is a rhombus lying in the first quadrant,  $O$  being the origin, find the equation of the line  $BC$ .



Watch Video Solution

62. Two sides of a rhombus lying in the first quadrant are given by  $3x - 4y = 0$  and  $12x - 5y = 0$ . If the length of the longer diagonal is 12, then find the equations of the other two sides of the rhombus.

 [Watch Video Solution](#)

63. If the line  $ax + by = 1$  passes through the point of intersection of  $y = x \tan \alpha + p \sec \alpha$ ,  $y \sin(30^\circ - \alpha) - x \cos(30^\circ - \alpha) = p$ , and is inclined at  $30^\circ$  with  $y = \tan \alpha x$ , then prove that  $a^2 + b^2 = \frac{3}{4p^2}$ .

 [Watch Video Solution](#)

64. Find the value of  $\lambda$ , if the lines  $3x - 4y - 13 = 0$ ,  $8x - 11y - 33 = 0$  and  $2x - 3y + \lambda = 0$  are concurrent.

 [Watch Video Solution](#)

65. If the lines  $a_1x + b_1y + 1 = 0$ ,  $a_2x + b_2y + 1 = 0$  and  $a_3x + b_3y + 1 = 0$  are concurrent, show that the points  $(a_1, b_1)$ ,  $(a_2, b_2)$  and  $(a_3, b_3)$  are collinear.

 [Watch Video Solution](#)

66. Show that the straight lines given by  $x(a + 2b) + y(a + 3b) = a + b$  for different values of  $a$  and  $b$  pass through a fixed point.

 [Watch Video Solution](#)

67. Let  $ax + by + c = 0$  be a variable straight line, where  $a$ ,  $b$  and  $c$  are the 1st, 3rd, and 7th terms of an increasing AP, respectively. Then prove that the variable straight line always passes through a fixed point. Find that point.

 [Watch Video Solution](#)

**68.** Prove that all the lines having the sum of the intercepts on the axes equal to half of the product of the intercepts pass through the point. Find the fixed point.



[Watch Video Solution](#)

**69.** Find the straight line passing through the point of intersection of  $2x + 3y + 5 = 0$ ,  $5x - 2y - 16 = 0$ , and through the point  $(-1, 3)$ .



[Watch Video Solution](#)

**70.** Consider a family of straight lines  $(x + y) + \lambda(2x - y + 1) = 0$ . Find the equation of the straight line belonging to this family that is farthest from  $(1, -3)$ .



[Watch Video Solution](#)

71. Let the sides of a parallelogram be  $U=a$ ,  $U=b$ ,  $V=a'$  and  $V=b'$ , where  $U=lx+my+n$ ,  $V=l'x+m'y+n'$ . Show that the equation of the diagonal through the point of intersection of

$$U = a, V = a' \text{ and } U = b, V = b' \text{ is given by } \begin{vmatrix} U & V & 1 \\ a & a' & 1 \\ b & b' & 1 \end{vmatrix} = 0.$$

 [Watch Video Solution](#)

72. Find the values of non-negative real number  $h_1, h_2, h_3, k_1, k_2, k_3$  such that the algebraic sum of the perpendiculars drawn from the points  $(2, k_1), (3, k_2), (7, k_3), (h_1, 4), (h_2, 5), (h_3, -3)$  on a variable line passing through  $(2, 1)$  is zero.

 [Watch Video Solution](#)

Example

1. Show that the lines  $4x + y - 9 = 0$ ,  $x - 2y + 3 = 0$ ,  $5x - y - 6 = 0$  make equal intercepts on any line of slope 2.

 [Watch Video Solution](#)

2. The equations of two sides of a triangle are  $3y - x - 2 = 0$  and  $y + x - 2 = 0$ . The third side, which is variable, always passes through the point  $(5, -1)$ . Find the range of the values of the slope of the third side, so that the origin is an interior point of the triangle.

 [Watch Video Solution](#)

3. Find the locus of the circumcenter of a triangle whose two sides are along the coordinate axes and the third side passes through the point of intersection of the line  $ax + by + c = 0$  and  $lx + my + n = 0$ .

 [Watch Video Solution](#)

4. Let  $ABC$  be a triangle with  $AB = AC$ . If  $D$  is the midpoint of  $BC$ ,  $E$  is the foot of the perpendicular drawn from  $D$  to  $AC$ , and  $F$  is the midpoint of  $DE$ , then prove that  $AF$  is perpendicular to  $BE$ .

 [Watch Video Solution](#)

5. A diagonal of rhombus  $ABCD$  is member of both the families of lines  $(x + y - 1) + \lambda(2x + 3y - 2) = 0$  and  $(x - y + 2) + \lambda(2x - 3y + 5) = 0$  and rhombus is  $(3, 2)$ . If the area of the rhombus is  $12\sqrt{5}$  sq. units, then find the remaining vertices of the rhombus.

 [Watch Video Solution](#)

6. Let  $ABC$  be a given isosceles triangle with  $AB = AC$ . Sides  $AB$  and  $AC$  are extended up to  $E$  and  $F$ , respectively, such that  $BE \times CF = AB^2$ . Prove that the line  $EF$  always passes through a fixed point.



Watch Video Solution

7. Let  $L_1 = 0$  and  $L_2 = 0$  be two fixed lines. A variable line is drawn through the origin to cut the two lines at  $R$  and  $S$ .  $P$  is a point on the line  $RS$  such that  $\frac{(m+n)}{OP} = \frac{m}{OR} + \frac{n}{OS}$ . Show that the locus of  $P$  is a straight line passing through the point of intersection of the given lines  $R, S, R$  are on the same side of  $O$ ).



Watch Video Solution

8. Let points  $A, B$  and  $C$  lie on lines  $y-x=0$ ,  $2x-y=0$  and  $y-3x=0$ , respectively. Also,  $AB$  passes through fixed point  $P(1,0)$  and  $BC$  passes through fixed point  $Q(0,-1)$ . Then prove that  $AC$  also passes through a fixed point and find that point.



Watch Video Solution



9. Consider two lines  $L_1$  and  $L_2$  given by  $x - y = 0$  and  $x + y = 0$ , respectively, and a moving point  $P(x, y)$ . Let  $d(P, L_i), i = 1, 2$ , represents the distance of point  $P$  from the line  $L_i$ . If point  $P$  moves in a certain region  $R$  in such a way that  $2 \leq d(P, L_1) + d(P, L_2) \leq 4$ , find the area of region  $R$ .

 [Watch Video Solution](#)

10. Let  $O(0, 0)$ ,  $A(2, 0)$ , and  $B\left(1, \frac{1}{\sqrt{3}}\right)$  be the vertices of a triangle. Let  $R$  be the region consisting of all those points  $P$  inside  $OAB$  which satisfy  $d(P, OA) \leq \min [d(P, OB), d(P, AB)]$ , where  $d$  denotes the distance from the point to the corresponding line. Sketch the region  $R$  and find its area.

 [Watch Video Solution](#)

11. A line through  $A(-5, -4)$  meets the lines  $x + 3y + 2 = 0$ ,  $2x + y + 4 = 0$  and  $x - y - 5 = 0$  at the points  $B$ ,  $C$  and  $D$  respectively, if  $\left(\frac{15}{AB}\right)^2 + \left(\frac{10}{AC}\right)^2 = \left(\frac{6}{AD}\right)^2$  find the equation of the line.

 [Watch Video Solution](#)

12. A rectangle PQRS has its side PQ parallel to the line  $y = mx$  and vertices  $P$ ,  $Q$ , and  $S$  on the lines  $y = a$ ,  $x = b$ , and  $x = -b$ , respectively. Find the locus of the vertex  $R$ .

 [Watch Video Solution](#)

## Concept Application Exercise 2 1

1. Find the equation of the right bisector of the line segment joining the points  $(3,4)$  and  $(-1,2)$ .



Watch Video Solution

2. about to only mathematics



Watch Video Solution

3. If the coordinates of the vertices of triangle  $ABC$  are  $(-1, 6)$ ,  $(-3, -9)$  and  $(5, -8)$ , respectively, then find the equation of the median through  $C$ .



Watch Video Solution

4. Find the equation of the line perpendicular to the line  $\frac{x}{a} - \frac{y}{b} = 1$  and passing through a point at which it cuts the x-axis.



Watch Video Solution

5. If the middle points of the sides  $BC$ ,  $CA$ , and  $AB$  of triangle  $ABC$  are  $(1, 3)$ ,  $(5, 7)$ , and  $(-5, 7)$ , respectively, then find the equation of the side  $AB$ .



[Watch Video Solution](#)

6. Find the equations of the lines which pass through the origin and are inclined at an angle  $\tan^{-1} m$  to the line  $y = mx + c$ .



[Watch Video Solution](#)

7. If  $(-2, 6)$  is the image of the point  $(4, 2)$  with respect to line  $L=0$ , then  $L$  is:



[Watch Video Solution](#)

8. Find the area bounded by the curves  $x + 2|y| = 1$  and  $x = 0$ .



[Watch Video Solution](#)

9. Find the equation of the straight line passing through the intersection of the lines  $x - 2y = 1$  and  $x + 3y = 2$  and parallel to  $3x + 4y = 0$ .

 [Watch Video Solution](#)

10. If the foot of the perpendicular from the origin to a straight line is at  $(3, -4)$ , then find the equation of the line.

 [Watch Video Solution](#)

11. A straight line through the point  $(2, 2)$  intersects the lines  $\sqrt{3}x + y = 0$  and  $\sqrt{3}x - y = 0$  at the point  $A$  and  $B$ , respectively. Then find the equation of the line  $AB$  so that triangle  $OAB$  is equilateral.

 [Watch Video Solution](#)

12. The equation of the straight line passing through the point  $(4, 3)$  and making intercepts on the co ordinate axes whose sum is  $-1$ , is

 [Watch Video Solution](#)

13. A straight line through the point  $A(3, 4)$  is such that its intercept between the axes is bisected at  $A$ . Its equation is :

 [Watch Video Solution](#)

14. A straight line  $L$  is perpendicular to the line  $5x - y = 1$ . The area of the triangle formed by line  $L$ , and the coordinate axes is 5. Find the equation of line  $L$ .

 [Watch Video Solution](#)

15. One side of a rectangle lies along the line  $4x + 7y + 5 = 0$ . Two of its vertices are  $(-3, 1)$  and  $(1, 1)$ . Find the equations of the other three

sides.

 [Watch Video Solution](#)

16. A line  $L_1 \equiv 3y - 2x - 6 = 0$  is rotated about its point of intersection with the y-axis in the clockwise direction to make it  $L_2$  such that the area formed by  $L_1, L_2$  the x-axis, and line  $x = 5$  is  $\frac{49}{3}$  sq units if its point of intersection with  $x = 5$  lies below the x-axis. Find the equation of  $L_2$ .

 [Watch Video Solution](#)

17. The diagonals  $AC$  and  $BD$  of a rhombus intersect at  $(5, 6)$ . If  $A \equiv (3, 2)$ , then find the equation of diagonal  $BD$ .

 [Watch Video Solution](#)

18. Find the equation of the straight line which passes through the origin and makes angle  $60^{\circ}$  with the line  $x + \sqrt{3}y + \sqrt{3} = 0$ .

 [Watch Video Solution](#)

19. A line intersects the straight lines  $5x - y - 4 = 0$  and  $3x - 4y - 4 = 0$  at  $A$  and  $B$ , respectively. If a point  $P(1, 5)$  on the line  $AB$  is such that  $AP : PB = 2 : 1$  (internally), find point  $A$ .

 [Watch Video Solution](#)

20. In the given figure,  $PQR$  is an equilateral triangle and  $OSPT$  is a square. If  $OT = 2\sqrt{2}$  units find the equation of lines  $OT, OS, SP, QR, PR,$  and  $PQ$ .

 [Watch Video Solution](#)



21. Two fixed points  $A$  and  $B$  are taken on the coordinates axes such that  $OA = a$  and  $OB = b$ . Two variable points  $A'$  and  $B'$  are taken on the same axes such that  $OA' + OB' = OA + OB$ . Find the locus of the point of intersection of  $AB'$  and  $A'B$ .

 [Watch Video Solution](#)

22. A regular polygon has two of its consecutive diagonals as the lines  $\sqrt{3}x + y - \sqrt{3}$  and  $2y = \sqrt{3}$ . Point  $(1, c)$  is one of its vertices. Find the equation of the sides of the polygon and also find the coordinates of the vertices.

 [Watch Video Solution](#)

23. Find the direction in which a straight line must be drawn through the point  $(1, 2)$  so that its point of intersection with the line  $x + y = 4$  may be at a distance of 3 units from this point.

 [Watch Video Solution](#)

 Watch Video Solution

## Concept Application Exercise 2.2

1. Two particles start from point  $(2, -1)$ , one moving two units along the line  $x + y = 1$  and the other 5 units along the line  $x - 2y = 4$ , If the particles move towards increasing  $y$ , then their new positions are:

 Watch Video Solution

2. The center of a square is at the origin and its one vertex is  $A(2, 1)$ . Find the coordinates of the other vertices of the square.

 Watch Video Solution

3. The straight line passing through  $P(x_1, y_1)$  and making an angle  $\alpha$  with  $x$ -axis intersects  $Ax + By + C = 0$  in  $Q$  then  $PQ =$  \_\_\_\_\_

 Watch Video Solution

4. The centroid of an equilateral triangle is  $(0,0)$ . If two vertices of the triangle lie on  $x+y = 2\sqrt{2}$ , then find all the possible vertices of triangle.

 [Watch Video Solution](#)

### Concept Application Exercise 2.3

1. Find the points on  $y$ -axis whose perpendicular distance from the line  $4x - 3y - 12 = 0$  is 3.

 [Watch Video Solution](#)

2. If  $p$  and  $p'$  are the distances of the origin from the lines  $x \sec \alpha + y \operatorname{cosec} \alpha = k$  and  $x \cos \alpha - y \sin \alpha = k \cos 2\alpha$ , then prove that  $4p^2 + p'^2 = k^2$ .

 [Watch Video Solution](#)

3. Prove that the lengths of the perpendiculars from the points  $(m^2, 2m)$ ,  $(mm', m + m')$ , and  $(m'^2, 2m')$  to the line  $x + y + 1 = 0$  are in GP.

 [Watch Video Solution](#)

4. The ratio in which the line  $3x+4y+2=0$  divides the distance between  $3x+4y+5=0$  and  $3x+4y-5=0$  is?

 [Watch Video Solution](#)

5. Find the incentre of a triangle formed by the lines  $x \cos \frac{\pi}{9} + y \sin \frac{\pi}{9} = \pi$ ,  $x \cos \frac{8\pi}{9} + y \sin \frac{8\pi}{9} = \pi$  and  $x \cos \frac{13\pi}{9} + y \sin \left( \frac{13\pi}{9} \right) = \pi$ .

 [Watch Video Solution](#)

6. Find the equations of lines parallel to  $3x - 4y - 5 = 0$  at a unit distance from it.

 [Watch Video Solution](#)

7. Find the equation of a straight line passing through the point  $(-5, 4)$  and which cuts off an intercept of  $\sqrt{2}$  units between the lines  $x + y + 1 = 0$  and  $x + y - 1 = 0$ .

 [Watch Video Solution](#)

### Concept Application Exercise 2 4

1. The point  $(8, -9)$  with respect to the lines  $2x + 3y - 4 = 0$  and  $6x + 9y + 8 = 0$  lies on (a) the same side of the lines (b) the different sides of the line (c) one of the line (d) none of these

 [Watch Video Solution](#)

2. How the following pairs of points are placed w.r.t the line  $3x-8y-7=0$ ?

(i)  $(-3, -4)$  and  $(1, 2)$       (ii)  $(-1, -1)$  and  $(3, 7)$

 [Watch Video Solution](#)

3. Find the range of  $(\alpha, 2 + \alpha)$  and  $\left(\frac{3\alpha}{2}, a^2\right)$  lie on the opposite sides of the line  $2x + 3y = 6$ .

 [Watch Video Solution](#)

4. If the point  $P(a^2, a)$  lies in the region corresponding to the acute angle between the lines  $2y = x$  and  $4y = x$ , then find the values of  $a$ .

 [Watch Video Solution](#)

5. If  $(a, 3a)$  is a variable point lying above the straight line  $2x+y+4=0$  and below the line  $x+4y-8=0$ , then find the values of  $a$ .



Watch Video Solution

6. Find the values of  $\alpha$  such that the variable point  $(\alpha, \tan\alpha)$  lies inside the triangle whose sides are

$$y = x + \sqrt{3} - \frac{\pi}{3}, x + y + \frac{1}{\sqrt{3}} + \frac{\pi}{6} = 0 \text{ and } x - \frac{\pi}{2} = 0$$



Watch Video Solution

7. Find the area of the region in which points satisfy

$$3 \leq |x| + |y| \leq 5.$$



Watch Video Solution

8. Find the area of the region formed by the points satisfying

$$|x| + |y| + |x + y| \leq 2.$$



Watch Video Solution

## Concept Application Exercise 2 5

1. Find the equation of the bisector of the obtuse angle between the lines  $3x - 4y + 7 = 0$  and  $12x + 5y - 2 = 0$ .

 [Watch Video Solution](#)

2. The incident ray is along the line  $3x - 4y - 3 = 0$  and the reflected ray is along the line  $24x + 7y + 5 = 0$ . Find the equation of mirrors.

 [Watch Video Solution](#)

3. If the two sides of rhombus are  $x + 2y + 2 = 0$  and  $2x + y - 3 = 0$ , then find the slope of the longer diagonal.

 [Watch Video Solution](#)



4. In triangle  $ABC$ , the equation of the right bisectors of the sides  $AB$  and  $AC$  are  $x + y = 0$  and  $y - x = 0$ , respectively. If  $A \equiv (5, 7)$ , then find the equation of side  $BC$ .

 [Watch Video Solution](#)

5. Show that the reflection of the line  $ax + by + c = 0$  on the line  $x + y + 1 = 0$  is the line  $bx + ay + (a + b - c) = 0$  where  $a \neq b$ .

 [Watch Video Solution](#)

6. The joint equation of two altitudes of an equilateral triangle is  $(\sqrt{3}x - y + 8 - 4\sqrt{3})(-\sqrt{3}x - y + 12 + 4\sqrt{3}) = 0$ . The third altitude has the equation

 [Watch Video Solution](#)



7.

Watch Video Solution

8. Two sides of a rhombus ABCD are parallel to the lines  $y = x + 2$  and  $y = 7x + 3$ . If the diagonals of the rhombus intersect at the point  $(1, 2)$  and the vertex A is on the y-axis, then vertex A can be



Watch Video Solution

## Concept Application Exercise 2.6

1. If  $a$  and  $b$  are two arbitrary constants, then prove that the straight line  $(a - 2b)x + (a + 3b)y + 3a + 4b = 0$  will pass through a fixed point. Find that point.



Watch Video Solution

2. If  $a, b, c$  are in harmonic progression, then the straight line  $\left(\left(\frac{x}{a}\right)\right)^{\frac{y}{b}} + \left(\frac{l}{c}\right) = 0$  always passes through a fixed point. Find that point.

 [Watch Video Solution](#)

3. A variable line passes through a fixed point P. The algebraic sum of the perpendiculars drawn from the points (2,0), (0,2) and (1,1) on the line is zero. Find the coordinate of the point P.

 [Watch Video Solution](#)

4. Consider the family of lines  $5x + 3y - 2 + \lambda_1(3x - y - 4) = 0$  and  $x - y + 1 + \lambda_2(2x - y - 2) = 0$ . Find the equation of a straight line that belongs to both the families.

 [Watch Video Solution](#)

5. If the straight lines  $x + y - 2 = 0$ ,  $2x - y + 1 = 0$  and  $ax + by - c = 0$  are concurrent, then the family of lines  $2ax + 3by + c = 0$  ( $a, b, c$  are nonzero) is concurrent at (a)  $(2, 3)$  (b)  $\left(\frac{1}{2}, \frac{1}{3}\right)$  (c)  $\left(-\frac{1}{6}, -\frac{5}{9}\right)$  (d)  $\left(\frac{2}{3}, -\frac{7}{5}\right)$



Watch Video Solution

### Exercise Single Correct Answer Type

1. Find the equations of the diagonals of the square formed by the lines  $x = 0$ ,  $y = 0$ ,  $x = 1$  and  $y = 1$ .

A.  $y=x, y+x=1$

B.  $y=x, x+y=2$

C.  $2y = x, y+x = 1/3$

D.  $y=2x, y+2x = 1$

Answer: A



Watch Video Solution

2. The coordinates of two consecutive vertices A and B of a regular hexagon ABCDEF are (1,0) and (2,0) respectively. The equation of the diagonal CE is:

A.  $\sqrt{3}x + y = 4$

B.  $x + \sqrt{3}y + 4 = 0$

C.  $x + \sqrt{3}y = 4$

D. none of these

**Answer: C**



Watch Video Solution

3. If each of the points  $(x_1, 4)$ ,  $(-2, y_1)$  lies on the line joining the points  $(2, -1)$  and  $(5, -3)$ , then the point  $P(x_1, y_1)$  lies on the line. (a)

$6(x + y) - 25 = 0$  (b)  $2x + 6y + 1 = 0$  (c)  $2x + 3y - 6 = 0$  (d)

$6(x + y) + 25 = 0$

A.  $6(x+y)-25 = 0$

B.  $2x+6y+1 = 0$

C.  $2x+3y-6=0$

D.  $6(x+y)+25=0$

**Answer: B**

 [Watch Video Solution](#)

4. The equation to the straight line passing through the point  $(a\cos^3\theta, a\sin^3\theta)$  and perpendicular to the line  $x\sec\theta + y\operatorname{cosec}\theta = a$  is

A.  $x\cos\theta - y\sin\theta = a\cos 2\theta$

B.  $x\cos\theta + y\sin\theta = a\cos 2\theta$

C.  $x\sin\theta + y\cos\theta = a\cos 2\theta$

D. none of these

**Answer: A**



Watch Video Solution

5. The line  $PQ$  whose equation is  $x - y = 2$  cuts the  $x$ -axis at  $P$ , and  $Q$  is  $(4, 2)$ . The line  $PQ$  is rotated about  $P$  through  $45^\circ$  in the anticlockwise direction. The equation of the line  $PQ$  in the new position is  $y = -\sqrt{2}$  (b)  $y = 2$  (c)  $x = 2$  (d)  $x = -2$

A.  $y = -\sqrt{2}$

B.  $y=2$

C.  $x=2$

D.  $x=-2$

Answer: C



Watch Video Solution

6. A line moves in such a way that the sum of the intercepts made by it on the axes is always  $c$ . The locus of the mid-point of its intercept between the

axes is (A)  $x + y = 2c$  (B)  $x + y = c$  (C)  $2(x + y) = c$  (D) None of these

A.  $x+y=2c$

B.  $x+y=c$

C.  $2(x+y)=c$

D.  $2x+y=c$

**Answer: C**



**Watch Video Solution**

7. If the x intercept of the line  $y = mx + 2$  is greater than  $\frac{1}{2}$  then the gradient of the line lies in the interval

A.  $(-1,0)$

B.  $\left(\frac{-1}{4}, 0\right)$

C.  $(-\infty, -4)$

D.  $(-4,0)$



**Answer: D**

 [Watch Video Solution](#)

**8.** The equation of a straight line on which the length of perpendicular from the origin is four units and the line makes an angle of  $120^\circ$  with the x-axis is

(a)  $x\sqrt{3} + y + 8 = 0$    (b)  $x\sqrt{3} - y = 8$    (c)  $x\sqrt{3} - y = 8$    (d)

$x - \sqrt{3}y + 8 = 0$

A.  $x\sqrt{3} + y + 8 = 0$

B.  $x\sqrt{3} - y = 8$

C.  $x\sqrt{3} - y = 8$

D.  $x - \sqrt{3} + 8 = 0$

**Answer: A**

 [Watch Video Solution](#)

9.  $ABCD$  is a square  $A \equiv (1, 2)$ ,  $B \equiv (3, -4)$ . If line  $CD$  passes through  $(3, 8)$ , then the midpoint of  $CD$  is (a)  $(2, 6)$  (b)  $(6, 2)$  (c)  $(2, 5)$  (d)  $\left(\frac{28}{5}, \frac{1}{5}\right)$

A.  $(2,6)$

B.  $(6,2)$

C.  $(2,5)$

D.  $(28/5,1/5)$

**Answer: D**

 [Watch Video Solution](#)

10. The equation of straight line which passes through the point  $(-4,3)$  such that the portion of the line between the axes is divided by the point in ratio  $5:3$  is -

A.  $9x-20y+96=0$

B.  $9x+20y=24$

C.  $20x+9y+53=0$

D. none of these

**Answer: A**

 [Watch Video Solution](#)

11. A square of side 'a' lies above the x-axis and has one vertex at the origin. The side passing through the origin makes an angle  $\alpha$  ( $0 < \alpha < \pi/4$ ) with the positive direction of x-axis and Find the equation of diagonal not passing through the origin ?

A.  $y(\cos\alpha + \sin\alpha) + x(\sin\alpha - \cos\alpha) = a$

B.  $y(\cos\alpha + \sin\alpha) + x(\sin\alpha + \cos\alpha) = a$

C.  $y(\cos\alpha + \sin\alpha) + x(\cos\alpha - \sin\alpha) = a$

D.  $y(\cos\alpha - \sin\alpha) - x(\sin\alpha - \cos\alpha) = a$

**Answer: C**

 [Watch Video Solution](#)

12. Let  $P = (-1, 0)$ ,  $Q = (0, 0)$  and  $R = (3, 3\sqrt{3})$  be three points. The equation of the bisector of the angle PQR

A.  $(\sqrt{3}/2)x + y = 0$

B.  $x + \sqrt{3}y = 0$

C.  $\sqrt{3}x + y = 0$

D.  $x + (\sqrt{3}/2)y = 0$

**Answer: C**

 [Watch Video Solution](#)

13. The equation of a line through the point  $(1, 2)$  whose distance from the point  $(3, 1)$  has the greatest value is (a)  $y = 2x$  (b)  $y = x + 1$  (c)  $x + 2y = 5$

(d)  $y = 3x - 1$

A.  $y=2x$

B.  $y=x+1$

C.  $x+2y=5$

D.  $y=3x-1$

**Answer: A**



**Watch Video Solution**

14. One diagonal of a square is along the line  $8x - 15y = 0$  and one of its vertex is  $(1, 2)$ . Then the equations of the sides of the square passing through this vertex are

(a)  $23x + 7y = 9, 7x + 23y = 53$

(b)  $23x - 7y + 9 = 0, 7x + 23y + 53 = 0$

(c)  $23x - 7y - 9 = 0, 7x + 23y - 53 = 0$  (d) none of these

A.  $7x-8y+9=0, 8x+7y-22=0$

B.  $9x-8y+7=0, 8x+9y-26=0$

C.  $23x-7y-9=0, 7x+23y-53=0$

D. none of these

**Answer: C**



[Watch Video Solution](#)

15. Prove that the parallelogram formed by the lines

$\frac{x}{a} + \frac{y}{b} = 1, \frac{x}{b} + \frac{y}{a} = 1, \frac{x}{a} + \frac{y}{b} = 2$  and  $\frac{x}{b} + \frac{y}{a} = 2$  is a rhombus.

A.  $\frac{\pi}{4}$

B.  $\frac{\pi}{2}$

C.  $\frac{\pi}{3}$

D.  $\frac{\pi}{6}$

**Answer: B**



[Watch Video Solution](#)

16. A line with positive rational slope, passes through the point A(6,0) and is at a distance of 5 units from B (1,3). The slope of line is

A.  $\frac{15}{8}$

B.  $\frac{8}{15}$

C.  $\frac{5}{8}$

D.  $\frac{8}{5}$

**Answer: B**



[Watch Video Solution](#)

17. A projectile A is projected from ground. An observer B running on ground with uniform velocity of magnitude  $v$  observes A to move along a straight line. The time of flight of A as measured by B is T. Then the range R of projectile on ground is

A.  $3x+3y-1=0$

B.  $x-3y+2=0$

C.  $5x+5y-3=0$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

**18.** Given  $A \equiv (1, 1)$  and  $AB$  is any line through it cutting the  $x$ -axis at  $B$ . If  $AC$  is perpendicular to  $AB$  and meets the  $y$ -axis in  $C$ , then the equation of the locus of midpoint  $P$  of  $BC$  is (a)  $x + y = 1$  (b)  $x + y = 2$  (c)  $x + y = 2xy$  (d)  $2x + 2y = 1$

A.  $x+y=1$

B.  $x+y=2$

C.  $x+y=2xy$

D.  $2x+2y=1$



**Answer: A**



**Watch Video Solution**

19. The number of possible straight lines passing through point(2,3) and forming a triangle with coordinate axes whose area is 12 sq. unit is: a. one b. two c. three d. four

A. one

B. two

C. three

D. four

**Answer: C**



**Watch Video Solution**

20. Two parallel lines lying in the same quadrant make intercepts  $a$  and  $b$  on  $x$  and  $y$  axes, respectively, between them. The distance between the lines is

(a)  $\frac{ab}{\sqrt{a^2 + b^2}}$  (b)  $\sqrt{a^2 + b^2}$  (c)  $\frac{1}{\sqrt{a^2 + b^2}}$  (d)  $\frac{1}{a^2} + \frac{1}{b^2}$

A.  $\sqrt{a^2 + b^2}$

B.  $\frac{ab}{\sqrt{a^2 + b^2}}$

C.  $\frac{1}{\sqrt{a^2 + b^2}}$

D.  $\frac{1}{a^2} + \frac{1}{b^2}$

Answer: B

 Watch Video Solution

21. The line  $L_1 \equiv 4x + 3y - 12 = 0$  intersects the  $x$ - and  $y$ -axes at  $A$  and  $B$ , respectively. A variable line perpendicular to  $L_1$  intersects the  $x$ - and the  $y$ -axis at  $P$  and  $Q$ , respectively. Then the locus of the circumcenter of triangle  $ABQ$  is (a)  $3x - 4y + 2 = 0$  (b)  $4x + 3y + 7 = 0$  (c)  $6x - 8y + 7 = 0$  (d) none of these

A.  $3x-4y+2 = 0$

B.  $4x+3y+7 = 0$

C.  $6x-8y+7=0$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

22. A beam of light is sent along the line  $x - y = 1$ , which after refracting from the x-axis enters the opposite side by turning through  $30^\circ$  towards the normal at the point of incidence on the x-axis. Then the equation of the refracted ray is  $(2 - \sqrt{3})x - y = 2 + \sqrt{3}$   $(2 + \sqrt{3})x - y = 2 + \sqrt{3}$   
 $(2 - \sqrt{3})x + y = (2 + \sqrt{3}) y = (2 - \sqrt{3})(x - 1)$

A.  $(2 - \sqrt{3})x - y = 2 + \sqrt{3}$

B.  $(2 + \sqrt{3})x - y = 2 + \sqrt{3}$

C.  $(2 - \sqrt{3})x + y = (2 + \sqrt{3})$

$$D. y = (2 + \sqrt{3})(x - 1)$$

**Answer: D**



**Watch Video Solution**

23. The number of integral values of  $m$  for which the x-coordinate of the point of intersection of the lines  $3x + 4y = 9$  and  $y = mx + 1$  is also an integer is 2 (b) 0 (c) 4 (d) 1

A. 2

B. 0

C. 4

D. 1

**Answer: A**



**Watch Video Solution**

24. If the sum of the distances of a point from two perpendicular lines in a plane is 1, then its locus is

(a) a square (b) a circle (c) a straight line (d) two intersecting lines

A. a square

B. a circle

C. a straight line

D. two intersecting lines

**Answer: A**



[Watch Video Solution](#)

25. The equation of set of lines which are at a constant distance 2 units from the origin is

A.  $x+y+2=0$

B.  $x+y+4=0$

$$C. x \cos \alpha + y \sin \alpha = 2$$

$$D. x \cos \alpha + y \sin \alpha = \frac{1}{2}$$

**Answer: C**



**Watch Video Solution**

**26.** The lines  $y = m_1x$ ,  $y = m_2x$  and  $y = m_3x$  make equal intercepts on the line  $x + y = 1$ . Then  $2(1 + m_1)(1 + m_3) = (1 + m_2)(2 + m_1 + m_3)$

$$(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$$

$$(1 + m_1)(1 + m_2) = (1 + m_3)(2 + m_1 + m_3)$$

$$2(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$$

A.  $2(1 + m_1)(1 + m_3) = (1 + m_2)(2 + m_1 + m_3)$

B.  $(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$

C.  $(1 + m_1)(1 + m_2) = (1 + m_3)(2 + m_1 + m_3)$

D.  $2(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$

**Answer: A**



**Watch Video Solution**

27. The condition on  $a$  and  $b$ , such that the portion of the line  $ax + by - 1 = 0$  intercepted between the lines  $ax + y = 0$  and  $x + by = 0$  subtends a right angle at the origin, is  $a = b$  (b)  $a + b = 0$   $a = 2b$  (d)  $2a = b$

A.  $a = b$

B.  $a + b = 0$

C.  $a = 2b$

D.  $2a = b$

**Answer: B**



**Watch Video Solution**

28. The area of the triangle formed by the lines  $y = ax$ ,  $x + y - a = 0$ , and  $y$ -axis is equal to

A.  $\frac{1}{2|1 + a|}$

B.  $\frac{a^2}{|1 + a|}$

C.  $\frac{1}{2} \frac{a}{|1 + a|}$

D.  $\frac{a^2}{2|1 + a|}$

**Answer: D**

 [Watch Video Solution](#)

29. The line  $\frac{x}{a} + \frac{y}{b} = 1$  meets the  $x$ -axis at  $A$ , the  $y$ -axis at  $B$ , and the line  $y = x$  at  $C$  such, that the area of  $\Delta AOC$  is twice the area of  $\Delta BOC$ . Then the coordinates of  $C$  are  $\left(\frac{b}{3}, \frac{b}{3}\right)$  (b)  $\left(\frac{2a}{3}, \frac{2a}{3}\right)$  (c)  $\left(\frac{2b}{3}, \frac{2b}{3}\right)$  (d) none of these

A.  $\left(\frac{b}{3}, \frac{b}{3}\right)$



B.  $\left(\frac{2a}{3}, \frac{2a}{3}\right)$

C.  $\left(\frac{2b}{3}, \frac{2b}{3}\right)$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

30. The line  $\frac{x}{3} + \frac{y}{4} = 1$  meets the y-axis and x-axis at A and B respectively. A square  $ABCD$  is constructed on the line segment  $AB$  away from the origin. The coordinates of the vertex of the square farthest from the origin are

A. 7,3

B. 4,7

C. 6,4

D. 3,8

**Answer: B**



**Watch Video Solution**

**31.** The area of a parallelogram formed by the lines  $ax \pm bx \pm c = 0$  is (a)

$\frac{c^2}{(ab)}$  (b)  $\frac{2c^2}{(ab)}$  (c)  $\frac{c^2}{2ab}$  (d) none of these

A.  $c^2 / (ab)$

B.  $2c^2 / (ab)$

C.  $c^2 / 2ab$

D. none of these

**Answer: B**



**Watch Video Solution**

**32.** One diagonal of a square is  $3x-4y+8=0$  and one vertex is  $(-1,1)$ , then the area of square is

A.  $\frac{1}{50}$ sq.unit

B.  $\frac{1}{25}$ sq.unit

C.  $\frac{3}{50}$ sq.unit

D.  $\frac{2}{25}$ sq.unit

**Answer: D**

 [Watch Video Solution](#)

**33.** In an isosceles triangle OAB , O is the origin and  $OA=OB=6$  . The equation of the side AB is  $x-y+1=0$  Then the area of the triangle is

A.  $2\sqrt{21}$

B.  $\sqrt{142}$

C.  $\sqrt{\frac{142}{2}}$

D.  $\sqrt{\frac{71}{2}}$

**Answer: D**



Watch Video Solution

34. about to only mathematics

A. 1 : 2

B. 3 : 4

C. 2 : 01

D. 4 : 3

Answer: B



Watch Video Solution

35. The coordinates of the foot of the perpendicular from the point (2, 3)

on the line  $-y + 3x + 4 = 0$  are given by  $\left(\frac{37}{10}, -\frac{1}{10}\right)$  (b)

$\left(-\frac{1}{10}, \frac{37}{10}\right)$   $\left(\frac{10}{37}, -10\right)$  (d)  $\left(\frac{2}{3}, -\frac{1}{3}\right)$

A. (37/10,-1/10)

B.  $(-1/10, 37/10)$

C.  $(10/37, -10)$

D.  $(2/3, -1/3)$

**Answer: B**

 [Watch Video Solution](#)

36. The straight lines  $7x - 2y + 10 = 0$  and  $7x + 2y - 10 = 0$  form an isosceles triangle with the line  $y = 2$ . The area of this triangle is equal to

$\frac{15}{7}$  sq units (b)  $\frac{10}{7}$  sq units  $\frac{18}{7}$  sq units (d) none of these

A.  $15/7$  sq. units

B.  $10/7$  sq. units

C.  $18/7$  sq. units

D. none of these

**Answer: C**

 [Watch Video Solution](#)

37. The equations of the sides of a triangle are  $x+y-5=0$ ,  $x-y+1=0$ , and  $y-1=0$ .

Then the coordinates of the circumcenter are

A. 2,1

B. 1,2

C. 2,-2

D. 1,-2

**Answer: A**

 [Watch Video Solution](#)

38. The equations of the sides of a triangle are

$x + y - 5 = 0$ ,  $x - y + 1 = 0$ , and  $x + y - \sqrt{2} = 0$  is

$\left(-\infty, -\frac{4}{3}\right) \cup \left(\frac{4}{3}, +\infty\right)$   $\left(-\frac{4}{3}, \frac{4}{3}\right)$  (c)  $\left(-\frac{3}{4}, \frac{4}{3}\right)$  none of these

A.  $(-\infty, -4/3) \cup (4/3, +\infty)$

B.  $(-4/3, 4/3)$

C.  $(-3/4, 4/3)$

D. none of these

**Answer: A**

 [Watch Video Solution](#)

**39.** The range of values of  $\theta$  in the interval  $(0, \pi)$  such that the points  $(3, 5)$  and  $(\sin \theta, \cos \theta)$  lie on the same side of the line  $x + y - 1 = 0$ , is

A.  $0 < \theta < \frac{\pi}{4}$

B.  $0 < \theta < \frac{\pi}{2}$

C.  $0 < \theta < \pi$

D.  $\frac{\pi}{4} < \theta < \frac{3\pi}{4}$

**Answer: B**



Watch Video Solution

40. Distance of origin from the line  $(1 + \sqrt{3})y + (1 - \sqrt{3})x = 10$  along the line  $y = \sqrt{3}x + k$  (1)  $\frac{2}{\sqrt{5}}$  (2)  $5\sqrt{2} + k$  (3) 10 (4) 5

A.  $\frac{5}{\sqrt{2}}$

B.  $5\sqrt{2} + k$

C. 10

D. 5

Answer: D



Watch Video Solution

41. Consider the points  $A(0, 1)$  and  $B(2, 0)$ , and  $P$  be a point on the line  $4x + 3y + 9 = 0$ . The coordinates of  $P$  such that  $|PA - PB|$  is maximum are (a)  $\left(-\frac{24}{5}, \frac{17}{5}\right)$  (b)  $\left(-\frac{84}{5}, \frac{13}{5}\right)$  (c)  $\left(\frac{31}{7}, \frac{31}{7}\right)$  (d)  $(-3, 0)$



Watch Video Solution



42. Consider the point  $A = (3, 4)$ ,  $B(7, 13)$ . If 'P' be a point on the line  $y = x$  such that  $PA + PB$  is minimum then coordinates of P is (A)  $\left(\frac{13}{7}, 13, 7\right)$  (B)  $\left(\frac{23}{7}, \frac{23}{7}\right)$  (C)  $\left(\frac{31}{7}, \frac{31}{7}\right)$  (D)  $\left(\frac{33}{7}, \frac{33}{7}\right)$

A.  $(12/7, 12/7)$

B.  $(-24/5, 17/5)$

C.  $(31/7, 31/7)$

D.  $(0,0)$

Answer: C

 Watch Video Solution

43. The area enclosed by  $2|x| + 3|y| \leq 6$  is

A. 3 sq. units

B. 4 sq. units

C. 12 sq. units

D. 24 sq. units

**Answer: C**

 [Watch Video Solution](#)

44.  $ABC$  is a variable triangle such that  $A$  is  $(1, 2)$ , and  $B$  and  $C$  on the line  $y = x + \lambda$  ( $\lambda$  is a variable). Then the locus of the orthocentre of triangle  $ABC$  is  $x + y = 0$  (b)  $x - y = 0$   $x^2 + y^2 = 4$  (d)  $x + y = 3$

A.  $x+y=0$

B.  $x-y=0$

C.  $x^2 + y^2 = 4$

D.  $x+y=3$

**Answer: D**

 [Watch Video Solution](#)

45. In  $ABC$ , the coordinates of the vertex  $A$  are  $(4, -1)$ , and lines  $x - y - 1 = 0$  and  $2x - y = 3$  are the internal bisectors of angles  $B$  and  $C$ . Then, the radius of the encircle of triangle  $ABC$  is (a)  $\frac{4}{\sqrt{5}}$  (b)  $\frac{3}{\sqrt{5}}$  (c)  $\frac{6}{\sqrt{5}}$  (d)  $\frac{7}{\sqrt{5}}$

A.  $4/\sqrt{5}$

B.  $3/\sqrt{5}$

C.  $6/\sqrt{5}$

D.  $7/\sqrt{5}$

**Answer: C**

 [Watch Video Solution](#)

46.  $P$  is a point on the line  $y + 2x = 1$ , and  $Q$  and  $R$  two points on the line  $3y + 6x = 6$  such that triangle  $PQR$  is an equilateral triangle. The length

of the side of the triangle is  $\frac{2}{\sqrt{5}}$  (b)  $\frac{3}{\sqrt{5}}$  (c)  $\frac{4}{\sqrt{5}}$  (d) none of these

A.  $2/\sqrt{15}$

B.  $3/\sqrt{5}$

C.  $4/\sqrt{5}$

D. none of these

**Answer: A**



**Watch Video Solution**

47. If the equation of base of an equilateral triangle is  $2x - y = 1$  and the vertex is  $(-1, 2)$ , then the length of the sides of the triangle is (a)  $\sqrt{\frac{20}{3}}$

(b)  $\frac{2}{\sqrt{15}}$  (c)  $\sqrt{\frac{8}{15}}$  (d)  $\sqrt{\frac{15}{2}}$

A.  $\sqrt{20/3}$

B.  $2/\sqrt{15}$

C.  $\sqrt{8/15}$

D.  $\sqrt{15/2}$

**Answer: A**

 [Watch Video Solution](#)

**48.** The locus of a point that is equidistant from the lines  $x + y - 2\sqrt{2} = 0$  and  $x + y - \sqrt{2} = 0$  is (a)  $x + y - 5\sqrt{2} = 0$  (b)  $x + y - 3\sqrt{2} = 0$  (c)  $2x + 2y - 3\sqrt{2} = 0$  (d)  $2x + 2y - 5\sqrt{5} = 0$

A.  $x + y - 5\sqrt{2} = 0$

B.  $x + y - 3\sqrt{2} = 0$

C.  $2x + 2y - 3\sqrt{2} = 0$

D.  $2x + 2y - 5\sqrt{2} = 0$

**Answer: C**

 [Watch Video Solution](#)

49. If the quadrilateral formed by the lines  $ax + by + c = 0$ ,  $a'x + b'y + c = 0$ ,  $ax + by + c' = 0$ ,  $a'x + b'y + c' = 0$  has perpendicular diagonals, then  $b^2 + c^2 = b'^2 + c'^2$   $c^2 + a^2 = c'^2 + a'^2$   $a^2 + b^2 = a'^2 + b'^2$  (d) none of these

A.  $b^2 + c^2 = b'^2 + c'^2$

B.  $c^2 + a^2 = c'^2 + a'^2$

C.  $a^2 + b^2 = a'^2 + b'^2$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

50. A line of fixed length 2 units moves so that its ends are on the positive x-axis and that part of the line  $x + y = 0$  which lies in the second quadrant. Then the locus of the midpoint of the line has equation.

A.  $x^2 + 5y^2 + 4xy - 1 = 0$

$$B. x^2 + 5y^2 + 4xy + 1 = 0$$

$$C. x^2 + 5y^2 - 4xy - 1 = 0$$

$$D. x^2 + 5y^2 - 4xy - 1 = 0$$

**Answer: A**

 [Watch Video Solution](#)

51. If the extremities of the base of an isosceles triangle are the points  $(2a, 0)$  and  $(0, a)$ , and the equation of one of the side is  $x = 2a$ , then the area of the triangle is  $5a^2$  sq units (b)  $\frac{5a^2}{2}$  sq units  $\frac{25a^2}{2}$  sq units (d) none of these

A.  $5a^2$ sq. units

B.  $5a^2 / 2$ sq. units

C.  $25a^2 / 2$ sq. units

D. none of these

**Answer: B**

 [Watch Video Solution](#)

52.  $A \equiv (-4, 0)$ ,  $B \equiv (4, 0)$  and  $N$  are the variable points of the y-axis such that  $M$  lies below  $N$  and  $MN = 4$ . Lines  $AM$  and  $BN$  intersect at  $P$ . The locus of  $P$  is (a)  $2xy - 16 - x^2 = 0$  (b)  $2xy + 16 - x^2 = 0$  (c)  $2xy + 16 + x^2 = 0$  (d)  $2xy - 16 + x^2 = 0$

A.  $2xy - 16 - x^2 = 0$

B.  $2xy + 16 - x^2 = 0$

C.  $2xy + 16 + x^2 = 0$

D.  $2xy - 16 + x^2 = 0$

**Answer: D**

 [Watch Video Solution](#)



53. The number of triangles that the four lines  $y = x + 3$ ,  $y = 2x + 3$ ,  $y = 3x + 2$ , and  $y + x = 3$  form is (a) 4 (b) 2 (c) 3 (d) 1

A. 4

B. 2

C. 3

D. 1

**Answer: C**



**Watch Video Solution**

54. A variable line  $\frac{x}{a} + \frac{y}{b} = 1$  moves in such a way that the harmonic mean of  $a$  and  $b$  is 8. Then the least area of triangle made by the line with the coordinate axes is (1) 8 sq. unit (2) 16 sq. unit (3) 32 sq. unit (4) 64 sq. unit

- A. 8 sq. unit
- B. 16 sq. unit
- C. 32 sq. unit
- D. 64 sq. unit

**Answer: C**

 [Watch Video Solution](#)

55. Given  $A(0, 0)$  and  $B(x, y)$  with  $x \in (0, 1)$  and  $y > 0$ . Let the slope of the line AB equals  $m_1$ . Point C lies on the line  $x = 1$  such that the slope of BC equals  $m_2$  where  $0 < m_2 < m_1$ . If the area of the triangle ABC can be expressed as  $(m_1 - m_2)f(x)$ , then largest possible value of  $f(x)$  is:

- A. 1
- B.  $1/2$
- C.  $1/4$
- D.  $1/8$

Answer: D

 Watch Video Solution

56. A triangle is formed by the lines  $x + y = 0$ ,  $x - y = 0$ , and  $lx + my = 1$ . If  $l$  and  $m$  vary subject to the condition  $l^2 + m^2 = 1$ , then the locus of its circumcenter is (a)  $(x^2 - y^2)^2 = x^2 + y^2$  (b)  $(x^2 + y^2)^2 = (x^2 - y^2)$  (c)  $(x^2 + y^2)^2 = 4x^2y^2$  (d)  $(x^2 - y^2)^2 = (x^2 + y^2)^2$

A.  $(x^2 - y^2)^2 = x^2 + y^2$

B.  $(x^2 - y^2)^2 = (x^2 - y^2)$

C.  $(x^2 - y^2) = 4x^2y^2$

D.  $(x^2 - y^2)^2 = (x^2 + y^2)^2$

Answer: A

 Watch Video Solution

57. Let  $P$  be  $(5, 3)$  and a point  $R$  on  $y = x$  and  $Q$  on the  $x$ -axis be such that  $PQ + QR + RP$  is minimum. Then the coordinates of  $Q$  are  $\left(\frac{17}{4}, 0\right)$  (b)  $(17, 0)$   $\left(\frac{17}{2}, 0\right)$  (d) none of these

A.  $(17/4, 0)$

B.  $(17, 0)$

C.  $(17/2, 0)$

D. none of these

**Answer: A**



[Watch Video Solution](#)

58. If a pair of perpendicular straight lines drawn through the origin forms an isosceles triangle with the line  $2x + 3y = 6$ , then area of the triangle so formed is  $36/13$  (b)  $12/17$  (c)  $13/5$  (d)  $17/14$

A.  $\frac{36}{13}$  sq. unit

B.  $\frac{12}{17}$  sq. unit

C.  $\frac{13}{5}$  sq. unit

D.  $\frac{17}{13}$  sq. unit

**Answer: A**

 [Watch Video Solution](#)

59. A point  $P(x,y)$  moves that the sum of its distance from the lines  $2x-y-3=0$  and  $x+3y+4=0$  is 7. The area bounded by locus  $P$  is (in sq. unit)

A. 70

B.  $70\sqrt{2}$

C.  $35\sqrt{2}$

D. 140

**Answer: B**

 [Watch Video Solution](#)

60. If AD, BE and CF are the altitudes of  $\triangle ABC$  whose vertex A is (-4,5). The coordinates of points E and F are (4,1) and (-1,-4), respectively. Equation of BC is

A.  $3x-4y+28=0$

B.  $4x+3y+28=0$

C.  $3x-4y-28=0$

D.  $x+2y+7=0$

**Answer: C**

 [Watch Video Solution](#)

61. The vertex A of  $\triangle ABC$  is (3,-1). The equation of median BE and angle bisector CF are  $x-4y+10=0$  and  $6x+10y-59=0$ , respectively. Equation of AC is

A.  $5x+18y=37$

B.  $15x+8y=37$

C.  $15x-8y=37$

D.  $15x+8y+37=0$

**Answer: B**

 [Watch Video Solution](#)

62. Suppose A, B are two points on  $2x - y + 3 = 0$  and  $P(1, 2)$  is such that  $PA=PB$ . Then the mid point of AB is

A.  $\left(\frac{-1}{5}, \frac{13}{5}\right)$

B.  $\left(\frac{-7}{5}, \frac{9}{5}\right)$

C.  $\left(\frac{7}{5}, \frac{-9}{5}\right)$

D.  $\left(\frac{-7}{5}, \frac{-9}{5}\right)$

**Answer: A**

 [Watch Video Solution](#)

63. Triangle formed by variable lines  $(a+b)x+(a-b)y-2ab=0$  and  $(a-b)x+(a+b)y-2ab=0$  and  $x+y=0$  is (where  $a, b \in R$ )

- A. (a) equilateral
- B. (b) Isosceles
- C. (c) scalene
- D. (d) none of these

Answer: D

 Watch Video Solution

64. A light ray coming along the line  $3x + 4y = 5$  gets reflected from the line  $ax + by = 1$  and goes along the line  $5x - 12y = 10$ . Then,

$$a = \frac{64}{115}, b = \frac{112}{15} \quad a = \frac{14}{15}, b = -\frac{8}{115} \quad a = \frac{64}{115}, b = -\frac{8}{115}$$
$$a = \frac{64}{15}, b = \frac{14}{15}$$



$$\text{A. } a = \frac{64}{115}, b = \frac{112}{15}$$

$$\text{B. } a = \frac{14}{15}, b = -\frac{18}{115}$$

$$\text{C. } a = \frac{64}{115}, b = -\frac{8}{115}$$

$$\text{D. } a = \frac{64}{15}, b = \frac{14}{15}$$

**Answer: C**

 [Watch Video Solution](#)

**65.** The point  $(2,1)$ , translated parallel to the line  $x - y = 3$  by the distance of 4 units. If this new position  $A'$  is in the third quadrant, then the coordinates of  $A'$  are-

$$\text{A. } (2 + 2\sqrt{2}, 1 + 2\sqrt{2})$$

$$\text{B. } (-2 + \sqrt{2}, -1 - 2\sqrt{2})$$

$$\text{C. } (2 - 2\sqrt{2}, 1 - 2\sqrt{2})$$

**D.** none of these

**Answer: C**



**Watch Video Solution**

66. One of the diagonals of a square is the portion of the line  $x/2+y/3=2$  intercepted between the axes. Then the extremities of the other diagonal are

- A. (5,5), (-1,1)
- B. (0,0), (4,6)
- C. (0,0),(-1,1)
- D. (5,5),(4,6)

**Answer: A**



**Watch Video Solution**

67. The point  $P(2,1)$  is shifted through a distance  $3\sqrt{2}$  units measured parallel to the line  $x+y=1$  in the direction of decreasing ordinates, to reach at  $Q$ . The image of  $Q$  with respect to given line is

- A. (3,-4)
- B. (-3,2)
- C. (0,-1)
- D. none of these

**Answer: A**



[Watch Video Solution](#)

68. Let  $O$  be the origin. If  $A(1, 0)$  and  $B(0, 1)$  and  $P(x, y)$  are points such that  $xy > 0$  and  $x + y < 1$ , then  $P$

- A.  $P$  lies either inside the triangle  $OAB$  or in the third quadrant
- B.  $P$  cannot lie inside the triangle  $OAB$

C. P lies inside the triangle OAB

D. P lies in the first quadrant only

**Answer: A**

 [Watch Video Solution](#)

69. In a triangle  $ABC$ , the bisectors of angles  $B$  and  $C$  lie along the lines  $x = y$  and  $y = 0$ . If  $A$  is  $(1, 2)$ , then the equation of line  $BC$  is  $2x + y = 1$

(b)  $3x - y = 5$  (c)  $x - 2y = 3$  (d)  $x + 3y = 1$

A.  $2x + y = 1$

B.  $3x - y = 5$

C.  $x - 2y = 3$

D.  $x + 3y = 1$

**Answer: B**

 [Watch Video Solution](#)

70. Line  $ax + by + p = 0$  makes angle  $\frac{\pi}{4}$  with  $x \cos \alpha + y \sin \alpha = p, p \in R^+$ . If these lines and the line  $x \sin \alpha - y \cos \alpha = 0$  are concurrent, then  $a^2 + b^2 = 1$  (b)  $a^2 + b^2 = 2$  2( $a^2 + b^2$ ) = 1 (d) none of these

A.  $a^2 + b^2 = 1$

B.  $a^2 + b^2 = 2$

C.  $2(a^2 + b^2) = 1$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

71. The equation of the line AB is  $y = x$ . If A and B lie on the same side of the line mirror  $2x - y = 1$ , then the equation of the image of AB is

A.  $x+y=2$

B.  $8x+y=9$

C.  $7x-y=6$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

72. The equation of the bisector of the acute angle between the lines

$2x - y + 4 = 0$  and  $x - 2y = 1$  is  $x - y + 5 = 0$   $x - y + 1 = 0$

$x - y = 5$  (d) none of these

A.  $x+y+5=0$

B.  $x-y+1=0$

C.  $x-y=5$

D. none of these

**Answer: B**



Watch Video Solution

73. The straight line  $4ax + 3by + c = 0$  passes through? , where

$a + b + c = 0$  (a)  $(4, 3)$  (b)  $\left(\frac{1}{4}, \frac{1}{3}\right)$   $\left(\frac{1}{2}, \frac{1}{3}\right)$  (d) none of these

A.  $(4,3)$

B.  $(1/4, 1/3)$

C.  $(1/2, 1/3)$

D. none of these

Answer: B



Watch Video Solution

74. If the lines  $ax + y + 1 = 0$ ,  $x + by + 1 = 0$  and  $x + y + c = 0$  ( $a, b, c$  being distinct and different from 1) are concurrent, then prove that

$$\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1.$$

A. 0

B. 1

C.  $1/(a+b+c)$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

75. If lines  $x + 2y - 1 = 0$ ,  $ax + y + 3 = 0$ , and  $bx - y + 2 = 0$  are concurrent, and  $S$  is the curve denoting the locus of  $(a, b)$ , then the least distance of  $S$  from the origin is  $\frac{5}{\sqrt{57}}$  (b)  $5/\sqrt{51}$   $5/\sqrt{58}$  (d)  $5/\sqrt{59}$

A.  $5/\sqrt{57}$

B.  $5/\sqrt{51}$

C.  $5/\sqrt{58}$

D.  $5/\sqrt{59}$



**Answer: C**

 [Watch Video Solution](#)

76. The straight lines  $x + 2y - 9 = 0$ ,  $3x + 5y - 5 = 0$ , and  $ax + by - 1 = 0$  are concurrent, if the straight line  $35x - 22y + 1 = 0$  passes through the point  $(a, b)$  (b)  $(b, a)$  (c)  $(-a, -b)$  (d) none of these

A.  $(a,b)$

B.  $(b,a)$

C.  $(-a,-b)$

D. none of these

**Answer: A**

 [Watch Video Solution](#)

77. If the straight lines  $2x + 3y - 1 = 0$ ,  $x + 2y - 1 = 0$ , and  $ax + by - 1 = 0$  form a triangle with the origin as orthocentre, then  $(a, b)$  is given by (a)  $(6, 4)$  (b)  $(-3, 3)$  (c)  $(-8, 8)$  (d)  $(0, 7)$

A. (6,4)

B. (-3,3)

C. (-8,8)

D. (0,7)

**Answer: C**

 [Watch Video Solution](#)

78. If  $\frac{a}{\sqrt{bc}} - 2 = \sqrt{\frac{b}{c}} + \sqrt{\frac{c}{b}}$ , where  $a, b, c > 0$ , then the family of lines  $\sqrt{ax} + \sqrt{by} + \sqrt{c} = 0$  passes through the fixed point given by (a)  $(1, 1)$  (b)  $(1, -2)$  (c)  $(-1, 2)$  (d)  $(-1, 1)$

A. (1,1)

B. (1,-2)

C. (-1,2)

D. (-1,1)

**Answer: D**

 [Watch Video Solution](#)

79. If it is possible to draw a line which belongs to all the given family of lines

$$y - 2x + 1 + \lambda_1(2y - x - 1) = 0, 3y - x - 6 + \lambda_2(y - 3x + 6) = 0,$$

$$ax + y - 2 + \lambda_3(6x + ay - a) = 0, \text{ then}$$

$$(a)a = 4 \text{ (b) } a = 3 \text{ (c) } a = -2 \text{ (d) } a = 2$$

A. a=4

B. a=3

C. a=-2

D.  $a=2$

**Answer: A**

 [Watch Video Solution](#)

80. If two members of family  $(2 + \lambda)x + (1 + 2\lambda)y - 3(1 + \lambda) = 0$  and line  $x+y=0$  make an equilateral triangle, the the incentre of triangle so formed is

- A.  $\left(\frac{1}{3}, \frac{1}{3}\right)$
- B.  $\left(\frac{7}{6}, -\frac{5}{6}\right)$
- C.  $\left(\frac{5}{6}, \frac{5}{6}\right)$
- D.  $\left(-\frac{3}{2}, -\frac{3}{2}\right)$

**Answer: A**

 [Watch Video Solution](#)

81. The set of lines  $x \tan^{-1} a + y \sin^{-1} \left( \frac{1}{\sqrt{1+a^2}} \right) + 2 = 0$  where  $a \in (0, 1)$  are concurrent at (a)  $\left( \frac{1}{\pi}, \frac{1}{\pi} \right)$  (b)  $\left( -\frac{4}{\pi}, -\frac{4}{\pi} \right)$  (c)  $(\pi, \pi)$  (d)

none of these

 [Watch Video Solution](#)

82. If  $\sin(\alpha + \beta)\sin(\alpha - \beta) = \sin \gamma(2 \sin \beta + \sin \gamma)$ , where  $\gamma \in (0, \pi)$

A. (1,1)

B. (-1,1)

C. (1,-1)

D. none of these

Answer: C

 [Watch Video Solution](#)

1. If  $P$  is a point  $(x, y)$  on the line  $y = -3x$  such that  $P$  and the point  $(3, 4)$  are on the opposite sides of the line  $3x - 4y = 8$ , then  $x > \frac{8}{15}$  (b)  $x > \frac{8}{5}$   $y < -\frac{8}{5}$  (d)  $y < -\frac{8}{15}$

A.  $x > 8/15$

B.  $x > 8/5$

C.  $x < -8/5$

D.  $y < -8/15$

**Answer: A:C**



**Watch Video Solution**

2. If  $(x, y)$  is a variable point on the line  $y = 2x$  lying between the lines  $2(x + 1) + y = 0$  and  $x + 3(y - 1) = 0$ , then  $x \in \left(-\frac{1}{2}, \frac{6}{7}\right)$  (b)  $x \in \left(-\frac{1}{2}, \frac{3}{7}\right)$   $y \in \left(-1, \frac{3}{7}\right)$  (d)  $y \in \left(-1, \frac{6}{7}\right)$

A.  $x \in (-1/2, 6/7)$

B.  $x \in (-1/2, 3/7)$

C.  $y \in (-1, 3/7)$

D.  $y \in (-1, 6/7)$

**Answer: B::D**

 [Watch Video Solution](#)

3. Let  $P(\sin \theta, \cos \theta)$  ( $0 \leq \theta \leq 2\pi$ ) be a point and let OAB be a triangle with vertices  $(0, 0)$ ,  $(\sqrt{\frac{3}{2}}, 0)$  and  $(0, \sqrt{\frac{3}{2}})$ . Find  $\theta$  if P lies inside  $\triangle OAB$

A.  $0 < \theta < \pi/12$

B.  $5\pi/2 < \theta < \pi/2$

C.  $0 < \theta < 5\pi/2$

D.  $5\pi/2 < \theta < \pi$

**Answer: A::B**

 Watch Video Solution

4. The lines  $x + 2y + 3 = 0$ ,  $x + 2y - 7 = 0$ , and  $2x - y - 4 = 0$  are the sides of a square. The equation of the remaining side of the square can be  $2x - y + 6 = 0$  (b)  $2x - y + 8 = 0$   $2x - y - 10 = 0$  (b)  $2x - y - 14 = 0$

A.  $2x - y + 6 = 0$

B.  $2x - y + 8 = 0$

C.  $2x - y - 10 = 0$

D.  $2x - y - 14 = 0$

Answer: A:D

 Watch Video Solution

5. Angle made with the x-axis by a straight line drawn through  $(1, 2)$  so that it intersects  $x + y = 4$  at a distance  $\frac{\sqrt{6}}{3}$  from  $(1, 2)$  is (a)  $105^\circ$  (b)  $75^\circ$  (c)  $60^\circ$  (d)  $15^\circ$



A.  $105^\circ$

B.  $75^\circ$

C.  $60^\circ$

D.  $15^\circ$

**Answer: B::D**



**Watch Video Solution**

6. The straight lines

$$2x + 11y - 5 = 0, 24x + 7y - 20 = 0 \text{ and } 4x - 3y - 2 = 0$$

A. they form a triangle

B. they are concurrent

C. one line bisects the angle between the other two

D. two of them are parallel

**Answer: C**



Watch Video Solution

7. A triangle is formed by the lines whose equations are AB:  $x+y-5=0$ , BC:  $x+7y-7=0$  and CA:  $7x+y+14=0$ .

Then

- A. angle at A is acute
- B. angle at C is acute
- C. internal angle bisector at angle B is  $3x+6y-16=0$
- D. external angle bisector at angle C is  $8x+8y+7=0$

**Answer: A,C,D**



Watch Video Solution

8. If the points  $\left(\frac{a^3}{(a-1)}\right)$ ,  $\left(\frac{(a^2-3)}{(a-1)}\right)$ ,  $\left(\frac{b^3}{(b-1)}\right)$ ,  $\left(\frac{(b^2-3)}{(b-1)}\right)$ , and  $\left(\frac{(c^2-3)}{(c-1)}\right)$ , where  $a, b, c$  are different from 1, lie on the

$$lx + my + n = 0 \quad , \quad \text{then} \quad a + b + c = -\frac{m}{l} \quad ab + bc + ca = \frac{n}{l}$$

$$abc = \frac{(m + n)}{l} abc - (bc + ca + ab) + 3(a + b + c) = 0$$

A.  $a + b + c = -\frac{m}{l}$

B.  $ab + bc + ca = \frac{n}{l}$

C.  $abc = \frac{(m + n)}{l}$

D.  $abc - (bc + ca + ab) + 3(a + b + c) = 0$

**Answer: A::B::D**



**Watch Video Solution**

9. Two sides of a rhombus OABC ( lying entirely in first quadrant or fourth quadrant) of area equal to 2 sq. units, are  $y = \frac{x}{\sqrt{3}}$ ,  $y = \sqrt{3}x$  Then possible coordinates of B is / are ('O' being the origin)

A.  $(1 + \sqrt{3}, 1 + \sqrt{3})$

B.  $(-1 - \sqrt{3}, -1 - \sqrt{3})$

C.  $(3 + \sqrt{3}, 3 + \sqrt{3})$

D.  $(\sqrt{3} - 1, \sqrt{3} - 1)$

**Answer: A::B**



**Watch Video Solution**

10. If  $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 1$  and  $\left(\frac{x}{c}\right) + \left(\frac{y}{d}\right) = 1$  intersect the axes at four concyclic points and  $a^2 + c^2 = b^2 + d^2$ , then these lines can intersect at,  $(a, b, c, d > 0)$  `

A. (1,1)

B. (1,-1)

C. (2,-2)

D. (3,3)

**Answer: A, B, C and D**



**Watch Video Solution**

11. The straight line  $3x + 4y - 12 = 0$  meets the coordinate axes at

$A$  and  $B$ . An equilateral triangle  $ABC$  is constructed. The possible

coordinates of vertex  $C$   $\left(2\left(1 - \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 - \frac{4}{\sqrt{3}}\right)\right)$

$\left(-2(1 + \sqrt{3}), \frac{3}{2}(1 - \sqrt{3})\right)$   $\left(2(1 + \sqrt{3}), \frac{3}{2}(1 + \sqrt{3})\right)$   
 $\left(2\left(1 + \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 + \frac{4}{\sqrt{3}}\right)\right)$

A.  $\left(2\left(1 - \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 - \frac{4}{\sqrt{3}}\right)\right)$

B.  $\left(-2(1 + \sqrt{3}), \frac{3}{2}(1 - \sqrt{3})\right)$

C.  $\left(2(1 + \sqrt{3}), \frac{3}{2}(1 + \sqrt{3})\right)$

D.  $\left(2\left(1 + \frac{3\sqrt{3}}{4}\right), \frac{3}{2}\left(1 + \frac{4}{\sqrt{3}}\right)\right)$

Answer: A::D

 Watch Video Solution

12. The equation of the lines passing through the point  $(1, 0)$  and at a distance  $\frac{\sqrt{3}}{2}$  from the origin is (a)  $\sqrt{3}x + y - \sqrt{3} = 0$  (b)  $x + \sqrt{3}y - \sqrt{3} = 0$  (c)  $\sqrt{3}x - y - \sqrt{3} = 0$  (d)  $x - \sqrt{3}y - \sqrt{3} = 0$

A.  $\sqrt{3}x + y - \sqrt{3} = 0$

B.  $x + \sqrt{3}y - \sqrt{3} = 0$

C.  $\sqrt{3}x - y - \sqrt{3} = 0$

D.  $x - \sqrt{3}y - \sqrt{3} = 0$

**Answer: A::C**

 [Watch Video Solution](#)

13. The sides of a triangle are the straight lines  $x + y = 1$ ,  $7y = x$ , and  $\sqrt{3}y + x = 0$ . Then which of the following is an interior point of the triangle? (a) Circumcenter (b) Centroid (c) Incenter (d) Orthocenter

A. Circumcenter

B. Centroid

C. Incenter

D. Orthocenter

**Answer: B::C**

 [Watch Video Solution](#)

14. If the straight line  $ax + cy = 2b$ , where  $a, b, c > 0$ , makes a triangle of area 2 sq. units with the coordinate axes, then (a)  $a, b, c$  are in GP (b)  $a, -b, c$  are in GP (c)  $a, 2b, c$  are in GP (d)  $a, -2b, c$  are in GP

A.  $a, b, c$  are in GP

B.  $a, -b, c$  are in GP

C.  $a, 2b, c$  are in GP

D.  $a, -2b, c$  are in GP

**Answer: A::B**

15. Consider the equation  $y - y_1 = m(x - x_1)$  . If  $m$  and  $x_1$  are fixed and different lines are drawn for different values of  $y_1$ , then (a) the lines will pass through a fixed point (b) there will be a set of parallel lines (c) all the lines intersect the line  $x = x_1$  (d) all the lines will be parallel to the line  $y = x_1$

- A. the lines will pass through a fixed point
- B. there will be a set of parallel lines
- C. all the lines intersect the line  $x = x_1$
- D. all the lines will be parallel to the line  $y = x_1$

**Answer: B::C**



16. Equation(s) of the straight line(s), inclined at  $30^\circ$  to the x-axis such that the length of its (each of their) line segment(s) between the coordinate axes is 10 units, is (are)  $x + \sqrt{3}y + 5\sqrt{3} = 0$   $x - \sqrt{3}y + 5\sqrt{3} = 0$   
 $x + \sqrt{3}y - 5\sqrt{3} = 0$   $x - \sqrt{3}y - 5\sqrt{3} = 0$

A.  $x + \sqrt{3}y + 5\sqrt{3} = 0$

B.  $x - \sqrt{3}y + 5\sqrt{3} = 0$

C.  $x + \sqrt{3}y - 5\sqrt{3} = 0$

D.  $x - \sqrt{3}y - 5\sqrt{3} = 0$

Answer: B::D

 Watch Video Solution

17. The lines  $x + y - 1 = 0$ ,  $(m - 1)x + (m^2 - 7)y - 5 = 0$ , and  $(m - 2)x + (2m - 5)y = 0$  are a.) concurrent for three values of  $m$  b.) concurrent for no value of  $m$  c.) parallel for one value of  $m$  d.) parallel for two value of  $m$

A. concurrent for three values of  $m$

B. concurrent for one value of  $m$

C. concurrent for no value of  $m$

D. parallel for  $m=3$

**Answer: C::D**

 [Watch Video Solution](#)

18. The equation of a straight line passing through the point  $(2, 3)$  and inclined at an angle of  $\tan^{-1}\left(\frac{1}{2}\right)$  with the line  $y + 2x = 5$  is  $y = 3$  (b)  $x = 2$  (c)  $3x + 4y - 18 = 0$  (d)  $4x + 3y - 17 = 0$

A.  $y=3$

B.  $x=2$

C.  $3x+4y-18=0$

D.  $4x+3y-17=0$

**Answer: B::C**



**Watch Video Solution**

**19.** Find the equation of a straight line on which the perpendicular from the origin makes an angle of  $30^\circ$  with  $x$ -axis and which forms a triangle of area  $50\sqrt{3}$  with the axes.

A.  $\sqrt{3}x + y - 10 = 0$

B.  $\sqrt{3}x + y + 10 = 0$

C.  $x + \sqrt{3}y - 10 = 0$

D.  $x - \sqrt{3}y - 10 = 0$

**Answer: A::B**



**Watch Video Solution**

20. A line is drawn perpendicular to line  $y = 5x$ , meeting the coordinate axes at  $A$  and  $B$ . If the area of triangle  $OAB$  is 10 sq. units, where  $O$  is the origin, then the equation of drawn line is (a)  $3x - y - 9 = 0$  (b)  $x + 5y = 10$  (c)  $x + 4y = 10$  (d)  $x - 4y = 10$

A. 12

B. -12

C. 10

D. -10

Answer: A::B

 Watch Video Solution

21. If  $x - 2y + 4 = 0$  and  $2x + y - 5 = 0$  are the sides of an isosceles triangle having area 10 sq units, the equation of the third side is (a)  $3x - y = -9$  (b)  $3x - y + 11 = 0$  (c)  $x - 3y = 19$  (d)  $3x - y + 15 = 0$

A.  $x+3y=-1$

B.  $x+3y=19$

C.  $3x-y=-9$

D.  $3x-y=11$

**Answer: A::B::C::D**

 [Watch Video Solution](#)

22. Find the value of  $a$  for which the lines  $2x + y - 1 = 0$ ,  $ax + 3y - 3 = 0$ ,  $3x + 2y - 2 = 0$  are concurrent.

A. -3

B. -1

C. 1

D. infinite value

**Answer: infinite**



Watch Video Solution

23. The lines  $px + qy + r = 0$ ,  $qx + ry + p = 0$ ,  $rx + py + q = 0$ , are concurrent then

A.  $p+q+r=0$

B.  $p^2 + q^2 + r^2 = pr + rp + pq$

C.  $p^3 + q^3 + r^3 = 3pqr$

D. none of these

Answer: A::B::C



Watch Video Solution

24.  $\theta_1$  and  $\theta_2$  are the inclination of lines  $L_1$  and  $L_2$  with the x-axis. If  $L_1$  and  $L_2$  pass through  $P(x_1, y_1)$ , then the equation of one of the angle bisector of these lines is

$$\text{A. } \frac{x - x_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\sin\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

$$\text{B. } \frac{x - x_1}{-\sin\left(\frac{\theta_1 - \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 - \theta_2}{2}\right)}$$

$$\text{C. } \frac{x - x_1}{\sin\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

$$\text{D. } \frac{x - x_1}{-\sin\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

**Answer: A::D**



**Watch Video Solution**

25. Consider the lines

$L_1 \equiv 3x - 4y + 2 = 0$  and  $L_2 \equiv 3y - 4x - 5 = 0$ . Now, choose the

correct statement(s).

(a) The line  $x+y=0$  bisects the acute angle between  $L_1$  and  $L_2$  containing the origin.

(b) The line  $x-y+1=0$  bisects the obtuse angle between  $L_1$  and  $L_2$  not containing the origin.

(c) The line  $x+y+3=0$  bisects the obtuse angle between  $L_1$  and  $L_2$

containing the origin.

(d) The line  $x-y+1=0$  bisects the acute angle between  $L_1$  and  $L_2$  not containing the origin.

A. The line  $x+y=0$  bisects the acute angle between  $L_1$  and  $L_2$  containing the origin.

B. The line  $x-y+1=0$  bisects the obtuse angle between  $L_1$  and  $L_2$  not containing the origin.

C. The line  $x+y+3=0$  bisects the obtuse angle between  $L_1$  and  $L_2$  containing the origin.

D. The line  $x-y+1=0$  bisects the acute angle between  $L_1$  and  $L_2$  not containing the origin.

**Answer: A::B**



**Watch Video Solution**



26. The sides of a rhombus are parallel to the lines  $x + y - 1 = 0$  and  $7x - y - 5 = 0$ . It is given that the diagonals of the rhombus intersect at  $(1, 3)$  and one vertex,  $A$  of the rhombus lies on the line  $y = 2x$ . Then the coordinates of vertex  $A$  are  $\left(\frac{8}{5}, \frac{16}{5}\right)$  (b)  $\left(\frac{7}{15}, \frac{14}{15}\right)$   $\left(\frac{6}{5}, \frac{12}{5}\right)$  (d)  $\left(\frac{4}{15}, \frac{8}{15}\right)$

A.  $(8/5, 16/5)$

B.  $(7/15, 14/15)$

C.  $(6/5, 12/5)$

D.  $(4/15, 8/15)$

Answer: A::C

 Watch Video Solution

27. Two straight lines  $u = 0$  and  $v = 0$  pass through the origin and the angle between them is  $\tan^{-1}\left(\frac{7}{9}\right)$ . If the ratio of the slope of  $v = 0$  and

$u = 0$  is  $\frac{9}{2}$ , then their equations are  $y + 3x = 0$  and  $3y + 2x = 0$

$2y + 3x = 0$  and  $3y + 2x = 0$   $2y = 3x$  and  $3y = x$   $y = 3x$  and  $3y = 2x$

A.  $y+3x=0$  and  $3y+2x=0$

B.  $2y+3x=0$  and  $3y+x=0$

C.  $2y=3x$  and  $3y=0$

D.  $y=3x$  and  $3y=2x$

**Answer: A::B::C::D**



**Watch Video Solution**

**28.** Let  $u \equiv ax + by + abz = 0$ ,  $v \equiv bx - ay + ba^3 = 0$ ,  $a, b \in R$ , be two straight lines. The equations of the bisectors of the angle formed by

$k_1u - k_2v = 0$  and  $k_1u + k_2v = 0$ , for nonzero and real  $k_1$  and  $k_2$  are

A.  $u=0$

B.  $k_2u + k_1v = 0$

C.  $k_2u - k_1v = 0$

D.  $v=0$

**Answer: A,D**



**Watch Video Solution**

29. Two sides of a triangle are parallel to the coordinate axes. If the slopes of the medians through the acute angles of the triangle are 2 and  $m$ , then  $m$  is (a)  $\frac{1}{2}$  (b) 2 (c) 4 (d) 8

A. a.  $1/2$

B. b. 2

C. c. 4

D. d. 8

**Answer: A::D**



**Watch Video Solution**

30. A line which makes an acute angle  $\theta$  with the positive direction of the x-axis is drawn through the point  $P(3, 4)$  to meet the line  $x = 6$  at  $R$  and  $y = 8$  at  $S$ . Then, (a)  $PR = 3\sec\theta$  (b)  $PS = 4\operatorname{cosec}\theta$  (c)

$$PR + PS = \frac{2(3\sin\theta + 4\cos\theta)}{\sin 2\theta} \quad \text{(d)} \quad \frac{9}{(PR)^2} + \frac{16}{(PS)^2} = 1$$

A.  $PR = 3\sec\theta$

B.  $PS = 4 \operatorname{cosec}\theta$

C.  $PR + PS = \frac{2(3\sin\theta + 4\cos\theta)}{\sin 2\theta}$

D.  $\frac{9}{(PR)^2} + \frac{16}{(PS)^2} = 1$

Answer: A::B::C::D

 Watch Video Solution

### Exercise Linked Comprehension Type

1. Let  $l$  be the line belonging to the family of straight lines  $(a + 2b)x + (a - 3b)y + a - 8b = 0$ ,  $a, b \in R$ , which is farthest from the

point  $(2, 2)$ , then area enclosed by the line  $L$  and the coordinate axes is

A.  $x+4y+7=0$

B.  $2x+3y+4=0$

C.  $4x-y-6=0$

D. none of these

**Answer: A**



[Watch Video Solution](#)

2. Let  $l$  be the line belonging to the family of straight lines  $(a + 2b)x + (a - 3b)y + a - 8b = 0$ ,  $a, b \in R$ , which is farthest from the point  $(2, 2)$ , then area enclosed by the line  $L$  and the coordinate axes is

A.  $4/3$  sq. units

B.  $9/2$  sq. units

C.  $49/8$  sq. units

D. none of these

**Answer: C**

 [Watch Video Solution](#)

3. Let  $L$  be the line belonging to the family of straight lines  $(a+2b)x + (a-3b)y + a - 8b = 0$ ,  $a, b \in \mathbb{R}$ , which is the farthest from the point  $(2, 2)$ .

If  $L$  is concurrent with the lines  $x - 2y + 1 = 0$  and  $3x - 4y + \lambda = 0$ , then the value of  $\lambda$  is

A. 2

B. 1

C. -4

D. 5

**Answer: D**

 [Watch Video Solution](#)

4. The equation of an altitude of an equilateral triangle is  $\sqrt{3}x + y = 2\sqrt{3}$  and one of its vertices is  $(3, \sqrt{3})$  then the possible number of triangles is a.

1 b. 2 c. 3 4. 4

A. 1

B. 2

C. 3

D. 4

**Answer: B**



[Watch Video Solution](#)

5. The equation of an altitude of an equilateral triangle is  $\sqrt{3}x + y = 2\sqrt{3}$  and one of its vertices is  $(3, \sqrt{3})$  then the possible number of triangles is

A. 0,0

B. 0,  $2\sqrt{3}$

C.  $3, -\sqrt{3}$

D. none of these

**Answer: D**

 [Watch Video Solution](#)

6. The equation of an altitude of an equilateral triangle is  $\sqrt{3}x + y = 2\sqrt{3}$ , and one of the vertices is  $(3, \sqrt{3})$ .

Which of the following is not one of the possible vertices of the triangle?

A.  $a. \sqrt{3}$

B.  $b. \sqrt{3}$

C.  $c. 2$

D.  $d. \text{none of these}$

**Answer: A**

 [Watch Video Solution](#)



7. A variable line  $L$  is drawn through  $O(0,0)$  to meet the lines  $L_1 : y-x-10=0$  and  $L_2 : y-x-20=0$  at the points  $A$  and  $B$  respectively. A point  $P$  is taken on  $L$  such that  $OP^2 = OA^2 + OB^2$  and  $P, A, B$  lies on same side of origin  $O$ . The locus of  $P$  is

A.  $3x+3y=40$

B.  $3x+3y+40=0$

C.  $3x-3y=40$

D.  $3y-3x=40$

**Answer: D**

 [Watch Video Solution](#)

8. A variable line  $L$  is drawn through  $O(0,0)$  to meet the line  $L_1$  and  $L_2$  given by  $y-x-10=0$  and  $y-x-20=0$  at Points  $A$  and  $B$ , respectively.

Locus of P, if  $OP^2 = OA \times OB$ , is a.  $(y + x)^2 = 50$  b.  $(y - x)^2 = 200$  c.

$(y - x)^2 = 100$  d. none of these

A.  $(y - x)^2 = 100$

B.  $(y + x)^2 = 50$

C.  $(y - x)^2 = 200$

D. none of these

**Answer: C**



**Watch Video Solution**

9. A variable line L drawn through  $O(0,0)$  to meet line  $l_1: y-x-10=0$  and  $L_2: y-x-20=0$  at the point A and B respectively then locus of point p is ' such that  $(OP)^2 = OA \cdot OB$ ,

A.  $(y - x)^2 = 80$

B.  $(y - x)^2 = 100$

C.  $(y - x)^2 = 64$

D. none of these

**Answer: A**

 [Watch Video Solution](#)

10. The line  $6x+8y=48$  intersects the coordinates axes at A and B, respectively. A line L bisects the area and the perimeter of triangle OAB, where O is the origin.

The number of such lines possible is a. 1 b. 2 c. 3 d. 4

A. 1

B. 2

C. 3

D. more than 3

**Answer: A**

 [Watch Video Solution](#)

11. if a line has direction ratio 2,-1,-2,determine its direction cosine

 [Watch Video Solution](#)

12. The line  $6x+8y=48$  intersects the coordinates axes at A and B, respectively.

A line L bisects the area and the perimeter of triangle OAB, where O is the origin.

Line L

- A. does not intersect AB
- B. does not intersect OB
- C. does not intersect OA
- D. can intersect all the sides

**Answer: C**

 [Watch Video Solution](#)

13.  $A(1, 3)$  and  $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$  are the vertices of a  $\Delta ABC$  and the equation of the angle bisector of  $\angle ABC$  is  $x + y = 2$ . find the equation of 'BC'

A.  $7x+3y-4=0$

B.  $7x+3y+4=0$

C.  $7x-3y+4=0$

D.  $7x-3y-4=0$

**Answer: B**

 [Watch Video Solution](#)

14.  $A(1, 3)$  and  $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$  are the vertices of a  $\Delta ABC$  and the equation of the angle bisector of  $\angle ABC$  is  $x + y = 2$ .

A. (A)  $(\frac{3}{10}, \frac{17}{10})$

B. (B)  $(\frac{17}{10}, \frac{3}{10})$

C. (C)  $(-5/2, 9/2)$

D. (D)  $(-1,1)$

**Answer: C**



[Watch Video Solution](#)

15.  $A(1, 3)$  and  $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$  are the vertices of a  $\triangle ABC$  and the equation of the angle bisector of  $\angle ABC$  is  $x + y = 2$ .

A.  $3x+7y=24$

B.  $3x+7y+24=0$

C.  $13x+7y+8=0$

D.  $13x-7y+8=0$

**Answer: A**



[Watch Video Solution](#)

16. Let ABCD be a parallelogram the equation of whose diagonals are  $AC: x + 2y = 3$ ;  $BD: 2x + y = 3$ . If length of diagonal  $AC = 4$  units and area of  $ABCD = 8$  sq. units. Find the length of the other diagonal is a.  $10/3$  b. 2 c.  $20/3$  d. None of these

A.  $10/3$

B. 2

C.  $20/3$

D. none of these

**Answer: C**



[Watch Video Solution](#)

17. ABCD is a parallelogram.  $x + 2y = 3$  and  $2x + y = 3$  are the equations of the diagonals AC and BD respectively.  $AC = 4$  units and area of parallelogram ABCD is 8 sq. units then The length of BC is equal to

A.  $\sqrt{232}/3$

B.  $4\sqrt{58}/9$

C.  $3\sqrt{58}/9$

D.  $4\sqrt{58}/9$

**Answer: A**

 [Watch Video Solution](#)

**18.** Let  $ABCD$  be a parallelogram the equation of whose diagonals are  $AC: x + 2y = 3$ ;  $BD: 2x + y = 3$ . If length of diagonal  $AC = 4$  units and area of  $ABCD = 8$  sq. units. Then

(i) The length of the other diagonal is

(ii) the length of side  $AB$  is equal to

 [Watch Video Solution](#)



19. Consider a triangle PQR with coordinates of its vertices as P(-8,5), Q(-15, -19), and R (1, -7). The bisector of the interior angle of P has the equation which can be written in the form  $ax+2y+c=0$ .

The distance between the orthocenter and the circumcenter of triangle PQR is

A.  $25/2$

B.  $29/2$

C.  $37/2$

D.  $51/2$

**Answer: A**



[Watch Video Solution](#)

20. Consider a triangle PQR with coordinates of its vertices as P(-8,5), Q(-15, -19), and R (1, -7). The bisector of the interior angle of P has the equation

which can be written in the form  $ax+2y+c=0$ .

The radius of the in circle of triangle PQR is

A. 4

B. 5

C. 6

D. 8

**Answer: B**



[Watch Video Solution](#)

21. Consider a triangle PQR with coordinates of its vertices as P(-8,5), Q(-15, -19), and R (1, -7). The bisector of the interior angle of P has the equation which can be written in the form  $ax+2y+c=0$ .

The radius of the in circle of triangle PQR is

The sum  $a + c$  is

A. 129

B. 78

C. 89

D. none of these

**Answer: C**

 [Watch Video Solution](#)

22. The base of an isosceles triangle measures 4 units base angle is equal to  $45^\circ$ . A straight line cuts the extension of the base at a point M at the angle  $\theta$  and bisects the lateral side of the triangle which is nearest to M.

The area of quadrilateral which the straight line cuts off from the given triangle is

A.  $\frac{3 + \tan\theta}{1 + \tan\theta}$

B.  $\frac{3 + 5\tan\theta}{1 + \tan\theta}$

C.  $\frac{3 + \tan\theta}{1 - \tan\theta}$

D.  $\frac{3 + 2\tan\theta}{1 + \tan\theta}$

**Answer: B**

 [Watch Video Solution](#)

23. The base of an isosceles triangle measures 4 units base angle is equal to  $45^\circ$ . A straight line cuts the extension of the base at a point M at the angle  $\theta$  and bisects the lateral side of the triangle which is nearest to M.

The possible range of values in which area of quadrilateral which straight line cuts off from the given triangle lie in (a)  $(\frac{5}{2}, \frac{7}{2})$  (b) (4,3) (c) (4,5) (d) (3,4)

A.  $\left(\frac{5}{2}, \frac{7}{2}\right)$

B. (4,3)

C. (4,5)

D. (3,4)

**Answer: D**

 [Watch Video Solution](#)

24. The base of an isosceles triangle measures 4 units base angle is equal to  $45^\circ$ . A straight line cuts the extension of the base at a point M at the angle  $\theta$  and bisects the lateral side of the triangle which is nearest to M.

The length of portion of straight line inside the triangle may lie in the range

A. (2,4)

B.  $\left(\frac{3}{2}, \sqrt{3}\right)$

C.  $(\sqrt{2}, 2)$

D.  $(\sqrt{2}, \sqrt{3})$

**Answer: C**



[Watch Video Solution](#)

25. Consider point A(6, 30), point B(24, 6) and line AB:  $4x+3y = 114$ .

Point  $P(0, \lambda)$  is a point on y-axis such that

$0 < \lambda < 38$  and point  $Q(0, \lambda)$  is a point on y-axis such that  $\lambda > 38$ .

For all positions of point P, angle APB is maximum when point P is

A. (0, 12)

B. (0, 15)

C. (0, 18)

D. (0, 21)

**Answer: C**



[Watch Video Solution](#)

**26.** Consider point A(6, 30), point B(24, 6) and line AB:  $4x+3y = 114$ .

Point  $P(0, \lambda)$  is a point on y-axis such that

$0 < \lambda < 38$  and point  $Q(0, \lambda)$  is a point on y-axis such that  $\lambda > 38$ .

The maximum value of angle APB is

A.  $\frac{\pi}{3}$

B.  $\frac{\pi}{2}$

C.  $\frac{2\pi}{3}$

D.  $\frac{3\pi}{3}$

**Answer: B**

 [Watch Video Solution](#)

27. Consider point A(6, 30), point B(24, 6) and line AB:  $4x+3y = 114$ .

Point  $P(0, \lambda)$  is a point on y-axis such that

$0 < \lambda < 38$  and point  $Q(0, \lambda)$  is a point on y-axis such that  $\lambda > 38$ .

For all positions of point Q, and AQB is maximum when point Q is

A. (0, 54)

B. (0, 58)

C. (0, 60)

D. (0, 1)

**Answer: B**

 [Watch Video Solution](#)

**Exercise Matrix Match Type**

1. Match the following lists:

List I	List II
<b>a.</b> Four lines $x + 3y - 10 = 0$ , $x + 3y - 20 = 0$ , $3x - y + 5 = 0$ , and $3x - y - 5 = 0$ form a figure which is	<b>p.</b> a quadrilateral which is neither a parallelogram nor a trapezium
<b>b.</b> The points $A(1, 2)$ , $B(2, -3)$ , $C(-1, -5)$ , and $D(-2, 4)$ in order are the vertices of	<b>q.</b> a parallelogram
<b>c.</b> The lines $7x + 3y - 33 = 0$ , $3x - 7y + 19 = 0$ , $3x - 7y - 10$ , and $7x + 3y - 4 = 0$ form a figure which is	<b>r.</b> a rectangle of area 10 sq. units
<b>d.</b> Four lines $4y - 3x - 7 = 0$ , $3y - 4x + 7 = 0$ , $4y - 3x - 21 = 0$ , $3y - 4x + 14 = 0$ form a figure which is	<b>s.</b> a square

6. Match the following lists:



## 2. Match the following lists:

List I	List II
<b>a.</b> The lines $y = 0$ ; $y = 1$ ; $x - 6y + 4 = 0$ , and $x + 6y - 9 = 0$ constitute a figure which is	<b>p.</b> a cyclic quadrilateral
<b>b.</b> The points $A(a, 0)$ , $B(0, b)$ , $C(c, 0)$ , and $D(0, d)$ are such that $ac = bd$ and $a, b, c, d$ are all positive. The points $A, B, C$ , and $D$ always constitute	<b>q.</b> a rhombus
<b>c.</b> The figure formed by the four lines $ax \pm by \pm c = 0$ , $a \neq b$ , is	<b>r.</b> a square
<b>d.</b> The line pairs $x^2 - 8x + 12 = 0$ and $y^2 - 14y + 45 = 0$ constitute a figure which is	<b>s.</b> a trapezium



Watch Video Solution

## 3. Match the following lists:

List I	List II
<b>a.</b> If lines $3x + y - 4 = 0$ , $x - 2y - 6 = 0$ , and $\lambda x + 4y + \lambda^2 = 0$ are concurrent, then the value of $\lambda$ is	<b>p.</b> $-4$
<b>b.</b> If the points $(\lambda + 1, 1)$ , $(2\lambda + 1, 3)$ , and $(2\lambda + 2, 2\lambda)$ are collinear, then the value of $\lambda$ is	<b>q.</b> $-1/2$
<b>c.</b> If the line $x + y - 1 -  \lambda/2  = 0$ , passing through the intersection of $x - y + 1 = 0$ and $3x + y - 5 = 0$ , is perpendicular to one of them, then the value of $\lambda$ is	<b>r.</b> $4$
<b>d.</b> If the line $y - x - 1 + \lambda = 0$ is equidistant from the points $(1, -2)$ and $(3, 4)$ , then $\lambda$ is	<b>s.</b> $2$

## 5. Match the following lists:



Watch Video Solution

4. Match the following lists:

List I	List II
a. If lines $3x + y - 4 = 0$ , $x - 2y - 6 = 0$ , and $\lambda x + 4y + \lambda^2 = 0$ are concurrent, then the value of $\lambda$ is	p. $-4$
b. If the points $(\lambda + 1, 1)$ , $(2\lambda + 1, 3)$ , and $(2\lambda + 2, 2\lambda)$ are collinear, then the value of $\lambda$ is	q. $-1/2$
c. If the line $x + y - 1 -  \lambda/2  = 0$ , passing through the intersection of $x - y + 1 = 0$ and $3x + y - 5 = 0$ , is perpendicular to one of them, then the value of $\lambda$ is	r. $4$
d. If the line $y - x - 1 + \lambda = 0$ is equidistant from the points $(1, -2)$ and $(3, 4)$ , then $\lambda$ is	s. $2$

5. Match the following lists:



Watch Video Solution

5. Match the following lists:

List I	List II
a. Four lines $x + 3y - 10 = 0$ , $x + 3y - 20 = 0$ , $3x - y + 5 = 0$ , and $3x - y - 5 = 0$ form a figure which is	p. a quadrilateral which is neither a parallelogram nor a trapezium
b. The points $A(1, 2)$ , $B(2, -3)$ , $C(-1, -5)$ , and $D(-2, 4)$ in order are the vertices of	q. a parallelogram
c. The lines $7x + 3y - 33 = 0$ , $3x - 7y + 19 = 0$ , $3x - 7y - 10 = 0$ , and $7x + 3y - 4 = 0$ form a figure which is	r. a rectangle of area 10 sq. units
d. Four lines $4y - 3x - 7 = 0$ , $3y - 4x + 7 = 0$ , $4y - 3x - 21 = 0$ , $3y - 4x + 14 = 0$ form a figure which is	s. a square

6. Match the following lists:

 Watch Video Solution

6. Match the following lists:

List I	List II
a. The lines $y = 0$ ; $y = 1$ ; $x - 6y + 4 = 0$ , and $x + 6y - 9 = 0$ constitute a figure which is	p. a cyclic quadrilateral
b. The points $A(a, 0)$ , $B(0, b)$ , $C(c, 0)$ , and $D(0, d)$ are such that $ac = bd$ and $a, b, c, d$ are all positive. The points $A, B, C$ , and $D$ always constitute	q. a rhombus
c. The figure formed by the four lines $ax \pm by \pm c = 0$ , $a \neq b$ , is	r. a square
d. The line pairs $x^2 - 8x + 12 = 0$ and $y^2 - 14y + 45 = 0$ constitute a figure which is	s. a trapezium



Watch Video Solution

7. Consider the lines given by

$$L_1: x + 3y - 5 = 0$$

$$L_2: 3x - ky - 1 = 0$$

$$L_3: 5x + 2y - 12 = 0$$

Match the following lists.

List I	List II
a. $L_1, L_2, L_3$ are concurrent if	p. $k = -9$
b. One of $L_1, L_2, L_3$ is parallel to at least one of the other two if	q. $k = -6/5$
c. $L_1, L_2, L_3$ form a triangle if	r. $k = 5/6$
d. $L_1, L_2, L_3$ do not form a triangle if	s. $k = 5$



Watch Video Solution

8. Consider a  $\triangle ABC$  in which sides AB and AC are perpendicular to  $x-y-4=0$  and  $2x-y-5=0$ , respectively. Vertex A is  $(-2, 3)$  and the circumcenter of  $\triangle ABC$  is  $(3/2, 5/2)$ .

The equation of the line in List I is of the form  $ax+by+c=0$ , where  $a, b, c \in I$ . Match it with the corresponding value of  $c$  in list II and then choose the correct code.

List I	List II
a. Equation of the perpendicular bisector of side $AB$	p. $-1$
b. Equation of the perpendicular bisector of side $AC$ .	q. $1$
c. Equation of side $AC$	r. $-16$
d. Equation of the median through $A$	s. $-4$

Codes :

$a \ b \ c \ d$

$r \ s \ p \ q$

$s \ r \ q \ p$

$q \ p \ s \ r$

$r \ p \ s \ q$

 [View Text Solution](#)

Exercise Numerical Value Type

1. about to only mathematics

 [Watch Video Solution](#)

2. The number of values of  $k$  for which the lines  $(k + 1)x + 8y = 4k$  and  $kx + (k + 3)y = 3k - 1$  are coincident is \_\_\_\_\_

 [Watch Video Solution](#)

3. about to only mathematics

 [Watch Video Solution](#)

4. The absolute value of the sum of the abscissas of all the points on the line  $x + y = 4$  that lie at a unit distance from the line  $4x + 3y - 10 = 0$  is \_\_\_\_\_

 [Watch Video Solution](#)

5. Two sides of a rectangle are  $3x+4y+5=0$ ,  $4x-3y+15=0$  and one of its vertices is  $(0, 0)$ . The area of rectangle is \_\_\_.

 Watch Video Solution

6. about to only mathematics

 Watch Video Solution

7. For all real values of  $a$  and  $b$  lines  $(2a + b)x + (a + 3b)y + (b - 3a) = 0$  and  $mx + 2y + 6 = 0$  are concurrent, then  $m$  is equal to (A) -2 (B) -3 (C) -4 (D) -5

 Watch Video Solution

8. The line  $3x + 2y = 24$  meets the  $y$ -axis at  $A$  and the  $x$ -axis at  $B$ . The perpendicular bisector of  $AB$  meets the line through  $(0, -1)$  parallel to

the x-axis at  $C$ . If the area of triangle  $ABC$  is  $A$ , then the value of  $\frac{A}{13}$

is \_\_\_\_\_

 Watch Video Solution

9. about to only mathematics

 Watch Video Solution

10. Triangle  $ABC$  with  $AB = 13$ ,  $BC = 5$ , and  $AC = 12$  slides on the coordinates axes with  $A$  and  $B$  on the positive x-axis and positive y-axis respectively. The locus of vertex  $C$  is a line  $12x - ky = 0$ . Then the value of  $k$  is \_\_\_\_\_

 Watch Video Solution

11. The line  $y = \frac{3x}{4}$  meets the lines  $x - y + 1 = 0$  and  $2x - y = 5$  at  $A$  and  $B$  respectively. Coordinates of  $P$  on  $y = \frac{3x}{4}$  such that  $PA \cdot PB = 25$ .



 [Watch Video Solution](#)

12. In a plane there are two families of lines  $y = x + r, y = -x + r$ , where  $r \in \{0, 1, 2, 3, 4\}$ . The number of squares of diagonals of length 2 formed by the lines is:

 [Watch Video Solution](#)

13. If  $5a + 5b + 20c = t$ , then find the value of  $t$  for which the line  $ax + by + c - 1 = 0$  always passes through a fixed point.

 [Watch Video Solution](#)

Archives Jee Main

1. The line L given by  $\frac{x}{5} + \frac{y}{b} = 1$  passes through the point (13,32). The line K is parallel to L and has the equation  $\frac{x}{c} + \frac{y}{3} = 1$  then the distance

between L and K is

A.  $\frac{23}{\sqrt{17}}$

B.  $\frac{23}{\sqrt{15}}$

C.  $\sqrt{17}$

D.  $\frac{17}{\sqrt{15}}$

**Answer: A**



**Watch Video Solution**

2. The line  $L_1: y - x = 0$  and  $L_2: 2x + y = 0$  intersect the line  $L_3: y + 2 = 0$  at P and Q respectively. The bisector of the acute angle between  $L_1$  and  $L_2$  intersects  $L_3$  at R. Statement-1 : The ratio  $PR:RQ$  equals  $2\sqrt{2}:\sqrt{5}$  Statement-2 : In any triangle, bisector of an angle divides the triangle into two similar triangles. Statement-1 is true, Statement-2 is true ; Statement-2 is correct explanation for Statement-1 Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for

Statement-1 Statement-1 is true, Statement-2 is false Statement-1 is false,  
Statement-2 is true

- A. Statement 1 is true, statement 2 is false.
- B. Statement 1 is true, statement 2 is true, statement 2 is the correct explanation of statement1.
- C. Statement 1 is true, statement 2 is true, statement 2 is not the correct explanation of statement 1.
- D. Statement 1 is false, statement 2 is true.

**Answer: A**

 [Watch Video Solution](#)

3. A line is drawn through the point (1, 2) to meet the coordinate axes at P and Q such that it forms a triangle OPQ, where O is the origin. If the area of the triangle OPQ is least, then the slope of the line PQ is

A.  $-\frac{1}{4}$

B. -4

C. -2

D.  $-\frac{1}{2}$

**Answer: C**

 [Watch Video Solution](#)

4. The x-coordinate of the incentre of the triangle that has the coordinates of mid points of its sides as  $(0, 1)$ ,  $(1, 1)$  and  $(1, 0)$  is

A.  $2 + \sqrt{2}$

B.  $2 - \sqrt{2}$

C.  $1 + \sqrt{2}$

D.  $1 - \sqrt{2}$

**Answer: B**

 [Watch Video Solution](#)

5. A ray of light along  $x + \sqrt{3}y = \sqrt{3}$  gets reflected upon reaching x-axis, the equation of the reflected ray is

A.  $y = x + \sqrt{3}$

B.  $\sqrt{3}y = x - \sqrt{3}$

C.  $y = \sqrt{3}x - \sqrt{3}$

D.  $\sqrt{3}y = x - 1$

**Answer: B**

 [Watch Video Solution](#)

6. Let  $a, b, c$  and  $d$  be non-zero numbers. If the point of intersection of the lines  $4ax + 2ay + c = 0$  and  $5bx + 2by + d = 0$  lies in the fourth quadrant and is equidistant from the two axes, then

A.  $2bc - 3ad = 0$

B.  $2bc+3ad=0$

C.  $3bc-2ad=0$

D.  $3bc+2ad=0$

**Answer: C**

 [Watch Video Solution](#)

7. Let PS be the median of the triangle with vertices  $P(2, 2)$ ,  $Q(6, -1)$  and  $R(7, 3)$ . The equation of the line passing through  $(1, -1)$  and parallel to PS is (1)  $4x - 7y - 11 = 0$  (2)  $2x + 9y + 7 = 0$  (3)  $4x + 7y + 3 = 0$  (4)  $2x - 9y - 11 = 0$

A.  $4x-7y-1=0$

B.  $2x+9y+7=0$

C.  $4x+7y+3=0$

D.  $2x-9y-11=0$

**Answer: B**

 [Watch Video Solution](#)

8. Locus of the image of the point  $(2, 3)$  in the line  $(2x - 3y + 4) + k(x - 2y + 3) = 0, k \in R$ , is a :

(1) straight line parallel to x-axis. (2) straight line parallel to y-axis (3) circle of radius  $\sqrt{2}$  (4) circle of radius  $\sqrt{3}$

A. Straight line parallel to x-axis

B. straight line parallel to y-axis

C. circle of radius  $\sqrt{2}$

D. circle of radius 3

**Answer: C**

 [Watch Video Solution](#)

9. Two sides of a rhombus are along the lines,  $x - y + 1 = 0$  and  $7x - y - 5 = 0$ . If its diagonals intersect at  $(-1, -2)$ , then which one of the following is a vertex of this rhombus ? (1)  $(-3, -9)$  (2)  $(-3, -8)$  (3)  $\left(\frac{1}{3}, -\frac{8}{3}\right)$  (4)  $\left(-\frac{10}{3}, -\frac{7}{3}\right)$

A.  $(-3, -8)$

B.  $\left(\frac{1}{3}, -\frac{8}{3}\right)$

C.  $\left(-\frac{10}{3}, -\frac{7}{3}\right)$

D.  $(-3, -9)$

**Answer: B**

 [Watch Video Solution](#)

Archives Jee Advanced

1. The locus of the orthocentre of the triangle formed by the lines  $(1 + p)x - py + p(1 + p) = 0$ ,  $(1 + q)x - qy + q(1 + q) = 0$  and  $y = 0$ ,



where  $p \neq q$ , is (A) a hyperbola (B) a parabola (C) an ellipse (D) a straight line

A. a hyperbola

B. a parabola

C. an ellipse

D. a straight line

**Answer: D**

 [Watch Video Solution](#)

2. A straight line L through the point (3,-2) is inclined at an angle  $60^\circ$  to the line  $\sqrt{3}x + y = 1$  If L also intersects the x-axis then the equation of L is

A.  $y + \sqrt{3}x + 2 - 3\sqrt{3} = 0$

B.  $y - \sqrt{3}x + 2 + 3\sqrt{3} = 0$

C.  $\sqrt{3}y - x + 3 + 2\sqrt{3} = 0$

$$D. \sqrt{3}y + x - 3 + 2\sqrt{3} = 0$$

**Answer: B**



**Watch Video Solution**

3. For  $a > b > c > 0$ , if the distance between  $(1, 1)$  and the point of intersection of the line  $ax + by - c = 0$  and  $bx + ay + c = 0$  is less than  $2\sqrt{2}$  then, (A)  $a + b - c > 0$  (B)  $a - b + c < 0$  (C)  $a - b + c > 0$  (D)  $a + b - c < 0$

A.  $a + b - c > 0$

B.  $a - b + c < 0$

C.  $a - b + c > 0$

D.  $a + b - c < 0$

**Answer: A**



**Watch Video Solution**

## Archives Numerical Value Type

1. For a point  $P$  in the plane, let  $d_1(P)$  and  $d_2(P)$  be the distances of the point  $P$  from the lines  $x - y = 0$  and  $x + y = 0$  respectively. The area of the region  $R$  consisting of all points  $P$  lying in the first quadrant of the plane and satisfying  $2 \leq d_1(P) + d_2(P) \leq 4$ , is



[Watch Video Solution](#)

## Single Correct Answer Type

1. In the  $xy$ -plane, how many straight lines whose  $x$ -intercept is a prime number and whose  $y$ -intercept is a positive integer pass through the point  $(4,3)$ ?

A. 1

B. 2

C. 3

D. 4

**Answer: B**

 [Watch Video Solution](#)

2. The condition that the equation  $lx + my + n = 0$  represents the equation of a straight line in the normal form is

A.  $l^2 + m^2 \pm 0, n > 0$

B.  $l^2 + m^2 \pm 0, n < 0$

C.  $l^2 + m^2 = 1, n < 0$

D.  $l^2 + m^2 = 1, n > 0$

**Answer: C**

 [Watch Video Solution](#)

3. In an isosceles triangle  $ABC$ , the coordinates of the points  $B$  and  $C$  on the base  $BC$  are respectively  $(1, 2)$  and  $(2, 1)$ . If the equation of the line  $AB$  is  $y = 2x$ , then the equation of the line  $AC$  is

A.  $y = \frac{1}{2}(x - 1)$

B.  $y = \frac{x}{2}$

C.  $y = x - 1$

D.  $2y = x + 3$

**Answer: B**



**Watch Video Solution**

4. If the coordinates of the points  $A, B, C$  be  $(-1, 5), (0, 0)$  and  $(2, 2)$  respectively, and  $D$  be the middle point of  $BC$ , then the equation of the perpendicular drawn from  $B$  to the line  $AD$  is

A.  $x + 2y = 0$

B.  $2x + y = 0$

C.  $x - 2y = 0$

D.  $2x - y = 0$

**Answer: C**

 [Watch Video Solution](#)

5. Two lines are drawn through  $(3,4)$ , each of which makes angle of  $45^\circ$  with the line  $x - y = 2$ . Then area of the triangle formed by these lines is (a) 9

(b)  $\frac{9}{2}$  (c) 2 (d)  $\frac{2}{9}$

A. 9

B.  $9/2$

C. 2

D.  $2/9$

**Answer: B**



[Watch Video Solution](#)

6. The line  $y = 2x + 4$  is shifted 2 units along  $+y$  axis, keeping parallel to itself and then 1 unit along  $+x$  axis direction in the same manner, then equation of the line in its new position is,

A.  $y = 2x + 6$

B.  $y = 2x + 5$

C.  $y = 2x + 4$

D. none of these

**Answer: C**



[Watch Video Solution](#)

7. A ray of light passing through the point  $A(2, 3)$  reflected at a point  $B$  on line  $x + y = 0$  and then passes through  $(5, 3)$ . Then the coordinates of  $B$  are

A.  $\left(\frac{1}{3}, -\frac{1}{3}\right)$

B.  $\left(\frac{2}{5}, -\frac{2}{5}\right)$

C.  $\left(\frac{1}{13}, -\frac{1}{13}\right)$

D. none of these

**Answer: C**



**Watch Video Solution**

8. If the transversal  $y = m_r x$  :  $r = 1, 2, 3$  cut off equal intercepts on the transversal  $x + y = 1$  then  $1 + m_1, 1 + m_2, 1 + m_3$  are in

A. A.P.

B. G.P.

C. H.P.

D. None of these

**Answer: C**





Watch Video Solution

9. The straight line  $y = x - 2$  rotates about a point where it cuts x-axis and become perpendicular on the straight line  $ax + by + c = 0$  then its equation is

A.  $ax + by + 20 = 0$

B.  $ax - by - 2a = 0$

C.  $bx + ay - 2b = 0$

D.  $ay - bx + 2b = 0$

**Answer: D**



Watch Video Solution

10. The two adjacent sides of parallelogram are  $y = 0$  and  $y = \sqrt{3}(x - 1)$ . If equation of one diagonal is  $\sqrt{3}y = (x + 1)$ , then equation of other diagonal is

A.  $\sqrt{3}y = (x - 1)$

B.  $y = \sqrt{3}(x + 1)$

C.  $y = -\sqrt{3}(x - 1)$

D.  $\sqrt{3}y = -(x + 1)$

**Answer: C**

 [Watch Video Solution](#)

11. A(3,0) and B(6,0) are two fixed points and  $U(x_1, y_1)$  is a variable point of the plane. AU and BU meet the y axis at C and D respectively and AD meets OU at V. Then for any position of U in the plane CV passes through fixed point (p,q) whose distance from origin is \_\_\_ units

A. 1units

B. 2 units

C. 3 units

D. 4 units

**Answer: B**



**Watch Video Solution**

12. If  $h$  denotes the A.M. and  $k$  denote G.M. of the intercept made on axes by the lines passing through  $(1, 1)$  then  $(h, k)$  lies on

A.  $y^2 = 2x$

B.  $y^2 = 4x$

C.  $y = 2x$

D.  $x + y = 2xy$

**Answer: A**



**Watch Video Solution**

13. Let  $A(a, 0)$  and  $B(b, 0)$  be fixed distinct points on the x-axis, none of which coincides with the  $O(0, 0)$ , and let  $C$  be a point on the y-axis. Let  $L$  be

a line through the  $O(0, 0)$  and perpendicular to the line AC. The locus of the point of intersection of the lines L and BC if C varies along is (provided  $c^2 + ab \neq 0$ )

A. (a)  $\frac{x^2}{a} + \frac{y^2}{b} = x$

B. (b)  $\frac{x^2}{a} + \frac{y^2}{b} = y$

C. (c)  $\frac{x^2}{b} + \frac{y^2}{a} = x$

D. (d)  $\frac{x^2}{b} + \frac{y^2}{a} = y$

**Answer: C**

 [Watch Video Solution](#)

14. If AD, BE and CF are the altitudes of a triangle ABC whose vertex A is the point  $(-4, 5)$ . The coordinates of the points E and F are  $(4, 1)$  and  $(-1, -4)$  respectively, then equation of BC is

A.  $3x - 4y - 28 = 0$

B.  $4x + 3y - 28 = 0$

C.  $3x - 4y + 28 = 0$

D.  $x + 2y + 7 = 0$

**Answer: A**



**Watch Video Solution**

15. Let P and Q be any two points on the lines represented by  $2x - 3y = 0$  and  $2x + 3y = 0$  respectively. If the area of triangle OPQ (where O is origin) is 5, then which of the following is not the possible equation of the locus of mid-point of PO?

A.  $4x^2 - 9y^2 + 30 = 0$

B.  $4x^2 - 9y^2 - 30 = 0$

C.  $9x^2 - 4y^2 - 30 = 0$

D. none of these

**Answer: C**



16. The acute angle between two straight lines passing through the point  $M(-6, -8)$  and the points in which the line segment  $2x + y + 10 = 0$  enclosed between the co-ordinate axes is divided in the ratio 1:2:2 in the direction from the point of its intersection with the x-axis to the point of intersection with the y-axis is: (a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{6}$  (d)  $\frac{\pi}{12}$

A.  $\pi/3$

B.  $\pi/4$

C.  $\pi/6$

D.  $\pi/12$

**Answer: B**

17. A variable line  $L$  is drawn through  $O(0,0)$  to meet the lines  $L_1: y-x-10=0$  and  $L_2: y-x-20=0$  at the points  $A$  and  $B$  respectively. A point  $P$  is taken on  $L$  such that  $OP^2 = OA^2 + OB^2$  and  $P, A, B$  lies on same side of origin  $O$ . The locus of  $P$  is

A.  $9x + 6y = 20$

B.  $6x - 9y = 20$

C.  $6x + 9y = 20$

D. none of these

**Answer: C**

 [Watch Video Solution](#)

18. The complete set of values of the parameter  $\alpha$  so that the point  $P(\alpha, (1 + \alpha^2)^{-1})$  does not lie outside the triangle formed by the lines  $L_1: 15y = x + 1$ ,  $L_2: 78y = 118 - 23x$  and  $L_3: y + 2 = 0$  is

A.  $(0, 5)$

B.  $[2, 5]$

C.  $[1, 5]$

D.  $[0, 2]$

**Answer: C**

 [Watch Video Solution](#)

19. if  $P, Q$  are two points on the line  $3x + 4y + 15 = 0$  such that  $OP = OQ = 9$  then the area of triangle  $OPQ$  is

A. 18 sq. units

B.  $18\sqrt{2}$  sq. units

C. 27 sq. units

D. none of these

**Answer: B**





Watch Video Solution

20. The number of points on the line  $3x + 4y = 5$ , which are at a distance of  $\sec^2 \theta + 2\cos \sec^2 \theta$ ,  $\theta \in R$ , from the point (1,3) is (a) 1 (b) 2 (c) 3 (d) infinite

A. 1

B. 2

C. 3

D. infinite

**Answer: B**



Watch Video Solution

21. ABC is an equilateral triangle whose centroid is origin and base BC is along the line  $11x + 60y = 122$ . Then

- A. Area of the triangle is numerically equal to the perimeter
- B. Area of triangle is numerically double the perimeter
- C. Area of triangle is numerically three times the perimeter
- D. Area of triangle is numerically half of the perimeter

**Answer: A**



**Watch Video Solution**

22. If the distance of a given point  $(\alpha, \beta)$  from each of two straight lines  $y = mx$  through the origin is  $d$ , then  $(\alpha\gamma - \beta x)^2$  is equal to (a)  $x^2 + y^2$   
(b)  $d^2(x^2 + y^2)$  (c)  $d^2$  (d) none of these

A.  $x^2 + y^2$

B.  $d^2(x^2 + y^2)$

C.  $d^2$

D. none of these

**Answer: B**



**Watch Video Solution**

23. The values of  $k$  for which lines  $kx + 2y + 2 = 0$ ,  $2x + ky + 3 = 0$ ,  $3x + 3y + k = 0$  are concurrent are

A.  $\{2, 3, 5\}$

B.  $\{2, 3, -5\}$

C.  $\{3, -5\}$

D.  $\{-5\}$

**Answer: C**



**Watch Video Solution**

24. The set of real values of  $k$  for which the lines  $x + 3y + 1 = 0$ ,  $kx + 2y - 2 = 0$  and  $2x - y + 3 = 0$  form a triangle is

A.  $R - \left\{ -4, \frac{2}{3} \right\}$

B.  $R - \left\{ -4, \frac{-6}{5}, \frac{2}{3} \right\}$

C.  $R - \left\{ \frac{-2}{3}, 4 \right\}$

D. R

**Answer: B**

 [Watch Video Solution](#)

25. Locus of the points which are at equal distance from  $3x + 4y - 11 = 0$  and  $12x + 5y + 2 = 0$  and which is near the origin is: (a)  $21x - 77y + 153 = 0$  (b)  $99x + 77y - 133 = 0$  (c)  $7x - 11y = 19$  (d) None of these

A.  $21x - 77y + 153 = 0$

B.  $99x + 77y - 133 = 0$

C.  $7x - 11y = 19$

D. None of these

**Answer: B**

 [Watch Video Solution](#)

26. Pair of lines through  $(1, 1)$  and making equal angle with  $3x - 4y = 1$  and  $12x + 9y = 1$  intersect x-axis at  $P_1$  and  $P_2$ , then  $P_1, P_2$  may be

A. (a)  $\left(\frac{8}{7}, 0\right)$  and  $\left(\frac{9}{7}, 0\right)$

B. (b)  $\left(\frac{6}{7}, 0\right)$  and  $(8, 0)$

C. (c)  $\left(\frac{8}{7}, 0\right)$  and  $\left(\frac{1}{8}, 0\right)$

D. (d)  $(8, 0)$  and  $\left(\frac{1}{8}, 0\right)$

**Answer: B**

 [Watch Video Solution](#)

27. The algebraic sum of distances of the line  $ax + by + 2 = 0$  from  $(1, 2), (2, 1)$  and  $(3, 5)$  is zero and the lines  $bx - ay + 4 = 0$  and

$3x + 4y + 5 = 0$  cut the coordinate axes at concyclic points. Then (a)

$a + b = -\frac{2}{7}$  (b) area of triangle formed by the line  $ax + by + 2 = 0$  with

coordinate axes is  $\frac{14}{5}$  (c) line  $ax + by + 3 = 0$  always passes through the

point  $(-1, 1)$  (d)  $\max\{a, b\} = \frac{5}{7}$

A.  $a + b = -\frac{2}{7}$

B. area of the triangle formed by the line  $ax + by + 2 = 0$  with

coordinate axes is  $\frac{14}{5}$

C. line  $ax + by + 3 = 0$  always passes through the point  $(-1, 1)$

D.  $\max\{a, b\} = \frac{5}{7}$

**Answer: C**

 [Watch Video Solution](#)

28. Equation of line which is equally inclined to the axis and passes through

a common points of family of lines

$$4acx + y(ab + bc + ca - abc) + abc = 0 \text{ (where } a, b, c > 0 \text{ are in } H. P. \text{)}$$

is

A.  $y - x = \frac{7}{4}$

B.  $y - x = -\frac{7}{4}$

C.  $y - x = \frac{1}{4}$

D.  $y - x = -\frac{1}{4}$

**Answer: A**



**Watch Video Solution**

29. The base  $BC$  of a  $ABC$  is bisected at the point  $(p, q)$  & the equation to the side  $AB$  &  $AC$  are  $px + qy = 1$  &  $qx + py = 1$ . The equation of the median through  $A$  is:  $(p - 2q)x + (q - 2p)y + 1 = 0$

$$(p + q)(x + y) - 2 = 0$$

$$(2pq - 1)(px + qy - 1) = (p^2 + q^2 - 1)(qx + py - 1) \text{ none of these}$$

A.  $qx - py = 0$

$$B. \frac{x}{p} + \frac{y}{q} = 2$$

$$C. (2pq - 1)(px + qy - 1) = (p^2 + q^2 - 1)(qx + py - 1)$$

$$D. (p - 2q)x + (q - 2p)y = p^2 + r^2$$

**Answer: C**

 [Watch Video Solution](#)

**30.**  $A(x_1, y_1), B(x_2, y_2), C(x_3, y_3)$  are three vertices of a triangle ABC.

$lx + my + n = 0$  is an equation of the line L. If the centroid of the triangle ABC is at the origin and algebraic sum of the lengths of the perpendicular from O the vertices of triangle ABC on the line L is equal to, then sum of the squares of reciprocals of the intercepts made by L on the coordinate axes is equal to

A. (a) 0

B. (b) 4

C. (c) 9



D. (d) 16

**Answer: C**

 [Watch Video Solution](#)

31. A straight line passes through the point of Intersection of the lines  $x - 2y - 2 = 0$  and  $2x - by - 6 = 0$  and the origin, then the set of values of 'b' for which the acute angle between this line and  $y = 0$  is less than  $45^\circ$  is

- A.  $(-\infty, 4) \cup (7, \infty)$
- B.  $(-\infty, 5) \cup (7, \infty)$
- C.  $(-\infty, 4) \cup (5, 7) \cup (7, \infty)$
- D.  $(-\infty, 4) \cup (4, 5) \cup (7, \infty)$

**Answer: D**

 [Watch Video Solution](#)

32. The locus of the foot of the perpendicular from the origin on each member of the family  $(4a + 3)x - (a + 1)y - (2a + 1) = 0$  (a)  $(2x - 1)^2 + 4(y + 1)^2 = 5$  (b)  $(2x - 1)^2 + (y + 1)^2 = 5$  (c)  $(2x + 1)^2 + 4(y - 1)^2 = 5$  (d)  $(2x - 1)^2 + 4(y - 1)^2 = 5$

A.  $(2x - 1)^2 + 4(y + 1)^2 = 5$

B.  $(2x - 1)^2 + (y + 1)^2 = 5$

C.  $(2x + 1)^2 + 4(y - 1)^2 = 5$

D.  $(2x - 1)^2 + 4(y - 1)^2 = 5$

**Answer: C**

 [Watch Video Solution](#)

### Comprehension Type

1. In a  $\triangle ABC$ ,  $A = (2, 3)$  and medians through B and C have equations  $x + y - 1 = 0$  and  $2y - 1 = 0$

Equation of median through A is

A.  $x + y = 4$

B.  $5x - 3y = 1$

C.  $5x + 3y = 1$

D.  $5x = 3y$

**Answer: B**



**Watch Video Solution**

2. In a  $\triangle ABC$ ,  $A = (2, 3)$  and medians through B and C have equations

$x + y - 1 = 0$  and  $2y - 1 = 0$

Equation of side BC is (a)  $5x + 13y + 11 = 0$  (b)  $5x - 3y = 1$  (c)  $5x = 3y$

(d)  $5x + 13y - 11 = 0$

A.  $5x + 13y + 11 = 0$

B.  $5x - 3y = 1$

C.  $5x = 3y$

D.  $5x + 13y - 11 = 0$

**Answer: A**



**Watch Video Solution**

3. Let A,B,C be angles of triangles with vertex  $A \equiv (4, -1)$  and internal angular bisectors of angles B and C be  $x - 1 = 0$  and  $x - y - 1 = 0$  respectively.

Slope of BC is

A.  $1/2$

B. 2

C. 3

D. 12

**Answer: B**



**Watch Video Solution**

## Multiple Correct Answers Type

1. The point  $P(\alpha, \alpha + 1)$  will lie inside the triangle whose vertices are  $A(0, 3)$ ,  $B(-2, 0)$  and  $C(6, 1)$  if

A.  $\alpha = -1$

B.  $\alpha = -\frac{1}{2}$

C.  $\alpha = \frac{1}{2}$

D.  $-\frac{6}{7} < \alpha < \frac{3}{2}$

**Answer: B::C::D**

 [Watch Video Solution](#)

2. A straight line through the point  $A(-2, -3)$  cuts the line  $x + 3y = 0$  and  $x + y + 1 = 0$  at  $B$  and  $C$  respectively. If  $AB \cdot AC = 20$  then equation of the possible line is

A.  $x - y = 1$

B.  $x - y + 1 = 0$

C.  $3x - y + 3 = 0$

D.  $3x - y = 3$

Answer: A::C

 Watch Video Solution

3. If  $A(3, 4)$  and  $B(-5, -2)$  are the extremities of the base of an isosceles triangle  $ABC$  with  $\tan C = 2$ , then point  $C$  can be

A.  $\left( \frac{3\sqrt{5} - 1}{2}, -(1 + 2\sqrt{5}) \right)$

B.  $\left( -\frac{(3\sqrt{5} + 5)}{2}, 3 + 2\sqrt{5} \right)$

C.  $\left( \frac{3\sqrt{5} - 1}{2}, 3 - 2\sqrt{5} \right)$

D.  $\left( -\frac{(3\sqrt{5} - 5)}{2}, -(1 - 2\sqrt{5}) \right)$

**Answer: A::B**



**Watch Video Solution**

4. If  $(a, b)$  be an end of a diagonal of a square and the other diagonals has the equation  $x - y = a$  then another vertex of the square can be

A.  $(a - b, a)$

B.  $(a, 0)$

C.  $(0, -a)$

D.  $(a + b, b)$

**Answer: B::D**



**Watch Video Solution**

5. The equation of the diagonals of a rectangle are  $y + 8x - 17 = 0$  and  $y - 8x + 7 = 0$ . If the area of the rectangle is  $8\text{squnits}$  then find the sides

of the rectangle

A.  $x = 1$

B.  $x + y = 1$

C.  $y = 9$

D.  $x - 2y = 3$

**Answer: A:C**



**Watch Video Solution**

6. If  $6a^2 - 3b^2 - c^2 + 7ab - ac + 4bc = 0$  then the family of lines  $ax + by + c = 0$ ,  $|a| + |b| \neq 0$  can be concurrent at concurrent (A) (-2,3) (B) (3,-1) (C) (2,3) (D) (-3,1)

A.  $(-2, -3)$

B.  $(3, -1)$

C.  $(2, 3)$



D.  $(-3, 1)$

**Answer: A::B**



**Watch Video Solution**

7. If graph of  $xy = 1$  is reflected in  $y = 2x$  to give the graph  $12x^2 + rxy + sy^2 + t = 0$ , then

A.  $r = 7$

B.  $s = -12$

C.  $t = 25$

D.  $r + s = -19$

**Answer: B::C::D**



**Watch Video Solution**

8. Let  $A, B, C$  be angles of triangles with vertex  $A \equiv (4, -1)$  and internal angular bisectors of angles  $B$  and  $C$  be  $x - 1 = 0$  and  $x - y - 1 = 0$  respectively.

If  $A, B, C$  are angles of triangle at vertices  $A, B, C$  respectively then

$$\cot\left(\frac{B}{2}\right)\cot\left(\frac{C}{2}\right) =$$

A. 2

B. 3

C. 4

D. 6

**Answer: D**

 [Watch Video Solution](#)