



CHEMISTRY

BOOKS - MTG CHEMISTRY (ENGLISH)

CHEMICAL KINETICS

Mcqs

1. The rate of chemical reaction

- A. keeps on increasing with time
- B. remains constant with time
- C. keeps on decreasing with time
- D. shows irregular trend with time.

Answer: C



Watch Video Solution

2. The minus sign in rate $= -\frac{d[A]}{dt}$ indicates the _____ in concentration of the _____ with time. The rate of a reaction is always _____ quantity. The rate of reaction increases with _____ in concentration of reactants. The blanks in the question corresponds to

- A. decrease, products, positive, increase
- B. increase, reactants, negative, decrease
- C. decrease, reactants, positive, increase
- D. increase, products, positive, increase

Answer: C

 [Watch Video Solution](#)

3. For a reaction $P + Q \rightarrow 2R + S$. Which of the following statements is incorrect?

A. Rate of disappearance of P = Rate of appearance of S

B. Rate of disappearance of Q = $2 \times$ Rate of appearance of R


C. Rate of disappearance of P = Rate of disappearance of Q

D. Rate of disappearance of Q = $\frac{1}{2} \times$ Rate of appearance of R

Answer: B

 [Watch Video Solution](#)

4. For this reaction



The relation between $\frac{d(NH_3)}{dt}$ and $-\frac{d(H_2)}{dt}$ is :-

A. $-\frac{1}{3} \frac{d[H_2]}{dt} = +\frac{1}{2} \frac{d[NH_3]}{dt}$

B. $-\frac{1}{2} \frac{d[H_2]}{dt} = +\frac{1}{3} \frac{d[NH_3]}{dt}$

C. $+\frac{1}{2} \frac{d[H_2]}{dt} = -\frac{1}{3} \frac{d[NH_3]}{dt}$

D. $+\frac{1}{3} \frac{d[H_2]}{dt} = -\frac{1}{2} \frac{d[NH_3]}{dt}$

Answer: A

 Watch Video Solution

5. For a reaction, $2N_2O_5(g) \rightarrow 4NO_2(g) + O_2(g)$ rate of reaction is:

$$\text{A. Rate} = -\frac{d[N_2O_5]}{dt} = -\frac{1}{4} \frac{d[NO_2]}{dt} = \frac{1}{2} \frac{d[O_2]}{dt}$$

$$\text{B. Rate} = -\frac{1}{2} \frac{d[n_2O_5]}{dt} = \frac{1}{4} \frac{d[NO_2]}{dt} = \frac{d[O_2]}{2}$$

$$\text{C. Rate} = -\frac{1}{4} \frac{d[N_2O_5]}{dt} = \frac{1}{2} \frac{d[NO_2]}{dt} = \frac{d[O_2]}{dt}$$

$$\text{D. Rate} = -\frac{1}{2} \frac{d[N_2O_5]}{dt} = \frac{1}{2} \frac{d[NO_2]}{dt} = \frac{1}{2} \frac{d[O_2]}{dt}$$

Answer: B

 Watch Video Solution

6. For the reaction $2N_2O_4 \rightleftharpoons 4NO_2$, given that

$$\frac{-d[N_2O_4]}{dt} = K \text{ and } \frac{d[NO_2]}{dt} = K, \text{ then}$$

A. $2k' = k$

B. $k' = 2k$

C. $k' = k$

D. $k = \frac{1}{4}k'$

Answer: B

 [Watch Video Solution](#)

7. For a reaction $2NH_3 \rightarrow N_2 + 3H_2$, it is observed that

$$\frac{-d(NH_3)}{dt} = k_1(NH_3), \quad \frac{d(N_2)}{dt} = k_2(NH_3), \quad \frac{d(H_2)}{dt} = k_3(NH_3)$$

What is the relation between k_1 , k_2 and k_3 ?

A. $k_1 = k_2 = k_3$

B. $k_1 = 3k_2 = 3k_3$

C. $1.5k_1 = 3k_2 = k_3$

D. $2k_1 = k_2 = 3k_3$

Answer: C

 [Watch Video Solution](#)

8. Consider the reaction, $2N_2O_5 \rightarrow 4NO_2 + O_2$ In the reaction NO_2 is being formed at the rate of $0.0125 \text{ mol } L^{-1}s^{-1}$. What is the rate of reaction at this time?

A. $0.0018 \text{ mol } L^{-1}s^{-1}$

B. $0.0031 \text{ mol } L^{-1}s^{-1}$

C. $0.0041 \text{ mol } L^{-1}s^{-1}$

D. $0.050 \text{ mol } L^{-1}s^{-1}$

Answer: B

 [Watch Video Solution](#)

9. For a reaction $R \rightarrow P$, the concentration of a reactant changes from 0.05 M to 0.04 M in 30 minutes. What will be average rate of reaction in minutes?

- A. $4 \times 10^{-4} M \text{min}^{-1}$
- B. $8 \times 10^{-4} \text{min}^{-1}$
- C. $3.3 \times 10^{-4} M \text{min}^{-1}$
- D. $2.2 \times 10^{-4} M \text{min}^{-1}$

Answer: C



[Watch Video Solution](#)

10. In a reaction $2HI \rightarrow H_2 + I_2$ the concentration of HI decreases from 0.5 mol L^{-1} to 0.4 mol L^{-1} in 10 minutes. What is the rate of reaction during this interval?

- A. $5 \times 10^{-3} M \text{min}^{-1}$

B. $2.5 \times 10^{-3} M \text{min}^{-1}$

C. $5 \times 10^{-1} M \text{min}^{-1}$

D. $2.5 \times 10^{-2} M \text{min}^{-1}$

Answer: A



Watch Video Solution

11. For the reaction $4NH_3 + 5O_2 \rightarrow 4NO + 6H_2O$, if the rate of disappearance of NH_3 is $3.6 \times 10^{-3} \text{ mol } L^{-1} s^{-1}$, what is the rate of formation of H_2O

A. $5.4 \times 10^{-3} \text{ mol } L^{-1} s^{-1}$

B. $3.6 \times 10^{-3} \text{ mol } L^{-1} s^{-1}$


C. $4 \times 10^{-4} \text{ mol } L^{-1} s^{-1}$

D. $0.6 \times 10^{-4} \text{ mol } L^{-1} s^{-1}$

Answer: A

[Watch Video Solution](#)

12. Nitrogen dioxide (NO_2) dissociates into nitric oxide (NO) and oxygen (O_2) as follows



If the rate of decrease of concentration of NO_2 is $6.0 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$. What will be the rate of increase of concentration of O_2 ?

A. $3 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

B. $6 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

C. $1 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

D. $1.5 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

Answer: A

[Watch Video Solution](#)

13. The rate of disappearance of SO_2 in the reaction $2SO_2 + O_2 \rightarrow 2SO_3$ is $1.28 \times 10^{-3} g/sec$ then the rate of formation of SO_3 is

A. $0.64 \times 10^{-5} gs^{-1}$

B. $0.32 \times 10^{-5} gs^{-1}$

C. $2.56 \times 10^{-5} gs^{-1}$

D. $1.28 \times 10^{-5} gs^{-1}$

Answer: D



[Watch Video Solution](#)

14. In a reaction $2X \rightarrow Y$, the concentration of X decreases from 3.0 mole/litre to 2.0 moles/litre in 5 minutes. The rate of reaction is

A. $0.1 \text{ mol } L^{-1} \text{ min}^{-1}$

B. $5 \text{ mol } L^{-1} \text{ min}^{-1}$

C. $1 \text{ mol L}^{-1}\text{min}^{-1}$

D. $0.5 \text{ mol L}^{-1}\text{min}^{-1}$

Answer: A

 [Watch Video Solution](#)

15. In a reaction, $2X \rightarrow Y$, the concentration of X decreases from 0.50M to 0.38 M in 10 min. What is the rate of reaction in M s^{-1} during this interval?

A. 2×10^{-4}

B. 4×10^{-2}

C. 2×10^{-2}

D. 1×10^{-2}

Answer: A

 [Watch Video Solution](#)

16. For the reaction, $2N_2O_5 \rightarrow 4NO_2 + O_2$ rate and rate constant are $1.02 \times 10^{-4} M \text{ sec}^{-1}$ and $3.4 \times 10^{-5} \text{ sec}^{-1}$ respectively, the concentration of N_2O_5 , at that time will be

A. 3.4×10^{-4}

B. 3

C. 5.2

D. 3.2×10^{-5}

Answer: B



[Watch Video Solution](#)

17. The rate constant of a reaction depends on

A. temperature or the reaction

B. extent of the reaction


C. initial concentration of the reactants

D. the time of completion of reaction.

Answer: A

 [Watch Video Solution](#)

18. The chemical reaction $2O_3 \rightarrow 3O_2$ proceeds as follows :



The rate law expression should be :

A. Rate = $k[O][O_3]$

B. Rate = $k[O_3]^2[O_2]^{-1}$

C. Rate = $k[O_3]^2$

D. Rate = $[O_2][O]$

Answer: B

 [Watch Video Solution](#)



Watch Video Solution

19. Rate constant in case of first order reaction is

- A. inversely proportional to the concentration units
- B. independent of concentration units
- C. directly proportional to concentration units
- D. inversely proportional to the square of concentration units.

Answer: B



Watch Video Solution

20. The rate law for a reaction $A + B \rightarrow C + D$ is given by the expression $k[A]$. The rate of reaction will be

- A. doubled on doubling the concentration of B
- B. halved on reducing the concentration of A to half

C. Decreased on increasing the temperature of the reaction

D. unaffected by any change in concentration or temperature.

Answer: B

 [Watch Video Solution](#)

21. A reaction in which reactants (R) are converted into products (P) follows second order kinetics. If concentration of R is increased by four times, what will be the increase in the rate formation of p?

A. 9 times

B. 4 times

C. 16 times

D. 8 times

Answer: C

 [Watch Video Solution](#)

22. The rate of a gaseous reaction is given by the expression $k[A]^2[B]^3$.

The volume of the reaction vessel is suddenly reduced to one-half of the initial volume. The reaction rate relative to the original rate will be

A. $\frac{1}{8}a$

B. $\frac{1}{2}a$

C. $2a$

D. $32a$

Answer: D



Watch Video Solution

23. The rate of formation of a dimer in a second order dimerisation reaction is $9.1 \times 10^{-6} \text{ mol L}^{-1} \text{ s}^{-1}$ at 0.01 mol L^{-1} monomer concentration. Calculate the rate constant for the reaction.

A. $9.1 \times 10^{-2} \text{ L mol}^{-1} \text{ s}^{-1}$

B. $9.1 \times 10^{-6} \text{mol}^{-1} \text{s}^{-1}$

C. $3 \times 10^{-4} \text{Lmol}^{-1} \text{s}^{-1}$

D. $27.3 \times 10^{-2} \text{Lmol}^{-1} \text{s}^{-1}$

Answer: A

 [Watch Video Solution](#)

24. The rate constant for the reaction, $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ is $3.0 \times 10^{-5} \text{s}^{-1}$. If the rate is $2.40 \times 10^{-5} \text{molL}^{-1} \text{s}^{-1}$, then the initial concentration of N_2O_5 (in molL^{-1}) is

A. 0.8

B. 0.7

C. 1.2

D. 1

Answer: B



[Watch Video Solution](#)

25. The order of reaction is decided by

A. temperature

B. mechanism of reaction as well as relative concentration of reactants

C. molecularity

D. pressure.

Answer: B



[Watch Video Solution](#)

26. Which of the following statements for the order of reaction is not correct ?





A. Order can be determined experimentally.

- B. order of reaction is equal to the sum of powers of concentration terms in rate law expression
- C. order cannot be fractional.
- D. order is not affected by stoichiometric coefficient of the reactants.

Answer: C

 [Watch Video Solution](#)

27. Which of the following is an example of a fractional order reaction?



Answer: D

 [View Text Solution](#)

28. What will be the rate equation for the reaction $2X + Y \rightarrow Z$, if the order of the reaction is zero?

A. Rate = $k[X][Y]$

B. Rate = k

C. Rate = $k[X]^0[Y]$

D. Rate = $k[X][Y]^0$

Answer: B



Watch Video Solution

29. For a reaction, $I^- + OCl^- \rightarrow IO^- + Cl^-$ in an aqueous medium, the rate of reaction is given by $\frac{d[IO^-]}{dt} = k \left(\frac{I^- [OCl^-]}{[OH^-]} \right)$. The overall order of reaction is

A. -1

B. 0

C. 1




D. 2

Answer: C



Watch Video Solution

30. For a reaction, $2NO + 2H_2 \rightarrow N_2 + 2H_2O$, the possible mechanism is



What is the rate law and order of the reaction?

A. Rate = $[N_2O_2]$, order = 1

B. Rate = $[N_2O_2][H_2]$, order = 2

C. Rate = $[N_2O_2]^2$ order = 2

$$\text{D. Rate} = [N_2O_2]^2[H_2], \text{ order} = 3$$

Answer: B

 [Watch Video Solution](#)

31. Rate constant of two reactions are given below. Identifying their order of reaction

$$k = 5.3 \times 10^{-2} \text{ L mol}^{-1} \text{ s}^{-1}$$

$$k = 3.8 \times 10^{-4} \text{ s}^{-1}$$

- A. (i) second order, (ii) first order
- B. (i) first order, second order
- C. (i) zero order, (ii) first order
- D. (i) second order, (ii) zero order

Answer: A

 [Watch Video Solution](#)

32. Find the values of A, B and C in the following table for the reaction

$X + Y \rightarrow Z$. The reaction is of first order w.r.t X and zero order w.r.t. Y.

Exp	$[X](\text{molL}^{-1})$	$[Y](\text{molL}^{-1})$	Initial rate $(\text{molL}^{-1}\text{s}^{-1})$
1	0.1	0.1	2×10^{-2}
2	A	0.2	4×10^{-2}
3	0.4	0.4	B
4	C	0.2	2×10^{-2}

A. $A = 0.2\text{molL}^{-1}$, $B = 8 \times 10^{-2}\text{molL}^{-1}\text{s}^{-1}$, $C = 0.1\text{molL}^{-1}$

B. $A = 0.4\text{molL}^{-1}$, $B = 4 \times 10^{-2}\text{molL}^{-1}\text{s}^{-1}$, $C = 0.02\text{molL}^{-1}$

C. $A = 0.2\text{molL}^{-1}$, $B = 2 \times 10^{-2}\text{molL}^{-1}\text{s}^{-1}$, $C = 0.4\text{molL}^{-1}$

D. $A = 0.4\text{molL}^{-1}$, $B = 2 \times 10^2\text{molL}^{-1}\text{s}^{-1}$, $C = 0.4\text{molL}^{-1}$

Answer: A

 [Watch Video Solution](#)

33. For a chemical reaction $A \rightarrow B$, the rate of reaction increases by a factor of 1.837 when the concentration of A is increased by 1.5 time. The order of reaction with respect to A is:

A. 1

B. 1.5

C. 2

D. 2.5

Answer: B



Watch Video Solution

34. For a reaction $A \rightarrow B$, the rate of reaction becomes twenty seven times when the concentration of A is increased three times. What is the order of the reaction?

A. 2

B. 1

C. 3

D. 0

Answer: C



Watch Video Solution

35. For the reaction $A + B \rightarrow$ products, what will be the order of reaction with respect to A and B?

Exp	$[A](\text{molL}^{-1})$	$[B](\text{molL}^{-1})$	$\text{Initial rate}(\text{molL}^{-1}\text{s}^{-1})$
1	2.5×10^{-4}	3×10^{-5}	5×10^{-4}
2	5×10^{-4}	6×10^{-5}	4×10^{-3}
3	1×10^{-3}	6×10^{-5}	1.6×10^{-2}

A. 1 with respect to A and 2 with respect to B

B. 2 with respect to A and 1 with respect to B

C. 1 with respect to A and 1 with respect to B


D. 2 with respect to A and 2 with respect to B

Answer: B



Watch Video Solution

36. The unit of rate constant for the reaction,



Which has rate = $k[H_2][NO]^2$, is

A. $\text{mol L}^{-1}\text{s}^{-1}$,

B. s^{-1}

C. $\text{mol}^{-2}\text{L}^2\text{s}^{-1}$

D. mol L^{-1}

Answer: C



Watch Video Solution

37. Match the rate law given in column I with the dimensions of rate constants given in column II and mark the appropriate choice.

Column I

(A) Rate = $k[NH_3]^0$

(B) Rate = $k[H_2O_2][I^-]$

(C) Rate = $k[CH_3CHO]^{3/2}$

(D) Rate = $k[C_5H_5Cl]$

Column II

(i) $\text{mol L}^{-1}\text{s}^{-1}$

(ii) $\text{L mol}^{-1}\text{s}^{-1}$

(iii) s^{-1}

(iv) $\text{L}^{1/2}\text{mol}^{-1/2}\text{s}^{-1}$

A. (A) \rightarrow (iv), (B) \rightarrow (iii), (C) \rightarrow (ii), (D) \rightarrow (i)

B. (A) \rightarrow (i), (B) \rightarrow (ii) (C) \rightarrow (iv) ,(D) \rightarrow (iii)

C. (A) \rightarrow (iii), (B) \rightarrow (ii) (C) \rightarrow (i) ,(D) \rightarrow (iv)

D. (A) \rightarrow (ii), (B) \rightarrow (i), (C) \rightarrow (vi), (D) \rightarrow (iii)

Answer: D

 [Watch Video Solution](#)

38. The decomposition of dimethyl ether is a fractional order reaction.

The rate of reaction is given by $\text{rate} = k(p_{\text{CH}_3\text{OCH}_3})^{3/2}$. If the pressure is measured in bar and time in minutes, then what are the units of rate and constant?

A. bar min^{-1} , $\text{bar}^2 \text{min}^{-1}$

B. bar min^{-1} , bar^2 , min^{-1}

C. $\text{bar}^{-1/2} \text{min}^{-1}$, bar^2 , min^{-1}

D. bar min^{-1} , $\text{bar}^{1/2} \text{min}^{-1}$

Answer: B

 [Watch Video Solution](#)

39. The unit of rate of reaction and rate of rate constant are same for a :

- A. zero order reaction
- B. first order reaction
- C. second order reaction
- D. third order reaction

Answer: A

 [Watch Video Solution](#)

40. The rate of the reaction:

$CH_3COOC_2H_5 + NaOH \rightarrow CH_3COONa + C_2H_5OH$ is given by the equation,

$$\text{rate} = k[\text{CH}_3\text{COOC}_2\text{H}_5][\text{NaOH}]$$

If concentration is expressed in mol/L the units of k are

A. $\text{mol}^{-2}\text{L}^2\text{s}^{-1}$

B. $\text{molL}^{-1}\text{s}^{-1}$

C. $\text{Lmol}^{-1}\text{s}^{-1}$

D. S^{-1}

Answer: C



Watch Video Solution

41. Fill in the blanks by choosing the correct option. Order of the reaction is the X of the powers to which concentration terms are raised in experimentally determined rate equation. The unit of first order rate constant is Y. The unit of first order rate constant when concentration measured in terms of pressure and time in minutes is Z

A. $X \rightarrow \text{product}, Y \rightarrow \text{molL}^{-1}\text{time}^{-1}, Z \rightarrow \text{atmmin}^{-1}$

B. $X \rightarrow \text{sum } Y \rightarrow \text{Lmol}^{-1}\text{time}^{-1}, Z \rightarrow \text{atmmin}^{-1}$

C. $X \rightarrow \text{product}, Y \rightarrow \text{Lmol}^{-1}, Z \rightarrow \text{atmmin}^{-1}$

D. $X \rightarrow \text{sum}, Y \rightarrow \text{tim}^{-1}, Z \rightarrow \text{min}^{-1}$

Answer: D



[Watch Video Solution](#)

42. The number of molecules of the reactants taking part in a single step of the reaction tells about. :

A. order of a reaction

B. molecularity of a reaction

C. fast step of the mechanism of a reaction

D. half-life of the reaction

Answer: B



[Watch Video Solution](#)

43. In any unimolecular reaction

- A. the order and molecularity of the slowest step are equal to one
- B. molecularity of the reaction can be zero, one or one step
- C. more than one reacting species are involved in one step
- D. molecularity of the reaction can be determined only experimentally.

Answer: A



[Watch Video Solution](#)

44. The overall rate of a reaction is governed by :

- A. the rate of fastest intermediate step
- B. the sum of the rates of all intermediate steps
- C. the average of the rates of all the intermediate steps
- D. the rate of slowest intermediate step.

Answer: D

 [Watch Video Solution](#)

45. For a reaction $X + Y \rightarrow Z$, rate $\propto [X]$. What is (i) molecularity and (ii) order of reaction ?

A. (i) 2, (ii) 1

B. (i) 2, (ii) 2

C. (i) 1, (ii) 1

D. (i) 1, (ii) 2

Answer: A

 [Watch Video Solution](#)


46. The reaction $2X \rightarrow Y + Z$ would be zero order reaction when

- A. rate remains unchanged at any concentration of Y and Z
- B. rate of reaction doubles if concentration of Y is doubled
- C. rate of reaction remains same at any concentration of X
- D. rate of reaction is directly proportional to square of concentration of X

Answer: C

 [Watch Video Solution](#)

47. Cyclopropane rearranges to form propene



This follows first order kinetics. The rate constant is $2.714 \times 10^{-3} \text{ sec}^{-1}$.

The unitial concentration of cyclopropane is 0.29 M. What will be the concentraion of cyclopropane after 100 sec?

A. 0.035 M

B. 0.22 M

C. 0.145 M

D. 0.0018 M

Answer: B

 [Watch Video Solution](#)

48. Derive an expression to calculate time required for completion of zero order reaction.

A. $t = \frac{[R_0]}{k}$

B. $t = [R] - [R_0]$

C. $t = \frac{k}{[R_0]}$

D. $t = \frac{[R_0] - [R]}{[R_0]}$

Answer: A

 [Watch Video Solution](#)

49. A plot of $\log(a-x)$ against time 't' is a straight line. This indicates that the reaction is of :

- A. zero order
- B. first order
- C. second order
- D. third order

Answer: B

 [Watch Video Solution](#)

50. Radioactive disintegration is an example of

- A. zero order reaction
- B. first order reaction

C. second order reaction

D. third order reaction

Answer: B

 [Watch Video Solution](#)

51. A first order reaction has a rate constant of $5 \times 10^{-3} \text{ s}^{-1}$. How long will 5.0g of this reaction take to reduce to 3.0g ?

A. 34.07 s

B. 7.57 s

C. 10.10 s

D. 15 g

Answer: A

 [Watch Video Solution](#)

52. In a first order reaction, the concentration of reactant decrease from 400 mol L^{-1} to 25 mol L^{-1} in 200 seconds. The rate constant for the reaction is

A. 1.01386 s^{-1}

B. $2 \times 10^{-4} \text{ s}^{-1}$

C. $1.386 \times 10^{-2} \text{ s}^{-1}$

D. $3.4 \times 10^{-4} \text{ s}^{-1}$

Answer: C



Watch Video Solution

53. A first order reaction is 20% complete in 10 minutes. What is the specific rate constant for the reaction?

A. 0.0970 min^{-1}

B. 0.009 min^{-1}

C. 0.0223 min^{-1}

D. 2.223 min^{-1}

Answer: C

 [Watch Video Solution](#)

54. The decomposition of a substance follows first order kinetics. If its concentration is reduced to $1/8$ th of its initial value in 12 minutes, the rate constant of the decomposition system is

A. $\left(\frac{2.303}{12} \frac{\log(1)}{8} \right) \text{ min}^{-1}$

B. $\left(\frac{2.303}{12} \log 8 \right) \text{ min}^{-1}$

C. $\left(\frac{0.693}{12} \right) \text{ min}^{-1}$

D. $\left(\frac{1}{12} \log 8 \right) \text{ min}^{-1}$

Answer: B

 [Watch Video Solution](#)

55. A first order reaction takes 40 min for 30 % decomposition. Calculate $t_{1/2}$. (Given $\log 7 = 0.845$)

A. 77.7 min

B. 52.5 min

C. 46.2min

D. 22.7 min

Answer: A



[Watch Video Solution](#)

56. The following data were obtained during the first order thermal decomposition of SO_2CI_2 at a constant volume.



Experiment	Time / s^{-1}	Total pressure/atm
1	0	0.5
2	100	0.6

What is the rate of reaction when total pressure is 0.65 atm?

A. 0.35atms^{-1}

B. $2.235 \times 10^{-3} \text{atms}^{-1}$

C. $7.8 \times 10^{-4} \text{atms}^{-1}$

D. $1.55 \times 10^{-4} \text{atms}^{-1}$

Answer: C

 [Watch Video Solution](#)

57. A first order reaction has a rate constant $1.15 \times 10^{-3} \text{s}^{-1}$. How long will 5g of this reactant take to reduce to 3g?

A. 444s

B. 400s

C. 528s

D. 669s

Answer: A

 [Watch Video Solution](#)

58. The decomposition of dinitrogen pentoxide (N_2O_5) follows first order rate law. Calculate the rate constant from the given data:

$$t = 800 \text{ sec} [N_2O_5] = 1.45 \text{ molL}^{-1} = [A]$$

$$t = 600 \text{ sec} [N_2O_5] = 0.88 \text{ molL}^{-1} = [A_2]$$

A. $3.12 \times 10^{-4} \text{ s}^{-1}$

B. $6.24 \times 10^{-4} \text{ s}^{-1}$



C. $2.84 \times 10^{-4} \text{ s}^{-1}$

D. $8.14 \times 10^{-4} \text{ s}^{-1}$

Answer: B

 [Watch Video Solution](#)

59. Two plots are shown below between concentration and time t . Which of the given orders are shown by the graphs respectively?







- A. Zero order and first order
- B. first order and second order
- C. zero order and second order
- D. first order and first order

Answer: C

 [Watch Video Solution](#)

60. Match the graphs given in column I with the order given in column II and mark the appropriate choice.

Column I		Column II	
(A)	 <p>Graph showing concentration $[A]$ on the y-axis versus time t on the x-axis. The plot is a straight line with a negative slope, indicating a linear decrease in concentration over time.</p>	(i)	Third order
(B)	 <p>Graph showing $\log [A]$ on the y-axis versus time t on the x-axis. The plot is a straight line with a negative slope, indicating a linear decrease in the logarithm of concentration over time.</p>	(ii)	First order
(C)	 <p>Graph showing $\frac{1}{[A]}$ on the y-axis versus time t on the x-axis. The plot is a straight line with a positive slope, indicating a linear increase in the reciprocal of concentration over time.</p>	(iii)	Zero order
(D)	 <p>Graph showing $\frac{1}{[A]^2}$ on the y-axis versus time t on the x-axis. The plot is a straight line with a positive slope, indicating a linear increase in the reciprocal of the square of concentration over time.</p>	(iv)	Second order

A. (A) \rightarrow (i), (B) \rightarrow (ii), (C) \rightarrow (iii), (D) \rightarrow (iv)

B. (A) \rightarrow (iii), (B) \rightarrow (ii), (C) \rightarrow (iii), (D) \rightarrow (iv)

C. (A) \rightarrow (ii), (B) \rightarrow (i), (C) \rightarrow (iii), (D) \rightarrow (iv)





D. (A) \rightarrow (iv), (B) \rightarrow (iii), (C) \rightarrow (i), (D) \rightarrow (ii)

Answer: B



Watch Video Solution

61. Match the plots in column I with their order in column II and mark the appropriate choice.

	Column I		Column II
(A)		(i)	Zero order
(B)		(ii)	First order
(C)		(iii)	Second order
(D)		(iv)	Third order

A. (A) \rightarrow (i), (B) \rightarrow (ii), (C) \rightarrow (iii), (D) \rightarrow (iv)

B. (A) \rightarrow (ii), (B) \rightarrow (i), (C) \rightarrow (iii), (D) \rightarrow (iv)

C. (A) \rightarrow (iv), (B) \rightarrow (iii), (C) \rightarrow (ii), (D) \rightarrow (i)

D. (A) \rightarrow (ii), (B) \rightarrow (i), (C) \rightarrow (iii), (D) \rightarrow (iv)

Answer: B

 [Watch Video Solution](#)

62. Half life of a first order reaction is 10 min. What % of reaction will be completed in 100 min ?`

A. 0.25

B. 0.5

C. 0.999

D. 0.75

Answer: C

 [Watch Video Solution](#)

63. The half life for radioactive decay of ^{14}C is 5730 years. An archaeological artifact containing wood had only 80% of the ^{14}C found in a living tree. Estimate the age of the sample.

A. 1485 years

B. 1845 years

C. 530 years

D. 4767 years

Answer: B



[Watch Video Solution](#)

64. What will be the half-life of the first order reaction for which the value of rate constant is 200s^{-1} ?

A. $3.46 \times 10^{-2}\text{s}$

B. $3.46 \times 10^{-3}\text{s}$

C. $4.26 \times 10^{-2} s$

D. $4.26 \times 10^{-3} s$

Answer: B

 [Watch Video Solution](#)

65. The rate constant of first order reaction is 10^{-2}min^{-1} . The half-life period of reaction is

A. 69.3 min

B. 34.65 min

C. 17.37 min

D. 3.46 min

Answer: B

 [Watch Video Solution](#)

66. In a first order reaction, The concentration of reactant is reduced to $\frac{1}{8}$ th of the initial concentration in 75 minutes at 298 K. What is the half period of the reaction in minutes?

- A. 50 min
- B. 15 min
- C. 30 min
- D. 25 min

Answer: D



[Watch Video Solution](#)

67. Calculate the half life of the reaction $A \rightarrow B$, when the initial concentration of A is 0.01molL^{-1} and initial rate is $0.00352\text{molL}^{-1}\text{min}^{-1}$. The reaction is of the first order

- A. 19.69 min

B. 1.969 min

C. 7.75 min

D. 77.5 min

Answer: B

 [Watch Video Solution](#)

68. In pseudo-unimolecular reactions :

A. both the reactants are present in low concentration

B. both the reactants are present in same concentration

C. one of the reactant is present in excess

D. one of the reactant is non-reactive.

Answer: C

 [Watch Video Solution](#)

69. The value of rate of a pseudo first order reaction depends upon

- A. the concentration of both the reactants present in the reaction
- B. the concentration of the reactant present small amount
- C. the concentration of the reactant present in excess
- D. the value of ΔH of the reaction.

Answer: B



[Watch Video Solution](#)


70. State one condition under which a bimolecular reaction may be kinetically of first order reactions.

- A. When both reactants have same concentration
- B. when one of the reacting species is in large excess
- C. when the reaction is at equilibrium
- D. when the activation energy of reaction is less

Answer: B

 [Watch Video Solution](#)

71. The hydrolysis of ethyl acetate is a reaction of :



- A. zero order
- B. pseudo first order
- C. second order
- D. third order

Answer: B

 [Watch Video Solution](#)

72. Which one of the following is wrongly matched?

A. Saponification of $CH_3COOC_2H_5$ - second order reaction

B. Hydrolysis of CH_3COOCH_3 - Pseudounimolecular reaction

C. Decomposition of H_2O_2 - First order reaction

D. Combination of H_2 and Br_2 to give HBr - Zero order reaction

Answer: D

 [Watch Video Solution](#)

73. In a pseudo first order hydrolysis of ester in water the following results were obtained:

t/s	0	30	60	90
[Ester]	0.55	0.31	0.17	0.085

(i) Calculate the average rate of reaction between the time interval 30 to 60 seconds.

(ii) Calculate the pseudo first order rate constant for the hydrolysis of ester.

A. $1.91 \times 10^{-2} s^{-1}$

B. $4.67 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

C. $1.98 \times 10^{-3} \text{ s}^{-1}$

D. $2.07 \times 10^{-2} \text{ s}^{-1}$

Answer: B



Watch Video Solution

74. The Activation energy for a chemical reaction mainly depends upon

- A. the difference in energies of reactants and products
- B. the sum of energies of reactants and products
- C. the difference in energy of intermediate complex with the average energy of reactants and products
- D. the difference in energy of intermediate complex and the average energy of reactants.

Answer: D



Watch Video Solution

75. The temperature dependence of the rate of a chemical reaction can be explained by Arrhenius equation which is

A. $k = Ae^{E_a/RT}$

B. $k = Ae^{-E_a/RT}$

C. $k = Ae \times \frac{E_a}{RT}$


D. $k = Ae \times \frac{RT}{E_a}$

Answer: B




Watch Video Solution


76. An endothermic reaction with high activation energy for the forward reaction is given by the diagram




A.



B.



C.



D.

Answer: C



Watch Video Solution

77. For an endothermic reaction, where ΔH represents the enthalpy of reaction in kJ mol^{-1} , the minimum value for the energy of activation will

be

- A. Less the zero
- B. equal to ΔH
- C. less than ΔH
- D. more than ΔH

Answer: D



Watch Video Solution

78. If hydrogen and oxygen are mixed and kept in the same vessel at room temperature, the reaction does not take place to form water because

- A. activation energy for the reaction is very high at room temperature
- B. molecules have no proper orientation to react to form water
- C. the frequency of collisions is not high enough for the reaction to take place

D. no catalysts present in the reaction mixture.

Answer: A

 [Watch Video Solution](#)

79. The temperature dependence of the rate of a chemical reaction is given by arrhenius equation, $k = Ae^{-E_a/RT}$. Which of the following graphs will be a straight line?

A. $\ln A$ vs $1/T$

B. $\ln A$ vs E_a

C. $\ln K$ vs $1/T$

D. $\ln k$ vs $-E_a/R$

Answer: C

 [Watch Video Solution](#)

80. Which of the following statements is not correct?

A. For a zero order reaction $t_{1/2}$ is proportional to initial concentration.

B. The relationship of variation of rate constant with temperature is given by

$$\frac{\log(k_2)}{k_1} = \frac{E_a}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

C. The unit of rate constant for a reaction is $\text{mol}^{1-n} \text{L}^{n-1} \text{s}^{-1}$ where n is order of the reaction.

D. The unit of rate of reaction changes with order of reaction.

Answer: D



Watch Video Solution

81. The decomposition of hydrocarbon follows the equation

$$k = (4.5 \times 10^{11} \text{ s}^{-1}) e^{-28000K/T}$$

Calculate E_a .

A. 669 kJ mol^{-1}

B. $232.79 \text{ kJ mol}^{-1}$

C. $4.5 \times 10^{11} \text{ kJ mol}^{-1}$

D. $28000 \text{ kJ mol}^{-1}$

Answer: B

 [Watch Video Solution](#)

82. The rate constant for a first order reaction at 300°C for which E_a is 35 kcal mol^{-1} and frequency constant is $1.45 \times 10^{11} \text{ s}^{-1}$ is

A. $10 \times 10^2 \text{ s}^{-1}$

B. $5.37 \times 10^{10} \text{ s}^{-1}$

C. $5 \times 10^{-4} \text{ s}^{-1}$

D. $7.94 \times 10^{-3} \text{ s}^{-1}$

Answer: D

 [Watch Video Solution](#)

83. Which of the following statement about the catalyst is/are true?

- A. A catalyst makes the reaction feasible by making ΔG more negative.
- B. A catalyst makes equilibrium constant more favourable for forward reaction
- C. A catalyst accelerate rate of reaction by bringing down the activation energy
- D. A catalyst always increases the rate of reaction.

Answer: C

 [Watch Video Solution](#)

84. When a catalyst is used an equilibrium process,

- A. it increases the rate of forward reaction
- B. it decreases the rate of backward reaction
- C. it decreases activation energy of forward process and decreases activation energy of backward process
- D. it fastens the attainment of equilibrium by lowering activation energy.

Answer: D



Watch Video Solution

85. Which of the following factors are responsible for the increase in the rate of a surface catalysed reaction?

- (i) A catalyst provides proper orientation for the reactant molecules to react.
- (ii) heat of adsorption of reactants on a catalyst helps reactant molecules to overcome activation energy.
- (iii) The catalyst increases the activation energy of the reactio.

A. (i) and (iii)

B. (i) and (ii)

C. (ii) and (iii)

D. (i),(ii) and (iii)


Answer: B



[Watch Video Solution](#)

86. For a reaction $A_2 + B_2 \rightleftharpoons 2AB$ the figure shows the path of the reaction in absence and presence of a catalyst. What will be energy of activation for forward (E_f) and backward (E_b) reaction in presence of a catalyst and ΔH for the reaction? The dotted curve is path of reaction in

presence of a catalyst.



- A. $E_f = 60\text{kJ/mol}$, $E_b = 70\text{kJ/mol}$, $\Delta H = 20\text{kJ/mol}$
- B. $E_f = 20\text{kJ/mol}$, $E_b = 2 - \text{kJ/mol}$, $\Delta H = 50\text{kJ/mol}$
- C. $E_f = 70\text{kJ/mol}$, $E_b = 20\text{kJ/mol}$, $\Delta H = 10\text{kJ/mol}$
- D. $E_f = 10\text{kJ/mol}$, $E_b = 20\text{kJ/mol}$, $\Delta H = -10\text{kJ/mol}$

Answer: D



Watch Video Solution

87. The increase in concentration of the reactants lead changes in

A. ΔH

B. collision frequency

C. activation energy

D. equilibrium constant.

Answer: B



Watch Video Solution

88. The rate constant is given by the equation $k = P \cdot Z e^{-E_a/RT}$. Which factor should register a decrease for the reaction to proceed more rapidly?

A. T

B. Z

C. E

D. pressure.

Answer: C



Watch Video Solution

89. Rate of a general reaction $A + B \rightarrow$ products can be expressed as follows on the basis of collision theory.

$$\text{Rate} = Z_{AB}e^{-E_a/RT}$$

Which of the following statements is not correct for the above expression?

A. Z is collision frequency and is equal to number of collisions per second per unit volume of the reaction mixture

B. $e^{E_a/RT}$ is the fraction of molecules with kinetic energy equal to or greater than E_a

C. E_a is activation energy of the reaction.

D. All the molecules which collide with one other are effective collisions.

Answer: D



[Watch Video Solution](#)

90. Threshold energy is equal to

- A. activation energy
- B. activation energy - energy of molecules
- C. activation energy + energy of molecules
- D. None of these

Answer: C



[Watch Video Solution](#)

91. For a certain reaction a large fraction of molecules has energy more than the threshold energy, still the rate of reaction is very slow. The possible reason for this could be that

- A. the colliding molecules could be large in size
- B. the colliding molecules must not be properly oriented for effective collisions
- C. the rate of reaction could be independent of the energy
- D. one of the reactant could be in excess.

Answer: B



[Watch Video Solution](#)

92. fill up the following with suitable terms

(i) Activation energy = Threshold energy - ____

(ii) Half-life period of zero order reaction = ____

(iii) Average rate of reaction = ___

(iv) Instantaneous rate of reaction

A. Potential energy $\frac{0.693}{k}, \frac{dx}{dt}, \frac{\Delta[A]}{\Delta t}$

B. Energy of reactants $\frac{1}{k}, \frac{\Delta[A]}{\Delta t}, \frac{dx}{dt}$

C. Energy of reaction $\frac{\log x}{t}, \frac{\Delta[A]}{\Delta t}, \frac{dx}{dt}$

D. Average kinetic energy of reactants $\frac{a}{2k}, \frac{\Delta[A]}{\Delta t}, \frac{dx}{dt}$


Answer: D



Watch Video Solution

Higher Order Thinking Skills

1. The reaction $2NO + Br_2 \rightarrow 2NOBr$, obeys the following mechanism:



expression of the above reaction can be written as

A. $r = k[NO]^2[Br_2]_2$

$$B. r = k[NO][Br_2]$$



$$C. r = k[NO][Br_2]$$

$$D. r = k[NOBr_2]$$

Answer: A

 [Watch Video Solution](#)

2. A hypothetical reaction $A_2 + B_2 \rightarrow 2AB$ follows the mechanism as given below:



The order of the overall reaction is

A. 2

B. 1

C. 1.5

D. zero

Answer: C

 Watch Video Solution

3. In acidic medium the rate of reaction between $[BrO_3^-]$ and $[Br^-]$ ions given by the expression

$$-\frac{d[BrO_3^-]}{dt} = k[BrO_3^-][Br^-][H^+]^2$$

It means

- (i) rate of constant of the reaction depends upon the concentration of H^+ ions
- (ii) rate of reaction is independent of the concentration of acid added
- (iii) the change in pH of the solution will affect the rate of reaction
- (iv) doubling the concentration of H^+ ions will increase the reactions rate by 4 items.

A. only (ii)

B. only (iii)

C. only (i) and (ii)

D. only (iii) and (iv)

Answer: D



Watch Video Solution

4. The rate of certain hypothetical reaction

$A + B + C \rightarrow$ Products, is given by

$$r = - \frac{dA}{dt} = k[A]^{1/2}[B]^{1/3}[C]^{1/4}$$

The order of a reaction is given by

A. 1

B. 2

C. 13/12

D. 5/4

Answer: D



Watch Video Solution



Watch Video Solution

5. For the reaction, $H_{2(g)} + Br_{2(g)} \rightarrow 2HBr_{(g)}$, the reaction rate $= k[H_2][Br_2]^{1/2}$. Which statement is true about this reaction?

- A. The reaction is of second order
- B. Molecularity of the reaction is $3/2$
- C. The unit of k is sec^{-1}
- D. Molecularity of the reaction is 2.

Answer: D



Watch Video Solution

6. For a first order reaction, the ratio of the time taken for $7/8^{\text{th}}$ of the reaction to complete to that of half of the reaction to complete is

- A. 3:1

B. 1:3

C. 2:3

D. 3:2

Answer: A

 [Watch Video Solution](#)

7. A first order reaction is 50% completed in 30 min at $27^{\circ}C$ and in 10 min at $47^{\circ}C$. Calculate the reaction rate constants at $27^{\circ}C$ and the energy of activation of the reaction in $kJmol^{-1}$.

A. $k = 0.0231\text{min}^{-1}$, $E_a = 43.848\text{kJmol}^{-1}$


B. $k = 0.017\text{min}^{-1}$, $E_a = 52.54\text{kJmol}^{-1}$

C. $k = 0.0693\text{min}^{-1}$, $E_a = 43.848\text{kJmol}^{-1}$


D. $k = 0.0231\text{min}^{-1}$, $E_a = 28.92\text{kJmol}^{-1}$

Answer: A


8. An exothermic chemical reaction proceeds by two stages reactants $\xrightarrow{\text{Stage I}}$ intermediate $\xrightarrow{\text{Stage II}}$ products, The activation energy of stage I is 50kJmol^{-1} . The overall enthalpy change of the reaction is -100kJ mol^{-1} . Which diagram could represent the energy level diagram for the reaction ?




A.



B.



C.



D.

Answer: C



Watch Video Solution

9. In a hypothetical reaction $X \rightarrow Y$, the activation energy for the forward backward reactions are 15 and 9 kJ mol^{-1} respectively. The potential energy of X is 10 kJ mol^{-1} . Which of the following statement is/are correct?

- (i) The threshold energy of the reaction is 25 kJ mol^{-1}
- (ii) The potential energy of Y is 16 kJ mol^{-1}
- (iii) Heat of reaction is 6 kJ mol^{-1}

The reaction is endothermic.

- A. Only (i)
- B. Only (i) and (ii)
- C. Only (ii) and (iii)
- D. All are correct

Answer: D

 [Watch Video Solution](#)

Ncert Exemplar Problems

1. The role of a catalyst is to change

- A. Gibbs energy of reaction
- B. enthalpy of reaction
- C. activation energy of reaction
- D. equilibrium constant.

Answer: C

 [Watch Video Solution](#)

2. In the presence of a catalyst, the heat evolved or absorbed during the reaction:

- A. increases
- B. decreases
- C. remains unchanged
- D. may increase or decrease

Answer: C



[Watch Video Solution](#)

3. Activation energy of a chemical reaction can be determined by


- A. determining the rate constant at standard temperature
- B. determining the rate constants at two temperatures
- C. determining probability of collision
- D. using catalyst

Answer: B



Watch Video Solution

4. Consider a first order gas phase decomposition reaction given below:



The initial pressure of the system before decomposition of A was p_i . After lapse of time t' . Total pressure of the system increased by x units and became P_t . the rate constant k for the reaction is given as

$$\text{A. } k = \frac{2.303}{t} \frac{\log(p_i)}{p_i - x}$$

$$\text{B. } k = \frac{2.303}{t} \frac{\log(p_i)}{2p_i - p_t}$$

$$\text{C. } k = \frac{2.303}{t} \frac{\log(p_i)}{2p_i + p_i}$$

$$\text{D. } k = \frac{2.303}{t} \frac{\log(p_i)}{p_i + x}$$


Answer: B




Watch Video Solution

5. According to Arrhenius equation rate constant k is equal to $Ae^{-E_a/RT}$.


Which of the following option. Represents the graph of $\ln k$ us $\frac{1}{T}$?




A.



B.



C.



D.

Answer: A



Watch Video Solution

6. Consider the Arrhenius equation given below and mark the correct option.

$$k = Ae^{-\frac{E_a}{RT}}$$

- A. Rate constant increases exponentially with increasing activation energy and decreasing temperature
- B. Rate constant decreases exponentially with increasing activation energy and decreasing temperature.
- C. Rate constant increases exponentially with decreasing activation energy and decreasing temperature.
- D. Rate Constant increases exponentially with decreasing activation energy and increasing temperature.

Answer: D



Watch Video Solution

7. Which of the following statement is not correct about order of a reaction ?

- A. The order of a reaction can be a fractional number.
- B. Order of a reaction is experimaentally determined quantity
- C. The order or a reaction is always equal to the sum of the stoichiometric coefficients of reactants in the balanced chemical equaiton for a reaction.
- D. The order of a reaction is the sum of the powers of molar concentration of the reactants in the rate law expression.

Answer: C



[View Text Solution](#)

8. Consider the graph given in Q.9 Which of the following options does not show instantaneous rate of reaction at 40^{th} second?

A. $\frac{V_5 - V_2}{50 - 30}$

B. $\frac{V_4 - V_2}{50 - 30}$

C. $\frac{V_3 - V_2}{40 - 30}$

D. $\frac{V_3 - V_1}{40 - 20}$

Answer: B

 [View Text Solution](#)


9. Which of the following statements is correct?

- A. The rate of a reaction decreases with passage of time as the concentration of reactants decreases.
- B. The rate of a reaction is same at any time during the reaction.
- C. The rate of a reaction is independent of temperature change.
- D. The rate of a reaction decreases with increases in concentration of reactant (s)

Answer: A

 [Watch Video Solution](#)

10. Which of the following expression is correct for the rate of reaction given below ?



A. $\frac{\Delta[\text{Br}^{-}]}{\Delta t} = 5 \frac{\Delta[\text{H}^{+}]}{\Delta t}$

B. $\frac{\Delta[\text{Br}^{-}]}{\Delta t} = \frac{6}{5} \frac{\Delta[\text{H}^{+}]}{\Delta t}$

C. $\frac{\Delta[\text{Br}^{-}]}{\Delta t} = \frac{5}{6} \frac{\Delta[\text{H}^{+}]}{\Delta t}$

D. $\frac{\Delta[\text{Br}^{-}]}{\Delta t} = 6 \frac{\Delta[\text{H}^{+}]}{\Delta t}$

Answer: C

 [Watch Video Solution](#)

11. Rate law for the reaction, $A + 2B \rightarrow C$ is found to be

$$\text{Rate} = k[A][B]$$

Concentration of reactant 'B' is doubled keeping the concentration of 'A' constant, the value of rate constant will be _____

- A. the same
- B. doubled
- C. quadrupled
- D. halved

Answer: A



[Watch Video Solution](#)

12. Which of the following statements is incorrect about the collision theory of chemical reaction?

- A. It considers reacting molecules or atoms to be hard spheres and ignores their structural features.
- B. Number of effective collisions determines the rate of reaction.
- C. Collision of atoms or molecules possessing sufficient threshold energy results into the product formation.
- D. Molecules should collide with sufficient threshold energy and proper orientation for the collision to be effective.

Answer: C

 [Watch Video Solution](#)

13. A first order reaction is 50 % completed in 1.26×10^{14} s. How much time would it take for 100 % completion?

A. 1.26×10^{15} s

B. 2.52×10^{14} s

C. $2.52 \times 10^{28} s$


D. Infinite

Answer: D



Watch Video Solution

14. Compound 'A' and B react according to the following chemical equation.



Concentration of either A or B were changed keeping the concentrations of one of the reactants constant and rates were measured as a function of initial concentration. Following results were obtained. Choose the correct option for the rate equations for this reaction.

Experiment	Initial concentration of $[A] / molL^{-1}$	Initial concentration of $[B]$
1	0.30	0.30
2	0.30	0.60
3	0.60	0.30

A. Rate = $k[A]^2[B]$

B. Rate = $k[A][B]^2$

C. Rate = $k[A][B]$

D. Rate = $k[A]^2[B]^0$

Answer: B



Watch Video Solution

15. Which of the following statements is not correct for the catalyst?

- A. It catalyze the forward and backward reaction to the same extent.
- B. It alters ΔG of the reaction.
- C. It is a substance that does not change the equilibrium constant of a reaction.
- D. It provides an alternate mechanism by reducing activation energy between reactants and products.

Answer: B



[Watch Video Solution](#)

16. The value of rate constant of a pseudo first order reaction


- A. depends on the concentration of reactants present in small amount
- B. depends on the concentrations of reactants present in excess
- C. is independent of the concentration of reactants
- D. depends only on temperature

Answer: B




[Watch Video Solution](#)


17. Consider the reaction $A \rightarrow B$. The concentration of both the reactants and the products varies exponentially with time. Which of the following figure correctly describes the change in concentration of reactants and products with time ?




A.



B.



C.



D.

Answer: B



Watch Video Solution

Assertion Reason

1. Assertion : Precipitation of silver chloride occurs instantaneously by mixing of aqueous solutions of silver nitrate and sodium chloride.

Reason : Ionic reactions occur very fast.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: A



[Watch Video Solution](#)

2. Assertion : The rate of reaction is the rate of change of concentration of a reactant or a product.


Reason : Rate of reaction remains constant during the complete reaction.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

3. Assertion : For the reaction



$$\text{Rate} = k[CHCl_3][Cl_2]^{1/2}$$

Reason : Rate of reaction is always equals to the sum of the stoichiometric coefficients of the reacting species in a balanced chemical equation.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

4. Assertion : Order of a reaction with respect to any reactant can be zero, positive negative of fractional.

Reason : Rate of a reaction cannot decrease with increase in concentration of a reactant or a product.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.

- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

5. Assertion : Molecularity greater than three is not observed.

Reason : The overall molecularity of complex reaction is equal to molecularity of the slowest step.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

6. Assertion : Complex reaction takes place in different steps and the slowest step determines the rate of reaction.

Reason : Order and molecularity of a reaction are always equal.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C



[Watch Video Solution](#)

7. Assertion : The decomposition of gaseous ammonia on a hot platinum surface is a zero order reaction at high pressure.

For a zero order reaction the rate of reaction is independent of initial concentration.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B



[Watch Video Solution](#)

8. Assertion : For a first order reaction, $t_{1/2}$ is independent of rate constant.

Reason : For a first reaction $t_{1/2} \propto [R]_0$.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.


D. If both assertion and reason are false.

Answer: D



Watch Video Solution

9. Assertion : The reaction



Reason : Change in concentration of H_2O is negligible.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A



Watch Video Solution

10. Assertion : The rate of a reaction sometimes does not depend on concentration.

Reason : Lower the activation energy, faster is the reaction

- A. If both assertion and reason are true and reason is the correct explanation of assertion.

- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

11. Assertion : For a chemical reaction with rise in temperature by 10° the rate constant is nearly doubled.

Reason : At $t + 10$, the fraction of molecules having energy equal to or greater than activation energy gets doubled.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: A



[Watch Video Solution](#)

12. Assertion : E_a of the forward reaction is higher than that of backward reaction in a reversible endothermic reaction.

Reason : Increasing the temperature of the substance increases the fraction of molecules which collide with energies greater than E_a .

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

13. Assertion : A catalyst increases the rate of reaction without itself undergoing any permanent chemical change.

Reason : A catalyst changes the Gibbs energy (ΔG) of the reaction and equilibrium constant of the reaction.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

14. Assertion : All molecules collisions lead to the formation of products.

Reason : Reactant molecules undergo chemical change irrespective of their collision.

Assertion : Rate of reaction increases with increase in temperature .

Reason : Number of effective collisions increases with increase in temperature.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: D



[View Text Solution](#)

15. Assertion : Rate of reaction increases with increase in temperature.

Reason : Number of effective collisions increases with increase in temperature.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A



[Watch Video Solution](#)

Rate Of A Chemical Reaction

1. The rate of chemical reaction

- A. keeps on increasing with time
- B. remains constant with time
- C. keeps on decreasing with time
- D. shows irregular trend with time.

Answer: C



Watch Video Solution

2. The minus sign in rate $= - \frac{d[A]}{dt}$ indicates the _____ in concentration of the _____ with time. The rate of a reaction is always _____ quantity. The rate of reaction increases with _____ in concentration of reactants. The blanks in the question corresponds to

- A. decrease, products, positive, increase
- B. increase, reactants, negative, decrease

C. decrease, reactants, positive, increase

D. increase, products, positive, increase


Answer: C



Watch Video Solution

3. For reversible reaction, $A + B \rightleftharpoons C + D$, the graph for rate of reaction with time is given below.

Mark the terms (p),(q) and (r)



- A. (p)-rate of backward reaction ,(q)-rate of forward reaction, (r)-
equilibrium,
- B. (p)-rate of forward reaction, (q)-rate of backward reaction, (r)-
equilibrium
- C. (p)-concentration of products (q)-concentration of reactants (r)-
rate of reaction

D. (p)-instantaneous rate of reaction, (q)-variation of rate, (r)-average rate of reaction

Answer: B

 [Watch Video Solution](#)

4. For a reaction $P + Q \rightarrow 2R + S$. Which of the following statements is incorrect?

A. Rate of disappearance of P = Rate of appearance of S

B. Rate of disappearance of Q = $2 \times$ Rate of appearance of R


C. Rate of disappearance of P = Rate of disappearance of Q

D. Rate of disappearance of Q = $\frac{1}{2} \times$ Rate of appearance of R

Answer: B

 [Watch Video Solution](#)

5. For this reaction



The relation between $\frac{d(NH_3)}{dt}$ and $-\frac{d(H_2)}{dt}$ is :-

A. $-\frac{1}{3} \frac{d[H_2]}{dt} = +\frac{1}{2} \frac{d[NH_3]}{dt}$

B. $-\frac{1}{2} \frac{d[H_2]}{dt} = +\frac{1}{3} \frac{d[NH_3]}{dt}$

C. $+\frac{1}{2} \frac{d[H_2]}{dt} = -\frac{1}{3} \frac{d[NH_3]}{dt}$

D. $+\frac{1}{3} \frac{d[H_2]}{dt} = -\frac{1}{2} \frac{d[NH_3]}{dt}$

Answer: A



Watch Video Solution

6. For a reaction, $2N_2O_5(g) \rightarrow 4NO_2(g) + O_2(g)$ rate of reaction is:

A. Rate = $-\frac{d[N_2O_5]}{dt} = -\frac{1}{4} \frac{d[NO_2]}{dt} = \frac{1}{2} \frac{d[O_2]}{dt}$

B. Rate = $-\frac{1}{2} \frac{d[n_2O_5]}{dt} = \frac{1}{4} \frac{d[NO_2]}{dt} = \frac{d[O_2]}{2}$

C. Rate = $-\frac{1}{4} \frac{d[N_2O_5]}{dt} = \frac{1}{2} \frac{d[NO_2]}{dt} = \frac{d[O_2]}{dt}$

$$\text{D. Rate} = -\frac{1}{2} \frac{d[N_2O_5]}{dt} = \frac{1}{2} \frac{d[NO_2]}{dt} = \frac{1}{2} \frac{d[O_2]}{dt}$$

Answer: B

 [Watch Video Solution](#)

7. For the reaction $2N_2O_4 \rightleftharpoons 4NO_2$, given that $-\frac{d[N_2O_4]}{dt} = K$ and $\frac{d[NO_2]}{dt} = K$, then

A. $2k' = k$

B. $k' = 2k$

C. $k' = k$

D. $k = \frac{1}{4}k'$

Answer: B

 [Watch Video Solution](#)

8. For a reaction $2NH_3 \rightarrow N_2 + 3H_2$, it is observed that $\frac{-d(NH_3)}{dt} = k_1(NH_3)$, $\frac{d(N_2)}{dt} = k_2(NH_3)$, $\frac{d(H_2)}{dt} = k_3(NH_3)$

What is the relation between k_1 , k_2 and k_3 ?

A. $k_1 = k_2 = k_3$

B. $k_1 = 3k_2 = 3k_3$

C. $1.5k_1 = 3k_2 = k_3$

D. $2k_1 = k_2 = 3k_3$

Answer: C



Watch Video Solution

9. Consider the reaction, $2N_2O_5 \rightarrow 4NO_2 + O_2$ In the reaction NO_2 is being formed at the rate of $0.0125 \text{ mol L}^{-1}\text{s}^{-1}$. What is the rate of reaction at this time?

A. $0.0018 \text{ mol L}^{-1}\text{s}^{-1}$

B. $0.0031 \text{ mol L}^{-1} \text{ s}^{-1}$

C. $0.0041 \text{ mol L}^{-1} \text{ s}^{-1}$

D. $0.050 \text{ mol L}^{-1} \text{ s}^{-1}$

Answer: B

 [Watch Video Solution](#)

10. For a reaction $R \rightarrow P$, the concentration of a reactant changes from 0.05 M to 0.04 M in 30 minutes. What will be average rate of reaction in minutes?

A. $4 \times 10^{-4} \text{ Mmin}^{-1}$

B. $8 \times 10^{-4} \text{ min}^{-1}$

C. $3.3 \times 10^{-4} \text{ Mmin}^{-1}$

D. $2.2 \times 10^{-4} \text{ Mmin}^{-1}$

Answer: C



Watch Video Solution

11. In a reaction $2HI \rightarrow H_2 + I_2$ the concentration of HI decreases from 0.5 mol L^{-1} to 0.4 mol L^{-1} in 10 minutes. What is the rate of reaction during this interval?

A. $5 \times 10^{-3} \text{ Mmin}^{-1}$

B. $2.5 \times 10^{-3} \text{ Mmin}^{-1}$

C. $5 \times 10^{-1} \text{ Mmin}^{-1}$

D. $2.5 \times 10^{-2} \text{ Mmin}^{-1}$

Answer: A



Watch Video Solution

12. For the reaction $4NH_3 + 5O_2 \rightarrow 4NO + 6H_2O$, if the rate of disappearance of NH_3 is $3.6 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$, what is the rate of formation of H_2O

A. $5.4 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

B. $3.6 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$


C. $4 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

D. $0.6 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

Answer: A

 [Watch Video Solution](#)

13. Nitrogen dioxide (NO_2) dissociates into nitric oxide (NO) and oxygen (O_2) as follows



If the rate of decrease of concentration of NO_2 is $6.0 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$. What will be the rate of increase of concentration of O_2 ?

A. $3 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

B. $6 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

C. $1 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$

$$D. 1.5 \times 10^{-12} \text{ mol L}^{-1} \text{ s}^{-1}$$

Answer: A



Watch Video Solution

14. The rate of disappearance of SO_2 in the reaction $2SO_2 + O_2 \rightarrow 2SO_3$ is $1.28 \times 10^{-3} \text{ g/sec}$ then the rate of formation of SO_3 is

A. $0.64 \times 10^{-5} \text{ gs}^{-1}$

B. $0.32 \times 10^{-5} \text{ gs}^{-1}$

C. $2.56 \times 10^{-5} \text{ gs}^{-1}$

D. $1.28 \times 10^{-5} \text{ gs}^{-1}$

Answer: D



Watch Video Solution

15. In a reaction $2X \rightarrow Y$, the concentration of X decreases from 3.0 mole/litre to 2.0 moles/litre in 5 minutes. The rate of reaction is

A. $0.1 \text{ mol L}^{-1}\text{min}^{-1}$

B. $5 \text{ mol L}^{-1}\text{min}^{-1}$

C. $1 \text{ mol L}^{-1}\text{min}^{-1}$

D. $0.5 \text{ mol L}^{-1}\text{min}^{-1}$

Answer: A



[Watch Video Solution](#)

16. In a reaction, $2X \rightarrow Y$, the concentration of X decreases from 0.50M to 0.38 M in 10 min. What is the rate of reaction in M s^{-1} during this interval?

A. 2×10^{-4}

B. 4×10^{-2}

C. 2×10^{-2}

D. 1×10^{-2}

Answer: A

 [Watch Video Solution](#)

17. For the reaction, $2N_2O_5 \rightarrow 4NO_2 + O_2$ rate and rate constant are $1.02 \times 10^{-4} M \text{ sec}^{-1}$ and $3.4 \times 10^{-5} \text{ sec}^{-1}$ respectively, the concentration of N_2O_5 , at that time will be

A. 3.4×10^{-4}

B. 3

C. 5.2

D. 3.2×10^{-5}

Answer: B

 [Watch Video Solution](#)

Factors Influencing Rate Of Reaction

1. The rate constant of a reaction depends on


- A. temperature of the reaction
- B. extent of the reaction
- C. initial concentration of the reactants
- D. the time of completion of reaction.

Answer: A



[Watch Video Solution](#)

2. The chemical reaction $2O_3 \rightarrow 3O_2$ proceeds as follows :



The rate law expression should be :

A. Rate = $k[O][O_3]$

B. Rate = $k[O_3]^2[O_2]^{-1}$

C. Rate = $k[O_3]^2$

D. Rate = $[O_2][O]$

Answer: B

 [Watch Video Solution](#)

3. Rate constant in case of first order reaction is

A. inversely proportional to the concentration units

B. independent of concentration units

C. directly proportional to concentration units

D. inversely proportional to the square of concentration units.

Answer: B

 [Watch Video Solution](#)

4. The rate law for a reaction $A + B \rightarrow C + D$ is given by the expression $k[A]$. The rate of reaction will be

- A. doubled on doubling the concentration of B
- B. halved on reducing the concentration of A to half
- C. Decreased on increasing the temperature of the reaction
- D. unaffected by any change in concentration or temperature.

Answer: B



Watch Video Solution

5. A reaction in which reactants (R) are converted into products (P) follows second order kinetics. If concentration of R is increased by four times, what will be the increase in the rate formation of p?

- A. 9 times

B. 4 times

C. 16 times

D. 8 times

Answer: C



Watch Video Solution

6. The rate of a gaseous reaction is given by the expression $k[A]^2[B]^3$.

The volume of the reaction vessel is suddenly reduced to one-half of the initial volume. The reaction rate relative to the original rate will be

A. $\frac{1}{8}a$

B. $\frac{1}{2}a$

C. $2a$

D. $32a$

Answer: D

[Watch Video Solution](#)

7. The rate of formation of a dimer in a second order dimerisation reaction is $9.1 \times 10^{-6} \text{ mol L}^{-1} \text{ s}^{-1}$ at 0.01 mol L^{-1} monomer concentration. Calculate the rate constant for the reaction.

A. $9.1 \times 10^{-2} \text{ L mol}^{-1} \text{ s}^{-1}$

B. $9.1 \times 10^{-6} \text{ mol}^{-1} \text{ s}^{-1}$

C. $3 \times 10^{-4} \text{ L mol}^{-1} \text{ s}^{-1}$

D. $27.3 \times 10^{-2} \text{ L mol}^{-1} \text{ s}^{-1}$

Answer: A

[Watch Video Solution](#)

8. The rate constant for the reaction, $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ is $3.0 \times 10^{-5} \text{ s}^{-1}$. If the rate is $2.40 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$, then the initial concentration of N_2O_5 (in mol L^{-1}) is

A. 0.8

B. 0.7

C. 1.2

D. 1

Answer: B



[Watch Video Solution](#)

9. The order of reaction is decided by

A. temperature

B. mechanism of reaction as well as relative concentration of reactants

C. molecularity

D. pressure.

Answer: B



Watch Video Solution

10. Which of the following statements for the order of reaction is not correct ?


- A. Order can be determined experimentally.
- B. order of reaction is equal to the sum of powers of concentration terms in rate law expression
- C. order cannot be fractional.
- D. order is not affected by stoichiometric coefficient of the reactants.




Answer: C



Watch Video Solution

11. Which of the following is an example of a fractional order reaction?








Answer: D

 [View Text Solution](#)

12. What will be the rate equation for the reaction $2X + Y \rightarrow Z$, if the order of the reaction is zero?



Answer: B

 [Watch Video Solution](#)



13. For a reaction, $I^- + OCl^- \rightarrow IO^- + Cl^-$ in an aqueous medium, the rate of reaction is given by $\frac{d[IO^-]}{dt} = k \left(\frac{I^- [OCl^-]}{[OH^-]} \right)$. The overall order of reaction is


- A. -1
- B. 0
- C. 1
- D. 2

Answer: C

 [Watch Video Solution](#)

14. For a reaction, $2NO + 2H_2 \rightarrow N_2 + 2H_2O$, the possible mechanism is





What is the rate law and order of the reaction?

A. Rate = $[N_2O_2]$, order = 1

B. Rate = $[N_2O_2][H_2]$, order = 2

C. Rate = $[N_2O_2]^2$ order = 2

D. Rate = $[N_2O_2]^2[H_2]$, order = 3

Answer: B



Watch Video Solution

15. Rate constant of two reactions are given below. Identifying their order of reaction

$$k = 5.3 \times 10^{-2} Lmol^{-1}s^{-1}$$

$$k = 3.8 \times 10^{-4} s^{-1}$$

A. (i) second order, (ii) first order

B. (i) first order, second order

C. (i) zero order, (ii) first order

D. (i) second order, (ii) zero order

Answer: A

 [Watch Video Solution](#)

16. Find the values of A, B and C in the following table for the reaction

$X + Y \rightarrow Z$. The reaction is of first order w.r.t X and zero order w.r.t. Y.

Exp	$[X](molL^{-1})$	$[Y](molL^{-1})$	Initial rate $(molL^{-1}s^{-1})$
1	0.1	0.1	2×10^{-2}
2	A	0.2	4×10^{-2}
3	0.4	0.4	B
4	C	0.2	2×10^{-2}

A. $A = 0.2molL^{-1}$, $B = 8 \times 10^{-2}molL^{-1}s^{-1}$, $C = 0.1molL^{-1}$

B. $A = 0.4molL^{-1}$, $B = 4 \times 10^{-2}molL^{-1}s^{-1}$, $C = 0.02molL^{-1}$

C. $A = 0.2molL^{-1}$, $B = 2 \times 10^{-2}molL^{-1}s^{-1}$, $C = 0.4molL^{-1}$

D. $A = 0.4molL^{-1}$, $B = 2 \times 10^2molL^{-1}s^{-1}$, $C = 0.4molL^{-1}$

Answer: A



[Watch Video Solution](#)

17. For a chemical reaction $A \rightarrow B$, the rate of reaction increases by a factor of 1.837 when the concentration of A is increased by 1.5 time. The order of reaction with respect to A is:

A. 1

B. 1.5

C. 2

D. 2.5

Answer: B



[Watch Video Solution](#)

18. For a reaction $A \rightarrow B$, the rate of reaction becomes twenty seven times when the concentration of A is increased three times. What is the order of the reaction?

A. 2

B. 1

C. 3

D. 0

Answer: C



[Watch Video Solution](#)

19. For the reaction $A + B \rightarrow$ products, what will be the order of reaction with respect to A and B?

Exp	$[A](\text{molL}^{-1})$	$[B](\text{molL}^{-1})$	$\text{Initialrate}(\text{molL}^{-1}\text{s}^{-1})$
1	2.5×10^{-4}	3×10^{-5}	5×10^{-4}
2	5×10^{-4}	6×10^{-5}	4×10^{-3}
3	1×10^{-3}	6×10^{-5}	1.6×10^{-2}

A. 1 with respect to A and 2 with respect to B

B. 2 with respect to A and 1 with respect to B


C. 1 with respect to A and 1 with respect to B

D. 2 with respect to A and 2 with respect to B

Answer: B

 [Watch Video Solution](#)

20. The unit of rate constant for the reaction,



Which has rate = $k[H_2][NO]^2$, is

A. $\text{mol L}^{-1}\text{s}^{-1}$,

B. s^{-1}

C. $\text{mol}^{-2}\text{L}^2\text{s}^{-1}$

D. mol L^{-1}

Answer: C

 Watch Video Solution

21. Match the rate law given in column I with the dimensions of rate constants given in column II and mark the appropriate choice.

Column I

Column II

$$(A) \text{Rate} = k[NH_3]^0$$

$$(i) \text{molL}^{-1}\text{s}^{-1}$$

$$(B) \text{Rate} = k[H_2O_2][I^-]$$

$$(ii) \text{Lmol}^{-1}\text{s}^{-1}$$

$$(C) \text{Rate} = k[CH_3CHO]^{3/2}$$

$$(iii) \text{S}^{-1}$$

$$(D) \text{Rate} = k[C_5H_5Cl]$$

$$(iv) \text{L}^{1/2}\text{mol}^{-1/2}\text{s}^{-1}$$

A. (A) \rightarrow (iv), (B) \rightarrow (iii), (C) \rightarrow (ii), (D) \rightarrow (i)

B. (A) \rightarrow (i), (B) \rightarrow (ii) (C) \rightarrow (iv) ,(D) \rightarrow (iii)

C. (A) \rightarrow (iii), (B) \rightarrow (ii) (C) \rightarrow (i) ,(D) \rightarrow (iv)

D. (A) \rightarrow (ii), (B) \rightarrow (i), (C) \rightarrow (vi), (D) \rightarrow (iii)

Answer: D

 Watch Video Solution

22. The decomposition of dimethyl ether is a fractional order reaction. The rate of reaction is given by $\text{rate} = k(p_{\text{CH}_3\text{OCH}_3})^{3/2}$. If the pressure is measured in bar and time in minutes, then what are the units of rate and constant?

- A. $\text{bar min}^{-1}, \text{bar}^2 \text{min}^{-1}$
- B. $\text{bar min}^{-1}, \text{bar}^2, \text{min}^{-1}$
- C. $\text{bar}^{-1/2} \text{min}^{-1}, \text{bar}^2, \text{min}^{-1}$
- D. $\text{bar min}^{-1}, \text{bar}^{1/2} \text{min}^{-1}$

Answer: B



[Watch Video Solution](#)

23. The unit of rate of reaction and rate of rate constant are same for a :

- A. zero order reaction
- B. first order reaction

C. second order reaction

D. third order reaction

Answer: A



Watch Video Solution

24. The rate of the reaction:

$CH_3COOC_2H_5 + NaOH \rightarrow CH_3COONa + C_2H_5OH$ is given by the equation,

$$\text{rate} = k[CH_3COOC_2H_5][NaOH]$$

If concentration is expressed in mol/L the units of k are

A. $mol^{-2}L^2s^{-1}$


B. $molL^{-1}s^{-1}$

C. $Lmol^{-1}s^{-1}$

D. S^{-1}

Answer: C

25. For a general reaction $X \rightarrow Y$, the plot of conc. of X vs time is given in the figure. What is the order of the reaction and what are the units of rate constant?



A. zero, $\text{mol L}^{-1}\text{s}^{-1}$

B. First, $\text{mol L}^{-1}\text{s}^{-1}$

C. First, s^{-1}

D. Zero, $L \text{ mol}^{-1} \text{ s}^{-1}$

Answer: A

 [Watch Video Solution](#)

26. Fill in the blanks by choosing the correct option. Order of the reaction is the X of the powers to which concentration terms are raised in experimentally determined rate equation. The unit of first order rate constant is Y. The unit of first order rate constant when concentration measured in terms of pressure and time in minutes is Z

A. $X \rightarrow$ product, $Y \rightarrow \text{molL}^{-1}\text{time}^{-1}$, $Z \rightarrow \text{atmmin}^{-1}$

B. $X \rightarrow$ sum $Y \rightarrow \text{Lmol}^{-1}\text{time}^{-1}$, $Z \rightarrow \text{atmmin}^{-1}$

C. $X \rightarrow$ product, $Y \rightarrow \text{Lmol}^{-1}$, $Z \rightarrow \text{atmmin}^{-1}$

D. $X \rightarrow$ sum, $Y \rightarrow \text{tim}^{-1}$, $Z \rightarrow \text{min}^{-1}$

Answer: D

 [Watch Video Solution](#)

27. The number of molecules of the reactants taking part in a single step of the reaction tells about. :

- A. order of a reaction
- B. molecularity of a reaction
- C. fast step of the mechanism of a reaction
- D. half-life of the reaction

Answer: B

 [Watch Video Solution](#)

28. In any unimolecular reaction

- A. the order and molecularity of the slowest step are equal to one
- B. molecularity of the reaction can be zero, one or one step

- C. more than one reacting species are involved in one step
- D. molecularity of the reaction can be determined only experimentally.

Answer: A

 [Watch Video Solution](#)

29. The overall rate of a reaction is governed by :

- A. the rate of fastest intermediate step
- B. the sum of the rates of all intermediate steps
- C. the average of the rates of all the intermediate steps
- D. the rate of slowest intermediate step.

Answer: D

 [Watch Video Solution](#)

30. For a reaction $X + Y \rightarrow Z$, rate $\propto [X]$. What is (i) molecularity and (ii) order of reaction ?

- A. (i) 2, (ii) 1
- B. (i) 2, (ii) 2
- C. (i) 1, (ii) 1
- D. (i) 1, (ii) 2

Answer: A



[Watch Video Solution](#)

Integrated Rate Equation

1. The reaction $2X \rightarrow Y + Z$ would be zero order reaction when

- A. rate remains unchanged at any concentration of Y and Z
- B. rate of reaction doubles if concentration of Y is doubled

C. rate of reaction remains same at any concentration of X

D. rate of reaction is directly proportional to square of concentration of X

Answer: C

 [Watch Video Solution](#)

2. Derive an expression to calculate time required for completion of zero order reaction.

A. $t = \frac{[R_0]}{k}$

B. $t = [R] - [R_0]$

C. $t = \frac{k}{[R_0]}$

D. $t = \frac{[R_0] - [R]}{[R_0]}$

Answer: A

 [Watch Video Solution](#)



3. A plot of $\log(a-x)$ against time 't' is a straight line. This indicates that the reaction is of :

- A. zero order
- B. first order
- C. second order
- D. third order

Answer: B

 [Watch Video Solution](#)

4. Observe the given graphs carefully.



Which of the given orders are shown by the graphs respectively?

- A. zero order, First order
- B. First order, Zero order
- C. First order, first order
- D. Second order, Zero order

Answer: C



[Watch Video Solution](#)

5. Radioactive disintegration is an example of

- A. zero order reaction
- B. first order reaction
- C. second order reaction
- D. third order reaction

Answer: B



Watch Video Solution

6. A first order reaction has a rate constant of $5 \times 10^{-3} \text{ s}^{-1}$. How long will 5.0 g of this reaction take to reduce to 3.0 g ?

A. 34.07 s

B. 7.57 s

C. 10.10 s

D. 15 g

Answer: A



Watch Video Solution

7. In a first order reaction, the concentration of reactant decrease from 400 mol L^{-1} to 25 mol L^{-1} in 200 seconds. The rate constant for the reaction is

A. $1.01386s^{-1}$

B. $2 \times 10^{-4}s^{-1}$

C. $1.386 \times 10^{-2}s^{-1}$

D. $3.4 \times 10^{-4}s^{-1}$

Answer: C

 [Watch Video Solution](#)

8. A first order reaction is 20% complete in 10 minutes. What is the specific rate constant for the reaction?

A. 0.0970 min^{-1}

B. 0.009 min^{-1}

C. 0.0223 min^{-1}

D. 2.223 min^{-1}

Answer: C

 [Watch Video Solution](#)

9. The decomposition of a substance follows first order kinetics. If its concentration is reduced to $1/8$ th of its initial value in 12 minutes, the rate constant of the decomposition system is

A. $\left(\frac{2.303}{12} \frac{\log(1)}{8} \right) \text{min}^{-1}$

B. $\left(\frac{2.303}{12} \log 8 \right) \text{min}^{-1}$

C. $\left(\frac{0.693}{12} \right) \text{min}^{-1}$

D. $\left(\frac{1}{12} \log 8 \right) \text{min}^{-1}$

Answer: B

 [Watch Video Solution](#)

10. A first order reaction takes 40 min for 30 % decomposition. Calculate $t_{1/2}$. (Given $\log 7 = 0.845$)

A. 77.7 min

B. 52.5 min

C. 46.2min

D. 22.7 min

Answer: A

 [Watch Video Solution](#)

11. The following data were obtained during the first order thermal decomposition of SO_2Cl_2 at a constant volume.



Experiment	Time / s^{-1}	Total pressure/atm
------------	-----------------	--------------------

1	0	0.5
---	---	-----

2	100	0.6
---	-----	-----

What is the rate of reaction when total pressure is 0.65 atm?

A. 0.35atms^{-1}

B. $2.235 \times 10^{-3} \text{atms}^{-1}$

C. $7.8 \times 10^{-4} \text{atms}^{-1}$

D. $1.55 \times 10^{-4} \text{atms}^{-1}$

Answer: C

 [Watch Video Solution](#)

12. A first order reaction has a rate constant $1.15 \times 10^{-3} \text{s}^{-1}$. How long will 5g of this reactant take to reduce to 3g?

A. 444s

B. 400s

C. 528s

D. 669s

Answer: A

 [Watch Video Solution](#)

13. The decomposition of dinitrogen pentoxide (N_2O_5) follows first order rate law. Calculate the rate constant from the given data:

$$t = 800 \text{ sec} [N_2O_5] = 1.45 \text{ molL}^{-1} = [A]$$

$$t = 600 \text{ sec} [N_2O_5] = 0.88 \text{ molL}^{-1} = [A_2]$$

A. $3.12 \times 10^{-4} \text{ s}^{-1}$

B. $6.24 \times 10^{-4} \text{ s}^{-1}$

C. $2.84 \times 10^{-4} \text{ s}^{-1}$

D. $8.14 \times 10^{-4} \text{ s}^{-1}$

Answer: B

 [Watch Video Solution](#)

14. Half life of a first order reaction in 10 min. What % of reaction will be completed in 100 min ?`

A. 0.25

B. 0.5

C. 0.999

D. 0.75

Answer: C



Watch Video Solution

15. The half life for radioactive decay of ^{14}C is 5730 years. An archaeological artifact containing wood had only 80% of the ^{14}C found in a living tree. Estimate the age of the sample.

A. 1485 years

B. 1845 years

C. 530 years

D. 4767 years

Answer: B



Watch Video Solution

16. What will be the half-life of the first order reaction for which the value of rate constant is 200s^{-1} ?

A. $3.46 \times 10^{-2}x$

B. $3.46 \times 10^{-3}s$

C. $4.26 \times 10^{-2}s$

D. $4.26 \times 10^{-3}s$

Answer: B



Watch Video Solution

17. The rate constant of first order reaction is 10^{-2}min^{-1} . The half-life period of reaction is

A. 69.3 min

B. 34.65 min

C. 17.37 min

D. 3.46 min

Answer: B



[Watch Video Solution](#)

18. In a first order reaction, The concentration of reactant is reduced to $\frac{1}{8}$ th of the initial concentration in 75 minutes at 298 K. What is the half period of the reaction in minutes?

A. 50 min

B. 15 min

C. 30 min

D. 25 min

Answer: D



Watch Video Solution

19. Calculate the half life of the reaction $A \rightarrow B$, when the initial concentration of A is 0.01molL^{-1} and initial rate is $0.00352\text{molL}^{-1}\text{min}^{-1}$. The reaction is of the first order

A. 19.69 min

B. 1.969 min

C. 7.75 min

D. 77.5 min

Answer: B



Watch Video Solution

20. In pseudo-unimolecular reactions :

A. both the reactants are present in low concentraion

- B. both the reactants are present in same concentration
- C. one of the reactant is present in excess
- D. one of the reactant is non-reactive.

Answer: C

 [Watch Video Solution](#)

21. The value of rate of a pseudo first order reaction depends upon

- A. the concentration of both the reactants present in the reaction
- B. the concentration of the reactant present small amount
- C. the concentration of the reactant present in excess
- D. the value of ΔH of the reaction.

Answer: B

 [Watch Video Solution](#)


22. State one condition under which a bimolecular reaction may be kinetically of first order reactions.

- A. When both reactants have same concentration
- B. when one of the reacting species is in large excess
- C. when the reaction is at equilibrium
- D. when the activation energy of reaction is less

Answer: B

 [Watch Video Solution](#)

23. The hydrolysis of ethyl acetate is a reaction of :



- A. zero order
- B. pseudo first order
- C. second order

D. third order

Answer: B

 [Watch Video Solution](#)

24. Which one of the following is wrongly matched?

- A. Saponification of $CH_3COOC_2H_5$ - second order reaction
- B. Hydrolysis of CH_3COOCH_3 - Pseudounimolecular reaction
- C. Decomposition of H_2O_2 - First order reaction
- D. Combination of H_2 and Br_2 to give HBr - Zero order reaction

Answer: D

 [Watch Video Solution](#)

25. In a pseudo first order hydrolysis of ester in water the following results were obtained:

t/s	0	30	60	90
[Ester]	0.55	0.31	0.17	0.085

(i) Calculate the average rate of reaction between the time interval 30 to 60 seconds.

(ii) Calculate the pseudo first order rate constant for the hydrolysis of ester.

A. $1.91 \times 10^{-2} s^{-1}$

B. $4.67 \times 10^{-3} mol L^{-1} s^{-1}$

C. $1.98 \times 10^{-3} s^{-1}$

D. $2.07 \times 10^{-2} s^{-1}$

Answer: B



[Watch Video Solution](#)

1. The Activation energy for a chemical reaction mainly depends upon

A. the difference in energies of reactants and products

B. the sum of energies of reactants and products

C. the difference in energy of intermediate complex with the average energy of reactants and products

D. the difference in energy of intermediate complex and the average energy of reactants.

Answer: D



[Watch Video Solution](#)

2. The temperature dependence of the rate of a chemical reaction can be explained by Arrhenius equation which is

A. $k = Ae^{E_a/RT}$

B. $k = Ae^{-E_a/RT}$


$$C. k = Ae \times \frac{E_a}{RT}$$

$$D. k = Ae \times \frac{RT}{E_a}$$

Answer: B

 Watch Video Solution

3. The energy diagram of a reaction $P + Q \rightarrow R + S$ is given. What are A and B in the graphs?



A. A \rightarrow activation energy, B \rightarrow heat of reaction

B. A \rightarrow threshold energy, B \rightarrow heat of reaction


C. A \rightarrow heat of reaction, B \rightarrow activation energy

D. A \rightarrow potential energy, B \rightarrow energy of reaction

Answer: A

 Watch Video Solution

4. The potential energy diagram for a reaction $X \rightarrow Y$ is given. A and C in the graph corresponding to



A. A \rightarrow activation energy, C $\rightarrow \Delta H^\circ$

B. A \rightarrow energy of reactants, C \rightarrow energy of products


C. $A \rightarrow \Delta H^\circ$, $C \rightarrow$ activation energy

D. $A \rightarrow$ activation energy, $C \rightarrow$ threshold energy


Answer: A

 [Watch Video Solution](#)


5. An endothermic reaction with high activation energy for the forward reaction is given by the diagram




A.



B.



C.



D.

Answer: C

[Watch Video Solution](#)

6. For an endothermic reaction, where ΔH represents the enthalpy of reaction in kJmol^{-1} , the minimum value for the energy of activation will be

- A. Less the zero
- B. equal to ΔH
- C. less than ΔH
- D. more than ΔH

Answer: D

[Watch Video Solution](#)

7. If hydrogen and oxygen are mixed and kept in the same vessel at room temperature, the reaction does not take place to form water because

- A. activation energy for the reaction is very high at room temperature
- B. molecules have no proper orientation to react to form water
- C. the frequency of collisions is not high enough for the reaction to take place
- D. no catalysts present in the reaction mixture.

Answer: A



Watch Video Solution

8. The temperature dependence of the rate of a chemical reaction is given by arrhenius equation, $k = Ae^{-E_a/RT}$. Which of the following graphs will be a straight line?

A. In A vs $1/T$

B. In A vs E_a

C. In K vs $1/T$

D. In k vs $-E_a/R$

Answer: C




Watch Video Solution

9. The temperature dependence of the rate constant k is expressed as

$$k = Ae^{-E_a/RT}$$

When a plot between $\log k$ and $1/T$ is plotted we get the graph as shown.



What is the value of slope in the graph?

- A. $\frac{E_a}{RT}$
- B. $-\frac{E_a}{2.303R}$
- C. $-\frac{E_a}{2.303RT} \log A$
- D. $-\frac{E_a}{2.303} \frac{R}{T}$

Answer: B



Watch Video Solution

10. Which of the following statements is not correct?

A. For a zero order reaction $t_{1/2}$ is proportional to initial concentration.

B. The relationship of variation of rate constant with temperature is given by

$$\frac{\log(k_2)}{k_1} = \frac{E_a}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

C. The unit of rate constant for a reaction is $\text{mol}^{1-n} \text{L}^{n-1} \text{s}^{-1}$ where n is order of the reaction.

D. The unit of rate of reaction changes with order of reaction.

Answer: D



[Watch Video Solution](#)

11. The decomposition of hydrocarbon follows the equation

$$k = (4.5 \times 10^{11} \text{ s}^{-1}) e^{-28000K/T}$$

Calculate E_a .

- A. 669 kJ mol^{-1}
- B. $232.79 \text{ kJ mol}^{-1}$
- C. $4.5 \times 10^{11} \text{ kJ mol}^{-1}$
- D. $28000 \text{ kJ mol}^{-1}$

Answer: B



[Watch Video Solution](#)

12. The rate constant for a first order reaction at 300°C for which E_a is 35

kcal mol^{-1} and frequency constant is $1.45 \times 10^{11} \text{ s}^{-1}$ is

- A. $10 \times 10^2 \text{ s}^{-1}$
- B. $5.37 \times 10^{10} \text{ s}^{-1}$

C. $5 \times 10^{-4} \text{ s}^{-1}$


D. $7.94 \times 10^{-3} \text{ s}^{-1}$

Answer: D

 Watch Video Solution

13. The graph of the effect of catalyst on activation energy is given below.

Fill up the blanks X and Y with appropriate statements



A. X = energy of activation without catalyst,

Y = energy of activation with catalyst

B. X = path of reaction with catalyst, Y = path of reaction without catalyst

C. X = energy of activation with catalyst,

Y = energy of activation without catalyst


D. X = energy of endothermic reaction, Y = energy of exothermic reaction

Answer: C



Watch Video Solution

14. Match the column I with column II and mark the appropriate choice.

Column I		Column II	
(A)	Zero order	(i)	$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$
(B)	First order	(ii)	
(C)	Endothermic reaction	(iii)	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$
(D)	Activation energy	(iv)	$k = \frac{1}{t} ([A]_0 - [A])$

A. (A) → (iv), (B) → (iii), (C) → (ii), (D) → (i)

B. (A) → (i), (B) → (ii) (C) → (iii), (D) → (iv)

C. (A) → (ii), (B) → (iii), (C) → (iv), (D) → (i)

D. (A) → (iii), (B) → (iv) (C) → (i), (D) → (ii)

Answer: A



Watch Video Solution

15. Which of the following statement about the catalyst is/are true?

- A. A catalyst makes the reaction feasible by making ΔG more negative.
- B. A catalyst makes equilibrium constant more favourable for forward reaction
- C. A catalyst accelerate rate of reaction by bringing down the activation energy
- D. A catalyst always increases the rate of reaction.

Answer: C

 [Watch Video Solution](#)

16. When a catalyst is used an equilibrium process,

- A. it increases the rate of forward reaction

B. it decreases the rate of backward reaction

C. it decreases activation energy of forward process and decreases activation energy of backward process

D. it fastens the attainment of equilibrium by lowering activation energy.

Answer: D

 [Watch Video Solution](#)

17. Which of the following factors are responsible for the increase in the rate of a surface catalysed reaction?

(i) A catalyst provides proper orientation for the reactant molecules to react.

(ii) heat of adsorption of reactants on a catalyst helps reactant molecules to overcome activation energy.

(iii) The catalyst increases the activation energy of the reactio.

A. (i) and (iii)

B. (i) and (ii)

C. (ii) and (iii)

D. (i),(ii) and (iii)

Answer: B



Watch Video Solution

Collision Theory Of Chemical Reactions

1. The increase in concentration of the reactants lead changes in

A. ΔH

B. collision frequency

C. activation energy

D. equilibrium constant.

Answer: B



[Watch Video Solution](#)

2. The rate constant is given by the equation $k = P \cdot Z e^{-E_a/RT}$. Which factor should register a decrease for the reaction to proceed more rapidly?

A. T

B. Z

C. E

D. pressure.

Answer: C



[Watch Video Solution](#)

3. Rate of a general reaction $A + B \rightarrow$ products can be expressed as follows on the basis of collision theory.

$$\text{Rate} = Z_{AB}e^{-E_a/RT}$$

Which of the following statements is not correct for the above expression?

A. Z is collision frequency and is equal to number of collisions per second per unit volume of the reaction mixture

B. $e^{E_a/RT}$ is the fraction of molecules with kinetic energy equal to or greater than E_a

C. E_a is activation energy of the reaction.

D. All the molecules which collide with one other are effective collisions.

Answer: D



Watch Video Solution

4. Threshold energy is equal to

- A. activation energy
- B. activation energy - energy of molecules
- C. activation energy + energy of molecules
- D. None of these

Answer: C



Watch Video Solution

5. For a certain reaction a large fraction of molecules has energy more than the threshold energy, still the rate of reaction is very slow. The possible reason for this could be that

- A. the colliding molecules could be large in size
- B. the colliding molecules must not be properly oriented for effective collisions

C. the rate of reaction could be independent of the energy

D. one of the reactant could be in excess.

Answer: B

 [Watch Video Solution](#)

6. fill up the following with suitable terms

(i) Activation energy = Threshold energy - _____

(ii) Half-life period of zero order reaction = _____

(iii) Average rate of reaction = _____

(iv) Instantaneous rate of reaction

A. Potential energy $\frac{0.693}{k}$, $\frac{dx}{dt}$, $\frac{\Delta[A]}{\Delta t}$

B. Energy of reactants $\frac{1}{k}$, $\frac{\Delta[A]}{\Delta t}$, $\frac{dx}{dt}$

C. Energy of reaction $\frac{\log x}{t}$, $\frac{\Delta[A]}{\Delta t}$, $\frac{dx}{dt}$

D. Average kinetic energy of reactants $\frac{a}{2k}$, $\frac{\Delta[A]}{\Delta t}$, $\frac{dx}{dt}$

Answer: D



[Watch Video Solution](#)

Ncert Exemplar

1. The role of a catalyst is to change

- A. Gibbs energy of reaction
- B. enthalpy of reaction
- C. activation energy of reaction
- D. equilibrium constant.

Answer: C



[Watch Video Solution](#)

2. In the presence of a catalyst, the heat evolved or absorbed during the reaction:

- A. increases
- B. decreases
- C. remains unchanged
- D. may increase or decrease

Answer: C

 [Watch Video Solution](#)


3. Activation energy of a chemical reaction can be determined by

- A. determining the rate constant at standard temperature
- B. determining the rate constants at two temperatures
- C. determining probability of collision
- D. using catalyst

Answer: B

 [Watch Video Solution](#)

4. Consider figure and mark the correct option.




- A. Activation energy of forward reaction is $E_1 + E_2$ and product is less stable than reactant.
- B. Activation energy of forward reaction is $E_1 + E_2$ and product is more stable than reactant.
- C. Activation energy of both forward and backward reaction is $E_1 + E_2$ and reactant is more stable than product.

D. Activation energy of backward reaction is E_1 and product is more stable than reactant.

Answer: A

 [Watch Video Solution](#)

5. Consider a first order gas phase decomposition reaction given below:



The initial pressure of the system before decomposition of A was p_i . After lapse of time t' . Total pressure of the system increased by x units and became P_t . the rate constant k for the reaction is given as

A. $k = \frac{2.303}{t} \frac{\log(p_i)}{p_i - x}$

B. $k = \frac{2.303}{t} \frac{\log(p_i)}{2p_i - p_t}$


C. $k = \frac{2.303}{t} \frac{\log(p_i)}{2p_i + p_i}$

D. $k = \frac{2.303}{t} \frac{\log(p_i)}{p_i + x}$


Answer: B

6. According to Arrhenius equation rate constant k is equal to $Ae^{-E_a/RT}$.


Which of the following option. Represents the graph of $\ln k$ us $\frac{1}{T}$?




A.



B.



C.



D.

Answer: A

7. Consider the Arrhenius equation given below and mark the correct option.

$$k = Ae^{-\frac{E_a}{RT}}$$

- A. Rate constant increases exponentially with increasing activation energy and decreasing temperature
- B. Rate constant decreases exponentially with increasing activation energy and decreasing temperature.
- C. Rate constant increases exponentially with decreasing activation energy and decreasing temperature.
- D. Rate Constant increases exponentially with decreasing activation energy and increasing temperature.

Answer: D



[Watch Video Solution](#)

8. Which of the following statement is not correct about order of a reaction ?

- A. The order of a reaction can be a fractional number.
- B. Order of a reaction is experimaentally determined quantity
- C. The order or a reaction is always equal to the sum of the stoichiometric coefficients of reactants in the balanced chemical equaiton for a reaction.
- D. The order of a reaction is the sum of the powers of molar concentration of the reactants in the rate law expression.


Answer: C



[View Text Solution](#)

9. A graph of volume of hydrogen released vs time for the reaction between zinc and dil. HCl is given in figure. On the basic of this mark the

correct option.



- A. Average rate upto 40 seconds is $\frac{V_3 - V_2}{40}$
- B. Average rate upto 40 seconds is $\frac{V_3 - V_2}{40 - 30}$
- C. Average rate upto 40 seconds is $\frac{v_3}{40}$
- D. Average rate upto 40 seconds is $\frac{V_3 - V_1}{40 - 20}$

Answer: C



View Text Solution

10. Consider the graph given in Q.9 Which of the following options does not show instantaneous rate of reaction at 40th second?

A. $\frac{V_5 - V_2}{50 - 30}$

B. $\frac{V_4 - V_2}{50 - 30}$

C. $\frac{V_3 - V_2}{40 - 30}$

D. $\frac{V_3 - V_1}{40 - 20}$

Answer: B



[View Text Solution](#)

11. Which of the following statements is correct?

A. The rate of a reaction decreases with passage of time as the concentration of reactants decreases.

B. The rate of a reaction is same at any time during the reaction.


C. The rate of a reaction is independent of temperature change.

D. The rate of a reaction decreases with increases in concentration of reactant (s)

Answer: A

 [Watch Video Solution](#)

12. Which of the following expression is correct for the rate of reaction given below ?



A. $\frac{\Delta[Br^{-}]}{\Delta t} = 5 \frac{\Delta[H^{+}]}{\Delta t}$




B. $\frac{\Delta[Br^{-}]}{\Delta t} = \frac{6}{5} \frac{\Delta[H^{+}]}{\Delta t}$

C. $\frac{\Delta[Br^{-}]}{\Delta t} = \frac{5}{6} \frac{\Delta[H^{+}]}{\Delta t}$

D. $\frac{\Delta[Br^{-}]}{\Delta t} = 6 \frac{\Delta[H^{+}]}{\Delta t}$

Answer: C

13. Which of the following graphs represents exothermic reaction?



A. (i) Only

B. (ii) Only

C. (iii) only

D. (i) and (ii)

Answer: A

14. Rate law for the reaction, $A + 2B \rightarrow C$ is found to be

$$\text{Rate} = k[A][B]$$

Concentration of reactant 'B' is doubled keeping the concentration of 'A' constant, the value of rate constant will be _____

- A. the same
- B. doubled
- C. quadrupled
- D. halved

Answer: A



[Watch Video Solution](#)

15. Which of the following statements is incorrect about the collision theory of chemical reaction?

- A. It considers reacting molecules or atoms to be hard spheres and ignores their structural features.
- B. Number of effective collisions determines the rate of reaction.
- C. Collision of atoms or molecules possessing sufficient threshold energy results into the product formation.
- D. Molecules should collide with sufficient threshold energy and proper orientation for the collision to be effective.

Answer: C

 [Watch Video Solution](#)

16. A first order reaction is 50 % completed in 1.26×10^{14} s. How much time would it take for 100 % completion?

A. 1.26×10^{15} s

B. 2.52×10^{14} s

C. $2.52 \times 10^{28} s$


D. Infinite

Answer: D



Watch Video Solution

17. Compound 'A' and B react according to the following chemical equation.



Concentration of either A or B were changed keeping the concentrations of one of the reactants constant and rates were measured as a function of initial concentration. Following results were obtained. Choose the correct option for the rate equations for this reaction.

Experiment	Initial concentration of $[A] / molL^{-1}$	Initial concentration of $[B]$
1	0.30	0.30
2	0.30	0.60
3	0.60	0.30

A. Rate = $k[A]^2[B]$

B. Rate = $k[A][B]^2$

C. Rate = $k[A][B]$

D. Rate = $k[A]^2[B]^0$

Answer: B



Watch Video Solution

18. Which of the following statements is not correct for the catalyst?

- A. It catalyts the forward and backward reaction to the same extent.
- B. It alters ΔG of the reaction.
- C. It is a substance that does not change the equilibrium constant of a reaction.
- D. It provides an alternate mechanism by reducing activation energy between reactants and products.

Answer: B



[Watch Video Solution](#)

19. The value of rate constant of a pseudo first order reaction


- A. depends on the concentration of reactants present in small amount
- B. depends on the concentrations of reactants present in excess
- C. is independent of the concentration of reactants
- D. depends only on temperature

Answer: B




[Watch Video Solution](#)


20. Consider the reaction $A \rightarrow B$. The concentration of both the reactants and the products varies exponentially with time. Which of the following figure correctly describes the change in concentration of reactants and products with time ?




A.



B.



C.



D.

Answer: B



Watch Video Solution

Assertion And Reason

1. Assertion : Precipitation of silver chloride occurs instantaneously by mixing of aqueous solutions of silver nitrate and sodium chloride.

Reason : Ionic reactions occur very fast.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: A



[Watch Video Solution](#)

2. Assertion : The rate of reaction is the rate of change of concentration of a reactant or a product.


Reason : Rate of reaction remains constant during the complete reaction.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

3. Assertion : For the reaction



$$\text{Rate} = k[CHCl_3][Cl_2]^{1/2}$$

Reason : Rate of reaction is always equals to the sum of the stoichiometric coefficients of the reacting species in a balanced chemical equation.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C



Watch Video Solution

4. Assertion : Order of a reaction with respect to any reactant can be zero, positive negative of fractional.

Reason : Rate of a reaction cannot decrease with increase in concentration of a reactant or a product.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

5. Assertion : Molecularity greater than three is not observed.

Reason : The overall molecularity of complex reaction is equal to molecularity of the slowest step.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

6. Assertion : Complex reaction takes place in different steps and the slowest step determines the rate of reaction.

Reason : Order and molecularity of a reaction are always equal.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C



[Watch Video Solution](#)

7. Assertion : The decomposition of gaseous ammonia on a hot platinum surface is a zero order reaction at high pressure.

For a zero order reaction the rate of reaction is independent of initial concentration.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: B



[Watch Video Solution](#)

8. Assertion : For a first order reaction, $t_{1/2}$ is independent of rate constant.

Reason : For a first reaction $t_{1/2} \propto [R]_0$.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: D

 Watch Video Solution

9. Assertion : The reaction

$C_{12}H_{22}O_{11} + H_2O \xrightarrow{H^+} C_6H_{12}O_6 + C_6H_{12}O_6$ is a first order reaction.
Cane sugar Glucose Fructose

Reason : Change in concentration of H_2O is negligible.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A



Watch Video Solution

10. Assertion : The rate of a reaction sometimes does not depend on concentration.

Reason : Lower the activation energy, faster is the reaction

- A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

11. Assertion : For a chemical reaction with rise in temperature by 10° the rate constant is nearly doubled.

Reason : At $t + 10$, the fraction of molecules having energy equal to or greater than activation energy gets doubled.

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: A

 [Watch Video Solution](#)

12. Assertion : E_a of the forward reaction is higher than that of backward reaction in a reversible endothermic reaction.

Reason : Increasing the temperature of the substance increases the fraction of molecules which collide with energies greater than E_a .

A. If both assertion and reason are true and reason is the correct explanation of assertion.

B. If both assertion and reason are true but reason is not the correct explanation of assertion.

C. If assertion is true but reason is false.

D. If both assertion and reason are false.

Answer: B

 [Watch Video Solution](#)

13. Assertion : A catalyst increases the rate of reaction without itself undergoing any permanent chemical change.

Reason : A catalyst changes the Gibbs energy (ΔG) of the reaction and equilibrium constant of the reaction.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: C

 [Watch Video Solution](#)

14. Assertion : All molecules collisions lead to the formation of products.

Reason : Reactant molecules undergo chemical change irrespective of their collision.

Assertion : Rate of reaction increases with increase in temperature .

Reason : Number of effective collisions increases with increase in temperature.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: D



[View Text Solution](#)

15. Assertion : Rate of reaction increases with increase in temperature.

Reason : Number of effective collisions increases with increase in temperature.

- A. If both assertion and reason are true and reason is the correct explanation of assertion.
- B. If both assertion and reason are true but reason is not the correct explanation of assertion.
- C. If assertion is true but reason is false.
- D. If both assertion and reason are false.

Answer: A



[Watch Video Solution](#)