



## MATHS

### BOOKS - NTA MOCK TESTS

### JEE MOCK TEST 2

#### Mathematics

1. Let  $z_1$  and  $z_2$  be  $n^{\text{th}}$  roots of unity which subtend a right angle at the origin. Then  $n$  must be of the form

A.  $4K + 1$

B.  $4K + 2$

C.  $4k+3$

D.  $4k$

**Answer: D**



**Watch Video Solution**

2. In  $(0, 2\pi)$ , the total number of points where  $f(x) = \max\{\sin x, \cos x, 1 - \cos x\}$  is not differentiable, are equal to

A. 3

B. 4

C. 5

D. 6

**Answer: A**



**Watch Video Solution**

3. The integral  $\int x \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) dx, (x > 0)$  is equal to

A.  $-x + (1 + x^2)\cot^{-1} x + c$

B.  $x - (1 + x^2)\cot^{-1} x + c$

C.  $x - (1 + x^2)\tan^{-1} x + c$

D.  $-x + (1 + x^2)\tan^{-1} x + c$

**Answer: D**



**Watch Video Solution**

4. The coefficient of  $x^n$  in the polynomial

$$(x + {}^{2n+1}C_0)(X + {}^{2n+1}C_1)(x + {}^{2n+1}C_2)\dots(X + {}^{2n+1}C_n)$$
 is

A.  $2^{n+1}$

B.  $2^{2n+1} - 1$

C.  $2^{2n-1}$

D.  $2^{2n}$

**Answer: D**

 [Watch Video Solution](#)

5. The mean of the data set comprising of 16 observations is 16. If one of the observation valued 16 is deleted and three new observations valued 3, 4 and 5 are added to the data, then the mean of the resultant data, is-

- A. 14.0
- B. 16.8
- C. 16.0
- D. 15.8

**Answer: A**

 [Watch Video Solution](#)

6. The lines represented by the equation

$$x^2 + 2\sqrt{3}xy + 3y^2 - 3x - 3\sqrt{3}y - 4 = 0, \text{ are}$$

- A. perpendicular to each other
- B. parallel
- C. inclined at  $45^\circ$  to each other
- D. None of these

**Answer: B**

 [Watch Video Solution](#)

7. At present, a firm is manufacturing 2000 items. It is estimated that the rate of change of production  $P$  w.r.t. additional number of workers  $x$  is given by  $\frac{dP}{dx} = 100 - 12\sqrt{x}$ . If the firm employs 25 more workers, then the new level of production of items is

- A. 3500

B. 4500

C. 2500

D. 3000

**Answer: A**



[Watch Video Solution](#)

8. Let  $f: R \rightarrow R$  be a periodic function such that  $f(T + x) = 1$

where  $T$  is a  $\left[1 - 3f(x) + 3(f(x))^2 - (f(x))^3\right]^{\frac{1}{3}}$  fixed positive

number, then period of  $f(x)$  is

A.  $T$

B.  $2T$

C.  $3T$

D. None of these

**Answer: B**

 [Watch Video Solution](#)

9. AB is vertical tower. The point A is on the ground and C is the middle point of AB. The part CB subtend an angle  $\alpha$  at a point P on the ground. If  $AP = nAB$ , then  $\tan \alpha =$

A.  $n = (n^2 + 1)\tan \alpha$

B.  $n = (2n^2 - 1)\tan \alpha$

C.  $n^2 = (2n^2 + 1)\tan \alpha$

D.  $n = (2n^2 + 1)\tan \alpha$

**Answer: D**

 [Watch Video Solution](#)

10. If  $f(x) = \begin{cases} x + 1 & x > 1 \\ 0, & x = 1 \\ 7 - 3x, & x < 1 \end{cases}$  then  $f'(0)$  equal to

A.  $-1$

B.  $-2$

C.  $-3$

D.  $-4$

**Answer: C**

 [Watch Video Solution](#)

11. If  $x = -1$  and  $x = 2$  are extreme points of  $f(x) = \alpha \log|x| + \beta x^2 + x$ , then

A.  $\alpha = 2, \beta = -\frac{1}{2}$

B.  $\alpha = 2, \beta = \frac{1}{2}$



$$C. \alpha = -6, \beta = \frac{1}{2}$$

$$D. \alpha = -6, \beta = -\frac{1}{2}$$

**Answer: A**

 [Watch Video Solution](#)

12. If  $x$  takes negative permissible value then  $\sin^{-1} x =$

A.  $-\cos^{-1} \sqrt{1 - x^2}$

B.  $\cos^{-1} \sqrt{x^2 - 1}$

C.  $\pi - \cos^{-1} \sqrt{1 - x^2}$

D.  $\cos^{-1} \sqrt{1 - x^2}$

**Answer: A**

 [Watch Video Solution](#)

13. If  $f(x) = \prod_{k=1}^{999} (x^2 - 47x + k)$ . then product of all real roots of  $f(x) = 0$  is

A. 550!

B. 551!

C. 552!

D. 999!

**Answer: C**



[Watch Video Solution](#)

14. In a certain town 25% families own a cellphone, 15% families own a scooter and 65% families own neither a cellphone nor a scooter. If 500 families own both a cellphone and scooter, then total number of families in the town is

A. 10000

B. 20000

C. 30000

D. 40000

**Answer: C**



[Watch Video Solution](#)

15. The number of ways of arranging 8 men and 5 women around a circular table such that no two women can sit together is

A.  $8!$

B.  $4!$

C.  $8!4!$

D.  $7! \times 8P_3$

**Answer: D**



16. Let  $A = \begin{bmatrix} -1 & 2 & -3 \\ -2 & 0 & 3 \\ 3 & -3 & 1 \end{bmatrix}$  be a matrix, then  $|A|adj(A^{-1})$  is equal to

A.  $O_{3 \times 3}$

B.  $\begin{bmatrix} -1 & 2 & -3 \\ -2 & 0 & 3 \\ 3 & -3 & 1 \end{bmatrix}$

C.  $I_3$

D.  $[-3, -3, 1), (3, 0, -2), (-1, 2, -3)]$

**Answer: B**

 Watch Video Solution

17. If the equation  $x^2 + 4 + 3 \sin(ax + b) - 2x = 0$  has at least one real solution, where  $a, b \in [0, 2\pi]$  then one possible value of  $(a + b)$

can be equal to

A.  $\frac{7\pi}{2}$

B.  $\frac{5\pi}{2}$

C.  $\frac{9\pi}{2}$

D. None of these

**Answer: A**



**Watch Video Solution**

**18.** Fifteen coupons are numbered 1, 2, 3, ..., 15 respectively. Seven coupons are selected at random one at a time with replacement. The probability that the largest number appearing on a selected coupon is 9 is :

A.  $\left(\frac{1}{15}\right)^7$

B.  $\left(\frac{3}{5}\right)^7$

C.  $\left(\frac{8}{15}\right)^7$

D.  $\left(\frac{2}{5}\right)^7$

**Answer: B**

 [Watch Video Solution](#)

19. For  $x \in R, x \neq 0$  if  $y(x)$  is a differentiable function such that  $x \int_1^x y(t) dt = (x + 1) \int_1^x ty(t) dt$ , then  $y(x)$  equals (where C is a constant)

A.  $Cx^3 e^{\frac{1}{x}}$

B.  $\frac{C}{x^2} e^{-\frac{1}{x}}$

C.  $\frac{C}{x} e^{-\frac{1}{x}}$

D.  $\frac{C}{x^3} e^{-\frac{1}{x}}$

**Answer: D**

 [Watch Video Solution](#)

20. If O is the origin and OP, OQ are distinct tangents to the circle  $x^2 + y^2 + 2gx + 2fy + c = 0$  then the circumcentre of the triangle OPQ is

- A.  $(-g, -f)$
- B.  $(g, f)$
- C.  $(-f, -g)$
- D. None of these

**Answer: D**

[Watch Video Solution](#)

21. The inradius of the triangle having sides 26,28,30 units is

[Watch Video Solution](#)

22. Let the sum  $\sum_{n=1}^9 \frac{1}{n(n+1)(n+2)}$  written in the rational form be  $\frac{p}{q}$  (where  $p$  and  $q$  are co-prime), then the value of  $\left[ \frac{q-p}{10} \right]$  is (where  $[.]$  is the greatest integer function)

 [Watch Video Solution](#)

23. In a  $\Delta ABC$  if  $\angle A = \angle B = \frac{1}{2} \left( \sin^{-1} \left( \frac{\sqrt{6}+1}{2\sqrt{3}} \right) + \sin^{-1} \left( \frac{1}{\sqrt{3}} \right) \right)$  and length of  $c = 6.3^{\frac{1}{4}}$ , then the area of  $\Delta ABC$

 [Watch Video Solution](#)

24.  $\lim_{x \rightarrow 0} \frac{\sin^{-1} x - \tan^{-1} x}{x^3}$  is equal to

 [Watch Video Solution](#)



25. A farmer  $F_1$  has a land in the shape of a triangle with vertices at  $P(0, 0)$ ,  $Q(1, 1)$  and  $R(2, 0)$ . From this land, a neighbouring farmer  $F_2$  takes away the region which lies between the side  $PQ$  and a curve of the form  $y = x^n$ , ( $n > 1$ ). If the area of the region taken away by the farmer  $F_2$  is exactly 30% of the area of  $\Delta PQR$ , then the value of  $n$  is \_\_\_\_\_.

 Watch Video Solution

26. If the integral

$$I = \int x^{\sin x} \left( \cos x \cdot nx + \frac{\sin x}{x} \right) dx, = (f(x))^{g(x)} + c (\forall x > 0)$$

then the range of  $y = g(x)$  is (where,  $c$  is an arbitrary constant)

A.  $[-1,1]$

B.  $[0,1]$

C.  $(0,1)$

D.  $(-1,1)$

Answer: A



Watch Video Solution

27. Let P and Q are two points in the xy plane on the curve  $y = x^{11} - 2x^7 + 7x^3 + 11x + 6$  such that  $\overrightarrow{OP} \cdot \hat{i} = 5, \overrightarrow{OQ} \cdot \hat{i} = -5$ , then the magnitude of  $\overrightarrow{OP} + \overrightarrow{OQ}$  is

A. 10

B. 12

C. 14

D. 8

Answer: B



Watch Video Solution

28. If the letters of the word REGULATIONS be arranged at random, find the probability that there will be exactly four letters between the  $R$  and the  $E$ .

A.  $\frac{6}{55}$

B.  $\frac{3}{55}$

C.  $\frac{5}{11}$

D.  $\frac{6}{11}$

Answer: A

 Watch Video Solution

29. If the point of intersection of the plane  $4x - 5y + 2z - 6 = 0$  with the line through the origin and perpendicular to the plane  $x - 2y - 4z = 4$  is P, then the distance of the point P from (1, 2, 3) is

A.  $\sqrt{63}$  units

B. 8 units

C.  $\sqrt{65}$  units

D.  $\sqrt{72}$  units

**Answer: C**



**Watch Video Solution**

**30.** The mean and variance of seven observations are 8 and 16 respectively. If five of these are 2,4,10,12 and 14, then find the remaining two observations.

A. 5,7

B. 3,5

C. 6,8

D. 4,2

**Answer: C**



**Watch Video Solution**

31. Let  $\triangle ABC$  is an isosceles triangle with  $AB = AC$ . If  $B = (0, a)$ ,  $C = (2a, 0)$  and the equation of AB is  $3x - 4y + 4a = 0$ , then the equation of side AC is

A.  $y = 8x - 16a$

B.  $3y = 4x - 8a$

C.  $x = 2a$

D.  $y + 8x = 16a$

**Answer: C**



**Watch Video Solution**

32. Let  $A(0,3)$  and  $B(0,12)$  be two vertices of a  $\triangle ABC$  where  $C = (x, 0)$ . If the circumcircle of  $\triangle ABC$  touches the x-axis, then the value of  $\cos 2\theta$  is (where  $\theta$  is angle ACB)

A.  $\frac{1}{3}$

B.  $\frac{1}{2}$

C.  $\frac{8}{15}$

D.  $\frac{7}{25}$

Answer: D

 [Watch Video Solution](#)

33. Consider three statements  $p$  : person 'A' passed in mathematics exam  $q$  : Person 'A' passed in physics exam  $r$  : Person 'A' passed in chemistry exam, Then the statement  $\sim((\sim(p \Rightarrow q) \Rightarrow r))$  is equivalent to

- A. Person A passed only in mathematics & physics & chemistry
- B. Person A failed only in mathematics & physics & chemistry
- C. Person A passed in all the three subjects mathematics & physics & chemistry
- D. Person A passed in chemistry but failed in mathematics & physics.

**Answer: A**

 [Watch Video Solution](#)

**34.** In equation  $(Z - 1)^n = Z^n = 1 (\forall n \in N)$  has solution, then n can be

- A. 4
- B. 12
- C. 15

**Answer: B** [Watch Video Solution](#)

35. The solution of the differential equation

$$\left(3x^2 \sin\left(\frac{1}{x}\right) + y\right)dx = x \cos\left(\frac{1}{x}\right)dx - xdy$$

is (where,  $c$  is an arbitrary constant)

A.  $\sin\left(\frac{1}{x}\right) = xy + c$

B.  $x^3 \sin\left(\frac{1}{x}\right) + xy = c$

C.  $x^3 \sin\left(\frac{1}{x}\right) = xy + c$

D.  $\sin\left(\frac{1}{x}\right) = x^3y + c$

**Answer: B** [Watch Video Solution](#)



36. The quadratic equation  $(1 - \sin \theta)x^2 + 2(1 - \sin \theta)x - 3 \sin \theta = 0$  has both roots complex for all  $\theta$  lying in the interval

A.  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

B.  $\left(0, \frac{3\pi}{2}\right)$

C.  $\left(\frac{\pi}{6}, \frac{7\pi}{6}\right)$

D.  $\left(\frac{7\pi}{6}, \frac{11\pi}{6}\right)$

Answer: D

 [Watch Video Solution](#)

37. The minimum value of the expression  $3x + 2y$  ( $\forall x, y > 0$ ), where  $xy^2 = 10$ , occurs when the value of  $y$  is equal to

A.  $\sqrt{10}$

B.  $\sqrt[3]{10}$

C.  $\sqrt[3]{30}$

D.  $\frac{1}{\sqrt{30}}$

**Answer: C**

 [Watch Video Solution](#)

38. If in the expansion of  $(1 + x)^m(1 - x)^n$  the coefficient of  $x$  and  $x^2$  are 3 and  $(-6)$  respectively, then the value of  $n$  is-

A. 6

B. 9

C. 12

D. 24

**Answer: B**



39. Number of words that can be formed with the letters of the word ALGEBRA so that all the vowels are separated (or no two vowels come together) is

- A. 720
- B. 2160
- C. 1440
- D. 1200

**Answer: A**

40. If  $f(k - x) + f(x) = \sin x$ , then the value of integral

$$I = \int_0^k f(x) dx \text{ is equal to}$$

A.  $\cos k$

B.  $2 \cos^2 \left( \frac{k}{2} \right)$

C.  $\sin^2 \left( \frac{k}{2} \right)$

D.  $\sin k$

**Answer: C**



**Watch Video Solution**

**41.** If the difference between the number of subsets of the sets  $A$  and  $B$  is 120, then choose the incorrect option.

A. Maximum value of  $n(A \cap B) = 3$

B. Minimum value of  $n(A \cap B) = 0$

C. Maximum value of  $n(A \cup B) = 21$

D. Minimum value of  $n(A \cup B) = 7$

**Answer: C**



**Watch Video Solution**

**42.** For the function  $f(x) = \sin(\pi[x]) \times \cos^{-1}([x])$ , choose the correct option.

(where  $[.]$  represents the greatest integer function)

- A. Domain of  $f(x)$  in  $[-1,1]$
- B. Range of  $f(x)$  contains exactly two elements
- C.  $f(x)$  is an identify function
- D.  $f(x)$  is a constant function

**Answer: D**



**Watch Video Solution**

43. The value of  $\lim_{x \rightarrow 0} \frac{\sin x}{3} \left[ \frac{5}{x} \right]$  is equal to

[where  $[.]$  represent the greatest integer function)

A.  $\frac{1}{3}$

B. 0

C.  $\frac{5}{3}$

D. 1

**Answer: C**

 [Watch Video Solution](#)

44. If  $y = \tan^{-1} \left( \frac{x}{1 + 6x^2} \right) + \tan^{-1} \left( \frac{2x - 1}{2x + 1} \right)$ , ( $\forall x > 0$ ) then

$\frac{dy}{dx}$  is equal to

A.  $\frac{3}{1 + 9x^2}$

B.  $\frac{1}{1 + 6x^2}$

C.  $\frac{1}{1 + 6x^2} + \frac{1}{1 + x^2}$

D.  $\frac{3}{1 + 6x^2}$

**Answer: A**

 [Watch Video Solution](#)

45. If  $|3x - 1|$ ,  $3$ ,  $|x - 3|$  are the first three terms of an arithmetic progression, then the sum of the first five terms can be

A. 5

B. 10

C. 20

D. 30

**Answer: A**

 [Watch Video Solution](#)

$$46. \text{ If } f(x) = \begin{cases} px + q & : x \leq 2 \\ x^2 - 5x + 6 & : 2 < x < 3 \\ ax^2 + bx + 1 & : x \geq 3 \end{cases}$$

is differentiable everywhere, then  $|p| + |q| + \left| \frac{1}{a} \right| + \left| \frac{1}{b} \right|$  is equal to

 [Watch Video Solution](#)

47. If the area bounded by  $y = \left| |x|^2 - 4|x| + 3 \right|$  and the x-axis from  $x = 1$  to  $x = 3$  is  $\frac{p}{q}$  (where, p & q are coprime) then the value of  $p + q$  is

A. 10

B. 9

C. 8

D. 7

**Answer: D**

 [Watch Video Solution](#)



48. Let  $M$  be a square matrix of order 3 whose elements are real number and  $adj(adjM) = \begin{bmatrix} 36 & 0 & -4 \\ 0 & 6 & 0 \\ 0 & 3 & 6 \end{bmatrix}$ , then the absolute value of  $Tr(M)$  is [Here,  $adj P$  denotes adjoint matrix of  $P$  and  $Tr(P)$  denotes trace of matrix  $P$  i.e., sum of all principal diagonal elements of matrix  $P$ ]

 [Watch Video Solution](#)

49. If common tangents of  $x^2 + y^2 = r^2$  and  $\frac{x^2}{16} + \frac{y^2}{9} = 1$  forms a square, then the length of diagonal of the square is

- A. 9
- B. 10
- C. 12
- D. 1

**Answer: B**



**Watch Video Solution**

**50.** The angular depression of the top and the foot of the chimney seen from the top of a tower on the same base level as the chimney are  $\tan^{-1}\left(\frac{4}{3}\right)$  and  $\tan^{-1}\left(\frac{5}{2}\right)$  respectively if the height of the tower is 150m. then the distance between the top of the chimney and the tower is



**Watch Video Solution**

**51.** Let  $P_1: 2x + y + z + 1 = 0$

$P_2: 2x - y + z + 3 = 0$  and  $P_3: 2x + 3y + z + 5 = 0$  be three planes, then the distance of the line of intersection of planes  $P_1 = 0$  and  $P_2 = 0$  from the plane  $P_3 = 0$  is

A.  $\frac{3}{\sqrt{14}}$  units

B.  $\frac{6}{\sqrt{14}}$  units

C.  $\frac{3}{\sqrt{7}}$  units

D.  $\frac{6}{\sqrt{7}}$  units

**Answer: B**

 [Watch Video Solution](#)

52. The parabolas  $C_1: y^2 = 4a(x - a)$  and  $C_2: y^2 = -4a(x - k)$  intersect at two distinct points A and B. If the slope of the tangent at A on  $C_1$  is same as the slope of the normal at B on  $C_2$ , then the value of k is equal to

A.  $3a$

B.  $2a$

C.  $a$

D.  $0$

**Answer: A**



**Watch Video Solution**

**53.** Let  $p, q$  and  $r$  be three statements, then  $(p \rightarrow q) \rightarrow r$  is equivalent to

A.  $(p \vee r) \wedge (q \vee r)$

B.  $(p \vee r) \wedge (\sim q \vee r)$

C.  $(p \wedge r) \vee (q \vee r)$

D.  $(p \vee r) \rightarrow r$

**Answer: B**



**Watch Video Solution**

54. Let two sides of a rectangle of area 20 sq. units are along lines  $x - y = 0$  and  $x + y = 2$ , then the locus of the point of intersection of diagonals is a

- A. pair of ellipse
- B. pair of straight lines
- C. pair of hyperbola having eccentricity 2 and  $\frac{2}{\sqrt{3}}$
- D. pair of hyperbola each having eccentricity  $\sqrt{2}$

Answer: D

 Watch Video Solution

55. Let  $2\vec{a} = \vec{b} \times \vec{c} + 2\vec{b}$  where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are three unit vectors, then sum of all possible values of  $\left| 3\vec{a} + 4\vec{b} + 5\vec{c} \right|$  is

- A. 10

B. 12

C. 14

D. 16

**Answer: C**

 [Watch Video Solution](#)

56. If  $f(1 + x) = f(1 - x) (\forall x \in R)$ , then the value of the integral

$$I = \int_{-7}^9 \frac{f(x)}{f(x) + f(2 - x)} dx \text{ is}$$

A. 0

B. 2

C. 8

D. 10

**Answer: C**

 [Watch Video Solution](#)

57. If  $f(x)$  is a real valued function such that

$f(x + 6) - f(x + 3) + f(x) = 0, \forall x \in R$ , then period of  $f(x)$  is

- A. 6
- B. 12
- C. 18
- D. 24

**Answer: C**

 [Watch Video Solution](#)

58. The value of  $\lim_{x \rightarrow 0} \frac{\tan^2 3x}{\sqrt{5} - \sqrt{4 + \sec x}}$  is equal to

- A.  $2\sqrt{5}$

B.  $-9\sqrt{5}$

C.  $9\sqrt{5}$

D.  $-36\sqrt{5}$

**Answer: D**



**Watch Video Solution**

**59.** If  $-\pi < \theta < \pi$ , the equation

$$(\cos 3\theta + 1)x^2 + 2(\cos 2\theta - 1)x + (1 - 2\cos \theta) = 0$$

has more than two roots for

A. no value of  $\theta$

B. one value of  $\theta$

C. two value of  $\theta$

D. all value of  $\theta$



**Answer: A**



**Watch Video Solution**

60. If  $\ln(2x^2 - 5)$ ,  $\ln(x^2 - 1)$  and  $\ln(x^2 - 3)$  are the first three terms of an arithmetic progression, then its fourth term is

A.  $\ln 8 - \ln 3$

B.  $\ln 3 - \ln 8$

C.  $\ln 24$

D.  $2 \ln 6$

**Answer: A**



**Watch Video Solution**

61. Image of line  $\frac{x-2}{3} = \frac{y-1}{1} = \frac{z-1}{-4}$  in the plane  $x + y + z = 7$  is

A.  $\frac{x-4}{1} = \frac{y-3}{1} = \frac{z-3}{1}$

B.  $\frac{x-3}{1} = \frac{y-4}{1} = \frac{z-3}{1}$

C.  $\frac{x-4}{3} = \frac{y-3}{1} = \frac{z-3}{-4}$

D.  $\frac{x-3}{1} = \frac{y-4}{1} = \frac{z-3}{-4}$

**Answer: C**

 [Watch Video Solution](#)

62. If  $\pi < \theta < \frac{3\pi}{2}$  and  $\cos \theta = -\frac{3}{5}$ , then  $\tan\left(\frac{\theta}{4}\right)$  is equal to

A.  $\frac{\sqrt{5}-1}{2}$

B.  $\frac{\sqrt{5}+1}{2}$

C.  $\frac{-\sqrt{5}-1}{4}$

D.  $\frac{-\sqrt{5} + 1}{4}$

**Answer: B**

 [Watch Video Solution](#)

**63.** If the value of integral

$$\int (x + \sqrt{x^2 - 1})^2 dx = ax^3 - x + b(x^2 - 1)^{\frac{1}{b}}, + C$$

(where, C is the constant of integration), then  $a \times b$  is equal to

A. 1

B.  $\frac{4}{9}$

C. 2

D.  $\frac{9}{4}$

**Answer: B**

 [Watch Video Solution](#)

64. The range of the function  $\sin^{-1}\left(\frac{x^2}{1+x^2}\right)$  is

A.  $[-\pi/2, \pi/2]$

B.  $[0, \pi/2)$

C.  $(0, \pi/2]$

D.  $(-\pi/2, \pi/2)$

**Answer: B**

 [Watch Video Solution](#)

65. The solution of the differential equation

$$\frac{1}{x^2} \left( \frac{dy}{dx} \right)^2 + 6 = \left( \frac{5}{x} \right) \frac{dy}{dx} \text{ is } y = \lambda x^2 + c$$

(where,  $c$  is an arbitrary constant). The sum of all the possible value of

$\lambda$  is

A.  $\frac{3}{2}$

B.  $\frac{5}{2}$

C.  $\frac{2}{5}$

D. 2

**Answer: B**

 [Watch Video Solution](#)

66. The number of tangents with positive slope that can be drawn from the origin to the curve  $y = \sin x$  is

A. 0

B. 2

C. 4

D. infinitely many

**Answer: D**

 [Watch Video Solution](#)

67. A complex number  $z$  is said to be unimodular if  $|z| = 1$ . Suppose  $z_1$  and  $z_2$  are complex numbers such that  $\frac{z_1 - 2z_2}{2 - z_1z_2}$  is unimodular and  $z_2$  is not unimodular. Then the point  $z_1$  lies on a : (1) straight line parallel to x-axis (2) straight line parallel to y-axis (3) circle of radius 2 (4) circle of radius  $\sqrt{2}$

- A. circle of radius  $\sqrt{2}$
- B. straight line parallel to x-axis
- C. straight line parallel to y-axis
- D. circle of radius 2

**Answer: D**

68. Equation of the straight line which meets the circle  $x^2 + y^2 = 8$  at two points where these points are at a distance of 2 units from the point A(2, 2) is

A.  $x + y = 2$

B.  $x + y = 3$

C.  $x + y = 1$

D.  $x + y = 0$

**Answer: B**

 [Watch Video Solution](#)

69. If  $x_1, x_2, \dots, x_n$  are  $n$  observations such that  $\sum_{i=1}^n (x_i)^2 = 400$

and  $\sum_{i=1}^n x_i = 100$  then possible values of  $n$  among the following is

A. 18

B. 20

C. 24

D. 27

**Answer: D**



[Watch Video Solution](#)

70. If the system of equation  $x - 2y + 5z = 3$

$2x - y + z = 1$  and  $11x - 7y + pz = q$  has infinitely many solution,

then

A.  $p + q = 2$

B.  $p + q = 10$

C.  $p - q = 2$

D.  $p - q = 5$



**Answer: C**



**Watch Video Solution**

**71.** A term of randomly chosen from the expansion of  $\left(\sqrt[6]{4} + \frac{1}{\sqrt[4]{5}}\right)^{20}$ .

If the probability that it is a rational term is P, then  $420P$  is equal to



**Watch Video Solution**

**72.** If a tangent of slope 2 of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{1} = 1$  passes through the point  $(-2, 0)$ , then the value of  $a^2$  is equal to



**Watch Video Solution**

**73.** The number obtained after dividing the number formed by the last three digits of  $17^{256}$  by 100 is



 Watch Video Solution

74. The area (in sq. units) bounded by  $y = 2 - |x - 2|$  and the x-axis is

 Watch Video Solution

75. Let  $y = x^3 - 6x^2 + 9x + 1$  be an equation of a curve, then the x-intercept of the tangent to this curve whose slope is least, is

 Watch Video Solution

76. Two intersecting lines lying in plane  $P_1$  have equations  $\frac{x-1}{1} = \frac{y-3}{2} = \frac{z-4}{3}$  and  $\frac{x-1}{2} = \frac{y-3}{3} = \frac{z-4}{1}$ . If the equation of plane  $P_2$  is  $7x - 5y + z - 6 = 0$ , then the distance between planes  $P_1$  and  $P_2$  is

A.  $\frac{11}{5\sqrt{3}}$

B.  $\frac{2}{\sqrt{3}}$

C.  $\frac{1}{\sqrt{3}}$

D.  $\frac{7}{5\sqrt{3}}$

**Answer: B**



[Watch Video Solution](#)

77. If  $t$  is real and  $\lambda = \frac{t^2 - 3t + 4}{t^2 + 3t + 4}$  then find number of the solution of the systems of equation  $3x - y + 4z = 0$ ,  $x + 2y - 3z = -2.6x + 5y + \lambda z = -3$  for a particular value of  $\lambda$ .

A. a unique solution

B. infinite solutions

C. no solution

D. 2 solutions

**Answer: A**



**Watch Video Solution**

**78.** The solution of the differential equation  $2ydx + xdy = 2x\sqrt{y}dx$  is (where,  $C$  is an arbitrary constant)

A.  $x\sqrt{y} = x + C$

B.  $x\sqrt{y} = \frac{x^2}{2} + C$

C.  $\frac{x}{\sqrt{y}} = x + C$

D.  $xy = C$

**Answer: B**



**Watch Video Solution**

79. The mean and variance of 10 observations are found to be 10 and 4 respectively. On rechecking it was found that an observation 8 was incorrect. If it is replaced by 18, then the correct variance is

A. 7

B. 8

C. 9

D.  $\frac{55}{6}$

**Answer: C**

 [Watch Video Solution](#)

80. The sum of the series

$3 + 8 + 16 + 27 + 41 \dots$  upto 20 terms is equal to

A. 4230

B. 4430

C. 4330

D. 4500

**Answer: B**



[Watch Video Solution](#)

**81.** The greatest integer less than or equal to  $(\sqrt{2} + 1)^6$  is

A. 196

B. 197

C. 198

D. 199

**Answer: B**



[Watch Video Solution](#)

82. If  $\cos x - \sin x = -\frac{5}{4}$ , where  $\frac{\pi}{2} < x < \frac{3\pi}{4}$ , then  $\cot\left(\frac{x}{2}\right)$  is equal to

A.  $\frac{4 - \sqrt{7}}{9}$

B. 8

C. -8

D.  $\frac{4 + \sqrt{7}}{9}$

Answer: D

 Watch Video Solution

83. In  $\Delta PQR$ , the equation of the internal angle bisector of angle Q is  $y = x$  and the equation of side PR is  $3x - y = 2$ . If coordinates of P are  $(3, 2)$  and  $2PQ = RQ$ , then the coordinates of Q are

A.  $(3, 3)$

B. (7, 7)

C. ( - 2, - 2)

D. (5, 5)

**Answer: B**



[Watch Video Solution](#)

**84.** Let the lines  $l_1$  and  $l_2$  be normals to  $y^2 = 4x$  and tangents to  $x^2 = -12y$  (where  $l_1$  and  $l_2$  are not x - axis). The absolute value of the difference of slopes of  $l_1$  and  $l_2$  is

A. 3

B. 2

C. 1

D.  $\frac{1}{2}$



Answer: C

 Watch Video Solution

85. The value of  $\lim_{x \rightarrow \infty} \frac{(\ln x)^2}{2 + 3x^2}$  is equal to

A.  $\frac{1}{3}$

B.  $\frac{2}{3}$

C. 1

D. 0

Answer: D

 Watch Video Solution

86. The value of  $\lim_{n \rightarrow \infty} \sum_{r=1}^n \left( \frac{2r}{n^2} \right) e^{\frac{r^2}{n^2}}$  is equal to

A.  $e$

B.  $2e$

C.  $e - 2$

D.  $e - 1$

**Answer: D**



[Watch Video Solution](#)

**87.** The direction cosines  $l$ ,  $m$  and  $n$  of two lines are connected by the relations  $l + m + n = 0$  and  $lm = 0$ , then the angle between the lines is

A.  $\frac{\pi}{3}$

B.  $\frac{\pi}{4}$

C.  $\frac{\pi}{2}$

D.  $0$

**Answer: A**



**Watch Video Solution**

**88.** The function  $f(x) = x^3 - ax$  has a local minimum at  $x = k$ , where  $k \geq 2$ , then a possible value of  $a$  is

A. 9

B. 11

C. 13

D. 8

**Answer: C**



**Watch Video Solution**

89. Let two circles having radii  $r_1$  and  $r_2$  are orthogonal to each other. If the length of their common chord is  $k$  times the square root of harmonic mean between the squares of their radii, then  $k^4$  is equal to

A. 13

B. 7

C. 4

D. 2

**Answer: C**



[Watch Video Solution](#)

90. The value of  $\int_0^{\frac{\pi}{2}} (\cos 2x \cos 2^2 x \cos 2^3 x \cos 2^4 x) dx$  is equal to

A. 0

B.  $\frac{1}{2}$

C.  $\frac{\pi}{2}$

D.  $\frac{\pi}{4}$

**Answer: A**



**Watch Video Solution**

**91.** Let  $A$  be a matrix of order  $3 \times 3$  such that  $|A| = 3$ . Let

$B = 3A^{-1}$  and  $C = \frac{\text{adj}A}{2}$ , then the value of  $|A^2B^3C^4|$  is

A.  $\frac{3^{16}}{2^{12}}$

B.  $\left(\frac{3}{2}\right)^{12}$

C.  $\frac{3^{10}}{2^8}$

D.  $\frac{3^{12}}{2^{14}}$

**Answer: A**



[Watch Video Solution](#)

92.

Let

$$\left(\hat{p} \times \vec{q}\right) \times \hat{p} + \left(\hat{p} \cdot \vec{q}\right) \vec{q} = (x^2 + y^2) \vec{q} + (14 - 4x - 6y) \hat{p}$$

where  $\hat{p}$  and  $\vec{q}$  are non - collinear vectors ( $\hat{p}$  is a unit vector) and  $x, y$  are scalars, then the value of  $x^2 + y^2$  is equal to

A. 10

B. 11

C. 12

D. 13

**Answer: D**[Watch Video Solution](#)

93. If  $p$  and  $q$  are two statements, then which of the following statement is a tautology

A.  $p \Rightarrow (p \vee \sim q)$

B.  $(p \vee q) \Rightarrow p$

C.  $p \Rightarrow (p \wedge q)$

D.  $p \Leftrightarrow (p \Rightarrow q)$

**Answer: A**

 [Watch Video Solution](#)

94. In an equilateral triangle  $ABC$ , equation of the sides  $BC$  is  $x + y - 2 = 0$  and the centroid of  $\Delta ABC$  is  $(0, 0)$ . If points  $A, B$  and  $C$  are in anticlockwise order, then the equation of side  $AC$  is

A.  $(y + 2) = (2 - \sqrt{3})(x + 2)$

B.  $(y + 2) = (2 + \sqrt{3})(x + 2)$

C.  $(y + 1) = (2 + \sqrt{3})(x + 1)$

D.  $x + 2 = 0$

**Answer: B**

 [Watch Video Solution](#)

95. The minimum distance between the curves

$y = \tan x, \forall x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  and  $\left(x - 2 - \frac{\pi}{4}\right)^2 + y^2 = 1$  is

A.  $\sqrt{2} - 1$

B.  $\sqrt{5} - 1$

C.  $\sqrt{5} + 1$

D. 2

**Answer: B**

 [Watch Video Solution](#)



 Watch Video Solution

96. A fair die is thrown  $n$  number of times. If the probability of always getting a number greater than the previous number is  $\frac{5}{54}$ , then the value of  $n$  is equal to ( $n \leq 6$ ).

 Watch Video Solution

97. How many 4 letter words can be formed from the word "MATHEMATICS" ?

A. 2500

B. 2454

C. 2400

D. 2254

**Answer: B**

 Watch Video Solution

98. If  $f(x) = \begin{cases} (1 + |\sin x|)^{\frac{p}{|\sin x|}}, & -\frac{\pi}{6} < x < 0 \\ q & : x = 0 \\ e^{\tan 3x \cdot \cot 5x} & : 0 < x < \frac{\pi}{6} \end{cases}$  is continuous at

$x = 0$ , then the value of  $2p + 10 \ln q$  is equal to

 Watch Video Solution

99. If  $f(x) = \sin x$ ,  $g(x) = \cos x$  and  $h(x) = \cos(\cos x)$ , then the integral  $I = \int f(g(x)) \cdot f(x) \cdot h(x) dx$  simplifies to  $-\lambda \sin^2(\cos x) + C$  (where,  $C$  is the constant of integration). The value of  $\lambda$  is equal to

 Watch Video Solution

100. If numerically greatest term in the expansion of  $(3 - 5x)^{11}$ , where  $x = \frac{1}{5}$ , is  $729\lambda$ , then the value of  $\frac{\lambda}{150}$  is

 [Watch Video Solution](#)

**101.** If a tower subtends equal angles at four points P, Q, R and S that lie in a plane containing the foot of the tower, then which of the following statements is always true (here, the tower is perpendicular to the plane containing the points P, Q, R, S)

A.  $\angle PQS = \angle PRS$

B.  $\angle PQR + \angle PSR = 180^\circ$

C.  $\angle PQS = 90^\circ \Rightarrow \angle PRS = 90^\circ$

D.  $(PQ)(RS) + (PS)(RQ) = (PR)(QS)$

**Answer: C**

 [Watch Video Solution](#)

102. The values of  $\lambda$  for which one root of the equation  $x^2 + (1 - 2\lambda)x + (\lambda^2 - \lambda - 2) = 0$  is greater than 3 and the other smaller than 2 are given by

A.  $2 < \lambda < 5$

B.  $1 < \lambda < 4$

C.  $1 < \lambda < 5$

D.  $2 < \lambda < 4$

Answer: D

 Watch Video Solution

103. Let a function  $f: (2, \infty) \rightarrow [0, \infty)$  defined as  $f(x) = \frac{|x - 3|}{|x - 2|}$ , then  $f$  is

A. injective & surjective

B. not injective but surjective

C. injective but not surjective

D. neither injective nor surjective

**Answer: B**



[Watch Video Solution](#)

**104.** Let  $n$  be a positive integer and  $z$  a complex number with unit modulus is a solution of the equation  $z^n + z + 1 = 0$ , then the value of  $n$  can be

A. 87

B. 97

C. 104

D. 222

Answer: C



Watch Video Solution

105. The value of  $\lim_{x \rightarrow 0} \frac{e^{-\left(\frac{x^2}{2}\right)} - \cos x}{x^3 \tan x}$  is equal to

A.  $\frac{1}{4}$

B.  $\frac{1}{8}$

C.  $\frac{1}{12}$

D.  $\frac{1}{16}$

Answer: C



Watch Video Solution

106. The value of  $\int \frac{(x-4)}{x^2 \sqrt{x-2}} dx$  is equal to (where , C is the constant of integration )

A.  $2x\sqrt{x-2} + C$

B.  $-\frac{2}{x}\sqrt{x-2} + C$

C.  $\frac{\sqrt{x-2}}{x} + C$

D.  $\frac{x}{\sqrt{x-2}} + C$

**Answer: B**



**Watch Video Solution**

**107.** The equation of the curve passing through the point (1,1) and satisfying the differential equation  $\frac{dy}{dx} = \frac{x + 2y - 3}{y - 2x + 1}$  is

A.  $x^2 - 4xy - y^2 + 6x + 2y - 4 = 0$

B.  $x^2 + 4xy - y^2 - 6x + 2y + 4 = 0$

C.  $x^2 + 4xy - y^2 - 6x - 2y + 4 = 0$

D.  $x^2 + 4xy + y^2 - 6x - 2y - 4 = 0$

**Answer: C**

 [Watch Video Solution](#)

**108.** Five different games are to be distributed among 4 children randomly. The probability that each child get at least one game is

A.  $\frac{1}{4}$

B.  $\frac{15}{64}$

C.  $\frac{21}{64}$

D.  $\frac{17}{632}$

**Answer: B**

 [Watch Video Solution](#)



109. Let the focus S of the parabola  $y^2 = 8x$  lies on the focal chord PQ of the same parabola . If PS = 6 , then the square of the slope of the chord PQ is

A.  $\frac{2}{\sqrt{5}}$

B.  $\frac{4}{5}$

C.  $\frac{5}{4}$

D.  $\frac{9}{4}$

**Answer: B**



**Watch Video Solution**

110. If  $p \rightarrow (q \vee r)$  is false, then the truth values of p,q,r are respectively

A. TFF

B. FFF

C. FTT

D. TTF

**Answer: A**

 [Watch Video Solution](#)

111.  $\frac{5}{3^2 7^2} + \frac{9}{7^2 11^2} + \frac{13}{11^2 15^2} + \dots \infty$

A.  $\frac{1}{8}$

B.  $\frac{1}{36}$

C.  $\frac{1}{54}$

D.  $\frac{1}{72}$

**Answer: D**

 [Watch Video Solution](#)

112. If  $13^{99} - 19^{93}$  is divided by 162, then the remainder is

A. 3

B. 6

C. 5

D. 0

**Answer: D**



**Watch Video Solution**

113. The  $\int_0^{\pi/2} \operatorname{sgn}\left(\sin^2 x - \sin x + \frac{1}{2}\right) dx$  is equal to , (where ,  $\operatorname{sgn}$

(x) denotes the sigum function of x)

A. 0

B. 1

C.  $\pi$

D.  $\frac{\pi}{2}$

**Answer: D**

 [Watch Video Solution](#)

114. If  $\vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$ ,  $\vec{a} \cdot \vec{b} = 2$  and  $\vec{a} \times \vec{b} = 2\hat{i} - \hat{k}$ , then  $\vec{b}$  is

A.  $(\hat{i} - 2\hat{j} + \hat{k})$

B.  $(4\hat{i} - 4\hat{j} + 2\hat{k})$

C.  $\frac{1}{2}(3\hat{i} + 7\hat{j} + 9\hat{k})$

D.  $\frac{1}{29}(7\hat{i} - 4\hat{j} + 14\hat{k})$

**Answer: D**

 [Watch Video Solution](#)

115. Equation of the plane passing through the point of intersection of

lines  $\frac{x-1}{3} = \frac{y-2}{1} = \frac{z-3}{2}$  &  $\frac{x-3}{1} = \frac{y-1}{2} = \frac{z-2}{3}$  and

perpendicular to the line  $\frac{x+5}{2} = \frac{y-3}{3} = \frac{z+1}{1}$  is

- A.  $2x + 3y + z + 7 = 0$
- B.  $2x - 3y - z + 22 = 0$
- C.  $2x + 3y + z - 22 = 0$
- D.  $2x + 3y + z + 13 = 0$

**Answer: C**

 [Watch Video Solution](#)

116. The equation of the tangent to the parabola  $y^2 = 4x$  whose slope is positive and which also touches  $x^2 + y^2 = \frac{1}{2}$  is

- A.  $y = x + 1$

B.  $y = 2x + 1$

C.  $x + y = 2$

D.  $y = 4x + \frac{1}{2}$

**Answer: A**



**Watch Video Solution**

**117.** If  $A$  is  $2 \times 2$  matrix such that  $A \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$  and  $A^2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ , then trace of  $A$  is (where the trace of the matrix is the sum of all principal diagonal elements of the matrix )

A. 1

B. 0

C. 2

D. 5

**Answer: A**

 [Watch Video Solution](#)

**118.** consider the planes  $P_1: 2x - y + z = 6$  and  $P_2: x + 2y - z = 4$  having normal  $\vec{N}_1$  and  $\vec{N}_2$  respectively . The distance of the origin from the plane passing through the point (1,1,1) and whose normal is perpendicular to  $N_1$  and  $N_2$  is

- A.  $\frac{7}{\sqrt{5}}$  units
- B.  $\sqrt{\frac{7}{5}}$  units
- C.  $\sqrt{\frac{3}{5}}$  units
- D.  $\frac{14}{\sqrt{35}}$  units

**Answer: B**

 [Watch Video Solution](#)

119. Let  $I_1 = \int_0^{\frac{\pi}{2}} \frac{dt}{1+t^6}$  and  $I_2 = \int_0^{\frac{\pi}{2}} \frac{x \cos x dx}{1+(x \sin x + \cos x)^6}$ , then

A.  $2I_1 = I_2$

B.  $I_1 = 2I_2$

C.  $I_1 = I_2$

D.  $I_1 = I_2 = 0$

**Answer: C**

 [Watch Video Solution](#)

120. A wire of length 28 cm is bent to form a circular sector, then the radius (in cm) of the circular sector such that the area of the circular sector is maximum is equal to

A. 5

B. 6



C. 7

D. 8

**Answer: C**

 [Watch Video Solution](#)

**121.** Let  $x^2 + y^2 = r^2$  and  $xy = 1$  intersect at  $A$  &  $B$  in first quadrant, If  $AB = \sqrt{14}$  then find the value of  $r$ .

 [Watch Video Solution](#)

**122.** If  $f(x) = \left\{ \frac{a + b \cos x + c \sin x}{x^2}, x > 0 \right\}, (9, x \geq 0)$  is continuous at  $x = 0$ , then the value of  $\frac{|a| + |b|}{5}$  is

 [Watch Video Solution](#)

**123.** Let  $p$  and  $q$  be the length of two chords of a circle which subtend angles  $36^\circ$  and  $60^\circ$  respectively at the centre of the circle. Then, the angle (in radian) subtended by the chord of length  $p + q$  at the centre of the circle is (use  $\pi = 3.1$ )

 [Watch Video Solution](#)

**124.** Let

$$a_r = r^4 C_r, b_r = (4 - r)^4 C_r, A_r = \begin{bmatrix} a_r & 2 \\ 3 & b_r \end{bmatrix} \text{ and } A = \sum_{r=0}^4 A_r \text{ then}$$

the value of  $|A|$  is equal to

 [Watch Video Solution](#)

**125.** The product of all the values of  $|\lambda|$ , such that the lines

$x + 2y - 3 = 0$ ,  $3x - y - 1 = 0$  and  $\lambda x + y - 2 = 0$  cannot form a triangle, is equal to

 [Watch Video Solution](#)

## Maths

1. The integral value of  $m$  for which the quadratic equation  $(2m - 3)x^2 - 4x + 2m - 3 = 0$  has both the roots negative is given by

[Watch Video Solution](#)

2. Let from a point  $A(h,k)$  chord of contacts are drawn to the ellipse  $x^2 + 2y^2 = 6$  such that all these chords touch the ellipse  $x^2 + 4y^2 = 4$ , then locus of the point  $A$  is

A.  $4x^2 + 9y^2 = 36$

B.  $x^2 + y^2 = 4$

C.  $x^2 - y^2 = 9$

D.  $x^2 + y^2 = 9$

Answer: D

 Watch Video Solution

3. If  $y(x)$  is the solution of the differential equation

$$\frac{dy}{dx} = -2x(y-1) \text{ with } y(0) = 1, \text{ then } \lim_{x \rightarrow \infty} y(x) \text{ equals}$$

 Watch Video Solution

4. 
$$\int \frac{\sin^2 x \cdot \sec^2 x + 2 \tan x \cdot \sin^{-1} x \cdot \sqrt{1-x^2}}{\sqrt{1-x^2}(1+\tan^2 x)} dx$$

A.  $(\sin^{-1} x)(\cos^2 x) + C$

B.  $(\sin^{-1} x)(\sin^2 x) + C$

C.  $(\cos^{-1} x)(\sin^2 x) + C$

D.  $-\sin^{-1} x(\sin^2 x) + C$

**Answer: B**



**Watch Video Solution**

5. The value of  $\lim_{x \rightarrow 0} \frac{x \cot(4x)}{\tan^2(3x)\cot^2(6x)}$  is equal to

A. 0

B. 4

C.  $\frac{2}{9}$

D. 1

**Answer: D**



**Watch Video Solution**

6. If  $n$  objects are arranged in a row, then the number of ways of selecting three of these objects so that no two of them are next to

each other is

A.  ${}^{n-3}C_3$

B.  ${}^{n-3}C_2$

C.  ${}^{n-2}C_2$

D.  ${}^{n-2}C_3$

**Answer: D**



**Watch Video Solution**

7. Solve  $\sin^{-1}(1 - x) - 2s \in^{-1} x = \frac{\pi}{2}$

A. 0

B.  $\frac{1}{2}$

C.  $0, \frac{1}{2}$

D.  $-\frac{1}{2}$

**Answer: A**



**Watch Video Solution**

8. If 1 , a, b and 4 are in harmonic progression , then the value of a + b is equal to

A.  $\frac{5}{4}$

B.  $\frac{10}{3}$

C.  $\frac{3}{10}$

D.  $\frac{4}{5}$

**Answer: B**



**Watch Video Solution**

9. fractional part of  $\frac{2^{78}}{31}$  is:

A.  $\frac{2}{31}$

B.  $\frac{4}{31}$

C.  $\frac{6}{31}$

D.  $\frac{8}{31}$

**Answer: D**



**Watch Video Solution**

**10.** Let  $f(x) = 10 - |x-5|$ ,  $x \in \mathbb{R}$ , then the set of all values of  $x$  at which  $f(x)$  is not differentiable is

A.  $\{0,5,10\}$

B.  $\{5,10\}$

C.  $\{0,5,10,15\}$

D.  $\{5,10,15\}$



**Answer: A**



**Watch Video Solution**

11. If two tangents drawn from the point P (h,k) to the parabola  $y^2 = 8x$  are such that the slope of one of the tangent is 3 times the slope of the other, then the locus of point P is

A.  $3y^2 = 16x$

B.  $3y^2 = 8x$

C.  $y^2 = 32x$

D.  $3y^2 = 32x$

**Answer: D**



**Watch Video Solution**

12. If  $I_1 = \int_{1-x}^k x \sin\{x(1-x)\}dx$  and  $I_2 = \int_{1-x}^k \sin\{x(1-x)\}dx$ ,

then

A. 2

B.  $\frac{1}{2}$

C. 1

D.  $\frac{1}{3}$

**Answer: B**

 [Watch Video Solution](#)

13. Let  $A$  is a matrix of order  $3 \times 3$  defined as  $A = [a_{ij}]_{3 \times 3}$ , where

$$a_{ij} = \lim_{x \rightarrow 0} \frac{1 - \cos(ix)}{\sin(ix)\tan(jx)} \quad (\forall 1 \leq i, j, \leq 3), \text{ then } A^2 \text{ is equal}$$

to

A.  $A$

B.  $\frac{3}{2}A$

C.  $\frac{2}{3}A$

D.  $\frac{1}{4}A$

**Answer: B**

 [Watch Video Solution](#)

14. If

$$\left[ \left( \vec{a} + 2\vec{b} + 3\vec{c} \right) \times \left( \vec{b} + 2\vec{c} + 3\vec{a} \right) \right] \cdot \left( \vec{c} + 2\vec{a} + 3\vec{b} \right) = 54$$

where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are 3 non - coplanar vectors, then the values of

$$\begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{c} \cdot \vec{a} & \vec{c} \cdot \vec{b} & \vec{c} \cdot \vec{c} \end{vmatrix} \text{ is equal to}$$

A. 9

B. 3

C. 6

D. 12

**Answer: A**

 [Watch Video Solution](#)

15. Let A be the point (1,2,3) and B be a point on the line  $\frac{x-1}{-2} = \frac{y+1}{3} = \frac{z-5}{4} = k$ . Then value of k such that line AB is perpendicular to the plane  $4x + 9y - 18z = 6$  is

A.  $-\frac{2}{5}$

B.  $\frac{1}{5}$

C.  $\frac{2}{5}$

D. no such value of k is possible

**Answer: C**

 [Watch Video Solution](#)

16. Let the circumcentre of  $\Delta ABC$  is S (-1,0) and the midpoints of sides AB and AC are E(1,-2) and F(-2,1) respectively , then the equation of the circumcircle of  $\Delta ABC$  is

A.  $(x + 1)^2 + y^2 = 5$

B.  $(x + 1)^2 + y^2 = 10$

C.  $(x + 1)^2 + y^2 = 15$

D.  $(x + 1)^2 + y^2 = 1$

**Answer: B**

 [Watch Video Solution](#)

17. If p and q are two statements , then which of the following statements is not equivalent to  $p \Leftrightarrow (p \Rightarrow q)$  ?

A.  $p \wedge q$

B.  $(p \Leftrightarrow q) \wedge (p \vee q)$

C.  $(p \Rightarrow q) \Leftrightarrow q$

D.  $(\neg p \Rightarrow q) \wedge (p \vee \neg q)$

**Answer: D**

 [Watch Video Solution](#)

18. Let  $F(n) = (\sin 1) \times (\sin 2) \times \dots \times \sin(n)$ ,  $\forall n \in \mathbb{N}$  then number of elements in the set  $A = \{f(1), f(2), \dots, f(6)\}$  that are positive are

 [Watch Video Solution](#)

19.  $a, b, c, \in \mathbb{N}$  and  $d = \begin{vmatrix} a & b & c \\ c & a & b \\ b & c & a \end{vmatrix}$ , then the least positive value of

D is

A. 4

B. 6

C. 3

D. 8

**Answer: A**



**Watch Video Solution**

20.  $f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = \lambda x + \sin x$  is onto if  $\lambda$  is an element of the set  $P$  and  $f(x)$  is one- one if  $\lambda$  is an element of the set  $Q$ , then (given ,  $\lambda$  is a real number )

A.  $P = Q$

B.  $P \subset Q$

C.  $P - Q = \{0\}$

D.  $Q \subset P$

**Answer: D**

 [Watch Video Solution](#)

**21.** Consider circles  $C_1$  &  $C_2$  touching both the axes and passing through (4,4), then the product of radii of these circles is

 [Watch Video Solution](#)

**22.** If  $P(z)$  is a variable point in the complex plane such that  $\text{IM}\left(-\frac{1}{z}\right) = \frac{1}{4}$ , then the value of the perimeter of the locus of  $P(z)$  is  
(use  $\pi = 3.14$ )

 [Watch Video Solution](#)

**23.** The probability of India winning a test match against Australia is  $\frac{1}{4}$ . Assuming the matches to be independent events, the probability



that in a 7 match series India's second win occurs at 4<sup>th</sup> test is P , then

256 P is equal to

A. 15

B. 12

C. 27

D. 40

**Answer: C**

 [Watch Video Solution](#)

**24.** The number of solutions of the equation  $|\cot x| = \cot x + \operatorname{cosec}x$  in  $[0, 10\pi]$  is /are

 [Watch Video Solution](#)

25. If  $\alpha$  is the only real root of  $x^3 + bx^2 + cx + 1 = 0$  ( $b < c$ ), then the value of  $|\lceil \alpha \rceil|$  is (where,  $\lceil \cdot \rceil$  represents the greatest integer function)



[Watch Video Solution](#)