



MATHS

BOOKS - NTA MOCK TESTS

JEE MOCK TEST 20

Mathematics

1. If the integral

$$I = \int x^{\sin x} \left(\cos x \cdot \ln x + \frac{\sin x}{x} \right) dx, = (f(x))^{g(x)} + c (\forall x > 0)$$

then the range of $y = g(x)$ is (where, c is an arbitrary constant)

A. [-1,1]

B. [0,1]

C. [0,1)

D. (-1,1)

Answer: A



Watch Video Solution

2. Let P and Q are two points in the xy plane on the curve

$y = x^{11} - 2x^7 + 7x^3 + 11x + 6$ such that

$\overrightarrow{OP} \cdot \hat{i} = 5, \overrightarrow{OQ} \cdot \hat{i} = -5$, then the magnitude of $\overrightarrow{OP} + \overrightarrow{OQ}$

is

A. 10

B. 12

C. 14

D. 8

Answer: B



Watch Video Solution

3. If the letters of the word REGULATIONS be arranged at random, find the probability that there will be exactly four letters between the R and the E .

A. $\frac{6}{55}$

B. $\frac{3}{55}$

C. $\frac{5}{11}$

D. $\frac{6}{11}$

Answer: A



Watch Video Solution

4. If the point of intersection of the plane $4x - 5y + 2z - 6 = 0$ with the line through the origin and perpendicular to the plane $x - 2y - 4z = 4$ is P, then the distance of the point P from (1, 2, 3) is

A. $\sqrt{63}$ units

B. 8 units

C. $\sqrt{65}$ units

D. $\sqrt{72}$ units

Answer: C



[Watch Video Solution](#)

5. The mean and variance of seven observations are 8 and 16 respectively. If five of these are 2, 4, 10, 12 and 14, then find the

remaining two observations.

A. 5,7

B. 3,5

C. 6,8

D. 4,2

Answer: C



Watch Video Solution

6. Let $\triangle ABC$ is an isosceles triangle with $AB = AC$. If $B = (0, a)$, $C = (2a, 0)$ and the equation of AB is $3x - 4y + 4a = 0$, then the equation of side AC is

A. $y = 8x - 16a$

B. $3y = 4x - 8a$

C. $x = 2a$

D. $y + 8x = 16a$

Answer: C



Watch Video Solution

7. Let $A(0,3)$ and $B(0,12)$ be two vertices of a $\triangle ABC$ where $C = (x, 0)$. If the circumcircle of $\triangle ABC$ touches the x-axis, then the value of $\cos 2\theta$ is (where θ is angle ACB)

A. $\frac{1}{3}$

B. $\frac{1}{2}$

C. $\frac{8}{15}$

D. $\frac{7}{25}$

Answer: D



Watch Video Solution

8. Consider three statements p : person 'A' passed in mathematics exam q : Person 'A' passed in physics exam r : Person 'A' passed in chemistry exam, Then the statement $\sim((\sim(p \Rightarrow q) \Rightarrow r))$ is equivalent to

- A. Person A passed only in mathematics & physics & chemistry
- B. Person A failed only in mathematics & physics & chemistry
- C. Person A passed in all the three subjects mathematics & physics & chemistry

D. Person A passed in chemistry but failed in mathematics & physics.

Answer: A



Watch Video Solution

9. In equation $(Z - 1)^n = Z^n = 1 (\forall n \in N)$ has solution, then n can be

A. 4

B. 12

C. 15

D. 21

Answer: B

10. The solution of the differential equation

$$\left(3x^2 \sin\left(\frac{1}{x}\right) + y\right)dx = x \cos\left(\frac{1}{x}\right)dx - xdy$$

is (where, c is an arbitrary constant)

A. $\sin\left(\frac{1}{x}\right) = xy + c$

B. $x^3 \sin\left(\frac{1}{x}\right) + xy = c$

C. $x^3 \sin\left(\frac{1}{x}\right) = xy + c$

D. $\sin\left(\frac{1}{x}\right) = x^3y + c$

Answer: B

11. The quadratic equation $(1 - \sin \theta)x^2 + 2(1 - \sin \theta)x - 3 \sin \theta = 0$ has both roots complex for all θ lying in the interval

A. $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

B. $\left(0, \frac{3\pi}{2}\right)$

C. $\left(\frac{\pi}{6}, \frac{7\pi}{6}\right)$

D. $\left(\frac{7\pi}{6}, \frac{11\pi}{6}\right)$

Answer: D



Watch Video Solution

12. The minimum value of the expression $3x + 2y$ ($\forall x, y > 0$), where $xy^2 = 10$, occurs when the value of y is equal to

A. $\sqrt{10}$

B. $\sqrt[3]{10}$

C. $\sqrt[3]{30}$

D. $\frac{1}{\sqrt{30}}$

Answer: C



Watch Video Solution

13. If in the expansion of $(1 + x)^m(1 - x)^n$ the coefficient of x and x^2 are 3 and (-6) respectively, then the value of n is-

A. 6

B. 9

C. 12

D. 24

Answer: B



Watch Video Solution

14. Number of words that can be formed with the letters of the word ALGEBRA so that all the vowels are separated (or no two vowels come together) is

A. 720

B. 2160

C. 1440

D. 1200

Answer: A



Watch Video Solution

15. If $f(k - x) + f(x) = \sin x$, then the value of integral

$I = \int_0^k f(x) dx$ is equal to

A. $\cos k$

B. $2 \cos^2\left(\frac{k}{2}\right)$

C. $\sin^2\left(\frac{k}{2}\right)$

D. $\sin k$

Answer: C



Watch Video Solution

16. If the difference between the number of subsets of the sets A and B is 120, then choose the incorrect option.

A. Maximum value of $n(A \cap B) = 3$

B. Minimum value of $n(A \cap B) = 0$

C. Maximum value of $n(A \cup B) = 21$

D. Minimum value of $n(A \cup B) = 7$

Answer: C



Watch Video Solution

17. For the function $f(x) = \sin(\pi[x]) \times \cos^{-1}([x])$, choose the correct option.

(where $[.]$ represents the greatest integer function)

A. Domain of $f(x)$ in $[-1,1]$

B. Range of $f(x)$ contains exactly two elements

C. $f(x)$ is an identity function

D. $f(x)$ is a constant function

Answer: D



Watch Video Solution

18. The value of $\lim_{x \rightarrow 0} \frac{\sin x}{3} \left[\frac{5}{x} \right]$ is equal to

[where $[\cdot]$ represent the greatest integer function)

A. $\frac{1}{3}$

B. 0

C. $\frac{5}{3}$

D. 1

Answer: C



Watch Video Solution

19. If $y = \tan^{-1}\left(\frac{x}{1+6x^2}\right) + \tan^{-1}\left(\frac{2x-1}{2x+1}\right)$, ($\forall x > 0$)
then $\frac{dy}{dx}$ is equal to

A. $\frac{3}{1+9x^2}$

B. $\frac{1}{1+6x^2}$

C. $\frac{1}{1+6x^2} + \frac{1}{1+x^2}$

D. $\frac{3}{1+6x^2}$

Answer: A

 [Watch Video Solution](#)

20. If $|3x-1|, 3, |x-3|$ are the first three terms of an arithmetic progression, then the sum of the first five terms can be

A. 5

B. 10

C. 20

D. 30

Answer: A



Watch Video Solution

21. If $f(x) = \begin{cases} px + q & : x \leq 2 \\ x^2 - 5x + 6 & : 2 < x < 3 \\ ax^2 + bx + 1 & : x \geq 3 \end{cases}$

is differentiable everywhere, then $|p| + |q| + \left| \frac{1}{a} \right| + \left| \frac{1}{b} \right|$ is equal to



Watch Video Solution

22. If the area bounded by $y = \left| |x|^2 - 4|x| + 3 \right|$ and the x-axis from $x = 1$ to $x = 3$ is $\frac{p}{q}$ (where, p & q are coprime) then the value of $p + q$ is

A. 10

B. 9

C. 8

D. 7

Answer: D



Watch Video Solution

23. Let M be a square matrix of order 3 whose elements are real

number and $adj(adjM) = \begin{bmatrix} 36 & 0 & -4 \\ 0 & 6 & 0 \\ 0 & 3 & 6 \end{bmatrix}$, then the absolute

value of $Tr(M)$ is [Here, adj P denotes adjoint matrix of P and $Tr(P)$ denotes trace of matrix P i.e., sum of all principal diagonal elements of matrix P]



[Watch Video Solution](#)

24. If common tangents of $x^2 + y^2 = r^2$ and $\frac{x^2}{16} + \frac{y^2}{9} = 1$ forms a square, then the length of diagonal of the square is

- A. 9
- B. 10
- C. 12
- D. 1

Answer: B



[Watch Video Solution](#)

25. The angular depression of the top and the foot of the chimney seen from the top of a tower on the same base level as the chimney are $\tan^{-1}\left(\frac{4}{3}\right)$ and $\tan^{-1}\left(\frac{5}{2}\right)$ respectively if the height of the tower is 150m. then the distance between the top of the chimney and the tower is



Watch Video Solution