



## MATHS

### BOOKS - NTA MOCK TESTS

#### JEE MOCK TEST 24

#### Maths

1. The integral value of  $m$  for which the quadratic equation  $(2m - 3)x^2 - 4x + 2m - 3 = 0$  has both the roots negative is given by

 [Watch Video Solution](#)

2. Let from a point  $A(h, k)$  chord of contacts are drawn to the ellipse  $x^2 + 2y^2 = 6$  such that all these chords touch the ellipse

$x^2 + 4y^2 = 4$ , then locus of the point A is

A.  $4x^2 + 9y^2 = 36$

B.  $x^2 + y^2 = 4$

C.  $x^2 - y^2 = 9$

D.  $x^2 + y^2 = 9$

**Answer: D**

 [Watch Video Solution](#)

3. If  $y(x)$  is the solution of the differential equation

$$\frac{dy}{dx} = -2x(y - 1) \text{ with } y(0) = 1, \text{ then } \lim_{x \rightarrow \infty} y(x) \text{ equals}$$

 [Watch Video Solution](#)

4. 
$$\int \frac{\sin^2 x \cdot \sec^2 x + 2 \tan x \cdot \sin^{-1} x \cdot \sqrt{1 - x^2}}{\sqrt{1 - x^2}(1 + \tan^2 x)} dx$$

A.  $(\sin^{-1} x)(\cos^2 x) + C$

B.  $(\sin^{-1} x)(\sin^2 x) + C$

C.  $(\cos^{-1} x)(\sin^2 x) + C$

D.  $-\sin^{-1} x(\sin^2 x) + C$

**Answer: B**



**Watch Video Solution**

5. The value of  $\lim_{x \rightarrow 0} \frac{x \cot(4x)}{\tan^2(3x)\cot^2(6x)}$  is equal to

A. 0

B. 4

C.  $\frac{2}{9}$

D. 1

**Answer: D**

 Watch Video Solution

6. If  $n$  objects are arranged in a row, then the number of ways of selecting three of these objects so that no two of them are next to each other is

A.  ${}^{n-3}C_3$

B.  ${}^{n-3}C_2$

C.  ${}^{n-2}C_2$

D.  ${}^{n-2}C_3$

**Answer: D**

 Watch Video Solution

7. Solve  $\sin^{-1}(1-x) - 2s \in^{-1} x = \frac{\pi}{2}$

A. 0

B.  $\frac{1}{2}$

C.  $0, \frac{1}{2}$

D.  $-\frac{1}{2}$

**Answer: A**



**Watch Video Solution**

**8.** If 1 , a, b and 4 are in harmonic progression , then the value of a + b is equal to

A.  $\frac{5}{4}$

B.  $\frac{10}{3}$

C.  $\frac{3}{10}$

D.  $\frac{4}{5}$

**Answer: B**

 [Watch Video Solution](#)

9. fractional part of  $\frac{2^{78}}{31}$  is:

A.  $\frac{2}{31}$

B.  $\frac{4}{31}$

C.  $\frac{6}{31}$

D.  $\frac{8}{31}$

**Answer: D**

 [Watch Video Solution](#)

10. Let  $f(x) = 10 - |x-5|$ ,  $x \in \mathbb{R}$ , then the set of all values of  $x$  at which  $f(x)$  is not differentiable is

A.  $\{0,5,10\}$

B.  $\{5,10\}$

C.  $\{0,5,10,15\}$

D.  $\{5,10,15\}$

**Answer: A**



**Watch Video Solution**

11. If two tangents drawn from the point P (h,k) to the parabola  $y^2 = 8x$  are such that the slope of one of the tangent is 3 times the slope of the other , then the locus of point P is

A.  $3y^2 = 16x$

B.  $3y^2 = 8x$

C.  $y^2 = 32x$

D.  $3y^2 = 32x$

**Answer: D**

 [Watch Video Solution](#)

12. If  $I_1 = \int_{1-x}^k x \sin\{x(1-x)\} dx$  and  $I_2 = \int_{1-x}^k \sin\{x(1-x)\} dx$ ,

then

A. 2

B.  $\frac{1}{2}$

C. 1

D.  $\frac{1}{3}$

**Answer: B**

 [Watch Video Solution](#)



13. Let  $A$  is a matrix of order  $3 \times 3$  defined as  $A = [a_{ij}]_{3 \times 3}$ , where

$$a_{ij} = \lim_{x \rightarrow 0} \frac{1 - \cos(ix)}{\sin(ix)\tan(jx)} (\forall 1 \leq i, j, \leq 3), \text{ then } A^2 \text{ is equal}$$

to

A.  $A$

B.  $\frac{3}{2}A$

C.  $\frac{2}{3}A$

D.  $\frac{1}{4}A$

**Answer: B**

 [Watch Video Solution](#)

14.

If

$$\left[ \left( \vec{a} + 2\vec{b} + 3\vec{c} \right) \times \left( \vec{b} + 2\vec{c} + 3\vec{a} \right) \right] \cdot \left( \vec{c} + 2\vec{a} + 3\vec{b} \right) = 54$$

where  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are 3 non - coplanar vectors, then the values of

$$\begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} & \vec{a} \cdot \vec{c} \\ \vec{b} \cdot \vec{a} & \vec{b} \cdot \vec{b} & \vec{b} \cdot \vec{c} \\ \vec{c} \cdot \vec{a} & \vec{c} \cdot \vec{b} & \vec{c} \cdot \vec{c} \end{vmatrix} \text{ is equal to}$$

A. 9

B. 3

C. 6

D. 12

**Answer: A**

 [Watch Video Solution](#)

15. Let A be the point (1,2,3) and B be a point on the line

$$\frac{x-1}{-2} = \frac{y+1}{3} = \frac{z-5}{4} = k \text{ Then value of } k \text{ such that line AB is}$$

perpendicular to the plane  $4x + 9y - 18z = 6$  is

A.  $-\frac{2}{5}$

B.  $\frac{1}{5}$

C.  $\frac{2}{5}$

D. no such value of  $k$  is possible

**Answer: C**



**Watch Video Solution**

**16.** Let the circumcentre of  $\triangle ABC$  is  $S(-1,0)$  and the midpoints of sides  $AB$  and  $AC$  are  $E(1,-2)$  and  $F(-2,1)$  respectively, then the equation of the circumcircle of  $\triangle ABC$  is

A.  $(x + 1)^2 + y^2 = 5$

B.  $(x + 1)^2 + y^2 = 10$

C.  $(x + 1)^2 + y^2 = 15$

D.  $(x + 1)^2 + y^2 = 1$

**Answer: B**



**Watch Video Solution**

17. If  $p$  and  $q$  are two statements , then which of the following statements is not equivalent to  $p \Leftrightarrow (p \Rightarrow q)$  ?

A.  $p \wedge q$

B.  $(p \Leftrightarrow q) \wedge (p \vee q)$

C.  $(p \Rightarrow q) \Leftrightarrow q$

D.  $(\neg p \Rightarrow q) \wedge (p \vee \neg q)$

**Answer: D**



**Watch Video Solution**

18. Let  $F(n) = (\sin 1) \times (\sin 2) \times \dots \times \sin(n)$ ,  $\forall n \in \mathbb{N}$  then number of elements in the set  $A = \{f(1), f(2), \dots, f(6)\}$  that are positive are

 [Watch Video Solution](#)

19.  $a, b, c, \in N$  and  $d = \begin{vmatrix} a & b & c \\ c & a & b \\ b & c & a \end{vmatrix}$ , then the least positive value of

D is

A. 4

B. 6

C. 3

D. 8

**Answer: A**

 [Watch Video Solution](#)

20.  $F: \mathbb{R} \rightarrow \mathbb{R}$ ,  $F(x) = \lambda x + \sin x$  is onto if  $\lambda$  is an element of the set  $P$  and  $f(x)$  is one- one if  $\lambda$  is an element of the set  $Q$ , then (given ,  $\lambda$  is a real number )

A.  $P = Q$

B.  $P \subset Q$

C.  $P - Q = \{0\}$

D.  $Q \subset P$

**Answer: D**

 [Watch Video Solution](#)

21. Consider circles  $C_1$  &  $C_2$  touching both the axes and passing through  $(4,4)$  , then the product of radii of these circles is

 [Watch Video Solution](#)

22. If  $P(z)$  is a variable point in the complex plane such that  $\operatorname{Im}\left(-\frac{1}{z}\right) = \frac{1}{4}$ , then the value of the perimeter of the locus of  $P(z)$  is (use  $\pi = 3.14$ )

 [Watch Video Solution](#)

23. The probability of India winning a test match against Australia is  $\frac{1}{4}$ . Assuming the matches to be independent events, the probability that in a 7 match series India's second win occurs at 4<sup>th</sup> test is  $P$ , then  $256P$  is equal to

A. 15

B. 12

C. 27

D. 40

**Answer: C**

 [Watch Video Solution](#)

**24.** The number of solutions of the equation  $|\cot x| = \cot x + \operatorname{cosec}x$  in  $[0, 10\pi]$  is /are

 [Watch Video Solution](#)

**25.** If  $\alpha$  is the only real root of  $x^3 + bx^2 + cx + 1 = 0 (b < c)$ , then the value of  $|\lceil \alpha \rceil|$  is (where,  $\lceil \cdot \rceil$  represents the greatest integer function)

 [Watch Video Solution](#)