



MATHS

BOOKS - NTA MOCK TESTS

JEE MOCK TEST 5

Mathematics

1. If the value of

$$(1 + \tan 1^\circ)(1 + \tan 2^\circ)(1 + \tan 3^\circ)\dots\dots\dots \text{Is}$$

2^λ , then the sum of the digits of the number λ is

A. 3

B. 6

C. 5

D. 4

Answer: C



Watch Video Solution

2. If $\frac{5 + 9 + 13 + \dots n \text{ terms}}{7 + 9 + 11 + \dots (n + 1) \text{ terms}} = \frac{17}{16}$,

then $n =$

A. 7

B. 12

C. 8

D. 15

Answer: A



Watch Video Solution

3. If $2f(xy) = (f(x))^y + (f(y))^x$ for all $x, y \in R$ and $f(1) = 3$, then the value of

$\sum_{t=1}^{10} f(t)$ is equal to

A. $\frac{3}{2}(3^{10} - 1)$

B. $\frac{3}{2}(3^9 - 1)$

C. $\frac{3^{10} - 1}{2}$

D. $\frac{1}{2}(3^{10} - 1)$

Answer: A



Watch Video Solution

4. If $\frac{1}{6}\sin\theta$, $\cos\theta$, $\tan\theta$ are in *G. P.* then θ is equal to ($n \in \mathbb{Z}$)

A. $\left\{ \theta : \theta = 2n\pi \pm \left(\frac{\pi}{6} \right), n \in I \right\}$

B. $\left\{ \theta : \theta = 2n\pi \pm \left(\frac{\pi}{3} \right), n \in I \right\}$

C. $\left\{ \theta : \theta = n\pi + (-1)^n \left(\frac{\pi}{3} \right), n \in I \right\}$

D. $\left\{ \theta : \theta = n\pi + \frac{\pi}{3}, n \in I \right\}$

Answer: B



Watch Video Solution

5. The value of integral $\int_0^1 \sqrt{\frac{1-x}{1+x}} dx$ is

A. -1

B. 1

C. $\frac{\pi}{2} - 1$

D. $\frac{\pi}{2} + 1$

Answer: C



Watch Video Solution

6. If $p, r, s > 0$ then

$$\lim_{x \rightarrow \infty} \left(\frac{p^{\frac{1}{x}} + q^{\frac{1}{x}} + r^{\frac{1}{x}} + s^{\frac{1}{x}}}{4} \right)^{3x}$$

A. $pqrs$

B. $(pqrs)^3$

C. $(pqrs)^{\frac{3}{2}}$

D. $(pqrs)^{\frac{3}{4}}$

Answer: D



Watch Video Solution

7. If $(27)^{999}$ is divided by 7, then the remainder is .

A. 1

B. 2

C. 3

D. 6

Answer: D



Watch Video Solution

8. If $A = \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix}$ and $A^2 - 4A + 10I = A$,

then k is equal to

A. 0

B. -4

C. 4

D. 1 or 4

Answer: C



Watch Video Solution

9. A piece of cheese is located at $(12, 10)$ in a coordinate plane. A mouse is at $(4, -2)$ and is running up the line $y = -5x + 18$. At the point (a, b) , the mouse starts getting farther from the cheese rather than closer to it. The value of $(a + b)$ is :

A. 6

B. 10

C. 18

D. 14

Answer: B



Watch Video Solution

10. Two medians drawn from the acute angles of a right angled triangle intersect at an angle $\frac{\pi}{6}$. If the length of the hypotenuse of the triangle is *3units*, then the area of the triangle (in sq. units) is

A. $\sqrt{3}$

B. 3

C. $\sqrt{2}$

D. 9

Answer: A



Watch Video Solution

11. If Z is a non-real complex number, then find

the minimum value of $\frac{\text{Im}z^5}{\text{Im}^5z}$

A. -2

B. -4

C. -5

D. -1

Answer: B



Watch Video Solution

12. If the coefficient of x^7 in $\left(ax^2 + \frac{1}{bx}\right)^{11}$ is equal to the coefficient of x^{-7} in

$\left(ax - \frac{1}{bx^2}\right)^{11}$ then

A. $ab = 1$

B. $\frac{a}{b} = 1$

C. $a + b = 1$

D. $a - b = 1$

Answer: A



Watch Video Solution

13. The average of five consecutive odd number is 61. Then the difference between the highest and lowest numbers is

A. 2

B. 5

C. 8

D. Cannot be determined

Answer: C



Watch Video Solution

14. The abscissa of A and B are the roots of the equation $x^2 + 2ax - b^2 = 0$ and their ordinates are the roots of the equation

$y^2 + 2py - q^2 = 0$. The equation of the circle

with AB as diameter is

A. $x^2 + y^2 + 2ax + 2py - b^2 - q^2 = 0$

B. $x^2 + y^2 + 2ax + py - b^2 - q^2 = 0$

C. $x^2 + y^2 + 2ax + 2py + b^2 + q^2 = 0$

D. None of these

Answer: A



Watch Video Solution

15. A tower T_1 of height 60 m is located exactly opposite to a tower T_2 of height 80 m on a straight road. From the top of T_1 , if the angle of depression of the foot of T_2 is twice the angle of elevation of the top of T_2 , then the width (in m) of the road between the feet of the towers T_1 and T_2 is

A. $20\sqrt{2}$

B. $10\sqrt{2}$

C. $10\sqrt{3}$

D. $20\sqrt{3}$

Answer: D



Watch Video Solution

16. The area between the curve $y = 2x^4 - x^2$, the x-axis, and the ordinates of the two minima of the curve is

A. $\frac{11}{60}$ sq. units

B. $\frac{7}{120}$ sq. units

C. $\frac{1}{30}$ sq. units

D. $\frac{7}{90}$ sq. units

Answer: B



Watch Video Solution

17. If a plane, intercepts on the coordinates axes at 8,4,4 then the length of the perpendicular from the origin to the plane is

A. $\frac{8}{3}$ units

B. $\frac{3}{8}$ units

C. 3 units

D. $\frac{4}{5}$ units

Answer: A



Watch Video Solution

18. If $f(x) = \begin{cases} \frac{\sqrt{4+ax} - \sqrt{4-ax}}{x} & -1 \leq x < 0 \\ \frac{3x+2}{x-8} & 0 \leq x \leq 1 \end{cases}$

continuous in $[-1, 1]$, then the value of a is

A. 1

B. -1

C. $\frac{1}{2}$

D. $-\frac{1}{2}$

Answer: D



Watch Video Solution

19. Check whether the relation R defined on the set $A = \{1, 2, 3, 4, 5, 6\}$ as $R = \{(a, b) : b = a + 1\}$ is reflexive, symmetric or transitive.

A. R is neither reflexive nor symmetric nor transitive

B. R is neither reflexive nor symmetric but transitive

C. R is not reflexive but symmetric and transitive

D. R is reflexive, symmetric and transitive

Answer: A



Watch Video Solution

20. $\int \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x} dx =$

A. $\frac{1}{2} \sin 2x + C$

B. $-\frac{1}{2} \sin 2x + C$

C. $-\frac{1}{2} \sin x + C$

D. $-\sin^2 x + C$

Answer: B



Watch Video Solution

21. Let $p(x) = x^2 + bx + c$, where b and c are integers. If $p(x)$ is a factor of both $x^4 + 6x^2 + 25$ and $3x^4 + 4x^2 + 28x + 5$, find the value of $p(1)$.



[Watch Video Solution](#)

22. There are eight rooms on the first floor of a hotel, with four rooms on each side of the corridor, symmetrical situated (that is each room is exactly opposite to one other room).

Four guests have to be accommodated in four of the eight rooms (that is, one in each) such that no two guests are in adjacent rooms or in opposite rooms. If N is the number of ways in which guests can be accommodated. Then the value of $\frac{N}{6}$ is



[Watch Video Solution](#)

23. Let $f: [1, \infty) \rightarrow [2, \infty)$ be a differentiable function such that $f(1) = \frac{1}{3}$. If

$$6 \int_1^x f(t) dt = 3x f(x) - x^3 \quad \text{for all } x \geq 1,$$

then the value of $3f(2)$ is



[Watch Video Solution](#)

24. If $y = \tan^{-1}(\sec x - \tan x)$, then $\frac{dy}{dx}$ is

equal to



[Watch Video Solution](#)

25. The volume of a cube is increasing at the rate of 18cm^3 per second. When the edge of

the cube is 12 cm, then the rate in cm^2 / s , at which the surface area of the cube increases, is



Watch Video Solution