



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 102

Mathematics

1. In the expansion of $(a + b)^n$, first three terms are 243, 810 and 1080 respectively, then the fourth term of the expansion is ($n \in \mathbb{N}$)

- A. 32
- B. 720
- C. 510
- D. 420

Answer: B



Watch Video Solution

2. If $z = x + iy$, $\forall x, y \in R$, $i^2 = -1$, $xy \neq 0$ and $|z| = 2$, then the imaginary part of $\frac{z+2}{z-2}$ cannot be

A. 1

B. 3

C. 2

D. 4

Answer: A



Watch Video Solution

3. The number of permutations of the alphabets of the word "GOOGLE" in which O's are together but G's are separated, is

A. 24

B. 48

C. 72

D. 36

Answer: D



Watch Video Solution

4. If B, C are square matrices of same order such that $C^2 = BC - CB$ and $B^2 = -I$, where I is an identity matrix, then the inverse of matrix $(C - B)$ is

A. C

B. $C + B$

C. $C - B$

D. I

Answer: B



[Watch Video Solution](#)

5. The tangent drawn to the hyperbola $\frac{x^2}{16} - \frac{y^2}{9} = 1$, at point P in the first quadrant whose abscissa is 5, meets the lines $3x - 4y = 0$ and $3x + 4y = 0$ at Q and R respectively. If O is the origin, then the area of triangle OQR is (in square units)

A. 6

B. 12

C. 3

D. 24

Answer: B



[Watch Video Solution](#)

6. Two natural numbers are randomly chosen and multiplied, then the chance that their product is divisible by 3 is

A. $\frac{4}{9}$

B. $\frac{5}{9}$

C. $\frac{2}{3}$

D. $\frac{1}{9}$

Answer: B



Watch Video Solution

7. If α and β are the roots of the equation $x^2 + x + c = 0$ such that $\alpha + \beta$, $\alpha^2 + \beta^2$ and $\alpha^3 + \beta^3$ are in arithmetic progression, then c is equal to

A. 1

B. $\frac{4}{3}$

C. $\frac{4}{5}$

D. $\frac{4}{7}$

Answer: D

 [Watch Video Solution](#)

8. In a harmonic progression t_1, t_2, t_3, \dots , it is given that $t_5 = 20$ and $t_6 = 50$. If S_n denotes the sum of first n terms of this, then the value of n for which S_n is maximum is

A. 6

B. 7

C. 9

D. 10

Answer: A

 [Watch Video Solution](#)

9. The locus of the centre of the circle which makes equal intercepts on the lines $x + y = 1$ and $x + y = 5$ is

A. $x - y = 2$

B. $x + y = 6$

C. $x + y = 3$

D. $x - y = 0$

Answer: C



Watch Video Solution

10. Consider the system of equations $\alpha x + y + z = p$, $x + \alpha y + z = q$ and $x + y + \alpha z = r$, then the sum of all possible distinct value(s) of α for which system does not possess a unique solution is

A. -2

B. 1

C. -1

D. 0

Answer: C



Watch Video Solution

11. The normal to the parabola $y^2 = 4x$ at $P(9, 6)$ meets the parabola again at Q. If the tangent at Q meets the directrix at R, then the slope of another tangent drawn from point R to this parabola is

A. 11

B. $\frac{11}{3}$

C. $\frac{3}{11}$

D. 3

Answer: B



Watch Video Solution

12. The number of points where $f(x) = |x^2 - 3|x| - 4|$ is non-differentiable is

A. 1

B. 2

C. 3

D. 4

Answer: C



Watch Video Solution

13. The complete set of values of α for which the lines $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and $\frac{x-3}{2} = \frac{y-5}{\alpha} = \frac{z-7}{\alpha+2}$ are

concurrent and coplanar is

A. $\{2, 3\}$

B. $\{0, 3\}$

C. $[-2, 3]$

D. R

Answer: D



Watch Video Solution

14. Let $f(x) = 2x + 1$ and $g(x) = \int \frac{f(x)}{x^2(x+1)^2} dx$. If $6g(2) + 1 = 0$ then $g\left(-\frac{1}{2}\right)$ is equal to

A. 4

B. -4

C. 3

D. 2

Answer: A



Watch Video Solution

15. Let $f(x)$ be a cubic function such that $f'(1) = f''(2) = 0$. If $x = 1$ is a point of local maxima of $f(x)$, then the local minimum value of $f(x)$ occurs at

A. $x = 0$

B. $x = 2$

C. $x = 4$

D. $x = 3$

Answer: D



Watch Video Solution

16. The maximum value of p for which the lines $3x - 4y = 2$, $3x - 4y = 12$, $12x + 5y = 7$ and $12x + 5y = p$ constitute the sides of a rhombus is

A. 33

B. 19

C. -19

D. 9

Answer: A



Watch Video Solution

17. The function $f: R \rightarrow R$ defined as $f(x) = \frac{x^2 - x + 1}{x^2 + x + 1}$ is

A. injective as well as surjective

B. injective but not surjective

C. surjective but not injective

D. neither injective nor surjective

Answer: D



Watch Video Solution

18. The value of $\lim_{x \rightarrow 0^+} \{x^{x^2} + x^{(x^x)}\}$ is equal to

A. 0

B. 1

C. 2

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

19. The area (in sq. units) bounded by $y = \ln x$, $y = \frac{x}{e}$ and y - axis is equal to

A. $\frac{e}{2} - 1$

B. $\frac{e}{2}$

C. $\frac{5e}{2}$

D. $\frac{3e}{2} - 1$

Answer: B



Watch Video Solution

20. Consider three vectors

$$\vec{p} = \hat{i} + \hat{j} + \hat{k}, \vec{q} = 3\hat{i} - \hat{j} + \hat{k} \text{ and } \vec{r} = \alpha\hat{i} + \beta\hat{j} + \lambda\hat{k}, \forall \alpha, \beta, \lambda \in R$$

. If $[\vec{p} \ \vec{q} \ \vec{r}]$ is maximum and $[\vec{r}] = 2\sqrt{6}$, then the value of $\alpha - \beta - \lambda$ is equal to

A. 8

B. 4

C. 0

D. -4

Answer: B

 [Watch Video Solution](#)

21. If $\sin \theta + \sin^2 \theta = 1$, then prove that $\cos^{12} \theta + 3 \cos^{10} \theta + 3 \cos^8 \theta + \cos^6 \theta - 1 = 0$

 [Watch Video Solution](#)

22. If $y = f(x)$ satisfies the differential equation $\frac{dy}{dx} + \frac{2x}{1+x^2}y = \frac{3x^2}{1+x^2}$ where $f(1) = 1$, then $f(2)$ is equal to

 [Watch Video Solution](#)

23. If the variance of the first 50 odd natural numbers is V_1 and the variance of next 50 odd natural numbers is V_2 , then $V_1 + V_2$ is equal to

 [Watch Video Solution](#)

24. If

$$I_1 = \int_0^{\frac{\pi}{2}} e^{\sin x} (1 + x \cos x) dx \quad \text{and} \quad I_2 = \int_0^{\frac{\pi}{2}} e^{\cos x} (1 - x \sin x) dx,$$

then $\left[\frac{I_1}{I_2} \right]$ is equal to (where $[x]$ denotes the greatest integer less than or equal to x)

 [Watch Video Solution](#)

25. The number of solution of $\cos^2 x + \cos^2 2x = 2$ in $[0, 20]$ is equal to

 [Watch Video Solution](#)