



## MATHS

### BOOKS - NTA MOCK TESTS

#### NTA JEE MOCK TEST 106

#### Mathematics

1. The value of the integral

$$\int_0^4 \{x^3 - 6x^2 + 12x - 4 + (x - 2)\cos(x - 2)\} dx \text{ is equal to}$$

A. 12

B. 16

C. 0

D. 10

**Answer: B**



**Watch Video Solution**

2. Let  $a, \lambda, \mu \in \mathbb{R}$ , Consider the system of linear equations  $ax + 2y = \lambda 3x - 2y = \mu$  Which of the following statement (s) is (are) correct?

- A. If  $a = -3$ , then the system has infinitely many solutions for all values of  $\lambda$  and  $\mu$
- B. If  $a \neq -3$ , then the system has a unique solution for all values of  $\lambda$  and  $\mu$
- C. If  $\lambda + \mu = 0$ , then the system has infinitely many solution for  $a = -3$
- D. IF  $\lambda + \mu \neq 0$ , then the system has no solution for  $a = -3$

**Answer:**



**Watch Video Solution**

3. Consider the statement p : If a cylinder is right circular cylinder then its volume is  $\frac{1}{3}\pi r^2 h$ .

Contrapositive of the statement p is

A. If the volume of a cylinder is  $\frac{1}{3}\pi r^2 h$  then it is not right circular cylinder

B. If the volume of a cylinder is not  $\frac{1}{3}\pi r^2 h$  then it is not right circular cylinder

C. If the volume of a cylinder is not  $\frac{1}{3}\pi r^2 h$  then it is not right circular cylinder

D. If a cylinder is not right circular cylinder then its volume is not  $\frac{1}{3}\pi r^2 h$

**Answer: B**



Watch Video Solution

4. Find the value of  $\sin^{-1}(\cos(\sin^{-1} x)) + \cos^{-1}(\sin(\cos^{-1} x))$

A. 0

B.  $\frac{\pi}{4}$

C.  $\frac{\pi}{2}$

D.  $\frac{3\pi}{4}$

Answer: C



Watch Video Solution

5. Let  $\int \frac{10 \ln x}{x^2} dx = f(x)$ , for all positive x. If  $f(e) = \frac{1}{e}$ , then  $f(2) + f(4)$  is equal to



Watch Video Solution

6. If  $f(x) = x^2 + \frac{x^2}{1+x^2} + \frac{x^2}{(1+x^2)^2} + \dots$  upto,  $\infty$ , then

A.  $\lim_{x \rightarrow 0} f(x)$  does not exist

B.  $f(x)$  is continuous but not differentiable at  $x = 0$

C.  $f(x)$  is discontinuous at  $x = 0$

D.  $f(x)$  is differentiable at  $x = 0$

**Answer: C**

 [Watch Video Solution](#)

7. The area (in square units) bounded by  $y = x^2 + x + 1$  and  $x + y = 2$  is

A.  $2\sqrt{2}$

B.  $\frac{4\sqrt{2}}{3}$

C.  $\frac{8\sqrt{2}}{3}$

D.  $\frac{\sqrt{2}}{3}$

**Answer: C**

 [Watch Video Solution](#)

8.

Suppose

$$\cos^2 y \cdot \frac{dy}{dx} = \sin(x + y) + \sin(x - y), |x| \leq \frac{\pi}{2} \text{ and } |y| \leq \frac{\pi}{2}. \quad \text{If}$$

$$y\left(\frac{\pi}{3}\right) = -\frac{\pi}{2}, \text{ then } y\left(\frac{\pi}{2}\right) \text{ is}$$

A. 0

B.  $\frac{\pi}{4}$

C.  $-\frac{\pi}{4}$

D.  $\frac{\pi}{2}$

**Answer: A**

 [Watch Video Solution](#)

9. If  $(1!)^2 + (2!)^2 + (3!)^2 + \dots + (99!)^2$  is divided by 100, the remainder is

A. 27

B. 28

C. 17

D. 14

**Answer: C**



[Watch Video Solution](#)

**10.** If a number of ellipse whose major axis is x - axis and the minor axis is y - axis be described having the same length of the major axis as  $2a$  but a variable minor axis, then the tangents at the ends of their latus rectum pass through fixed points whose distance from the centre is equal to

A.  $\frac{1}{2}$  units

B. 1 unit

C.  $\frac{3}{2}$  units

D. 2 units

**Answer: B**



**Watch Video Solution**

**11.** Find the set of all values of  $a$  for which the roots of the equation

$$x^2 - 2ax + a^2 + a - 3 = 0 \text{ are less than } 3,$$

A.  $a < 2$

B.  $2 \leq a \leq 3$

C.  $3 < a \leq 4$

D.  $a > 4$

**Answer: A**



**Watch Video Solution**

**12.**

Let

$$F(x) = \left(1 + \sin\left(\frac{\pi}{2k}\right)\right) \left(1 + \sin\left(k - 1\right)\frac{\pi}{2k}\right) \left(1 + \sin(2k + 1)\frac{\pi}{2k}\right) \left(1 + \sin\left(\dots\right)\right)$$



. The value of  $F(1) + F(2) + F(3)$  is equal to

A.  $\frac{3}{16}$

B.  $\frac{1}{4}$

C.  $\frac{5}{16}$

D.  $\frac{7}{16}$

**Answer: D**



**Watch Video Solution**

13. Number of solutions of the equation

$\sin^4 x - \cos^2 x \sin x + 2 \sin^2 x + \sin x = 0 \in 0 \leq x \leq 3\pi$  is \_\_\_\_\_

A.  $\frac{3\pi}{4}$

B.  $2\pi$

C.  $4\pi$

D.  $6\pi$

Answer: D



Watch Video Solution

14.

Let

$$\vec{a} = (\cos \theta)\hat{i} - (\sin \theta)\hat{j}, \vec{b} = (\sin \theta)\hat{i} + (\cos \theta)\hat{j}, \vec{c} = \hat{k} \text{ and } \vec{r} = 7\hat{i} +$$

. IF  $\vec{r} = x\vec{a} + y\vec{b} + z\vec{c}$ , then the value of  $\frac{x^2 + y^2}{z}$  is equal to

A. 3

B. 5

C. 50

D. 7

Answer: B



Watch Video Solution

15. Two numbers are selected randomly from the set  $S = \{1, 2, 3, 4, 5, 6\}$  without replacement one by one. The probability that minimum of the two numbers is less than 4 is

A.  $\frac{1}{15}$

B.  $\frac{14}{15}$

C.  $\frac{1}{5}$

D.  $\frac{4}{5}$

**Answer: D**



**Watch Video Solution**

16. A line is drawn from the point  $P(1, 1, 1)$  and perpendicular to a line with direction ratio  $(1, 1, 1)$  to intersect the plane  $x + 2y + 3z = 4$  at  $Q$ . The locus of point  $Q$  is :

A.  $\frac{x}{1} = \frac{y - 5}{-2} = \frac{z + 2}{1}$

$$B. \frac{x}{-2} = \frac{y-5}{1} = \frac{z+2}{1}$$

$$C. x = y = z$$

$$D. \frac{x}{2} = \frac{y}{3} = \frac{z}{5}$$

**Answer: A**



**Watch Video Solution**

17. If  $\alpha, \beta, \gamma$  are the roots of  $px^2 + qx^2 + r = 0$ , then the value of the determinant  $|\alpha\beta\beta\gamma\gamma\alpha\beta\gamma\gamma\alpha\alpha\beta\gamma\alpha\alpha\beta\beta\gamma|$  is  $p$  b.  $q$  c.  $0$  d.  $r$

A.  $pq$

B.  $qr$

C.  $0$

D.  $pr$

**Answer: C**



**Watch Video Solution**

18. If  $f(x) = \sin^{-1} \frac{2 \cdot (3)^x}{1 + 9^x}$ , then  $f' \left( -\frac{1}{2} \right)$  is equal to

A.  $\sqrt{3} \log_e \sqrt{3}$

B.  $-\sqrt{3} \log_e \sqrt{3}$

C.  $-\sqrt{3} \log_e 3$

D.  $\sqrt{3} \log_e 3$

**Answer: A**



**Watch Video Solution**

19. The line  $2x - y + 1 = 0$  is tangent to the circle at the point  $(2, 5)$  and the centre of the circle lies on  $x - 2y = 4$ . The square of the radius of the circle is

A. 45

B. 75

C. 20

D. 50

**Answer: A**



[Watch Video Solution](#)

**20.** The number of ways in which four different letters can be put in their four addressed envelopes such that atleast two of them are in the wrong envelopes are

A. 23

B. 25

C. 6

D. 7

**Answer: A**



[Watch Video Solution](#)

21. The maximum number of points on the parabola  $y^2 = 16x$  which are equidistant from a variable point P (which lie inside the parabola) are

 [Watch Video Solution](#)

22. The slope of the tangent to the curve  $y = x^3 + x + 54$  which also passes through the origin is

 [Watch Video Solution](#)

23. In an arithmetic progression containing 99 terms, the sum of all the odd numbered terms, the sum of all the odd numbered terms is 2550. If the sum of all the 99 terms of the arithmetic progression is  $k$ , then  $\frac{k}{100}$  is equal to

 [Watch Video Solution](#)

24. Consider the system of equations  $ax + y + bz = 0$ ,  $bx + y + az = 0$  and  $ax + by + abz = 0$  where  $a, b \in \{0, 1, 2, 3, 4\}$ . The number of ordered pairs  $(a, b)$  for which the system has non-trivial solutions is

 [Watch Video Solution](#)

25. Let  $A(z_1)$ ,  $B(z_2)$  and  $C(z_3)$  be complex numbers satisfying the equation  $|z| = 1$  and also satisfying the relation  $3z_1 = 2z_2 + 2z_3$ . Then  $|z_2 - z_3|^2$  is equal to

 [Watch Video Solution](#)