



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 30

Mathematics

1. The number of solutions of the equation

$$z^3 + \frac{3(\bar{z})^2}{|z|} = 0 \text{ (where, } z \text{ is a complex number)}$$

are

A. 2

B. 3

C. 6

D. 5

Answer: D



Watch Video Solution

2. If $f(x) = (x^3 + x + 1)\tan(\pi[x])$ (where, $[x]$ represents the greatest integer part of x), then

A. domain of $f(x)$ is $R - (2n + 1)\frac{\pi}{2}, n \in I$

B. range of $f(x) \in R$

C. $f(x)$ is an even function

D. $f(x)$ is a non - periodic function

Answer: C



Watch Video Solution

3. A five digit number is chosen at random. The probability that all the digits are distinct and digits at odd places are odd and digits at even places are even, is

A. ${}^5P_2 \times \frac{{}^5P_3}{9 \times 10^4}$

B. $\frac{{}^5P_2 \times {}^5P_3}{10^5}$

C. $\frac{{}^5C_2 \times {}^5C_3 \times 2}{10^4 \times 9}$

D. $\frac{{}^5C_2 \times {}^5C_3}{9 \times 10^4}$

Answer: A



Watch Video Solution

4. The length of the focal chord of the parabola

$y^2 = 4x$ at a distance of 0.4 units from the origin

is

A. 22 units

B. 23 units

C. 24 units

D. 25 units

Answer: D



Watch Video Solution

5. If p, q, r in \mathbb{N} , then the number of points having position vector $p\hat{i} + q\hat{j} + r\hat{k}$ such that $8 \leq p + q + r \leq 12$ are

A. 110

B. 185

C. 144

D. 108

Answer: B



Watch Video Solution

6. Let α, β and γ be the roots of the equation $x^3 + 6x^2 - px - 42 = 0$. If α, β and γ are in arithmetic progression then $|\alpha| + |\beta| + |\gamma| =$

A. 10

B. 11

C. 12

D. 13

Answer: C



Watch Video Solution

7. The solution of the differential equation

$$\frac{ydx - xdy}{xy} = xdx + ydy \text{ is (where, } C \text{ is an}$$

arbitrary constant)

A. $\frac{x}{y} = x + y + C$

B. $\frac{x}{y} = \frac{x^2 + y^2}{2} + c$

C. $\ln\left(\frac{x}{y}\right) = x^2 + y^2 + C$

D. $2 \ln\left(\frac{x}{y}\right) = x^2 + y^2 + C$

Answer: D



Watch Video Solution

8. If a circle drawn by assuming a chord parallel to the transverse axis of hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ as diameter always passes through $(2, 0)$, then

A. $|a| = |b| = 2$

B. $|b| \neq |a|$

C. $|b| = |a| = 1$

D. $|b| = |a| = 3$

Answer: A



Watch Video Solution

9. If A is a square matrix of order 2×2 and

$B = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, such that $AB = BA$, then A can be

A. $\begin{bmatrix} 1 & 4 \\ 6 & 7 \end{bmatrix}$

B. $\begin{bmatrix} 1 & 4 \\ 7 & 6 \end{bmatrix}$

C. $\begin{bmatrix} 2 & 2 \\ 2 & 4 \end{bmatrix}$

D. $\begin{bmatrix} 3 & 4 \\ 4 & 9 \end{bmatrix}$

Answer: A



Watch Video Solution

10. Let \oplus and \otimes are two mathematical operators. If $p \oplus (p \otimes q)$ is not a tautology, then \oplus and \otimes can be

A. \vee and \Rightarrow respectively

B. \Rightarrow and \wedge respectively

C. \Rightarrow and \vee respectively

D. None of these

Answer: B



Watch Video Solution

11. The coefficient of x^{10} in the expansion of $(1+x)^{15} + (1+x)^{16} + (1+x)^{17} + \dots + (1+x)^{30}$

A. ${}^{31}C_{10} - {}^{15}C_{10}$

B. ${}^{31}C_{11} - {}^{15}C_{11}$

C. ${}^{30}C_{10} - {}^{15}C_{10}$

D. ${}^{31}C_{10} - {}^{14}C_{11}$

Answer: B



Watch Video Solution

12. A vector \vec{r} is equally inclined with the vectors $\vec{a} = \cos \theta \hat{i} + \sin \theta \hat{j}$, $\vec{b} = -\sin \theta \hat{i} + \cos \theta \hat{j}$ and $\vec{c} = \hat{k}$, then the angle between \vec{r} and \vec{a} is

A. $\cos^{-1} \left(\frac{1}{\sqrt{2}} \right)$

B. $\cos^{-1} \left(\frac{1}{3} \right)$

C. $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$

D. $\cos^{-1}\left(\frac{1}{2}\right)$

Answer: C



Watch Video Solution

13. Let $f(x)$ be a non - constant polynomial such that $f(a) = f(b) = f(c) = 2$. Then the minimum number of roots of the equation $f''(x) = 0$ in $x \in (a, c)$ is/are

A. 2

B. 1

C. 0

D. 3

Answer: B



Watch Video Solution

14. Let the incentre of $\triangle ABC$ is $I(2, 5)$. If $A = (1, 13)$ and $B = (-4, 1)$, then the coordinates of C are

A. $(1, 10)$

B. (10, 1)

C. (8, 2)

D. (9, 2)

Answer: B



Watch Video Solution

15. Lying in the plane $x + y + z = 6$ is a line L passing through $(1, 2, 3)$ and perpendicular to the line of intersection of planes $x + y + z = 6$ and $2x - y + z = 4$, then the equation of L is

$$\text{A. } \frac{x-1}{4} = \frac{y-2}{-7} = \frac{z-3}{3}$$

$$\text{B. } \frac{x-1}{2} \frac{y-2}{1} = \frac{z-3}{-3}$$

$$\text{C. } \frac{x-1}{4} = \frac{y-2}{-5} = \frac{z-3}{1}$$

$$\text{D. } \frac{x-1}{3} = \frac{y-2}{1} = \frac{z-3}{-4}$$

Answer: C



Watch Video Solution

16. Let PQ and RS be tangents at the extremities of the diameter PR of a circle of radius r . If PS and RQ intersect at a point X on the circumference of the circle, then $2r$ equals

A. $\sqrt{PQ \cdot RS}$

B. $\frac{PQ + RS}{2}$

C. $\frac{2PQ \cdot RS}{PQ + RS}$

D. $\sqrt{\frac{PQ^2 + RS^2}{2}}$

Answer: A



Watch Video Solution

17. Let A and B are two matrices of order 3×3 ,

where $|A| = -2$ and $|B| = 2$, then

$|A^{-1} \text{adj}(B^{-1}) \text{adj}(2A^{-1})|$ is equal to

A. 2

B. -2

C. 4

D. 8

Answer: B



Watch Video Solution

18. The sum to infinite terms of the arithmetic -
gemoetric progression $3, 4, 4, \frac{32}{9}, \dots$ is equal
to

A. 27

B. 30

C. 24

D. 25

Answer: A



Watch Video Solution

19. The value of $\int_{-1}^1 \left(\sin^{-1} x + \frac{x^5 + x^3 - 1}{\cos x} \right) dx$

is equal to

A. $\tan 1$

B. 0

C. $2 \tan 1$

D. $-2 \tan 1$

Answer: D



Watch Video Solution

20. If $E = \cos^2 71^\circ + \cos^2 49^\circ + \cos 71^\circ \cos 49^\circ$,

then the value of $10E$ is equal to

A. 7.5

B. 2.5

C. 3.5

D. 4.5

Answer: A



Watch Video Solution

21. If $x = 3 \cos t$ and $y = 5 \sin t$, where t is a parameter, then $9 \frac{d^2 y}{dx^2}$ at $t = -\frac{\pi}{6}$ is equal to



Watch Video Solution

22. The area (in sq. units) of the region bounded by the curves $y = 2 - x^2$ and $y = |x|$ is k , then the value of $3k$ is



[Watch Video Solution](#)

23. A number equal to 2 times the mean and with a frequency equal to k is inserted in a data having n observation. If the new mean is $\frac{4}{3}$ times the old mean, then the value of $\frac{k}{n}$ is



[Watch Video Solution](#)

$$24. \text{ If } f(x) = \begin{cases} \frac{\sqrt{x}}{\sqrt{4+\sqrt{x}}-a} & x > 0 \\ c & x = 0 \\ \frac{4e^{\frac{2}{x}} + 3e^{\frac{1}{x}}}{e^{\frac{2}{x}} + be^{\frac{1}{x}}} & x < 0 \end{cases}$$

continuous at $x = 0$ for some constants a , b and c ,

then the value of $\frac{50b}{a}$ is equal to

 [Watch Video Solution](#)

25. The integral $I = \int (\sin(x^2) + 2x^2 \cos(x^2)) dx$

(where $I = xh(x) + c$, C is the constant of

integration). If the range of $H(x)$ is $[a, b]$, then

the value of $a + 2b$ is equal to

 [Watch Video Solution](#)

