



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 32

Mathematics

1. Two vertical poles of height 10 m and 40 m stand apart on a horizontal plane. The height (in meters) of the point of intersection of the line joining the top of intersection of the lines joining the top of

each pole to the foot of the other, from this horizontal plane is

A. 8

B. 10

C. 6

D. 4

Answer: A



Watch Video Solution

2. The total number of solution(s) of the equation

$$2x + 3 \tan x = \frac{5\pi}{2} \text{ in } x \in [0, 2\pi] \text{ is/are equal to}$$

A. 1

B. 2

C. 3

D. 4

Answer: C



Watch Video Solution

3. If $y = |\tan x - |\sin x ||$, then the value of $\frac{dy}{dx}$ at $x = \frac{5\pi}{4}$ is

A. $\frac{2\sqrt{2} + 1}{\sqrt{2}}$

B. $\frac{2\sqrt{2} - 1}{\sqrt{2}}$

C. $\frac{\sqrt{2} + 1}{2}$

D. $\frac{\sqrt{2} - 1}{2}$

Answer: B



Watch Video Solution

4. The value of k for which the sum of the squares of the roots of $2x^2 - 2(k - 2)x - (k + 1) = 0$ is least is

A. 1

B. $\frac{3}{2}$

C. 2

D. $\frac{5}{2}$

Answer: B



Watch Video Solution

5. If $\lim_{x \rightarrow 0} \frac{\sin 2x - a \sin x}{x^3}$ exists finitely, then the value of a is

A. 0

B. 2

C. 1

D. 4

Answer: B



Watch Video Solution

6. If $f(x) = \tan^{-1} \sqrt{x^2 + 4x} + \sin^{-1} \sqrt{x^2 + 4x + 1}$

- A. domain of $f(x)$ contains 3 integers only
- B. range of $f(x)$ has two elements only
- C. $f(x)$ is a constant function $\forall x \in R$
- D. $f(x)$ contains only two elements in its domain

Answer: D



Watch Video Solution

7. The function

$$f(x) = \max \{ (1 - x), (1 + x), 2 \}, x \in (-\infty, \infty)$$

is

A. discontinuous at exactly two points

B. differentiable $\forall x \in \mathbb{R}$

C. differentiable $\forall x \in \mathbb{R} - \{-1, 1\}$

D. continuous $\forall x \in \mathbb{R} - \{0, 1, -1\}$

Answer: C



Watch Video Solution

8. A nine - digit number is formed using the digits 1, 2, 3, 5 and 7. The probability that the product of all digits is always 1920 is

A. $\frac{1}{5^9}$

B. $\frac{7}{5^8}$

C. $\frac{72}{5^9}$

D. $\frac{1}{7!}$

Answer: C



Watch Video Solution

9. The order of the differential equation of the family of curves $y = a3^{bx+c} + d \sin(x + e)$ is (where, a, b, c, d, e are arbitrary constants)

A. 5

B. 4

C. 3

D. 2

Answer: B



Watch Video Solution

10. Focus of hyperbola is $(\pm 3, 0)$ and equation of tangent is $2x + y - 4 = 0$, find the equation of hyperbola is

A. $4x^2 - 5y^2 = 20$

B. $5x^2 - 4y^2 = 20$

C. $4x^2 - 5y^2 = 1$

D. $5x^2 - 4y^2 = 1$

Answer: A



Watch Video Solution

11. Which of the following statement is not a fallacy?

A. $p \wedge (\sim(\sim p \Rightarrow q))$

B. $\sim((p \wedge q) \Rightarrow p)$

C. $\sim(p \Rightarrow (p \vee q))$

D. $\sim p \vee (\sim p \Rightarrow q)$

Answer: D



Watch Video Solution

12. The value of $\int \frac{e^{\sqrt{x}}}{\sqrt{x}(1 + e^{2\sqrt{x}})} dx$ is equal to

(where, C is the constant of integration)

A. $\tan^{-1}(2e^{\sqrt{x}}) + C$

B. $\ln\left(\frac{1 + e^x}{1 - e^{\sqrt{x}}}\right) + C$

C. $2 \tan^{-1}(e^{\sqrt{x}}) + C$

D. $(\tan^{-1} x)e^{\sqrt{x}} + C$

Answer: C



Watch Video Solution

13. A plane passes through $(1, -2, 1)$ and is perpendicular to two planes $2x - 2y + z = 0$ and $x - y + 2z = 4$. The distance of the plane from the point $(0, 2, 2)$ is

A. $\frac{3}{\sqrt{2}}$ units

B. $4\sqrt{2}$ units

C. $3\sqrt{2}$ units

D. $2\sqrt{2}$ units

Answer: A



Watch Video Solution

14. Consider $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$

such that $AB = BA$. then the value of

$$\frac{a_{12}}{a_{21}} + \frac{a_{11}}{a_{22}} \text{ is}$$

A. 2

B. 4

C. $\frac{3}{2}$

D. $\frac{1}{\sqrt{2}}$

Answer: C



Watch Video Solution

15. A line passing through the point $(2, 2)$ encloses an area of 4 sq. units with coordinate axes. The sum of intercepts made by the line on the x and y axis is equal to

A. -2

B. 4

C. -4

D. 2

Answer: C



Watch Video Solution

16. If 2, 7, 9 and 5 are subtracted respectively from four numbers in geometric progression, then the resulting numbers are in arithmetic progression. The smallest of the four numbers is

A. -24

B. -12

C. 6

D. 3

Answer: A



Watch Video Solution

17. The coefficient of x^6 in the expansion of $(1 + x + x^2 + x^3)(1 - x)^6$ is

A. -10

B. 10

C. 9

D. -9

Answer: D



Watch Video Solution

18. The acute angles between the curves $y = 2x^2 - x$ and $y^2 = x$ at $(0, 0)$ and $(1, 1)$ are α and β respectively, then

A. $\alpha - \beta = 0$

B. $\alpha + \beta = 0$

C. $\alpha > \beta$

D. $\alpha < \beta$

Answer: A



Watch Video Solution

19. The slope of the tangent of the curve

$$y = \int_x^{x^2} (\cos^{-1} t^2) dt \text{ at } x = \frac{1}{\sqrt{2}} \text{ is equal to}$$

A. $\cos^{-1} \left(\frac{1}{4} \right) - \frac{\pi}{3}$

B. $\cos^{-1} \left(\frac{1}{4} \right) + \frac{\pi}{3}$

C. $\sqrt{2} \cos^{-1} \left(\frac{1}{4} \right) - \frac{\pi}{3}$

D. $\sqrt{2} \cos^{-1} \left(\frac{1}{4} + \frac{\pi}{3} \right)$

Answer: C



Watch Video Solution

20. The value of $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} e^{\sec^2 x} \frac{\sin x}{\cos^3 x} dx$ is equal to

A. $\frac{1}{2}e^4$

B. $\frac{1}{2}e^{\frac{4}{3}}$

C. $\frac{1}{2}(e^{\frac{4}{3}} - e^4)$

D. $\frac{1}{2}(e^2 - 1)$

Answer: C



Watch Video Solution

21. Let PQ be a focal chord of the parabola $y^2 = 4ax$. If the centre of a circle having PQ as its diameter lies on the line $\sqrt{5}y + 4 = 0$, then length of the chord PQ, is



Watch Video Solution

22. If $\lambda \in \mathbb{R}$ such that the origin and the non-real roots of the equation $2z^2 + 2z + \lambda = 0$ form the vertices of an equilateral triangle in the argand plane, then $\frac{1}{\lambda}$ is equal to



Watch Video Solution

23. Let $|A| = 1$, $|\vec{b}| = 4$ and $\vec{a} \times \vec{r} + \vec{b} = \vec{r}$. If the projection of \vec{r} along \vec{a} is 2, then the projection of \vec{r} along \vec{b} is



Watch Video Solution

24. Let a, x, y, z be real numbers satisfying the equations

$$ax + ay = z$$

$$x + ay = z$$

$x + ay = az$, where x, y, z are not all zero, then the number of the possible values of a is



Watch Video Solution

25. If the radius of the circle passing through the origin and touching the line $x + y = 2$ at $(1, 1)$ is

r units, then the value of $3\sqrt{2}r$ is



Watch Video Solution