



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 59

Mathematics

1. Consider the matrix $A = \begin{bmatrix} 3 & 1 \\ -6 & -2 \end{bmatrix}$, then

$(I + A)^{40}$ is equal to

A. $I + 2^{38}A$

B. $I + 2^{39}A$

C. $I + (2^{40} + 1)A$

D. $I + (2^{40} - 1)A$

Answer: D



Watch Video Solution

2. The lines joining the origin to the points of intersection of the line $4x + 3y = 24$ with the circle $(x - 3)^2 + (y - 4)^2 = 25$ are

A. coincident

B. perpendicular

C. equally inclined to x - axis

D. None of these

Answer: B



Watch Video Solution

3. The value of the integral $I = \int_0^{\frac{\pi}{2}} \frac{\cos x - \sin x}{10 - x^2 + \frac{\pi x}{2}} dx$

is equal to

A. $\frac{\pi}{2}$

B. π

C. 0

D. 4π

Answer: C



Watch Video Solution

4. If $\sin x + \cos x = \frac{\sqrt{7}}{2}$, where $x \in \left[0, \frac{\pi}{4}\right]$, then the value of $\tan \frac{x}{2}$ is equal to

A. $\frac{3 - \sqrt{7}}{3}$

B. $\frac{\sqrt{7} - 2}{3}$

C. $\frac{4 - \sqrt{7}}{4}$

D. $\frac{5 - \sqrt{3}}{2}$

Answer: B



Watch Video Solution

5. The equation $\tan^4 x - 2\sec^2 x + a^2 = 0$ will have at least one solution, if

A. $|a| \leq 4$

B. $|a| \leq 2$

C. $|a| \leq \sqrt{3}$

D. $|a| > 2$

Answer: C



Watch Video Solution

6. The statement $\sim p \rightarrow (q \rightarrow p)$ is equivalent to

A. $p \rightarrow (p \rightarrow q)$

B. $p \rightarrow (p \vee q)$

C. $q \rightarrow p$

D. $q \rightarrow (p \rightarrow q)$

Answer: C



Watch Video Solution

7. If the standard deviation of n observation

x_1, x_2, \dots, x_n is 5 and for another set of n

observation y_1, y_2, \dots, y_n is 4, then the standard deviation of n observation

$x_1 - y_1, x_2 - y_2, \dots, x_n - y_n$ is

A. 1

B. $\frac{\sqrt{5}}{2}$

C. 5

D. Data insufficient

Answer: D



Watch Video Solution

8. The domain of the function

$$f(x) = \log_2 [1 - \log_{12}(x^2 - 5x + 16)] \text{ is}$$

A. $(1, 4)$

B. $(-\infty, 4]$

C. $[1, \infty)$

D. $[1, 4]$

Answer: A



Watch Video Solution

9. The length of the portion of the common tangent to $x^2 + y^2 = 16$ and $9x^2 + 25y^2 = 225$ between the two points of contact is

A. $\frac{9}{4}$ units

B. $\frac{\sqrt{3}}{4}$ units

C. $\frac{3}{4}\sqrt{7}$ units

D. $\frac{5}{4}\sqrt{7}$ units

Answer: C



Watch Video Solution

10. The equation of the curve lying in the first quadrant, such that the portion of the x - axis cut - off between the origin and the tangent at any point P is equal to the ordinate of P, is (where, c is an arbitrary constant)

A. $y = ce^{\frac{x}{y}}$

B. $ye^{\frac{x}{y}} = c$

C. $ye^{\frac{y}{x}} = c$

D. $y = ce^{\frac{y}{x}}$

Answer: B



Watch Video Solution

11. $\sum_{r=1}^n = \frac{r}{r^4 + r^2 + 1}$ is equal to

A. $\frac{n^2 + n}{2(n^2 + n + 1)}$

B. $\frac{n^2 + 2n}{2(n^2 + n + 1)}$

C. $\frac{2n^2 + n}{2(n^2 + n + 1)}$

D. $\frac{n^2 + n}{(n^2 + n + 1)}$

Answer: A



Watch Video Solution

12. If the integral $I = \int e^{x^2} x^3 dx = e^{x^2} f(x) + c$, where c is the constant of integration and $f(1) = 0$, then

the value of $f(2)$ is equal to

A. 4

B. $\frac{5}{2}$

C. $\frac{3}{2}$

D. 3

Answer: C



Watch Video Solution

13. The points on the curve $y = x^2$ which are closest to the point $P(0, 1)$ are

A. $(\pm \sqrt{2}, 2)$

B. $\left(\pm \frac{1}{2}, \frac{1}{4}\right)$

C. $\left(\pm \frac{1}{\sqrt{2}}, \frac{1}{2}\right)$

D. $\left(\pm \frac{1}{4}, \frac{1}{16}\right)$

Answer: C



Watch Video Solution

14. Let ΔOAB be an equilateral triangle with side length unity (O being the origin). Also, M and N being closer to A and N being closer to B. position vectors

of A, B, M and N are \vec{a} , \vec{b} , \vec{m} and \vec{n} respectively, then the value of $\vec{m} \cdot \vec{n}$ is equal to

A. $\frac{1}{2}$

B. $\frac{2}{3}$

C. $\frac{13}{18}$

D. $\frac{4}{9}$

Answer: C



Watch Video Solution

15. If A and B are two independent events such that

$$P(A) > \frac{1}{2}, P(A \cap B^C) = \frac{3}{25} \text{ and } P(A^C \cap B) = \frac{8}{25}$$

, then $P(A)$ is equal to (where, A^c and B^c represent the complement of events A and B respectively)

A. $\frac{1}{5}$

B. $\frac{3}{5}$

C. $\frac{3}{4}$

D. $\frac{4}{5}$

Answer: B



Watch Video Solution

16. If A , B and C are square matrices of order 3 and $|A| = 2$, $|B| = 3$ and $|C| = 4$, then the value of $|3(\text{adj}A)BC^{-1}|$ is equal to (where, $\text{adj} A$ represents the adjoint matrix of A)

A. 27

B. $\frac{27}{4}$

C. $\frac{81}{2}$

D. 81

Answer: D



Watch Video Solution

17. $\sum_{r=0}^n (n-r) \binom{n}{r}^2$ is equal to

A. $n^2 \binom{2n-1}{n}$

B. $n^2 \binom{2n}{n-1}$

C. $n^2 \binom{2n-1}{n-1}$

D. $m^2 \binom{2n-2}{n}$

Answer: D



Watch Video Solution

18. For a complex number z , the equation

$z^2 + (p + iq)zr + is = 0$ has a real root (where p ,

q, r, s are non-zero real numbers and $i^2 = -1$), then

A. $pqr = r^2 + p^2s$

B. $prs = q^2 + r^2p$

C. $prs = p^2 + s^2q$

D. $pqs = s^2 + q^2r$

Answer: D



Watch Video Solution

19. The length of the normal chord which subtends an angle of 90° at the vertex of the parabola $y^2 = 4x$ is

A. $6\sqrt{3}$ units

B. $7\sqrt{2}$ units

C. $8\sqrt{2}$ units

D. $9\sqrt{2}$ units

Answer: A



Watch Video Solution

20. Let $f(x) = \begin{cases} \frac{2^{\frac{1}{x}} - 1}{2^{\frac{1}{x}} + 1} & : x \neq 0 \\ 0 & : x = 0 \end{cases}$, then $f(x)$ is

A. continuous and differentiable at $x = 0$

B. continuous but not differentiable at $x = 0$

C. differentiable but not continuous at $x = 0$

D. none of these

Answer: D



Watch Video Solution

21. If the total number of positive integral solution of $15 < x_1 + x_2 + x_3 \leq 20$ is k, then the value of $\frac{k}{100}$ is equal to



Watch Video Solution

22. If $3 \tan^{-1} \left(\frac{1}{2 + \sqrt{3}} \right) - \tan^{-1} \cdot \frac{1}{3} = \tan^{-1} \cdot \frac{1}{x}$,

then the value of x is equal to



Watch Video Solution

23. If the straight lines

$$x + 2y = 3, 2x + 3y = 5 \text{ and } k^2x + ky = -1$$

represent a triangle which is right - angled, then the

value of k are k_1 and k_2 . The value of $\left| \frac{k_1 + k_2}{k_1 - k_2} \right|$ is

 [Watch Video Solution](#)

24. Two lines $\frac{x - 1}{2} = \frac{y - 2}{3} = \frac{z - 3}{4}$ and

$\frac{x - 4}{5} = \frac{y - 1}{2} = \frac{z}{1}$ intersect at a point P. If the

distance of P from the plane $2x - 3y + 6z = 7$ is λ

units, then the value of 49λ is equal to

 [Watch Video Solution](#)

25. The area (in sq. units) bounded by $y = 2^x$ and $y = 2x - x^2$ from $x = 1$ to $x = 2$ is $k \log_2 e - l$, then the value of $\left| \frac{k}{l} \right|$ is equal to



Watch Video Solution