

**MATHS****BOOKS - NTA MOCK TESTS****NTA JEE MOCK TEST 76****Mathematics**

1. The domain of the function $f(x) = 2 \sin^{-1} \left\{ \log_2 \left(\frac{1}{2} x^2 \right) \right\}$ is

A. $[-2, -1] \cup (1, 2]$

B. $(-2, -1] \cup [1, 2]$

C. $[-2, -1] \cup [1, 2]$

D. $(-2, -1) \cup (1, 2)$

Answer: C



[Watch Video Solution](#)

2. Which of the following is a function whose graph is symmetrical about the origin ?

A. $f(x) = (2^x + 2^{-x})$

B. $f(x) = \left[\log \left(x + \sqrt{1 + x^2} \right) \right]^2$

C. $f(x + y) = f(x) + f(y) \forall x, y \in R$

D. None of these

Answer: C

[Watch Video Solution](#)

3. Let the function $f(x) = x^2 \sin\left(\frac{1}{x}\right)$, $\forall x \neq 0$ is continuous at $x = 0$.

Then, the value of the function at $x = 0$ is

A. 0

B. -1

C. 1

D. indeterminate

Answer: A



Watch Video Solution

4. All the students of a class performed poorly in Mathematics. The teacher decided to give grace marks of 12 to the entire class. Which of the following statistical measures will not change even after the grace marks were given?

A. Median

B. Mode

C. Variance

D. Mean

Answer: C

 [Watch Video Solution](#)

5. If p, q and r are simple propositions with truth values T, F and T , respectively, then the truth value of $(\sim p \vee q) \wedge \sim r \rightarrow p$ is

- A. True if the truth values of p, q, r are T, F, T respectively
- B. False if the truth values of p, q, r are T, F, T respectively
- C. False if the truth values of p, q, r are T, F, F respectively
- D. True if the truth values of p, q, r are F, T, F respectively

Answer: A

 [Watch Video Solution](#)

6. A bag contains 5 white balls, 3 black balls, 4 yellow balls. A ball is drawn from the bag, its colour is noted and put back into the bag with 5

additional balls of the same colour. The process is repeated. The probability that a yellow ball is drawn in the 1st draw given that a white ball is drawn in the 2nd draw is

A. $\frac{9}{55}$

B. $\frac{4}{17}$

C. $\frac{1}{5}$

D. $\frac{2}{11}$

Answer: B



Watch Video Solution

7. Let $P(1, 7, \sqrt{2})$ be a point and the equation of the line L is $\frac{x-1}{\sqrt{2}} = \frac{y-7}{1} = \frac{z-\sqrt{2}}{-1}$. If PQ is the distance of the plane $\sqrt{2}x + y - z = 1$ from the point P measured along a line inclined at an angle 60° with L, then the length of PQ is

A. 3 units

B. $3\sqrt{2}$ units

C. 6 units

D. 8 units

Answer: C

 [Watch Video Solution](#)

8. Let $x = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix}$ be a matrix. If $X^6 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then the number of divisors of $(a + b + 2020c + d)$ is equal to

A. 7

B. 14

C. 21

D. 28

Answer: B

 [Watch Video Solution](#)

9. Let $A = \begin{bmatrix} x & 2 & -3 \\ -1 & 3 & -2 \\ 2 & -1 & 1 \end{bmatrix}$ be a matrix and $|\text{adj}(\text{adj}A)| = (12)^4$, then

the sum of all the values of x is equal to

A. -24

B. 24

C. -18

D. 1

Answer: C

 [Watch Video Solution](#)

10. Let the coordinates of two points P and Q be $(1, 2)$ and $(7, 5)$ respectively. The line PQ is rotated through 315° in clockwise direction about the point of trisection of PQ which is nearer to Q . The equation of the line in the new position is

A. $2x - y - 6 = 0$

B. $x - y - 1 = 0$

C. $3x - y - 11 = 0$

D. $3x - y - 9 = 0$

Answer: C



Watch Video Solution

11. If the circles $(x - 3)^2 + (y - 4)^2 = 16$ and $(x - 7)^2 + (y - 7)^2 = 9$ intersect at points A and B, then the area (in sq. units) of the quadrilateral C_1AC_2B is equal to (where, C_1 and C_2 are centres of the given circles)

A. 6

B. 12

C. 18

D. 24

Answer: B



Watch Video Solution

12. If the eccentricity of the hyperbola $x^2 - y^2 \sec^2 \alpha = 5$ is $\sqrt{3}$ times the eccentricity of the ellipse $x^2 \sec^2 \alpha + y^2 = 25$, then $\tan^2 \alpha$ is equal to

A. 2

B. 1

C. 3

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

13. Number of complex numbers satisfying equation

$z^3 = \bar{z}$ & $\arg(z + 1) = \frac{\pi}{4}$ simultaneously is

A. i

B. $1 + 2i$

C. $2 + 3i$

D. $3 + 4i$

Answer: A



Watch Video Solution

14. The coefficient of x^6 in $\{(1+x)^6 + (1+x)^7 + \dots + (1+x)^{15}\}$

is

A. ${}^{16}C_9$

B. ${}^{16}C_5 - {}^{16}C_5$

C. ${}^{16}C_6 - 1$

D. ${}^{16}C_6$

Answer: A

 [Watch Video Solution](#)

15. The position vector of a point P is $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ where $x, y, z \in \mathbb{N}$ and $\vec{a} = \hat{i} + \hat{j} + \hat{k}$. If $\vec{r} \cdot \vec{a} = 10$, then the number of possible position of P is

A. 36

B. 72

C. 66

D. 54

Answer: A

 [Watch Video Solution](#)

16. If the equation $2x^2 - 7x + 9 = 0$ and $ax^2 + bx + 18 = 0$ have a common root, then $(a, b \in \mathbb{R})$

A. $a = 2, b = -7$

B. $a = \frac{-7}{2}, b = 1$

C. $a = 4, b = -14$

D. $a = 4, b = -7$

Answer: C



Watch Video Solution

17. Consider the integral $I_n = \int_0^{\frac{n}{4}} \frac{\sin(2n-1)x}{\sin x} dx$, then the value of $I_{20} - I_{19}$ is

A. $\frac{1}{20}$

B. $\frac{-1}{19}$

C. $\frac{-1}{25}$

D. $\frac{1}{19}$

Answer: B



Watch Video Solution

18. The area (in sq. units) bounded between $y = 3 \sin x$ and $y = -4 \sin^3 x$ from $x = 0$ to $x = \pi$ is

A. 4π

B. 34π

C. 4

D. $\frac{34}{3}$

Answer: D



Watch Video Solution

19. If $f(x)$ is a differentiable function satisfying $|f'(x)| \leq 2 \forall x \in [0, 4]$ and $f(0) = 0$, then

A. $f(x) = 18$ has no solution in $x \in [0, 4]$

B. $f(x) = 18$ has more than 2 solutions in $x \in [0, 4]$

C. $f(x) = 14$ has 3 solutions in $x \in [0, 4]$

D. $f(x) = 20$ has 2 solutions in $x \in [0, 4]$

Answer: A



Watch Video Solution

20. The equation of the curve satisfying the differential equation

$\frac{dy}{dx} + \frac{y}{x^2} = \frac{1}{x^2}$ and passing through $\left(\frac{1}{2}, e^2 + 1\right)$ is

A. $y = e^x + 1$

B. $y = e^{\frac{1}{x}} - 1$

C. $y = 1 + e^{\frac{1}{x}}$

D. $y = 1 + e^{-x}$

Answer: C



Watch Video Solution

21. If from the top of a tower, 60 meters high, the angles of depression of the top and floor of a house are 3° and 60° respectively, then the height (in meters) of the house is

 [Watch Video Solution](#)

22. Let $\vec{p}, \vec{q}, \vec{r}, \vec{s}$ are non-zero vectors in which no two of them are perpendicular and no three of them are coplanar. If $(\vec{p} \times \vec{r}) \cdot (\vec{p} \times \vec{s}) + (\vec{r} \times \vec{p}) \cdot (\vec{q} \times \vec{s}) = k [(\vec{p} \times \vec{q}) \cdot (\vec{s} \times \vec{r})]$, then the value of $\frac{k}{2}$ is equal to

 [Watch Video Solution](#)

23. A point P on the parabola $y^2 = 4x$, the foot of the perpendicular from it upon the directrix and the focus are the vertices of an equilateral triangle. If the area of the equilateral triangle is β sq. units, then the value of β^2 is



[Watch Video Solution](#)

24. Let the sets $A = \{2, 4, 6, 8, \dots\}$ and $B = \{3, 6, 9, 12, \dots\}$ such that $n(A) = 200$ and $n(B) = 250$. If $n(A \cup B) = k$, then $\frac{k}{100}$ is equal to



[Watch Video Solution](#)

25. The area (in sq. units) bounded by the curve $e^x y - 2 = 0$ with the x-axis from $x = 0$ to $x = \ln 2$ is



[Watch Video Solution](#)