



MATHS

BOOKS - NTA MOCK TESTS

NTA JEE MOCK TEST 79

Mathematics

1. Three balls are marked with 2, 4 and 6. They are placed in a box and a ball is drawn, its number is noted and the ball drawn is returned to the box. The process is repeated two more times. If the probability to the sum of all three numbers is 12 is λ , then 108λ is equal to

A. 28

B. 14

C. 20

D. 10

Answer: A



Watch Video Solution

2. The number of 2×2 matrices A with real entries, such that $A + A^T = 3I$ and $AA^T = 5I$, is equal to

A. 0

B. 1

C. 2

D. infinite

Answer: C



Watch Video Solution

3. The sum of the intercepts of the plane on the coordinate axes, passing through the intersection of the planes

$$2x + 3y + 3z - 5 = 0 \quad \text{and} \quad 2x - 5y + 3z + 1 = 0 \quad \text{and}$$

parallel to the line $\frac{x - 1}{2} = \frac{y - 2}{-5} = \frac{z - 3}{-7}$, is

A. $\frac{2}{5}$

B. $\frac{11}{105}$

C. $\frac{11}{102}$

D. $\frac{3}{101}$

Answer: C



Watch Video Solution

4. If $A_r = \begin{bmatrix} \frac{1}{r(r+1)} & \frac{1}{3^r} \\ 2 & 3 \end{bmatrix}$, then $\lim_{n \rightarrow \infty} \sum_{r=1}^n |A_r|$ is equal to

A. 0

B. 2

C. 4

D. 8

Answer: B



Watch Video Solution

5. The solution of the equation $\cos^{-1} x + \cos^{-1} 2x = \frac{2\pi}{3}$

is

A. $x = \pm \frac{\sqrt{3}}{\sqrt{28}}$

B. $x = \pm \frac{\sqrt{3}}{\sqrt{26}}$

C. $x = \pm \frac{1}{\sqrt{2}}$

D. $x = \pm \frac{\sqrt{3}}{\sqrt{38}}$

Answer: A



Watch Video Solution

6. Negation of "Monu is in class X or Anu is in class XII" is

A. Monu is not in class X and Anu is in class XII

B. Monu is not in class X and Anu is not in class XII

C. Either Monu is not in class X or Anu is not in class XII

D. None of these

Answer: B



Watch Video Solution

7. The value of $\lim_{x \rightarrow 0} (\cos x + \sin bx)^{\frac{a}{x}}$ is equal to

A. e^{-ab}

B. e^{ab^2}

C. e^{ab}

D. e^{-b^2a}

Answer: C

 [Watch Video Solution](#)

8. If $f(x) = [x]\tan(\pi x)$ then $f'(k^+)$ is equal to (where k is some integer and $[.]$ denotes greatest integer function)

 [Watch Video Solution](#)

9. Consider the integrals

$$I = \int \frac{\sin x}{3 \cos x + 4 \sin x} dx \quad \text{and} \quad J = \int \frac{\cos x}{3 \cos x + 4 \sin x} dx.$$

Then, the integral $4I - 3J$ is equal to (where, c is the constant of integration)

A. $\ln|2 \cos x - 4 \sin x| + c$

B. $\ln|3 \cos x + 4 \sin x| + c$

C. $\ln|3 \sin x + 4 \cos x| + c$

D. $\ln|3 \sin x - 4 \cos x| + c$

Answer: B



Watch Video Solution

10. If the area bounded by $f(x) = \tan^3 x + \tan x$ from $x = 0$ to $x = \frac{\pi}{4}$ is k square units, then the maximum value of

$g(x) = k \sin x$ is $\left(\forall x \in \left[0, \frac{\pi}{4} \right] \right)$

A. 2

B. $\frac{1}{2\sqrt{2}}$

C. 4

Answer: B



Watch Video Solution

11. For the function $f(x) = \sin^3 x - 3 \sin x + 4 \forall x \in \left[0, \frac{\pi}{2}\right]$, which of the following is true?

A. Greatest value of the function is 2π

B. Greatest value of the function is 4

C. Rolle's Theorem is applicable to $f(x)$ in $x \in \left[0, \frac{\pi}{2}\right]$

D. LMVT is not applicable to $f(x)$ in $x \in \left[0, \frac{\pi}{2}\right]$

Answer: B



Watch Video Solution

12. The curve satisfying the differential equation $\sin(x^3)e^y dy + 3x^2 \cos(x^3)e^y dx = x \sin(x^2) dx$ C is the constant of integration is $\lambda \sin(x^3)e^y + \cos(x^2) = C$.

Then, the value of λ is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

13. A possible three digit even number which can be formed with the condition that If 5 is one of the digit, then 7 is next digit is :

A. 5

B. 325

C. 345

D. 265

Answer: D



[Watch Video Solution](#)

14. The discriminant of the quadratic $(2x + 1)^2 + (3x + 2)^2 + (4x + 3)^2 + \dots n$ terms

$= 0, \forall n > 3, x \in R$ is

A. positive

B. zero

C. negative

D. depends on n

Answer: C



Watch Video Solution

15.

Simplify

$$P = \frac{1}{2\sqrt{1} + \sqrt{2}} + \frac{1}{3\sqrt{2} + 2\sqrt{3}} + \dots + \frac{1}{100\sqrt{99} + 99\sqrt{100}}$$

A. $\frac{1}{10}$

B. $\frac{3}{10}$

C. $\frac{9}{10}$

D. $\frac{1}{2}$

Answer: C



Watch Video Solution

16. The general solution of the equation

$\sin^2 x + \cos^2 3x = 1$ is equal to :

(where $n \in I$)

A. $x = \frac{n\pi}{3}$

B. $x = n\pi + \frac{\pi}{4}$

C. $x = \frac{n\pi}{4}$

D. $x = n\pi + \frac{\pi}{2}$

Answer: C

 Watch Video Solution

17. The line $L_1: \frac{x}{5} + \frac{y}{b} = 1$ passes through the point (13, 32) and is parallel to $L_2: \frac{x}{c} + \frac{y}{3} = 1$. Then, the distance between L_1 and L_2 is

- A. $\frac{17}{\sqrt{15}}$ units
- B. $\frac{23}{\sqrt{17}}$ units
- C. $\frac{23}{\sqrt{17}}$ units
- D. $\frac{23}{\sqrt{15}}$ units

Answer: C

 Watch Video Solution

18. B & C are fixed points having co-ordinates $(3, 0)$ and $(-3, 0)$ respectively. If the vertical angle BAC is 90° , then the locus of the centroid of the triangle ABC has the equation :

A. $x^2 + y^2 = 1$

B. $x^2 + y^2 = 2$

C. $9(x^2 + y^2) = 1$

D. $9(x^2 + y^2) = 4$

Answer: A



Watch Video Solution

19. The product of the slopes of the common tangents of the ellipse $x^2 + 4y^2 = 16$ and the parabola $y^2 - 4x - 4 = 0$ is

A. $-\frac{1}{15}$

B. $\frac{1}{16}$

C. $-\frac{1}{3}$

D. $-\frac{1}{2}$

Answer: B



Watch Video Solution

20. For a complex number Z , if

$|Z - i| \leq 2$ and $Z_1 = 5 + 3i$, then the maximum value of

$|iZ + Z_1|$ is (where, $i^2 = -1$)

A. $7 + \sqrt{13}$

B. $7 + \sqrt{12}$

C. 7

D. $\sqrt{34} - 2$

Answer: C



Watch Video Solution

21. Consider a parabola $y^2 = 4x$ with vertex at A and focus at S. PQ is a chord of the parabola which is normal at point P. If the abscissa and the ordinate of the point P are equal, then the square of the length of the diameter of the circumcircle of triangle PSQ is



Watch Video Solution

22. if $\sum \frac{k}{k+1} \cdot {}^{100}C_k = \frac{a \cdot 2^{100} + b}{c}$ where $a, b, c \in N$

then find the least value of $a + b + c$



Watch Video Solution

23. The value of the integral $I = \int_0^{\frac{\pi}{2}} \frac{\sin^3 x - \cos^3 x}{1 + \sin^6 x \cos^6 x} dx$ is

equal to



Watch Video Solution

24. If $\sqrt{x+y} + \sqrt{y-x} = 2$, then the value of $\frac{d^2y}{dx^2}$ is

equal to



Watch Video Solution

25. Let \vec{a} and \vec{b} are vectors such that

$$|\vec{a}| = 2, |\vec{b}| = 3 \text{ and } \vec{a} \cdot \vec{b} = 4. \quad \text{If}$$

$$\vec{c} = \left(3\vec{a} \times \vec{b} \right) - 4\vec{b}, \text{ then } |\vec{c}| \text{ is equal to}$$



Watch Video Solution