

BIOLOGY

BOTANY AND ZOOLOGY FOR NEET AND AIIMS

PLANT GROWTH AND DEVELOPMENT

Exercise I General Growth

1. The maximum growth phase is

- A. Lag phase
- B. Steady phase
- C. Log phase
- D. Senescent phase



- **2.** Plant growth is .
 - A. Irreversible

- B. Increase in size
- C. Localised
- D. All the above



Watch Video Solution

3. Correcet sequence of differnet phase of growth is

A. Cell division, Cell differentiaiton Cell elongation B. Cell differntiation, Cell division Cell elongation

C. Cell elongation , Cell division , Cell differntiation

D. Cell division, Cell elongation, Cell differentation

Answer: D



4. The correct sequence of stages , through which the plants passes during its life span.

A. Seed
$$\;
ightarrow\;$$
 plantlet $\;
ightarrow\;$ seedling $\;
ightarrow\;$ mature plants

B. Seedling
$$ightarrow$$
 seed $ightarrow$ plantlet $ightarrow$ mature plant

C. Plantlet
$$ightarrow$$
 seeding $ightarrow$ seed $ightarrow$

mature plant

D. Seed ightarrow seedling ightarrow plantlet ightarrow mature plant

Answer: D



Watch Video Solution

5. Identify the worng statement form the following .

A. Both growth and differenation are open in plants

- B. Plants body in made up of derivatives of meristems
- C. Secondary growth of leaves is due to the activity of lateral meristemg
- D. Apical meristems of shoot and root tip cause primary growth



6. Which of the following can not be takne as parmeter of measuring the growth in plant body

- A. Fresh (or) dry weight of plants body
- B. Thickness of leaf lamina
- C. Surface area of leaf lamina
- D. Length of root (or) stem

Answer: B



7. The Shape of growth curve obtained when the length of an organ is plotted against the time is

A. Parabola

B. S' Shape

C. Linear

D. Step wise

Answer: C



8. The exponential growth can be expressed as

A.
$$Lt = L_0 + rt$$

B.
$$W_1=W_0e^{rt}$$

C.
$$W_1 = W_1 + rt$$

D.
$$\log_e\!\left(rac{W_0}{W_1}
ight) = rt$$

Answer: B



9. The absolute growth rate and relative growth rates of living systems are.

A. Quantiative comparisons

B. Qualitative comparison

C. Both 1 and 2

D. None

Answer: A



10. During exponential phase of geometrical growth

A. Cell elongate but do not divide

B. Cells divide but do not elongate

C. Both the progeny cells of mitotic

division retain the ability of cell division

D. Cell degenerate due to deficiency of nutrients

Answer: C



- 11. Water is essential for growth for .
- (A) Englargement of cells.
- (B) Turgidity of cells for extension of growth.
- (C) Providing medium for enzymeatic activites .
- (D) The source of energy.
 - A. All correct expept 'A'
 - B. All correct expept 'B'
 - C. All correct expept 'C'
 - D. All correct expept 'D'



- **12.** $W_1=W_0$ ert is the expression of exponential growth , in which r denotes .
 - A. Initial growth
 - B. Final growth
 - C. Relative growth rate
 - D. Base of natural logarithms



- **13.** The very first even in the sequential stage of development of a higher plant cell is
 - A. Elongation
 - B. Plasmatic growth
 - C. Maturation
 - D. Senescence

Answer: B



- **14.** The ability of producing differnet kinds of structures at differnet phases of life is called
 - A. Plasticity
 - B. Re-differnatiation
 - C. Plasmacity
 - D. Maturation

Answer: A



Watch Video Solution

15. Heterophyllous aquatic plants of the following is

- A. Cotton
- B. Coriander
- C. Larkspur
- D. Buttercup



Watch Video Solution

16. Pick out the correct statement from the following .

A. The parenchyma cells undergo differnetiation during plant tissue culture

- B. The tissues arising out of same meristem have always same structures at maturity
- C. The products of meristematic cells retain

 the power of cell division by

 redifferentiation
- D. Epidermins is a product of differntiation



17. Intracellular intrinsic factor that influence the development in plants is

- A. Genetical
- **B.** Chemical
- C. Physical
- D. Nutritional

Answer: A



18. Exponential phase of growth among the following is .

A. Lag phase

B. Senescence phase

C. Logarithmic phase

D. Steady state

Answer: C



19. The kind of growth in unicelluar organims like bacteria is .

- A. Logarithmic
- B. Arithmietic
- C. Geometric
- D. Plasticity

Answer: C



20. The growth in plants in measured by .
A. Pfeiffer's auxanometer
B. Osmometer
C. Manometer
D. Respirometer
Answer: A



Watch Video Solution

21. Development is the sum of

- A. Cell division and cell elongation
- B. Cell division and cell differenation
- C. Growth and differentiation
- D. Growth and de differentiation



Watch Video Solution

22. Growth is measured by the following parameter

- A. Increase in fresh wieght, dry weight & length
- B. Increase in cell number , cell volume and area
- C. Increase in cell volume and dry wieght
- D. Increase in cell number, volume, area length fresh weight and dry weight.



23. Wound healing in the matured part of the secondary stem occurs due to .

- A. Differentiation
- B. De-differneation
- C. Re differenatiation
- D. none of the above

Answer: B



- 24. Plasticity refers to

 A. Ability of the plants to undergo different
 - pathways of metabolism suitable to the enbironments
 - B. Ability to exhibit different phase of life to produce different types of structures
 - C. Ability of plants to grow through out their life
 - D. 1 & 2

Exercise I Plant Growth Regulartors General

1. The important of coleoptile tip in phototropic curvautre was first experimentally confirmed by

A. Julius von Sachs

B. Charles Darwn and Francis Darwin

C. Boysen - Jenson

D. F.W.Went

Answer: B



Watch Video Solution

2. Which one of the following pairs is not correctly matched ?

A. Adenine derivative - kinetin

B. Carotenoid derivat

C. Terpense - IAA

D. Indole compounds - IBA

Answer: C



Watch Video Solution

3. Growth promoter hormones are

A. IAA, ABA and CK

B. IAA, GA and ABA

C. IAA, GA and CK

D. ABA, Ck and GA



Watch Video Solution

4. A green plants bends towards the source of light when exposed to the light on only one side . Which of the following is the best explanation of the phenomenon?

A. The apices of the stems are attached by light

- B. The plants needs light for photosynthesis
- C. Auxin accumulates on the shaded side to induce great cell elongation on that side
- D. Ligh reduces length.



- **5.** All the following pairs are growth promoting activites except
 - A. Tropic growth and pattern formation
 - B. Flowering and fruiting
 - C. Seed formation and seed germination
 - D. Dormancy and Abscission



- 6. FW. Went is associated with
 - A. Isolation of auxins from the coleptiles tips of oat seedlings
 - B. Isolation of auxins form the coleoptiles tips of canary grass seedling
 - C. Isolution of Gibberellins form Gibberalla fuzikuroi
 - D. Isolation of auxins form the coleoptiles of both canary grass and oat seedlings

Answer: A



Watch Video Solution

7. Which one of the following is not a natural polymer

- A. ABA
- B. Zeatin
- C. 2,4-D
- D. N6 furfuryl amino purine



- **8.** Choose the incorrect statement form the following
 - A. Cytokinis show specific effect on cytokinesis
 - B. ABA is antagonistic phytohomone to

 Gibberellic acids

C. Natural cytokinins are sythesized in root apices only

D. Ethylene increase absorption surface of root system

Answer: C



Watch Video Solution

9. Plants hormomes are .

A. Growth regulators

- B. Growth promotors
- C. Growth inhibitors
- D. All of these

Answer: D



Watch Video Solution

10. Which one of the following is a natural growth inhibitor?

A. NAA

- B. ABA
- C. IAA
- D. GA

Answer: B



Watch Video Solution

11. Which one of the following is not correctly matched?

A. Promote cell division - kinetin

- B. Promote senescence GA
- C. Promote floweing Auxins
- D. Promote Abscission Ethylene

Answer: B



- 12. Identify the mis mathc of following?
 - A. Adenine derivative kinetin
 - B. Carotenoid derivative ABA

C. Terpenes - IAA

D. Indole compounds - IBA

Answer: C



Watch Video Solution

Exercise I Auxins

1. Phytohormone which inhibits flowering in serveral plants but stimulates flowering in Pineapple plants is .

- A. 2-4 D
- B. IBA
- C. GA_3
- D. Ethylene

Answer: D



Watch Video Solution

2. Identify a pair of hormoes whose effets are opposite to each other regarding apical dominace

- A. Auxins and ABA
- B. Auxins and Cytokinins
- C. Gibberellins and Cytokinins
- D. Ethylene and ABA

Answer: B



- 3. Removal of apical bud makes the plants
 - A. Bushy

- B. Grow rapidly
- C. Grow slowly
- D. Dermant



- **4.** 2,4-D stands for
 - A. 2,4 Dichlorophenoxy Acetic Acid
 - B. 2,4- D Dichloro Butryic Acid

- C. 2,4- Dichloronaphthoxy Acetic Acid
- D. 2,4- Dichloronaphthalene Acetic Acid



- **5.** Auxins were first isolated from
 - A. Coleoptiles of canary grass
 - B. Coleoptiles of canary Avena
 - C. Rice seedling

D. Human urine

Answer: D



Watch Video Solution

6. Which of the following physiological activity is not induced by Auxins in plants

- A. Rooting in stem cutting
- B. Xylen differnation
- C. Induction of parthencarpy

D. Induction of abscission of yong leaves and furits

Answer: D



Watch Video Solution

7. Which of the following physiological activity is not induced by Auxins in plants?

A. Indole - 3 -butyric acid

B. Napthalene acetic acid

- C. 2,3-5 tri chloro phenoxy acetic acid
- D. 2,4 dichloro phenoxy acetic acid



- **8.** Which of the following is not a functional attribute of auxins ?
 - A. Promote premature fruit and leaf fall
 - B. Promote mature fruit and leaf fall

- C. Parthenocarpy
- D. Controls xylen differentiaion



Watch Video Solution

9. The harmone widely used in tea plantations for hedge making

- A. ABA
- **B.** Auxins

 $\mathsf{C}.\,GA_3$

D. Ethylene

Answer: B



Watch Video Solution

10. The modified form of adenine is

A. 2,4 - Dichlorophenoxy acetic acid

B. N6 - furfuryl amino purine

C. 2-chloro ethyl phosphonic acid

D. Dichloro phenyl methyl urea

Answer: B



Watch Video Solution

11. Removal of apical bud during prusing results in the

- A. Death of plant
- B. Early flowering
- C. Promotion of lateral branching

D. Promotion of lateral branching

Answer: C



Watch Video Solution

12. A high concentration of synthetic auxins is generally used for .

- A. Regulating the growth of roots
- B. Weed control
- C. Controlling cell enlargement

D. Preventing the growth of lateral shoots

Answer: B



Watch Video Solution

13. Which one prevents premature fall of fruits

A. IAA

B. Ethylene

 $\mathsf{C}.\,GA_3$

D. Zeatin



Watch Video Solution

14. Which of the following is not an influence of auxins?

- A. Apical dominace
- B. Parthenocarpy
- C. Tropic movements
- D. Bolting

Answer: D



Watch Video Solution

15. For plants tissue culture which among the following is required ?

- A. Trypsin
- B. Kinetin
- C. Caffeine
- D. Cocumarin

Answer: B



Watch Video Solution

Exercise I Gibberellins

- **1.** Bolting is associated with physiological function of
 - A. Gibberellic acid
 - B. Cytokinins
 - C. IAA

D. ABA

Answer: A



Watch Video Solution

2. Bakane disease is caused by

A. Fungus

B. Alga

C. Bacterium

D. Virus



Watch Video Solution

- 3. Gibberellins were first extracted from
 - A. Coeloptile tip
 - B. Root tip
 - C. Fungus
 - D. Bacterium

Answer: C

4. Gibberellic acid was discovered in connection with a disease in rice seedling, caused by a

A. Virus

B. Bacterium

C. Fungus

D. Nematoda

Answer: C

5. Which of the following phytohormones is used to speed up the malting process in brewing industry?

A. GA_1

B. GA_2

C. GA_3

D. GA_4

Answer: C



6. The 'PGR' which induces increase in the length of grapes stalks

A. IAA

B. Gibberellins

C. ABA

D. Ethylene

Answer: B



Watch Video Solution

7. Which of the following is called Antigibberellin?

A. Auxin

B. ABA

C. Ethylene

D. Cytokinin

Answer: B



8. Foolish seedling disease of Rice led to the discovery of

A. GA

B. ABA

C. 2,4-D

D. IAA

Answer: A



9. Which of the PGRs	induces parthenocarpy in
Tomatoes ?	

- A. Auxin
- B. Gibberellins
- C. Cytokinin
- D. Ethylene



10. Cell elongation in internodal regions of the green plants takes place due to

- A. Indole acetic acid
- B. Gibberellic acid
- C. Cytokinins
- D. Ethylene

Answer: B



11. The rosette habit of cabbage can be changed by application of

- A. IAA
- B. GA
- C. ABA
- D. Ethephon

Answer: B



12. The Phenomenon of elongation of stem in the plants with rosette habit is know as

- A. Rosetting
- B. Bolting
- C. Phase of elongation
- D. 1 and 3

Answer: B



13. The hormones which are found to be more effective in inducing parthenocarpy are

- A. Cytokinis
- **B.** Auxins
- C. Ethylene
- D. Ethephon

Answer: B



Exercise I Cytokinins

- 1. Cytokinins
 - A. Promote abscission
 - B. Influence water movement
 - C. inhibit protoplasmic streaming
 - D. help retain chlorophyll

Answer: D



=: The hist hataral cytokinin or plants is	2.	The	first	natural	cyto	okinin	of	plants	is
---	----	-----	-------	---------	------	--------	----	--------	----

- A. zeatin
- B. kinetin
- C. dihydrooxyzeation
- D. riboxylzeatin



3. Cytokinins are mostly produced in

A. Shoot apex

B. Root apex

C. old leaves

D. Ripe fruits

Answer: B



- 4. Cytokinins are generally
 - A. Acids
 - B. Aminopurines
 - C. Phenols
 - D. Glucosides

Answer: B



5. PGR which induces the formation of new leaves and chloroplast in leaves is

- A. Cytokinins
- B. Ethylene
- C. Gibberellins
- D. Abscisic acid

Answer: D



6. One hormone helps to produce new leaves, while the other promotes abscission of leaves .

A. ABA and Auxins

These are respectively

B. Cytokinins

C. Gibberellins and cytokinins

D. Etylene and Gibbenllins

Answer: B



7. Natural cytokinis are synthesized in tissue that one

- A. Senescent
- B. Dividing rapidly
- C. Storing food material
- D. Differentating

Answer: B



8. Hormone primarily connected with cell division in

- A. IAA
- B. NAA
- C. Cytokinin/zeation
- D. Gibberellic acid

Answer: C



9. Which of the following is a cytokin

A. Phytochrome

B. Leucine

C. Ethylene

D. Zeatin

Answer: D



10. Cut or excised leaves remain green for long if induced to root or dippped in .

- A. Gibberellins
- B. Cytokinins
- C. Auxins
- D. Ethylene

Answer: B



Exercise I Ethylene

1. Name the gaseous hromone which stimulates transverse or isodiametirc growth

A. Ethylene

B. ABA

C. Sodium salt of NAA

D. Ethephon

Answer: A



2. Ethylene is

- A. Neither a growth promoter nor growth inhibitor
- B. A nitrogenous phytohormone
- C. A gaseous phytohormone that hastens fruits ripening
- D. A volatile phytohormone that caused wounds in plants body

Answer: C



Watch Video Solution

3. Phytohormone responsible for 'Respiratory climactic' is .

A. ABA

B. Ethylene

 $\mathsf{C}.\,GA_3$

D. IAA

Answer: B



Watch Video Solution

- 4. Ethylene is highly effective in
 - A. Root growth and root hair formation
 - B. Abscission of palnts organs
 - C. Apical hook formation in dicot seedlings
 - D. Fruit ripening

Answer: D

- **5.** Which of the followign is not the physiological effect of Ethylene?
 - A. Increasing absorption surface of root
 - B. Elongation of apple fruits
 - C. Enahances rate of respiration during furit ripening
 - D. Petiole elongation in deep water rice

plants

Answer: B



- **6.** On hormone helps in ripening of fruits, while the other sitmulates closure of stomata. These are respectively.
 - A. ABA and Auxin
 - B. Ethylene and ABA
 - C. ABA and Ethylene
 - D. GA and ABA

Answer: B



Watch Video Solution

- 7. The gaseous furit ripening phytohromone is
 - A. Ethylene
 - B. Kinetin
 - C. GA
 - D. ABA

Answer: A

8. Ethylene is liberated during

A. Developing seeds

B. Ripening in some furits

C. Developing ovules

D. Germinating seeds

Answer: B



Exercise I Aba

1. A grwoth inhibitor responsible for stomatal closure is

A. Zeatin

B. Abscisic acid

C. Ethylene

D. GA_3

Answer: B

- 2. Abcisic acid promotes
 - A. Cell division
 - B. Dormacy
 - C. Shoot elongation
 - D. Cell elongation and wall formation

Answer: B



3. Which one is antagonist to Gas

A. NAA

B. Zetain

C. ABA

D. Ethylene

Answer: C



4. The dormancy of potato tubers can be increased by the spray of .

A. Auxins and Gibberellins

B. Gibberellins

C. Auxins and cytokinins

D. ABA

Answer: D



5. Closure of stomata is brought about by.

A. Abscisic acid

B. Kinetin

C. Giberellin

D. IBA

Answer: A



Watch Video Solution

Exercise I Dormancy

1. A seed unable to germinate because the external conditions are not favourable is called .

A. Cilmacteric C

B. Quiescence

C. Dormancy

D. Scarification

Answer: B



2. Seed dormancy is due to the presence of hard seed coats, in the members of .

A. Fabaceae

B. Solanaceae

C. Liliacae

D. Malvaceae

Answer: B



Exercise I Photoperiodism

1. In some plants, flowering is either Quantitatively or Qualitatively dependent on exposure to low temperature and it is called

A. Vernalisation

B. Dormancy

C. Quiescence

D. Photoperiodism

Answer: A

2. The phenomenon of the influence of relative duration of day night on flowering is called

A. Vernalisation

B. Photobleaching

C. Photoropism

D. Photoperidism

Answer: D



3. In flowering plants, the site of perception of light/ dark duration is

- A. Leaves
- B. Buds
- C. Branches
- D. Stem

Answer: A



4. Which is not affected by light?

A. Photosynthesis

B. Flowering

C. Fertilization

D. Transpiration

Answer: C



5. Effect of day length duration duration of plant development is .

A. Chemotropism

B. Phototropism

C. Photoperidism

D. Photonasty

Answer: C



6.	Long	day	plants	blossom	in
----	------	-----	--------	---------	----

- A. Summer
- B. Autumn
- C. Spring
- D. 1 and 3

Answer: A



Watch Video Solution

Exercise I Vernalization

1. F	lowering	dependent	on	low	temperature
ехр	osure is				

- A. Vernalisation
- B. Thermotropy
- C. Cryoscopy
- D. Cryostat

Answer: A



- 2. Vernalisation is
 - A. Growth curve related to light
 - B. Effect of photoperiods of plants growth
 - C. Speeding up ability to flower by low temperature treatment
 - D. Diurnal photoperiodicity

Answer: C



Exercise Ii Auxins

1. Who experimentally proved that the auxins were responsible for the apical dominace

A. L.G. Paleg

B. Thimann and Skoog

C. Skoog and Miller

D. Burg and thiman

Answer: B



2. The importance of coleoptile tip in phototropic curvature was first experimentally confirmed by

A. Juslius von Sachs

B. Charles Darwin and Francis Darwin

C. Boysen - Jenson

D. F.W.Went

Answer: B



3. Which one of the following statement concerning growth regulators is incorrect.

A. Gibberellins cause the formation of more male flowers in Cannabis

B. Abscisic acid manitains seeds in a dormant state

C. Cytokinins increase the K + concentration in guard cells

D. Some synthetic auxins control grass weeds in pulse crops

Answer: D



Watch Video Solution

4. Study the following statements and select the correct ones

A. Auxin is transported from base to tip in large quantities

- B. Cells on the dark side expand farther than those on the light side
- C. Apical dominace can be removed by the decapitation
- D. Tropic movements in pants are under the control of gibberellins

Answer: C



5. Which one of the following pairs is not correctly matched ?

A. Adenine derivative - kinetin

B. Carotenoid derivative - ABA

C. Terpenes - IAA

D. Indole compounds - IBA

Answer: C



6. Darwing discovered that canary grass coleoptile bends towards light only when .

A. It is 5 cm long

B. The seedling is growing in blue light

C. Coeloptile tip is intact and exposed to

light

D. Cooler nights

Answer: C



7. Removal of apical bud	makes the	plants
---------------------------------	-----------	--------

- A. Bushy
- B. Grow rapidly
- C. Grow slowly
- D. Dormant

Answer: A



- 8. Bioassay for auxin is
 - A. Dwarf maize test
 - B. Avena curvautre test
 - C. Cell division test
 - D. Green leaf test

Answer: B



- 9. 2,4-D stands for
 - A. 2,4-Dichlorophenoxy Acetic Acid
 - B. 2,4-Dichloro Butyric Acid
 - C. 2,4 Dichlornaphthoxy Acetic Acid
 - D. 2,4-D-chloronaphthalene Acetic Acid

Answer: A



10. Hormone that break dormancy of potato tuber is

- A. IAA
- B. ABA
- C. Zeatin
- D. Gibberellin

Answer: A



11. Abscission layeris formed when the concentration of

- A. Auxin increase
- B. Auxin decreases
- C. Gibberellins decreases
- D. Gibberellins increases

Answer: B



Exercise Ii Gibberellins

1. Bolting is associated with physiological function of

A. Gibberelic acid

B. Cytokinins

C. IAA

D. ABA

Answer: A



2. Physiological function of the phytohormone , that is synthesized from acetly co - enzyme A molecules is

A. Promotion of seed germination

B. Promotion of apical dominance

C. Removal of apical dominance

D. Promotion of triple response growth

Answer: A



3. In generminating cereals, amylase synthesis is stimulated by

A. Cytokinins

B. Abscisic acid

C. Gibberellins

D. Auxin

Answer: C



4. Which of the following is not related to Gibberelin

A. Bolting of rosette plants

B. Replacing long day requirement

C. Removal of genetic dwarfism

D. Bending movement of coleoptile

Answer: D



5. Genetic Dwarfness can be controlled by treating plants with

- A. Cytokinin
- B. Gibberellic acid
- C. Auxin
- D. Antigibberellin

Answer: B



6. The Phenomenon of elongation of stem in the plants with rosette habit is know as

- A. Rosetting
- B. Bolting
- C. Phase of elongation
- D. 1 and 3

Answer: B



- 7. Which one is the test for gibberellin?
 - A. Bolting in cabbage
 - B. Morphogenesis in carrot cells
 - C. Rapid divisions in carrt cells
 - D. Elongation of oat coleoptile

Answer: A



8. Exogenous GA induce formation of male flowers on genetically female plants of

- A. Cucurbita
- B. Luffa
- C. Cucumis
- D. Carica

Answer: C



9. The hormones which are found to be more effective in inducing parthenocarpy are

- A. Cytokinins
- B. Gibberellims
- C. Ethylene
- D. Ethephon

Answer: B



10. Gibberellic acid induces flowering in

A. Some plants only

B. In long day plants under short day conditions

C. In short day plants under long day condition

D. Day neutral plants

Answer: B



11. Bakane disease is caused by

- A. Fungus
- B. Alga
- C. Bacterium
- D. Virus

Answer: A



12. Gibberellins were first extracted from

- A. Coeloptile tip
- B. Root tip
- C. Fungus
- D. Bacterium

Answer: C



Watch Video Solution

Exercise Ii Cytokinins

Watch Video Solution							
-							

2. The first cytokinin was isolated by

1. Richmond'- Long effect is related to .

- A. Skoog (1970)
- B. Evins, 1971
- C. Miller etal (1954)
- D. Leopoid (1969)

Answer: C



Watch Video Solution

3. Cytokinins

A. Promote abscission

- B. Influence water movement
- C. inhibit protoplasmic streaming
- D. help retain chlorophyll

Answer: D



- **4.** The first natural cytokinin of plants is
 - A. Zeatin
 - B. kinetin

- C. dihydrooxyzeation
- D. Riboxylzeatin

Answer: A



- 5. Cytokinins are mostly produced in
 - A. Shoot apex
 - B. Root apex
 - C. Young leaves

D. Lateral buds

Answer: B



Watch Video Solution

6. Cytokinins

- A. Induce cell division and inhibit ageing
- B. Maintain dormancy
- C. Induce abission
- D. Inhibit cell division

Answer: A



Watch Video Solution

- 7. Cytokinins are generally
 - A. Acids
 - **B.** Aminopurines
 - C. Phenols
 - D. Glucosides

Answer: B

8. Knowledge about cytokinins was gained after skoog' work on

A. Tomato pith

B. Tobacco pith

C. Rice seedling

D. Avena curvature test

Answer: B



Watch Video Solution

9. Hormone primarily connected with cell division in

A. IAA

B. NAA

C. Cyokinin/zeatin

D. Gibberllic acid

Answer: C



10. Which of the following is a cytokin

- A. Phytochrome
- B. Leucine
- C. Ethylene
- D. Zeatin

Answer: D



11. The compounds having a highly specific hydro-phhilic group are named as

- A. Auxins
- **B.** Cytokinins
- C. Gibberellins
- D. Ethylene

Answer: B



12. Cut or excised leaves remain green for long if induced to root or dippped in .

- A. Gibberellins
- B. Cytokinins
- C. Auxins
- D. Ethylene

Answer: B



13. Maximum concentration of cytokinins occurs in

A. Grwoing embryos an fruits

B. Apical buds

C. Lateral buds and fruits

D. Root tips

Answer: A



14. Morphogenesis is controlled by an interaction between

- A. Auxins and gibberllins
- B. Auxins and cytokinins
- C. Gibberellins and cytokinins
- D. None of these above

Answer: B



15. The phenomenon of delay of senescence by cytokinins is known as

- A. Richmond lang effect
- B. Senescence of effect
- C. Loomis and Torry effect
- D. Sorokin et al effect, 1962

Answer: A



16. The regular which retards ageing of plants parts is

- A. Cytokinin
- B. Auxin decreases
- C. Gibberellins
- D. Abscisis acid

Answer: A



17. The hormones which regulate phloem transport are

- A. Auxins
- **B.** Cytokinins
- C. Gibberellins
- D. Ethylene

Answer: B



18.	For	plants	tissue	culture	which	among	the
fol	lowi	ng is re	quired	?			

- A. Trypsin
- B. Kinetin
- C. Caffeine
- D. Cocumarin

Answer: B



- A. Roots
- **B.** Leaves
- C. Shoot tip
- D. Fruit

Answer: A



20. Phytohormone which inhibits flowering in serveral plants but stimulates flowering in Pineapple plants is .

- A. 2-4, D
- B. IBA
- C. GA3
- D. Ethylene

Answer: D



21. Name the gaseous hromone which stimulates transverse or isodiametirc growth

- A. Ethylene
- B. ABA
- C. Sodium salt of NAA
- D. Ethephon

Answer: A



22.	The	gaseous	furit	ripening	phytohromone	3
is						

- A. Ethylene
- B. kinetin
- C. GA
- D. ABA

Answer: A



23. Ethylene is a by product of

- A. Developing seeds
- B. Ripening in some furits
- C. Developing ovules
- D. Germinating seeds

Answer: B



24. Which is not affected by light?

A. Photosynthesis

B. Flowering

C. Fertilization

D. Transpiration

Answer: C



25. Ethylene is liberated during

- A. Developing seeds
- B. Ripening in some furits
- C. Developing ovules
- D. Germinating seeds

Answer: B



26. Ethylene gas

- A. Is saturated hydrocarbon
- B. Slows down ripening of apples
- C. Retards ripening of tomatoes
- D. Speeds up maturation of furits

Answer: A



1. Part of the plant body where Abscisic acid is not synthesized but Cytokinins are sytheszied is

A. Leaves

B. Flower

C. Roots

D. Seeds

Answer: C



2. A grwoth inhibitor responsible for stomatal closure is

A. Zeatin

B. Abscisic acid

C. Giberellin

D. IBA

Answer: B



3. Abcisic acid promotes

- A. Cell division
- B. Leaf fall senescence and dormacy
- C. Shoot elongation
- D. Cell elongation and formation

Answer: B



4.	Closure	of stomata	is broug	ht about by.
----	---------	------------	----------	--------------

A. Abscisic acid

B. kinetin

C. Giberellin

D. IBA

Answer: A



5. Hormone responsible for ageing is

- A. GA
- B. IAA
- C. ABA
- D. Cytokinin

Answer: C



1. A long day plant is

- A. Wheat
- B. Soyabeam
- C. Tobacco
- D. Xanthium

Answer: C



- 2. Pick up the correct explanation
 - A. Xanthium Long dry plant
 - B. Sunflower short day plant
 - C. Wheat Short day plant
 - D. Tomato Day neutral plant

Answer: D



- 3. In short day plants, flowering is induced by
 - A. Photoperiod less than 12 hours
 - B. Photoperiod below a critical length and uninterrupted long night
 - C. Long night
 - D. Long day

Answer: B



4. DNF	is '
---------------	------

- A. Cestrum
- B. Cucumber
- C. Potato
- D. Radish

Answer: B



Watch Video Solution

5. Short day plants flower in

- A. Spring
- B. Summer
- C. Autumn
- D. 1 and 2

Answer: A



Watch Video Solution

6. Effect of day length duration duration of plant development is .

- A. Chemotropism
- B. Phototropism
- C. Photoperidism
- D. Photperiodism

Answer: B



Watch Video Solution

7. The most effetive wavelenght of light for flowering are

- A. Blue and red
- B. Red and far red
- C. Green
- D. Orange and red

Answer: B



- **8.** Phytochrome is
 - A. Light receptor

- B. Dark receptor
- C. Photosynthetic pigment
- D. A light sensitive hormone

Answer: A



Watch Video Solution

9. Phytcohrome is involved in picking stimulus in

A. Phototropism

- B. Phototropism
- C. Photorespiration
- D. Photosynthesis

Answer: C



- **10.** Phytochrome is
 - A. Protein
 - B. Pigment

- C. light sensitive movement
- D. Hormone induced process

Answer: A



Watch Video Solution

11. Pfr interconversion is regulated by.

A. ABA

B. 2,4-D

C. IAA

D. GA

Answer: D



Watch Video Solution

12. Phytochrome in plants is sensitive to

- A. Gree light
- B. Blue light
- C. Red light
- D. Red and far red light

Answer: D



Watch Video Solution

13. Photoperiodism is probably due to the sythesis of

- A. Cytokinins
- B. Gibberllins
- C. Auxin
- D. Florigen

Answer: D



Watch Video Solution

14. The hormone capable of replacing the requirment of long photoperiods for flowering is

- A. Ethylene
- B. Auxin
- C. Gibberellin
- D. Cytokinin

Answer: C



Watch Video Solution

Exercise Ii Vernalization

- **1.** Flowering dependent on low temperature exposure is
 - A. Vernalisation
 - B. Thermotropy
 - C. Cryoscopy

D. Cryostat

Answer: A



- 2. The term " vernalization" was coined by
 - A. Garner and Allard
 - B. Darwin
 - C. Geoffery
 - D. Lysenko

Answer: D



Watch Video Solution

3. Vernalisation is

- A. Growth curve related to light
- B. Effect of photoperiods of plants growth
- C. Speeding up ability to flower by low temperature treatment
- D. Diurnal photoperiodicity

Answer: C



Watch Video Solution

- 4. Vernalisation can often be replaced by
 - A. Auxin
 - **B.** Cytokinins
 - C. Gibberellins
 - D. Ethylenis

Answer: C

- 5. The stimulus responsible for vernalization is
 - A. Phytochrome
 - B. Vernalin
 - C. Florigen
 - D. None of these

Answer: B



6. The perception of stimulus of cold treatment/ vernalization is received by

- A. Leaves
- B. Shoot apex
- C. Axillary buds
- D. Stem

Answer: B



7. Low temperature required for vernalization is usually between .

A.
$$10-10^{\circ}$$
 C

B.
$$5-15^{\circ}C$$

C.
$$1-30^{\circ}C$$

D.
$$1-5^{\circ}C$$

Answer: D



8. The plant substance which become inhibitory on accmulation are .

A. Abscisic acid

B. Phenolic inhibitors

C. Auxins

D. Cytokinis

Answer: B



9. In autumn, leaf fall occurs because

A. Formation , of abscission layer at the base of leaves

B. Leaf becomes heavy

C. Leaf does not remain green

D. Of low temperature

Answer: A




Exercise li Pgr S

- 1. Growth promoter hormones are
 - A. IAA, ABA and CK
 - B. IAA, GA and ABA
 - C. IAA, GA and CK
 - D. ABA, CK and GA

Answer: A



2. Match the following.



The correct match is .

Answer: C



Watch Video Solution

3. Which amongst the following is a natural growth regulator?

A. NAA

B. Ethylene

C. 2,4-D

D. Benzaldehyde

Answer: B



- **4.** Sprouting of potato can be prevented in storage by
 - A. Malic hydrazide
 - B. Gibberellins
 - C. Indole acetic acid
 - D. Cytokinins

Answer: A



Watch Video Solution

Exercise Iii Previous Aipmt Neet Questions

- 1. Root hairs develop from the rgion of :
 - A. Maturation
 - B. Elongation
 - C. Root cap

D. Meristematic activity

Answer: A



Watch Video Solution

2. Fruit and leaf drop at early states can be prevented by the application of ::

A. Cytokinins

B. Ethylene

C. Auxins

D. Gibberellic acid

Answer: C



Watch Video Solution

3. The avena curvature is used for biosassay of

A. Ethylene

B. ABA

 $\mathsf{C}.\,GA_3$

D. IAA

Answer: D



Watch Video Solution

- 4. Auxins can be bioassayed by
 - A. Potometer
 - B. Lettuce hypocotly elongation
 - C. Avean coleoptile curvautre
 - D. Hydroponics

Answer: C

5. What causes a green plant exposed to the light, on only one side, to bend towards the source of light as it grows?

A. Light stimlates plant cells on the lighted side to grow faster

B. Auxin accumulates on the shaded side, stimulating

- C. Green plants need light to perform photosynthesis
- D. Green plants seek light because the are phototropic .

Answer: B



Watch Video Solution

6. Typical growth curve in plants is

A. Stair -steps

- B. Parabolic
- C. Sigmoid
- D. Linear

Answer: C



Watch Video Solution

7. Which one of the following growth regulators is know as strees hormone?

A. Abscisic acid

B. Ethylene

 $\mathsf{C}.\,GA_3$

D. Indole acetic acid

Answer: A



Watch Video Solution

8. Dr. F.Went note that if coleoptile tips were removed and planced on agar for one hour, the agar would produce a bending when placed on one side of freshly -cut coleoptile

stumps. Of what significance is this experiment?

A. It made possible the isolation and exact identification of auxin

B. It is the basic for quantiative determination of small amounts of growth - promoting substance .

C. It supports the hydpothesis that IAA is auxin

D. It demonstrated polar movement of auxins

Answer: A



Watch Video Solution

9. During seed germination its stored food is mobilized by

A. ABA

B. Gibberllin

- C. Ethylene
- D. Cytokinin

Answer: B



- 10. Vernalisation stimulates flowering in
 - A. Zamikand
 - B. Turmeric
 - C. Carrot

D. Ginger

Answer: C



Watch Video Solution

11. Throught their effects on plants growth regulator, what do the temperature and light control in the plants?

A. Apical dominace

B. Flowering

- C. Closure of stomata
- D. Fruit elongation .

Answer: B



Watch Video Solution

12. Which one of the following generally acts as an antagonist to gibberllins?

- A. Zeatin
- B. Ethylene

C. ABA

D. IAA

Answer: C



Watch Video Solution

13. Root development is promoted by

A. Abscisic acid

B. Auxin

C. Gibberellin

D. Ethylen

Answer: D



Watch Video Solution

14. One of the commoly used plant growth hormone is tea plantation is

A. Ethylene

B. Abscisic acid

C. Zeatin

D. Indole - 3 - acetic acid

Answer: D



Watch Video Solution

15. Coiling of graden pea tendrils around any support is an example of

A. Thigmotaxis

B. Thigmonasty

C. Thigmotropism

D. Thermotaxis

Answer: C



Watch Video Solution

16. Phototropic curvature is the result of uneven distribution of

- A. Gibberellin
- B. Phytochrome
- C. Cytokinins

D. Auxin

Answer: D



Watch Video Solution

17. Which one of the following acids is a derivative of carotenoids?

A. Indole - 3-acetic acid

B. Gibberellic acid

C. Abscisic acid

D. Indole butyric acid

Answer: C



Watch Video Solution

18. One of the synthetic auxn is

A. IAA

B. GA

C. IBA

D. NAA

Answer: D



- **19.** Senescence as an active developmental celluar process in the growth , and funcitoning of a flowering plant, is indicated in
 - A. Annual plants
 - B. Floral parts
 - C. Vessesl and tracheid differenatation
 - D. Leaf abscission

Answer: C



Watch Video Solution

20. Importance of day length in flowering of plants was first shown in

- A. Cotton
- B. Petunia
- C. Lemna
- D. tobacco

Answer: D



Watch Video Solution

21. The wavelenght of light absorbed by P_r from of phytochrome is

- A. 680 nm
- B. 720 nm
- C. 620 nm
- D. 640 nm

Answer: A



- **22.** Which one of the following pairs , is not correctly matched ?
 - A. Gibberellic acid Leaf fall
 - B. Cytokinin Cell division
 - C. IAA -Cell wall elongation
 - D. Abscisic acid Stomatal closure

Answer: A



Watch Video Solution

23. Opening of floral buds into flowers, is a type of

- A. autonomic movement of variation
- B. paratonic movement of growth
- C. autonomic movement of growth
- D. autonomic movement of locomotion

Answer: C



Watch Video Solution

24. An enzyme that can stimulate germination of barley seeds is

A. invertase

B. lpha - amylase

C. lipase

D. protease

Answer: B



Watch Video Solution

25. Farmers in a particular region were concerned that premature yellowing of leaves of a pulse corp mightcause decrease in the yield. Which treatment could be most beneficial to obtain maximum seed yield?

A. Application of iron magnesium to promote synthesis of chlorophyll

- B. Frequent irrigation of the crop
- C. Treatment of the plans with cytokinins along with a small does of nitrogenous fertilizer
- D. Removal of all yellow leaves and spraying the remaining green leaes with 2,4,5-trichlorophenozy acetic acid .

Answer: C



- **26.** How does prunning help in making the hedge dense?
 - A. It relases wound hormones
 - B. It induces the differentiation of new shoots from the rootstock
 - C. It frees axillary buds from apical dominace.
 - D. The apical shoot grow faster after pruing.

Answer: C



Watch Video Solution

27. Treatment of seeds at low temperature under moist conditions to break its dormancy is called .

A. stratification

B. scarification

C. vernalization

D. chelation

Answer: A

