



## PHYSICS

### BOOKS - U-LIKE PHYSICS (HINGLISH)

### CBSE SAMPLE QUESTION PAPER 2019-20 (SOLVED)

#### Section A

1. A charge  $q$  is placed at the point of intersection of body diagonals of a cube. The electric flux passing through any one of its face is

A.  $\frac{q}{6 \epsilon_0}$

B.  $\frac{3q}{\epsilon_0}$

C.  $\frac{6q}{\epsilon_0}$

D.  $\frac{q}{3 \epsilon_0}$

**Answer: A**



[View Text Solution](#)

2. The electric potential of earth is taken to be zero because earth is a good

A. insulator.

B. conductor.


C. semiconductor.

D. dielectric.

**Answer: B**

 [View Text Solution](#)

3. If the ammeter in the given circuit shown in the diagram reads 2 A, the resistance R is



A.  $1\Omega$

B.  $2\Omega$

C.  $3\Omega$

D.  $4\Omega$

**Answer: A**



[View Text Solution](#)

4. The heat producted by 100 W heater in 2 mintues is equal to

A. 10.5 kj

B. 16.3 kj

C. 12.0 kj

D. 14.2 kJ

**Answer: C**



**View Text Solution**

5. Time period of a charged particle undergoing a circular motion in a uniform magnetic field is independent of

- A. speed of the particle.
- B. mass of the particle.
- C. charge of the particle.
- D. magnetic field.

**Answer: A**



**View Text Solution**

6. The final image formed in an astronomical refracting telescope with respect to the object is

- A. real inverted.
- B. real erect.
- C. virtual erect.
- D. virtual inverted.

**Answer: D**



**View Text Solution**

7. The shape of the interference fringes in Young's double - slit experiment when  $D$  (distance between slit and screen) is very large as compared to fringe width is nearly

A. straight line.

B. parabolic.

C. circular

D. hyperbolic

**Answer: A**



**View Text Solution**

8. Unpolarised light is incident on a plane surface having refractive index  $\sqrt{3}$ . The angle of incidence at which reflected and refracted rays would become perpendicular to each other is

A.  $15^\circ$

B.  $30^\circ$

C.  $45^\circ$

D.  $60^\circ$

**Answer: D**



[View Text Solution](#)



9. Photoelectric emission from a given surface of metal can take place when the value of a 'physical quantity' is less than the energy of incident photon. The physical quantity is

- A. Threshold frequency.
- B. Work function of surface.
- C. Threshold wavelength.
- D. Stopping potential.

**Answer: B**



[View Text Solution](#)

10. A photon beam of energy 12.1 eV is incident on a hydrogen atom. The orbit to which electron of H-atom be excited is

A. 2nd orbit.

B. 3rd orbit

C. 4th orbit.

D. 5th orbit.

**Answer: B**



**View Text Solution**

**11.** Note : Fill in the blanks with appropriate answer :

Horizontal and vertical components of earth's magnetic field at a place are equal. The angle of dip at that place is \_).



[View Text Solution](#)

**12.** A free floating magnetic needle at North pole is \_\_\_\_\_ to the surface of earth.



[View Text Solution](#)

13. The magnetic flux linked with a coil changes by  $2 \times 10^{-2}$  Wb when the current by 0.01 A.

The self - inductance of the coil is \_\_\_\_\_.



[View Text Solution](#)

14. If the angular speed of the armature of a dynamo is doubled then the amplitude of the induced e.m.f. will become \_\_\_\_\_.



[View Text Solution](#)

15. An electron is accelerated through a potential difference of 100 V, then de-Broglie wavelength associated with it is approximately \_\_\_\_\_ Å.

 [View Text Solution](#)

16. An equilateral prism is made up of material of refractive index  $\sqrt{3}$ . The angle of minimum deviation of light passing through the prism is \_\_\_\_\_.

 [View Text Solution](#)

17. Which physical quantity in a nuclear reaction is considered equivalent to the  $Q$  - value of the reaction?

 [View Text Solution](#)

18. Zener diode is used in reverse bias. When its reverse bias is increased, how does the thickness of the depletion layer change?

 [View Text Solution](#)

19. The initial concentration of a radioactive substance is  $N_0$  and its half life is 12 hours. What will be its

concentration after 36 hours?



[View Text Solution](#)

20. Work function of Sodium is 2.75 eV. What will be K.E. of emitted electron when photon of energy 3.54 eV is incident on the surface of sodium ?



[View Text Solution](#)

21. From the information of energy band gaps of diodes, how do you decide which can be light emitting diodes?



[View Text Solution](#)

22. Give any one advantage of LEDs over conventional incandescent low power lamps.



[View Text Solution](#)

## Section B


1. Derive the expression from drift velocity of free electron in terms of relaxation time and electric field applied across a conductor.



[View Text Solution](#)



2. Find total energy stored in capacitors given in the circuit.



[View Text Solution](#)

3. An  $\alpha$  - particle and a proton are accelerated through same potential difference. Find the ratio.  $(v_\alpha / v_p)$  of velocities acquired by two particles.

[View Text Solution](#)

4. What is Brewster's angle ? Derive relation between Brewster angle and refractive index of medium which produces Plane Polarised light.



[View Text Solution](#)

5. The work function of Cs is 2.14 eV. Find

(a) threshold frequency for Cs.

(b) Wavelength of incident light if the photo current is brought to zero by stopping potential of 0.6V.



[View Text Solution](#)

6. Derive the expression for the radius of  $n$ th Bohr's orbit in Hydrogen atom.

 [View Text Solution](#)

7. Energy of electron in first excited state in Hydrogen atom is  $-3.4\text{eV}$ . Find K.E. and P.E. of electron in the ground state.

 [View Text Solution](#)

8. Draw energy band diagram of p and n type semiconductors. Also write two difference between p

and n type semiconductors.



[View Text Solution](#)

9. Energy gap in p - n photodiode is 2.8 eV. Can it detect a wavelength of 6000 nm ? Justify your answer.




[View Text Solution](#)

## Section C

1. State working principle of potentiometer. Explain how the balance point shifts when value of resistor R

increases in the circuit of potentiometer, given below.



[View Text Solution](#)

2. Using Biot - Savart's law, derive an expression for magnetic field at any point on axial line of a current carrying circular loop. Hence, find magnitude of magnetic field intensity at the centre of circular coil.

[View Text Solution](#)

3. Obtain the resonant frequency and  $Q$  - factor of a series LCR circuit with  $L = 3H$ ,  $C = 27\mu F$ ,  $R = 7.4\Omega$ .

It is desired to improve the sharpness of resonance of circuit by reducing its full width at half maximum by a factor of 2. Suggest a suitable way.



[View Text Solution](#)


4. State the conditions of total internal reflection.

Refractive indices of the given prism material for Red,

Blue and Green colours are respectively 1.39, 1.48 and

1.42 respectively. Trace the path of rays through the

prism.



[View Text Solution](#)

5. Define resolving power of an astronomical refracting telescope and write expression for it in normal adjustment. Assume that light of wavelength  $6000\text{\AA}$  is coming from a star, what is the limit of resolution of a telescope whose objective has a diameter of 2.54 m?



[View Text Solution](#)

6. Write the basic assumptions used in the derivation of lens- maker's formula and hence derive this expression.



[View Text Solution](#)

7. Show that  ${}_{92}^{238}U$  cannot spontaneously emit a proton.

Given :

$${}_{92}^{238}U = 238.05079u, {}_{91}^{237}Pa = 237.05121u, {}_1^1H = 1.00783u$$



[View Text Solution](#)



8. Suggest an ideal to convert a full wave bridge rectifier to a half wave rectifier by changing the connecting wire/s. Draw the diagram and explain your answer.




[View Text Solution](#)

## Section D

1. (a) Using Gauss's law, derive expression for intensity of electric field at any point near the infinitely long straight uniformly charged wire.

(b) The electric field components in the following figure are  $E_x = \alpha x$ ,  $E_y = 0$ ,  $E_z = 0$ , in which

$\alpha = 400\text{N/Cm}$ . Calculate (i) the electric flux through the cube, and (ii) the charge within the cube assume that  $a = 0.1\text{m}$ .




[View Text Solution](#)

2. (a) Define electrostatic potential at a point. Write its SI unit.

Three charges  $q_1, q_2$  and  $q_3$  are kept respectively at points A, B and C as shown in figure Write the expression for electrostatic potential energy of the

system.




(b) Depict the equipotential surfaces due to

(i) an electric dipole.

(ii) two identical negative charges separated by a small distance.

 [View Text Solution](#)

3. In the following diagram, the arm PQ of the rectangular conductor is moved from  $x = 0$ , outwards.



The uniform magnetic field is perpendicular to the plane and extends from  $x = 0$  to  $x = b$  and is zero for  $x > b$ . Only the arm PQ possesses substantial resistance 'r'. Consider the situation when the arm PQ is pulled outwards from  $x = 0$  to  $x = 2b$ , and then moved back to  $x = 0$  with constant speed 'v'. Obtain expressions from the (i) magnetic flux, (ii) the induced emf, (iii) the force necessary to pull the arm and (iv) the power dissipated as joule heat.

Sketch the variation of these quantities with distance.



[View Text Solution](#)

4. Write working principle of cyclotron and with a suitable diagram explain its working. Give any two application of cyclotron.



[View Text Solution](#)

5. Derive mirror equation for a convex mirror. Using it, show that a convex mirror always produces a virtual image, independent of the location of object.



[View Text Solution](#)

6. (a) Draw a ray diagram for final image formed at distance of distinct vision ( $D$ ) by a compound microscope and write expression for its magnifying power.

(b) An angular magnification (magnifying power) of  $30\times$  is desired for a compound microscope using as objective of focal length  $1.25\text{ cm}$  and eye piece of focal length  $5\text{ cm}$ . How will you set up the compound microscope?



[View Text Solution](#)