

## CHEMISTRY

### BOOKS - NARENDRA AWASTHI

#### DILUTE SOLUTION

##### Exercise

1. The vapour pressure of a give liquid will decrease if :

- A. surface area of liquid decreased
- B. the volume of liquid in the container is decreased
- C. the volume of the vapour phase is increased

D. the temperature is decreased

**Answer: d**

 [Watch Video Solution](#)

2. The normal boiling point of water is  $373\text{K}$  . Vapour pressure of water at temperature  $T$  is  $19\text{ mmHg}$  . If enthalpy of vaporization is  $40.67\text{kJ/mol}$ , then temperature  $T$  would be

- A.  $250\text{ K}$
- B.  $291.4\text{ K}$
- C.  $230\text{ K}$
- D.  $290\text{ K}$

**Answer: B**

 [Watch Video Solution](#)

3. A sample of the liquid  $H_2O$  at 18.0 g is injected into an evacuated 7.6 L flask maintained at  $27.0^\circ C$ . If vapour pressure of  $H_2O$  at  $27.0^\circ C$  is 24.63 mm Hg, what weight percentage of the water will be vaporised when the system comes to equilibrium? Assume water vapours behaves as an ideal gas. The volume occupied by the liquid water is negligible compared to the volume of the container:

A. 0.01

B. 0.1

C. 0.18

D. 0.2

**Answer: a**

 [Watch Video Solution](#)

4. Raoult's law is obeyed by each constituent of a binary liquid solution when :

A. the forces of attraction between like molecules are greater than those between unlike molecules

B. the forces of attraction between like molecules are smaller than those between unlike molecules

- C. the forces of attraction between like molecules are identical with those between unlike molecules
- D. the volume occupied by unlike molecules are different

**Answer: c**

 [Watch Video Solution](#)

5. For a binary ideal liquid solution, the total vapour of the solution is given as:

A.  $P_{\text{total}} = P_A^\circ + (P_A^\circ - P_B^\circ)X_B$

B.  $P_{\text{total}} = P_B^\circ + (P_A^\circ - P_B^\circ)X_A$

$$C. P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) X_A$$

$$D. P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) X_B$$

**Answer: b**

 [Watch Video Solution](#)

6. For an ideal liquid solution with  $P_A^\circ > P_B^\circ$ , which relation between  $X_A$  ((mole fraction of A in liquid phase) and  $Y_A$  (mole fraction of A in vapour phase) is correct ?

A.  $Y_a < Y_b$

B.  $X_A > X_B$

C.  $\frac{Y_A}{Y_B} > \frac{X_A}{X_B}$

$$D. \frac{Y_A}{Y_B} < \frac{X_A}{X_B}$$

**Answer: C**

 **Watch Video Solution**

7.  $X_A$  and  $X_B$  are the mole fraction of A and B respectively in liquid phase  $y_A$  and  $y_B$  are the mole fraction of A and B respective in vapour phase. Find out the slope of straight line if a graph is plotted  $\frac{1}{y_A}$  along Y-axis against  $\frac{1}{x_A}$  along X-axis gives straight line [ $p_A^\circ$  and  $p_B^\circ$  are vapour pressure of pure components A and B].

A.  $\frac{P_B^\circ}{P_A^\circ}$

B.  $\frac{P_A^\circ}{P_B^\circ}$

C.  $P_B^\circ - P_A^\circ$

D.  $P_A^\circ - P_B^\circ$

**Answer: a**

 [Watch Video Solution](#)

**8.** For a dilute solution, Raoult's law states that:

A. the lowering of vapour pressure is equal to the mole fraction solute

B. the relative lowering of vapour pressure is equal to the mole fraction of solute



C. the relative lowering of vapour pressure is proportional to the amount of solute in solution

D. the vapour pressure of the solution is equal to the mole fraction of solvent

**Answer: b**

 [Watch Video Solution](#)

9. The solubility of a specific non-volatile salt is 4 g in 100 g of water at  $25^{\circ}C$ . If 2.0g, 4.0g and 6.0 g of the salt added of 100 g of water at  $25^{\circ}$ , in system X, Y and Z. The vapour pressure would be in the order:

A.  $X < Y < Z$

B.  $X > Y > Z$

C.  $Z > X = Y$


D.  $X > Y = Z$

Answer: d



Watch Video Solution


10. The pair of boiling point and compound are given as,






$80^\circ C$     $65^\circ C$     $212^\circ C$     $184^\circ C$

*I*   *II*   *III*   *IV*

Which will show lowest vapour pressure at room temperature





**Answer: B**



**Watch Video Solution**

11. 6.0 g of urea (molecular mass = 60) was dissolved in 9.9 moles of water. If the vapour pressure of pure water is  $P^\circ$ , the vapour pressure of solution is :

A.  $0.10 P^\circ$

B.  $1.10 P^\circ$

C.  $0.90 P^\circ$

D.  $0.99 P^\circ$

**Answer: d**

 [Watch Video Solution](#)

**12.** An ideal solution was found to have a vapour pressure of 80 torr when the mole fraction of a non-volatile solute was 0.2. What would be the vapour pressure of the pure solvent at the same temperature?

A. 64 torr

B. 80 torr

C. 100 torr

D. 400 torr

**Answer: C**



**Watch Video Solution**

**13.** If the vapor pressure of a dilute aqueous solution of glucose is  $750\text{mm}$  of Hg at  $373\text{K}$ , then molality of solute is

A. 0.26

B. 0.73

C. 0.74

D. 0.039

**Answer: c**

 [Watch Video Solution](#)

14. Estimate the lowering of vapour pressure due to the solute (glucose) in a 1.0 M aqueous solution at  $100^{\circ}C$ :

- A. 10 torr
- B. 18 torr
- C. 13.45 torr
- D. 24 torr

**Answer: c**

 [Watch Video Solution](#)

15. Calculate the weight of non – volatile solute having molecular weight 40, which should be dissolved in 57gm octane to reduce its vapour pressure to 80 % :

A. 47.2 g

B. 5 g

C. 106.2 g

D. None of these

**Answer: b**



**Watch Video Solution**

16. Equal mass of a solute are dissolved in equal mass of two solvents A and B and formed very dilute solution. The relative lowering of vapour pressure for the solution B has twice the relative lowering of vapour pressure for the solution A. If  $m_A$  and  $M_B$  are the molecular mass of solvents A and B respectively, then :

A.  $M_A = M_B$

B.  $M_B = 2 \times M_A$

C.  $M_A = 4M_B$

D.  $M_A = 2M_B$

**Answer: b**



**Watch Video Solution**



17. For an ideal solution of two components A and B, if  $x_A$  and  $y_A$  are mole fractions of component 'A' in solution and vapour phase respectively, then the slope of linear line in the graph drawn between  $1/x_A$  and  $1/y_A$  is

A.  $X_A = Y_A$

B.  $X_A > Y_A$

C.  $X_A < Y_A$

D. Data insufficient

**Answer: c**



**Watch Video Solution**

18. At  $25^{\circ}C$ , the vapour pressure of pure liquid A (mol. Mas = 40) is 100 torr, (mol. = 80). The vapour pressure at  $25^{\circ}C$  of a solution containing 20 g of each A and B is :

- A. 80 torr
- B. 59.8 torr
- C. 68 torr
- D. 48 torr

**Answer: a**



[Watch Video Solution](#)

19. Two liquids A and B form an Ideal solution. At 300K, the V.P of solution containing one mole of 'A' and 4 mole

'B' is 560mm Hg. At the same temp. If one mole of 'B' is taken out from the solution the V.P of the solution has decreased by 10mm Hg, the V.P, of pure A & B are (in min)

A. 400600

B. 500 , 500

C. 600 , 400

D. none of these

**Answer: a**



[Watch Video Solution](#)

**20.** Two liquids A and B have vapour pressure in the ratio

$P_A^0 : P_B^0 = 1 : 3$  at a certain temperature . Assume A and B

from an ideal solution and the ratio of mole fraction of A to B in the vapour phase is 4 : 3 . Then the mole fraction of B in the solution at the same temperature is :

A.  $\frac{1}{5}$

B.  $\frac{2}{3}$

C.  $\frac{4}{5}$

D.  $\frac{1}{4}$

**Answer: a**



[Watch Video Solution](#)

**21.** Two liquids A and B have  $P_A^\circ$  and  $P_B^\circ$  in the ratio of 1 : 3 and the ratio of number of moles of A and B in liquid

phase are 1 : 3 then mole fraction of 'A' in vapour phase in equilibrium with the solution is equal to :

A. 0.1

B. 0.2

C. 0.5



D. 1

**Answer: a**



[Watch Video Solution](#)

**22.** Which represents correct difference when non-volatile solute is present in an ideal solution?



A. I, II, III

B. I, III

C. II, III

D. I, II

**Answer: a**



**Watch Video Solution**

**23. Select correct statement :**

- A. Solution has more molecules randomness than a pure solvent. The entropy change between solution and solid is larger than the entropy change between pure solvent and solid
- B. Heat of fusion of solution are positive
- C. Solution containing sugar freezes at a lower temperature than pure water
- D. All are correct statements

**Answer: d**



**Watch Video Solution**

24. Select correct statement :

A. Heats of vaporisation for a pure solvent and for a solution are similar because similar intermolecular forces between solvent molecules must be overcome in both cases

B. Entropy change between solution and vapour is smaller than the entropy change between pure solvent and vapour

C. Boiling point of the solution is larger than that of the pure solvent


D. All are correct statements



Answer: d

 Watch Video Solution

25. The vapour pressure curves of the same solute in the same solvent are shown below. The curves are parallel to each other and does not intersect. The concentrations of solutions are in order of :



A.  $I < II < III$

B.  $I = II = III$

C.  $I > II > III$

D.  $I > III > II$


**Answer: a**



**Watch Video Solution**

**26.** Boiling point composition diagram of the liquid-vapour equilibrium for A and B is shown in the figure. If a binary liquid mixture of A and B is distilled fractionally, which of

the following would be correct observation ?



$X_A = 1$  % Composition  $X_B = 1$

- A. Composition of the still (residue) will approach pure liquid B only
- B. composition of the distillate will approach pure A only
- C. Composition of distillate and residue will approach pure A and B respectively

D. Neither of the component can be obtained in pure state

**Answer: c**

 [Watch Video Solution](#)

27. The boiling point of an azeotropic mixture of water and ethyl alcohol is less than that of the theoretical value of water and alcohol mixture. Hence the mixture shows

A. the mixture will show negative deviation from Raoult's law

- B. the mixture will show positive deviation from Raoult's law
- C. the mixture can be considered as pure solution
- D. this mixture can be considered as pure solution

**Answer: b**

 [Watch Video Solution](#)

**28.** Formation of a solution from two components can be considered as :

(i) pure solvent  $\rightarrow$  separated solvent molecules,  $\Delta H_1$

(ii) Pure solute  $\rightarrow$  separated molecules,  $\Delta H_2$

(iii) separated solvent and solute molecules  $\rightarrow$  solution,

$$\Delta H_3$$

solution so formed will be ideal if :

A.  $\Delta H_{\text{soln}} = \Delta H_1 + \Delta H_2 + \Delta H_3$

B.  $\Delta H_{\text{soln}} = \Delta H_1 + \Delta H_2 - \Delta H_3$

C.  $\Delta H_{\text{soln}} = \Delta H_1 - \Delta H_2 - \Delta H_3$

D.  $\Delta H_{\text{soln}} = \Delta H_3 - \Delta H_1 - \Delta H_2$

**Answer: a**

 [Watch Video Solution](#)

**29.** Total vapour pressure of mixture of 1 mol X ( $P_X^\circ = 150$  torr) and 2 mol Y ( $P_Y^\circ = 300$  torr) is 240 torr. In this case :

- A. there is a negative deviation from Raoult's law
- B. there is a positive deviation from Raoult's law
- C. there is no deviation from Raoult's law
- D. can not be decided

**Answer: a**



**Watch Video Solution**

**30.** In a mixture of A and B, components show positive deviation when:

A. A-B interaction is stronger than A-A and B-B interaction

B. A-B interaction is weaker than A-A and B-B interaction

C.  $\Delta V_{\text{mix}} < 0$ ,  $\Delta S_{\text{mix}} > 0$

D.  $\Delta V_{\text{mix}} = 0$ ,  $\Delta S_{\text{mix}} > 0$


**Answer: b**

 [Watch Video Solution](#)

**31.** A liquid mixture having composition corresponding to point Z in the figure shown is subjected to distillation at constant pressure. Which of the following statements



is correct about the process?



- A. The composition of distillate differs from the mixture
- B. The boiling point goes on changing
- C. The mixture has highest vapour pressure than for any other composition

D. Composition of an azeotrope alters on changing the external pressure

**Answer: d**

 [Watch Video Solution](#)

**32. Which will form maximum boiling azeotrope ?**

A.  $C_6H_6 + C_6H_5CH_3$  solution

B.  $HNO_3 + H_2O$  solution

C.  $C_2H_5OH + H_2O$  solution

D. n-hexane and n-heptane

**Answer: B**



Watch Video Solution

33. Total vapour pressure of mixture of 1 mole of volatile components A ( $P_{A^{\circ}}=100$  mm Hg) and 3 mole of volatile component B ( $P_{B^{\circ}} = 80$  mm Hg) is 90 mm Hg. For such case:

- A. There is positive deviation from Raoult's law
- B. boiling point has been lowered
- C. force of attraction between A and B is weaker than that between A and A or between B and B
- D. All the above statements are correct

**Answer: d**

34. The azeotropic mixture of water ( $B. P. = 100^{\circ}C$ ) and  $HCl$  ( $B. P. = 86^{\circ}C$ ) boils at about  $120^{\circ}C$ . During fractional distillation of this mixture it is possible to obtain :

- A. pure  $HCl$
- B. pure  $H_2O$
- C. pure  $H_2O$  as well as pure  $HCl$
- D. Neither  $C_2H_5OH$  nor  $HCl$

Answer: d

35. Azeotropic mixture of water and  $C_2H_5OH$  boils at 351

K. By distilling the mixture it is possible to obtain

A. pure  $C_2H_5OH$  only

B. Pure water only

C. Neither  $C_2H_5OH$  nor water

D. Both water and  $C_2H_5OH$  in pure state

**Answer: c**



**Watch Video Solution**

**36.** An azeotropic mixture of two liquids has a boiling point higher than either of them when it :

- A. shows positive deviation from Raoult's law
- B. shows negative deviation from Raoult's law
- C. shows ideal behaviour
- D. is saturated

**Answer: b**



**Watch Video Solution**

**37.** If two liquids A ( $P_A^\circ = 100\text{torr}$ ) and ( $P_B^\circ = 200\text{ torr}$ ) which are completely immiscible with each other (each

one will behave independently of the other) are present in a closed vessel, the total vapour pressure of the system will be :

- A. less than 100 torr
- B. greater than 200 torr
- C. between 100 to 200 torr
- D. 300 torr

**Answer: d**



[Watch Video Solution](#)

**38.** When a liquid that is immiscible with water was steam distilled at  $95.2^{\circ}C$  at a total pressure of 748 torr, the

distilled contained 1.25 g of the liquid per gram of water .

The vapour pressure of water is 648 torr at  $95.2^{\circ}C$ , what

is the molar mass of liquid?

A. 7.975 g/mol

B. 166 g/ mol

C. 145.8 g/mol

D. None of these

**Answer: c**



**Watch Video Solution**

**39.** Which of the following is a colligative property



- A. Vapour pressure
- B. Depression in f.pt.
- C. Elevation in b.pt.
- D. Osmotic pressure

**Answer: a**



**Watch Video Solution**

**40.** The degree of an electrolyte is  $a$  and its Van't Hoff factor is  $i$ . The number of ions obtained by complete dissociation of 1 molecules of electrolyte as :

A. 
$$\frac{i + a - 1}{a}$$

B.  $i = a - 1$

C.  $\frac{i - 1}{a}$

D.  $i = 1 + \frac{a}{1 - a}$

**Answer: a**



**Watch Video Solution**

**41.** One mole of a solute A is dissolved in a given volume of a solvent. The association of the solute takes place as follows:

A.  $i = 1 - a$

B.  $i = 1 + \frac{a}{n}$

C.  $i = \frac{1 - a + \frac{a}{n}}{1}$

D.  $i = 1$

**Answer: C**

 [Watch Video Solution](#)

42. The van't Hoff factor  $i$  for a compound which undergoes dissociation in one solvent and association in other solvent is respectively

A. greater than one and less than one

B. less than one and greater than one

C. less than one and less than one

D. greater than one and greater than one

**Answer: A**

 [Watch Video Solution](#)

**43.** Which solution has the highest vapour pressure ?

A. 0.02 M NaCl at  $50^{\circ}\text{C}$

B. 0.03 M sucrose at  $15^{\circ}\text{C}$

C. 0.005 m  $\text{CaCl}_2$  at  $500^{\circ}\text{C}$

D. 0.005 M  $\text{CaCl}_2$  at  $25^{\circ}\text{C}$

**Answer: c**

 [Watch Video Solution](#)

44. An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase ?

A. addition of water

B. addition of NaCl

C. addition of  $Na_2SO_4$

D. Addition of 1.0 molal KI

**Answer: a**



**Watch Video Solution**

45. Four solutions of  $K_2SO_4$  with the concentrations 0.1m, 0.01, 0.001 m, and 0.0001 m are available. The maximum value of colligative property corresponds to :

- A. 0.0001 m solution
- B. 0.001 m solution
- C. 0.01 m solution
- D. 0.1 m solution

**Answer: d**



**Watch Video Solution**

46. Moles of  $K_2SO_4$  to be dissolved in 12 mol water to lowest its vapour pressure by 10 mm of Hg at a temperature at which vapour pressure of pure water is 50 mm is :

A. 1.5 mole

B. 2 mole

C. 1 mole

D. 3 mole

**Answer: D**



**Watch Video Solution**

47. A very diluted saturated solution of a sparingly soluble salt  $X_3Y_4$  has a vapour pressure of 20 mm Hg at temperature T, while pure water exerts a pressure of 20.0126 mm Hg at the same temperature. Calculate molality (m) at temperature T :

A.  $6.3 \times 10^{-4}$

B.  $3.5 \times 10^{-2}$

C.  $5 \times 10^{-3}$

D. None of these

**Answer: c**



**Watch Video Solution**



48. When 1 mole of a solute is dissolved in 1 kg of  $H_2O$ , boiling point of solution was found to be  $100.5^\circ C$ .  $K_b$  for  $H_2O$  is :

- A. 0.5
- B. 100
- C. 100.5
- D. 95.5

**Answer: a**



**Watch Video Solution**

49. Chloroform,  $CHCl_3$ , boils at  $61.7^\circ C$ . If the  $K_b$  for chloroform is  $3.63^\circ C / \text{molal}$ , what is the boiling point of a solution of 15.0 kg of  $CHCl_3$  and 0.616 kg of acenaphthalene,  $C_{12}H_{10}$ ?

A. 61.9

B. 62

C. 52.2





D. 62.67

Answer: d



Watch Video Solution

50. A compound has the empirical formula  $C_{10}H_8Fe$ . A solution of 0.26 g of the compound in 11.2 g of benzene ( $C_6H_6$ ) boils at  $80.26^\circ C$ . The boiling point of benzene is  $80.10^\circ C$ , the  $K_b$  is  $2.53^\circ C/molal$ . What is the molecules formula of the compound?



Answer: d



Watch Video Solution

51. A solution of 0.640 g of azulene in 100.0 g of benzene is  $80.23^\circ\text{C}$ . The boiling point of benzene is  $80.10^\circ\text{C}$ , and  $K_b$  is  $2.53^\circ\text{C/molal}$ . What is the molecular mass of azulene?

A. 108

B. 99

C. 125

D. 134

**Answer: c**



**Watch Video Solution**

52. One molal solution of a carboxylic acid in benzene shows the elevation of boiling point of 1.518 K. The degree of association for dimerization of the acid in benzene is (  $K_b$  for benzene =  $2.53 \text{ K kg mol}^{-1}$  ):

A. 0.6

B. 0.7

C. 0.75

D. 0.8

**Answer: d**



**Watch Video Solution**

53. The boiling point elevation for toluene is  $3.32 \text{ K kg mol}^{-1}$ . The normal boiling point of toluene is  $110.7^\circ \text{C}$ . The enthalpy of vapourization of the toluene would be nearly :

A.  $17.0 \text{ kJ mol}^{-1}$

B.  $34.0 \text{ kJ mol}^{-1}$

C.  $51.0 \text{ kJ mol}^{-1}$

D.  $68.0 \text{ kJ mol}^{-1}$

**Answer: b**



**Watch Video Solution**

54. Which of the following aqueous solutions should have the highest boiling point ?

A. 0.015 M urea

B. 0.01 M  $KNO_3$

C. 0.10M  $Na_2SO_4$

D. 0.015 m glucose

Answer: c



Watch Video Solution

55. Calculate the percentage degree of dissociation of an electrolyte  $XY_2$  (Normal molar mass = 164) in water if the

observed molar mass by measuring elevation in boiling point is 65.6

A. 75%

B. 25%

C. 65%

D. None of these

**Answer: a**



**Watch Video Solution**

**56.** if the elevation in boiling point of a solution of non-volatile, non-electrolytic and non-associating solute in solvent ( $K_b = x K kg mol^{-1}$ ) is y K, then the depression in



freezing point of solution of same concentration would be

$(K_f)$  of the solvent =  $z k. kgmol^{-1}$ )

A.  $2x \frac{y}{y}$

B.  $y \frac{z}{x}$

C.  $x \frac{z}{y}$

D.  $y \frac{z}{2x}$

**Answer: b**



**Watch Video Solution**

57. When a solution containing non- volatile solute freezes, which equilibrium would exist?

A. solid solvent  $\Leftrightarrow$  liquid solvent

B. solid solute  $\Leftrightarrow$  liquid solution

C. solid solute  $\Leftrightarrow$  liquid solvent

D. solid solvent  $\Leftrightarrow$  liquid solution

**Answer: d**



**Watch Video Solution**

**58.** Bromoform has a normal has freezing point of  $7.734^{\circ}C$  and  $K_f = 14.4^{\circ}C/m$ . a solution of 2.60 g of an unknown substance in 100 g of freezes at  $5.43^{\circ}C$ . What is the molecules mass of the unknown substance ?

A. 16.25

B. 162.5

C. 100

D. none of these

**Answer: b**



**Watch Video Solution**

59.  $C_6H_6$  freezes at  $5.5^\circ C$ . At what temperature will a solution of 10.44 g of  $C_4H_{10}$  in 200 g of  $C_6H_6$  freeze  $K_f(C_6H_6) = 5.12^\circ C/m$

A.  $4.608^\circ C$

B.  $0.892^\circ C$

C.  $5.5^{\circ}C$

D. none of these

**Answer: b**

 [Watch Video Solution](#)

**60.** How much ethyl alcohol must be added to  $1.00L$  of water so that the solution will not freeze at  $-4^{\circ}F$ ?

A.  $>20\text{ g}$

B.  $<10.75\text{ g}$

C.  $<494.5\text{ g}$

D.  $>494.5\text{ g}$

**Answer: d**



**Watch Video Solution**

**61.** The freezing point of a solution of 2.40 g of biphenyl( $C_{12}H_{10}$ ) in 75.0 g of benzene ( $C_6H_6$ ) is  $4.40^\circ C$ . The normal freezing point of benzene is  $5.50^\circ C$ . What is the molal freezing point constant for benzene ?

A. 5.3

B. 5.1

C. 4.6





D. 4.8

**Answer: a**



Watch Video Solution

62. A solution containing 1.8 g of a compound (empirical formula  $CH_2O$ ) in 40 g of water is observed to freeze at  $-0.465^\circ\text{C}$ . The molecules formula of the compound is ( $K_f$  of water  $= 1.86\text{K mol}^{-1}$ ):



Answer: d



Watch Video Solution

63. Freezing point of the following equilibrium, liquid solvent  $\rightleftharpoons$  solid solvent is :

A.  $\frac{\Delta H - \Delta G}{T \Delta S}$

B.  $\frac{\Delta H}{\Delta S}$

C.  $\frac{\Delta G}{\Delta S}$

D.  $\frac{\Delta S}{\Delta H}$

Answer: b



Watch Video Solution

64. Freezing point of a solution is smaller than freezing point of a solvent. It is due to :

A.  $\Delta H$  of solution and solvent is almost identical since intermolecular force between solvent molecules are involved

B.  $\Delta S$  solution (between solution and solid ) is larger than that of the  $\Delta S$  of solvent (between solvent and solid)

C.  $\Delta S$  of the solution is smaller than that of the solvent





D.  $\Delta H$  of the solution is much higher than of solvent but  $\Delta S$  of solvent than that of the solvent



**Answer: b**

 **Watch Video Solution**

65. When 36g of a non-volatile, non-electrolytic solute having the empirical formula  $CH_2O$  is dissolved in 1.2 kg of water, the solution freezes at  $-0.93^\circ C$ . The molecular formula of the solute is ( $K_f$  of water =  $1.86 \text{ K kg mol}^{-1}$ )



**Answer: d**



Watch Video Solution

66. When 36g of a non-volatile, non-electrolytic solute having the empirical formula  $CH_2O$  is dissolved in 1.2 kg of water, the solution freezes at  $-0.93^\circ C$ . The molecular formula of the solute is ( $K_f$  of water =  $1.86 \text{ K kg mol}^{-1}$ )

A.  $140 \text{ gmol}^{-1}$

B.  $150.5 \text{ gmol}^{-1}$

C.  $160 \text{ gmol}^{-1}$

D.  $155 \text{ gmol}^{-1}$

Answer: b



Watch Video Solution

**67.** Assertion: Camphor is used as solvent in the experimental determination of molecular mass of naphthalene and anthracene

Reason: Camphor has high cryoscopic constant

- A. it is readily available
- B. it has a very high cryoscopic constant
- C. it is volatile
- D. it is solvent for organic substances

**Answer: b**



**Watch Video Solution**

68. How is molar mass related to the depression in freezing point of a solution ?

- A.  $[Fe(H_2O)_6]Cl_3$
- B.  $[Fe(H_2O)_5Cl]Cl_2 \cdot H_2O$
- C.  $[Fe(H_2O)_4Cl_2]Cl \cdot 2H_2O$
- D.  $[Fe(H_2O)_3Cl_3] \cdot 3H_2O$

**Answer: a**



**Watch Video Solution**

69. Which of the following solutions (*1molal*) will have the maximum freezing point, assuming equal ionization in each case?

- A.  $[Fe(H_2O)_6Cl]Cl_3$
- B.  $[Fe(H_2O)_5Cl]Cl_2 \cdot H_2O$
- C.  $[Fe(H_2O)_4Cl_2]Cl \cdot 2H_2O$
- D.  $[Fe(H_2O)_3Cl_3] \cdot 3H_2O$

**Answer: d**






**Watch Video Solution**

**70.** Calculate depression of freezing point for 0.56 molal aq. Solution of KCl.

(Given :  $K_{f(H_2O)} = 1.8 \text{ kg mol}^{-1}$ ).

- A.  $[pt(H_2O)_6]Cl_4$






Answer: c



Watch Video Solution

71. A complex is represented as  $CoCl_3 \cdot xNH_3$ . Its 0.1 molal solution in water shows  $\Delta T_f = 0.558k$ .  $K_f$  for  $H_2O$  is  $1.86K/mol$ . Assuming 100% ionisation of complex and coordination number of Co is six. The formula of complex is








D. none of these


**Answer: b**



**Watch Video Solution**

**72.** The freezing point of equimolal aqueous solution will be highest for





**Answer: d**



**Watch Video Solution**

**73.** The freezing point of equimolal aqueous solution will be highest for

A. 160

B. 90

C. 45

D. 180

**Answer: a**





Watch Video Solution

74. Depression in freezing point of 0.01 molal aqueous HCOOH solution is 0.02046. one molal aqueous urea solution freezes at  $-1.86^{\circ}C$ , assuming molality equal to molarity, pH of HCOOH solution is :

A. 2

B. 3

C. 4

D. 5

**Answer: b**



Watch Video Solution

75. When mercuric iodide is added to the aqueous solution of potassium iodide, the

- A. freezing point is raised
- B. Freezing point is lowered
- C. freezing point does not change
- D. boilingpoint does not change

**Answer: a**



**Watch Video Solution**

76. The solution of acetic acid in benzene conains

- A. freezing point of the solution reduces
- B. average molar mass of solute increases
- C. boiling point of solution increases
- D. molar mass of solute decreases

**Answer: b**



**Watch Video Solution**

**77.** The temperature of a city was found to be  $-9.3^{\circ}\text{C}$ . A car used, whose radiator was filled with 5 L of water. What minimum quantity of antifreezing agent ethylene glycol were added to water of radiator in order to use the car for travelling? ( $K_f$  of water  $1.86\text{ k mol}^{-1}$ )

A. 3200 g

B. 1670 g

C. 1550 g

D. 2100 g

**Answer: c**



**Watch Video Solution**

**78.** The cryoscopic constant of water is  $1.86 \text{ K kg mol}^{-1}$ . A 0.01 molal acetic acid solution produces a depression of  $0.0194^\circ \text{C}$  in the freezing point. The degree of dissociation of acetic acid is :

A. zero

B. 0.043

C. 0.43

D. 1

**Answer: b**



**Watch Video Solution**

**79.** In a 0.5 molal solution KCl, KCl is 50% dissociated. The freezing point of solution will be ( $K_f = 1.86 \text{ K kg mol}^{-1}$ ):

A. 274.674 K

B. 271.60 K

C. 273 K

D. none of these

**Answer: b**

 **Watch Video Solution**

**80.** A 1.0 g sample of  $\text{Co}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_3\text{Cl}_3$  is dissolved in 25.0 g of water and the freezing point of the solution is  $-0.87^\circ\text{C}$ . How many ions are produced per mole of compound? The  $K_f$  of water is  $1.86^\circ\text{C}/\text{molal}$

A. 2

B. 3

C. 4

D. 5

**Answer: c**



**Watch Video Solution**

**81.** An aqueous solution contain 3% and 1.8% by mass. Urea and glucose respectively. What is the freezing point of solution ? ( $K_f = 1.86^\circ C/m$ )

A.  $-1.172^\circ C$

B.  $-2.27^\circ C$

C.  $-1.5^\circ C$

D. none of these

**Answer: a**



**Watch Video Solution**

82. phenol associates in benzene to a certain extent in dimerisation reaction. A solution containing 0.02 kg of phenol in 1.0 kg of benzene has its freezing point depressed 0.69 k Hence degree of dissociation of phenol dimerises will be. [ $K_f(C_6H_6) = 5.12\text{ k mol}^{-1}$ ]

A. 0.63

B. 0.73

C. 0.83

D. 0.93



**Answer: b**



**Watch Video Solution**



83. The first ionisation potential is maximum for



Answer: d



Watch Video Solution

84. In a 0.2 molal aqueous solution of a weak acid HX the degree of dissociation is 0.25. The freezing point of the

solution will be nearest to: ( $K_f = 1.86 \text{ K kg mol}^{-1}$ )

A.  $-0.26^\circ \text{ C}$

B.  $0.465^\circ \text{ C}$

C.  $-0.48^\circ \text{ C}$

D.  $-0.465^\circ \text{ C}$

**Answer: d**



**Watch Video Solution**

**85.** An aqueous solution of 0.01 M KCl cause the same elevation in boiling point as an aqueous solution of urea.

The concentration of urea solution is :

A. 0.01 m

B. 0.005 M

C. 0.02 M

D. 0.04 M

**Answer: c**



**Watch Video Solution**

**86.** when some NaCl was dissolved in water, the freezing point depression was numerically equal to twice the molal f.p. depression constant. The relative lowering of vapour pressure of the solution is nearly :

A. 0.036

B. 0.018

C. 0.0585


D. 0.072

**Answer: a**

 [Watch Video Solution](#)

**87.** Which one of the following statement is false ?

A. The correct order of osmotic pressure for 0.01 M aqueous solution of each follows



B. Isotonic solutions are those solutions which have the same osmotic pressure

C. Raoult's law states that the vapour pressure of a component over a solution is proportional to its mole fraction in liquid state

D. Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression

**Answer: d**



**Watch Video Solution**

88. 0.1 molal aqueous solution of an electrolyte  $AB_3$  is 90% ionised. The boiling point of the solution at 1 atm is (

$$K_b(H_2O) = 0.52 \text{ kg mol}^{-1})$$

A. 273.19 K

B. 374.92 K

C. 376.4 K




D. 373.19 K

**Answer: d**



**Watch Video Solution**

89. Which of the following aqueous solutions has osmotic pressure nearest to pure solvent ?



Answer: d



Watch Video Solution

90. 0.1 M NaCl and 0.05 M  $BaCl_2$  solutions are separated by a semi-permeable membrane in a container. For this

system, choose the correct answer

A. There is no movement of any solution across the membrane

B. Water flows from  $BaCl_2$  solution towards NaCl solution

C. Water flows from NaCl solution towards  $BaCl_2$  solution

D. Osmotic pressure of 0.1 M NaCl is lower than the osmotic pressure of  $BaCl_2$ (assume complete dissociation)

**Answer: b**



**Watch Video Solution**



**91.** Two aqueous solutions, A and B, are separated by a semi-permeable membrane. The osmotic pressure of solution A immediately begins to decrease. Which of the following statements is true?

- A. The solvent molecules are moving from the solution of higher osmotic pressure to that of lower osmotic pressure
- B. The initial osmotic pressure of solution B is greater than that of solution A.
- C. Solvent molecules are moving from solution B into solution A.

D. Both (a) and (b) are true statements.

**Answer: c**



**Watch Video Solution**

**92.** Which one of the following pair of solutions is an isotonic?

A. 0.1 M urea and 0.1 M NaCl

B. 0.1 M urea and 0.2 M  $MgCl_2$

C. 0.1 M NaCl and 0.1 M  $Na_2SO_4$

D. 0.1 M  $C(NO_3)_2$  and 0.1 M  $Na_2SO_4$




**Answer: d**



Watch Video Solution

93. The empirical formula of a non-electrolyte is  $CH_2O$ . A solution containing  $3 \text{ g L}^{-1}$  of the compound exerts the same osmotic pressure as that of  $0.05 \text{ M}$  glucose solution.

The molecules formula of the compound is :



Answer: b



Watch Video Solution

94. A semipermeable membrane used in the measurement of osmotic pressure of a solution allows the passage of

- A. solute molecular through it
- B. solvent molecules though it
- C. both solvent and solute molecules
- D. either solvent or solute

**Answer: b**



[Watch Video Solution](#)

95. In osmosis the water always moves towards ....

- A. higher vapour pressure to lower vapour pressure
- B. higher concentration to lower concentration
- C. lower vapour pressure to higher vapour pressure
- D. higher osmotic pressure to lower osmotic pressure

**Answer: a**



**Watch Video Solution**

**96.** The osmotic pressures of equimolar solutions of urea,

$BaCl_2$  and  $AlCl_3$  will be in the order :

- A.  $AlCl_3 > BaCl_2$  gt urea
- B.  $BaCl_2 > AlCl_3$  gt urea

C. urea  $\text{gt}$   $\text{BaCl}_2 > \text{AlCl}_3$

D.  $\text{BaCl}_2 > \text{urea} > \text{AlCl}_3$

**Answer: a**

 [Watch Video Solution](#)

**97.** Assuming each salt to be 90% dissociated which of the following will have the highest osmotic pressure?

A. decimolar aluminium sulphate

B. decimolar barium chloride solution

C. decimolar sodium sulphate solution

D. solution of valume of decimolar barium choride and  
decimolar sodium suphate solutions

**Answer: a**

 [Watch Video Solution](#)

**98.** consider 0.1 M solutions of two solutes X and Y. The behaves as a univalent electrolyte while the solute Y dimerises in solution. Which of the following statement are correct regarding these solutions?

(1) The boiling point of the solution of X will be higher than that of Y

(2) The osmotic pressure of the solution of Y will be lower than that of X

(3) The freezing point of the solution of X will be lower than that of Y

(4) The relative lowering of vapour pressure of both the solutions will be the same It brgt Select the correct answer from the option given below

A. 1, 2 and 3

B. 2, 3 and 4

C. 1, 2 and 4

D. 1, 3 and 4

**Answer: a**



**Watch Video Solution**



99. If  $M_{\text{normal}}$  is the normal molecular mass and  $\alpha$  is the degree of ionisation of  $K_3[Fe(CN)_6]$ , then the abnormal molecular mass of the complex in the solution will be

A.  $M_{\text{normal}}(1 + 2\alpha)^{-1}$

B.  $M_{\text{normal}}(1 + 3\alpha)^{-1}$

C.  $M_{\text{normal}}(1 + \alpha)^{-1}$

D. equal to  $M_{\text{normal}}$

**Answer: b**



**Watch Video Solution**

**100.** Equal volumes of 0.1 M urea and 0.1 M glucose are mixed. The mixture will have :

- A. lower osmotic pressure
- B. same osmotic pressure
- C. higher osmotic pressure
- D. none of these

**Answer: b**

 [Watch Video Solution](#)

**101.** A 5% (w/V ) solution of cane sugar (molecular mass = 342) is isotonic with 1% (w/V) solution of a substance X.

The molecular mass of X is :

A. 34.2

B. 171.2

C. 68.4

D. 136.8

**Answer: c**



**Watch Video Solution**

**102.** Insulin  $(C_2H_{10}O_5)_n$  is dissolved in a suitable solvent and the osmotic pressure ( $\pi$ ) of solutions of various concentrations  $(g/cm^3)C$  is measured at  $20^\circ C$ . The

slope of a plot of  $\pi$  against  $C$  is found to be  $4.65 \times 10^{-3}$ .

The molecular weight of insulin is:

A.  $3 \times 10^5$

B.  $9 \times 10^5$

C.  $4.5 \times 10^5$

D.  $5.16 \times 10^6$

**Answer: d**



[Watch Video Solution](#)

**103.** An aqueous solution of sucrose ( $C_{12}H_{22}O_{11}$ ) having a concentration of 34.2gram/ litra has an osmotic pressure of 2.38 atmospheres at  $17^\circ\text{C}$ . For an aqurous

solution of glucose ( $C_6H_{12}O_6$ ) to be isotonic with this solution , its concentration should be :

- A. 34.2 gram per liter
- B. 17.1 gram per liter
- C. 18.0 gram per liter
- D. 36.0 gram per liter

**Answer: c**

 [Watch Video Solution](#)

**104.** Which of the following experimental methods is adopted to determine osmotic pressure?

A. Berkley- Hartely's method

B. Beckmann's method

C. Landsberger's method

D. Differential method

**Answer: a**



**Watch Video Solution**

**105.** Based upon the technique of reverse osmosis the approximate pressure required to desalinate sea water containing 2.5% (mass/volume)  $KNO_3$  at  $27^\circ C$  will be

A. 10.5 atm

B. 21 atm

C. 12.2 atm

D. 6.09 atm

**Answer: c**



**Watch Video Solution**

**106.** A 1% (mass/vol) KCl solution is ionised to the extent of 80%. The osmotic pressure at  $27^{\circ}\text{C}$  of the solution will be :

A. 6.95 atm

B. 5.94 atm

C. 2.71 atm

D. 3.30 atm

**Answer: b**

 [Watch Video Solution](#)

**107.** The osmotic pressure of blood is 7.40 atm. at  $27^{\circ}C$  .

Number of mole of glucose to be used per liter for an intravenous injection that is to have the same osmotic pressure as blood is

A. 0.3

B. 0.2

C. 0.1



D. 0.4

**Answer: a**



**Watch Video Solution**

**108.** The relationship between osmotic pressure ( $\pi_1$ ,  $\pi_2$  and  $\pi_3$ ) at a definite temperature when 1 g glucose, 1 g urea and 1 g sucrose are dissolved in 1 liter of water is (assume  $i = 1$  for all):

A.  $\pi_1 > \pi_2 > \pi_3$

B.  $\pi_3 > \pi_1 > \pi_2$


C.  $\pi_2 > \pi_1 > \pi_3$

D.  $\pi_2 > \pi_3 > \pi_1$


Answer: c

 Watch Video Solution

109. van't Hoff proved that osmotic pressure ( $\pi$ ) is a colligative property. For an ideal solution, osmotic pressure ( $\pi$ ) is helpful to determine that molecular mass of solute using  $M_B = \frac{W_B RT}{\pi \cdot V}$  Relation can expressed by the curve (C = concentration) :




A.




B.

C.



D.



**Answer: a**

 [Watch Video Solution](#)

**110.** A solution containing 4.0 g of PVC in 2 litre of dioxane (industrial solvent ) was found to have an osmotic pressure  $3.0 \times 10^{-4}$  atm at  $27^\circ C$  The molecular mass of the polymer will be :

A.  $1.6 \times 10^4$

B.  $1.6 \times 10^5$

C.  $1.6 \times 10^3$

D.  $1.6 \times 10^2$

**Answer: b**



**Watch Video Solution**

**111.** The osmotic pressure of 0.010 M solutions of KI and of sucrose ( $C_{12}H_{22}O_{11}$ ) are 0.432 atm and 0.24 atm respectively . The van't Hoff factor for KI is :

A. 1.8

B. 0.8

C. 1.2

D. 1

**Answer: a**

 [Watch Video Solution](#)

**112.** What is the correct sequence of osmotic pressure of 0.01M aq. solution of :

(a)  $Al_2(SO_4)_3$       (b)  $Na_3PO_4$       (c)  $BaCl_2$       (d) *Glucose*

A.  $\pi_4 > \pi_2 > \pi_3 > \pi_1$

B.  $\pi_3 > \pi_4 > \pi_2 > \pi_1$

C.  $\pi_3 > \pi_4 > \pi_1 > \pi_2$

D.  $\pi_1 > \pi_2 > \pi_3 > \pi_4$

**Answer: d**

 **Watch Video Solution**

**113.** 1.0 molar solution of the complex of the salt,  $CrCl_3 \cdot 6H_2O$ , displays an osmotic pressure of  $3RT$ . 0.5 L of the same solution on treatment with excess of  $AgNO_3$  solution will yield (assume  $a = 1$ ):

A. 0.5 mole of AgCl

B. 1.0 mole of AgCl

C. 1.5 mole of AgCl

D. 3.0 mole of AgCl

**Answer: b**

 [Watch Video Solution](#)

**114.** A 0.010 g sample of  $Cr(NH_3)_4(SO_4)Cl$  is dissolved in 25.0 mL of water and the osmotic pressure of the solution is 59.1 torr at  $25^\circ C$ . How many moles of ions are produced per mole of compound?

A. 1

B. 4

C. 2

D. 3

**Answer: c**



Watch Video Solution

115. Which of the following aqueous solutions should have the highest osmotic pressure?

A. 0.011 M  $AlCl_3$  at  $50^\circ C$

B. 0.03 m NaCl at  $25^\circ C$

C. 0.012 m  $(NH_4)_2SO_4$  at  $25^\circ$

D. 0.03 m NaCl at  $50^\circ C$


Answer: d



Watch Video Solution



116.  $X_3Y_2(i = 5)$  when reacted with  $A_2B_3(i = 5)$  in aqueous solution gives brown colour. These are separated by a semipermeable membrane  $AB$  as shown. Due to osmosis there is :







- A. brown colour formation in side X
- B. brown colour formation in side Y
- C. formation in both of the sides X and Y
- D. no brown colour formation

Answer: d



Watch Video Solution

117. Which of the following curves represents the Henry's law ?



**Answer: a**



**Watch Video Solution**

**118.** Assertion: The solubility of the gas in a liquid increases with increase of pressure.

Reason : The solubility of a gas in a liquid is directly proportional to the pressure of the gas.

- A. temperature
- B. pressure
- C. Both (a) and (b)
- D. none of these

**Answer: b**



Watch Video Solution

119. At  $300K$ ,  $40mL$  of  $O_3(g)$  dissolves in  $100g$  of water at  $1.0atm$ . What mass of ozone dissolved in  $400g$  of water at a pressure of  $4.0atm$  at  $300K$  ?

A.  $0.1\text{ g}$

B.  $1.24\text{ g}$

C.  $0.48\text{ g}$

D.  $4.8\text{ g}$

Answer: b



Watch Video Solution

120. 1 kg of water under a nitrogen pressure of 1 atmosphere dissolves 0.02 gm of nitrogen at 293 K.

Calculate Henry's law constant :

A.  $7.2 \times 10^{-4}$  L/atm

B.  $7.7 \times 10^3$  atm

C.  $2 \times 10^{-5}$  atm

D.  $2 \times 10^{-2}$  atm

**Answer: a**



**Watch Video Solution**

121. According to Henry's Law , the partial pressure of gas ( $P_g^1$ ) is directly , propotional to molefraction of gas in dissolved state , i.e. ,  $P_{\text{gas}}^1 = K_H \cdot X_{\text{gas}}$ . Which are correct .

- A.  $K_H$  is characteristic constant for a given gas-solvent system
- B. Higher is the value of  $K_H$ , lower is solubility of gas for a given partial pressure of gas
- C.  $K_H$  has temperature dependence
- D.  $K_H$  decreases with increase of temprature

Answer: d



Watch Video Solution

122. At 760 torr pressure and  $20^{\circ}C$  temperature, 1 L of water dissolves 0.04 gm of pure oxygen or 0.02 gm of pure nitrogen. Assuming that dry air is compound of 20% oxygen and 80% nitrogen (by volume), the masses (in g/L) of oxygen and nitrogen dissolved by 1 L of water at  $20^{\circ}C$  exposed to air at a total pressure of 706 torr are respectively :

A. 0.008, 0.016

B. 0.016, 0.008

C. 0.16, 0.08

D. 0.04, 0.02

**Answer: a**



**Watch Video Solution**

123. The plot of  $1/x_A$  versus  $1/y_A$  (where  $X_A$  and  $Y_A$  are the mole fraction of A in liquid and vapour phases, respectively) is linear with slope and intercept respectively are given as (y-axis =  $1/y_A$ , x-axis =  $1/x_A$ )

A.  $\frac{P_A^\circ}{P_B^\circ}, \frac{P_B^\circ - P_A^\circ}{P_B^\circ}$

B.  $\frac{P_B^\circ}{P_A^\circ}, \frac{P_A^\circ - P_B^\circ}{P_A^\circ}$

C.  $\frac{P_B^\circ}{P_A^\circ}, \frac{P_B^\circ}{P_B^\circ - P_A^\circ}$

D.  $P_A^\circ - P_B^\circ, \frac{P_A^\circ}{P_B^\circ}$

Answer: b



Watch Video Solution



**124.** At  $48^{\circ}C$ , the vapour pressure of pure  $CS_2$  is 850 torr .  
A solution of 2.0 g of sulphur in 100g of  $CS_2$  has a vapour pressure 844.9 torr. Determine the atomicity of sulphur molecule :

A. 1

B. 2

C. 4

D. 8

**Answer: d**



**Watch Video Solution**

125. If two liquids A ( $P_A^\circ = 100\text{torr}$ ) and ( $P_B^\circ = 200\text{ torr}$ ) which are completely immiscible with each other (each one will behave independently of the other) are present in a closed vessel, the total vapour pressure of the system will be :





- A. 150
- B. 180
- C. 188.88
- D. 198.88

**Answer: c**



**Watch Video Solution**

126. Which of the following represents correctly the changes in thermodynamic properties during the formation of 1 mole of an ideal binary solution :



Answer: c



Watch Video Solution

127. A certain non-volatile electrolyte contains 40% carbon, 6.7% hydrogen and 53.3% oxygen. An aqueous solution containing 5% by mass of the solute boils at  $100.15^{\circ}\text{C}$ . Determine the molecular formula of the compound ( $K_b = 0.51^{\circ}\text{C}/m$ ):

A.  $\text{HCHO}$

B.  $\text{CH}_3\text{OH}$

C.  $\text{C}_2\text{H}_5\text{OH}$

D.  $\text{C}_6\text{H}_{12}\text{O}_6$

Answer: d



Watch Video Solution

128. An aqueous solution boils at  $101^{\circ}C$ . What is the freezing point of the same solution?

(Gives :  $K_f = 1.86^{\circ}C/m$  and  $K_b = 0.51^{\circ}C/m$ )

A.  $3.647^{\circ}C$

B.  $-3.647^{\circ}C$

C.  $-0.199^{\circ}C$

D. none of these

**Answer: b**



**Watch Video Solution**

129. An industrial waste water I found to contain 8.2%  $Na_3PO_4$  and 12%  $MgSO_4$  by mass in solution. If %

ionisation of  $Na_3PO_4$  and  $MgSO_4$  are 50 and 60 respectively then its normal boiling point is [ $K_b(H_2O) = 0.50 K kg mol^{-1}$ ]:

- A.  $102.3^\circ C$
- B.  $103.35^\circ C$
- C.  $101.785^\circ$
- D. none of these

**Answer: c**

 [Watch Video Solution](#)

130. Ratio of  $\frac{\Delta T_b}{K_b}$  of 10 g  $AB_2$  and 14 g  $A_2B$  per 100 g of solvent in their respective, solution ( $AB_2$  and  $A_2B$

both are non-electrolytes ) is 1 mole/ kg in both cases.

Hence, atomic wt. of A and B are respectively :

A. 100, 40

B. 60, 20

C. 20, 60

D. None of these

**Answer: b**



[Watch Video Solution](#)

**131.** The freezing point of solution containing  $0.2g$  of acetic acid in  $20.0g$  of benzene is lowered by  $0.45^{\circ}C$ .

Calculate the degree of association of acetic acid in

benzene.

$$(K_f = 5.12K^{\circ} mol^{-1} kg^{-1})$$

A. 0.527

B. 0.8

C. 0.945

D. None of these

**Answer: c**



**Watch Video Solution**

**132.** If the boiling point of an aqueous solution containing a non-volatile solute is  $100.15^{\circ}C$ . What is its freezing



point? Given latent heat of fusion and vapourization of water  $80\text{calg}^{-1}$  and  $540\text{calg}^{-1}$ , respectively.

A.  $0.361^\circ\text{C}$

B.  $-0.361^\circ\text{C}$

C.  $-3.61^\circ\text{C}$

D. None of these

**Answer: b**



[Watch Video Solution](#)

**133.** 1.0 g of a monobasic acid HA in 100 g water lowers the freezing point by 0.155 K. IF 0.75 g, of same acid requires 25 mL of N/5 NaOH solution for complete

neutralisation then %, degree of ionization of acid is (

$K_f \text{ of } H_2O = 1.86 K kg mol^{-1}$ ):

A. 0.2

B. 0.25

C. 0.4

D. 0.5

**Answer: b**



[Watch Video Solution](#)

**134.** 0.1 M KI and 0.2 M  $AgNO_3$  are mixed in 3 : 1 volume ratio. The depression of freezing point of the resulting solution will be [ $K_b(H_2O) = 1.86 K kg mol^{-1}$ ]:

A. 3.72 K

B. 1.86 K

C. 0.93 K

D. 0.279 K

**Answer: d**



**Watch Video Solution**

**135.** If 0.1 M  $H_2SO_4$ (aq.) solution shows freezing point  $-0.3906^\circ C$  then what is the  $K_{a2}$  for  $H_2SO_4$ ? (Assume  $m = M$  and  $K_f(H_2O) = 1.86 K kg mol^{-1}$ )

A. 0.122

B. 0.0122

C.  $1.11 \times 10^{-3}$

D. None of these

**Answer: b**



**Watch Video Solution**

**136.** A living cell contains a solution which is isotonic with 0.2 M glucose solution. What osmotic pressure develops when the cell is placed in 0.05 M  $BaCl_2$  solution at 300 K ?

A. 1.23 atm

B. 3.69 atm

C. 6.15 atm

D. None of these

**Answer: a**

 [Watch Video Solution](#)

**137.** What is the osmotic pressure of 0.2 M HX (aq.) solution at 300 K ?

A. 4.926 atm

B. 0.5024 atm

C. 5.024 atm

D. None of these

**Answer: c**

 [Watch Video Solution](#)

**138.** A solution contain 8 g of a carbohydrate in 100 g of water has a density 1.025 g/mL and an osmotic pressure of 5 atm at  $27^{\circ}C$ . What is the molar mass of the carbohydrate?

A. 387

B. 374

C. 3740


D. None of these

**Answer: b**



Watch Video Solution

139. Study the following figure and choose the correct options. Assuming complete dissociation of electrolyte:



- A. There will be net movement of any substance across the membrane
- B.  $MgCl_2$  will flow towards the  $Al_2(SO_4)_3$  solution
- C.  $Al_2(SO_4)_3$  will flow towards the  $MgCl_2$  solution

D. The  $\pi$  (osmotic pressure ) of 0.1 M  $MgCl_2$  is higher than the  $\pi$  of 0.05 M  $Al_2(SO_4)_3$

**Answer: d**

 [Watch Video Solution](#)

**140.** Two pure liquids A & B which can form ideal solution have vapour pressures 300 torr & 800 torr . A mixture of the vapour of A & B for which the mole fraction of A is 0.25 is slowly compressed at temperature T.


The total pressure applied when only last bubble of vapour remains

 [Watch Video Solution](#)



**141.** Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.



Loss in mass of solvent ( $w_{II}$ ) will be proportional to :

A.  $P^\circ - P$

B.  $P - P^\circ$

C.  $\frac{P}{P^\circ}$

D.  $P \times P^\circ$


**Answer: a**

 [Watch Video Solution](#)

**142.** Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and

the solvent is kept constant.



Gain in mass of anhydrous  $\text{CaCl}_2$  is proportional to :

- A.  $P$
- B.  $P^\circ$
- C.  $P - P^\circ$
- D.  $P^\circ - P$


**Answer: b**



**Watch Video Solution**

**143.** Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.



$\frac{P^\circ - P}{P^\circ}$  is equal to :

A.  $\frac{w_I}{w_{II} + w_I}$

B.  $\frac{w_{II}}{w_I + w_{II}}$

C.  $\frac{w_I}{w_{II} - w_I}$

D.  $\frac{w_{II}}{w_I}$


**Answer: b**

 [Watch Video Solution](#)

**144.** Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and

the solvent is kept constant.



Dry air was passed thorough 9.24 g of solute in 108 g of water and then through pure water. The loss in mass of solution was 3.2 g and that of pure water 0.08 g . The molecular mass (g/mol) of solute is nearly :


- A. 50
- B. 62
- C. 70
- D. 80

**Answer: b**



**Watch Video Solution**

145. A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant  $K_b$ . The solute dimerises in the solution according to the following equation. The degree of association is (a) :



The van't Hoff factor will be:

A.  $i = 1 - 2a$

B.  $i = 1 - \frac{a}{2}$

C.  $i = 1 + \frac{a}{2}$

D.  $i = 1 + a$

**Answer: b**



**Watch Video Solution**

**146.** A dilute solutions "x" moles of A in 1kg of solvent with molal elevation constant  $K_b$  . The solution dimerises in the solution .  $2A \rightleftharpoons A_2$  ( $\alpha$  be degree of association)

The molecular weight observed will be :

- A. greater than actual molecular mass
- B. lesser than actual molecular mass
- C. equal to the actual molecular mass
- D. cannot be predicted by the date given

**Answer: a**



**Watch Video Solution**



**147.** A dilute solution having one mole of solute 'S' in 1 kg solvent with molal elevation constant  $K_b$ . Solute undergoes association as follows.  $2S \rightleftharpoons S_2$  The degree of association of solute  $\alpha$  is given by the following expression .

$$\alpha = \frac{1 - i}{1 - 1/n}$$

Where,  $n$  is the number of molecular of solute undergoing association .

The degree of association can be given as :

A.  $a = \frac{(K_b x - \Delta T_b)}{\Delta T_b 2}$

B.  $a = \frac{2(K_b x - \Delta T_b)}{K_b x}$

C.  $a = 2 + \frac{2 \Delta T_b}{K_b x}$

D.  $a = \frac{\Delta T_b}{2K_b x}$

**Answer: b**



**Watch Video Solution**

**148.** When non-volatile solute is added to a pure solvent, the:

- A. vapour pressure of the solution becomes lower than the vapour pressure of the pure solvent
- B. rate of evaporation of solvent is reduced
- C. solute does not affect the rate of condensation
- D. none of these

**Answer: a,b,c**





**149.** The total vapour pressure of a binary solution is given by

$$P = (100X_A + 260X_B)\text{mm Hg}$$

where,  $X_A$  and  $X_B$  are the mole fractions of components A and B. This indicates that the:

- A. vapour pressure of solution is less than the pure B component
- B. vapour pressure of solution is less than the pure A component
- C. vapour pressure of pure A is 100 mm Hg and that of pure B is 260 mm Hg

D. the vapour pressure of pure A and B are 260 mm Hg  
and 100 mm hg respectively

**Answer: a,b,c**

 [Watch Video Solution](#)

**150.** Two compounds form an ideal solution at room temperature. Which of the following are correct for this ideal solution ?

(a)  $\Delta G = +ve$

(b)  $\Delta S = +ve$  surrounding

(c)  $\Delta S = +ve$  system

(d)  $\Delta_{mix}H = 0$

A.  $\Delta H_{\text{mix}} = 0$  and  $\Delta V_{\text{mix}} = 0$

B.  $\Delta V_{\text{mix}} = 0$  and  $\Delta S_{\text{mix}} > 0$

C.  $\Delta H_{\text{mix}} > 0$  and  $\Delta S_{\text{mix}} > 0$

D.  $\Delta G_{\text{mix}} < 0$  and  $\Delta S_{\text{mix}} > 0$

**Answer: a,b,d**



**Watch Video Solution**

**151.** Which of the following is correct for a non-ideal solution of liquids A and B showing negative deviation?

A.  $\Delta H_{\text{mix}} = -ve$

B.  $\Delta V_{\text{mix}} = -ve$

C.  $\Delta S_{\text{mix}} = -ve$

D.  $\Delta G_{\text{mix}} = -ve$

**Answer: a,b,d**

 **Watch Video Solution**

**152.** A binary solution of liquids A and B will show positive deviation from Raoult's law if it fulfils the following condition:

A.  $P_A > X_A P_{A^\circ}$  and  $P_B > X_B P_{B^\circ}$

B. The intermolecular forces of A - B lt A - A, B - B

C.  $\Delta H_{\text{mixing}}$  is positive

D.  $\Delta V$  mixing is negative

**Answer: a,b,c**

 **Watch Video Solution**

**153.** Which of the following statement is/are correct about acetone and trichloromethane mixture?

A. Mixtures of acetone and trichloromethane show positive deviation from Raoult's law

B. The forces of attraction acting between molecules of acetone and trichloromethane in a mixture are

stronger than those acting between the molecules  
in pure acetone

C. Pure acetone can be obtained by the careful  
fractional distillation of any mixture of acetone and  
trichloromethane

D. When acetone and trichloromethane are mixed, the  
enthalpy change is negative

**Answer: b,d**



**Watch Video Solution**

**154.** The azeotropic solution of two miscible liquids:



A. can be separated by simple distillation

B. may show positive or negative deviation from

Raoult's law

C. are supersaturated solution

D. behave like a single component and boil at a

constant temperature

**Answer: b,d**

 [Watch Video Solution](#)

**155.** For exact determination of molecular mass through colligative properties measurement :

A. solute must be volatile

B. solution must be vary dilute

C. solution must be formed by similar nature of substances

D. solute must not be dissociated

**Answer: b,d**



**Watch Video Solution**

**156.** In the depression of freezing point experiment, it is found that

(a) The vapour pressure of the solution is less than that of pure solvent

(b) The vapour pressure of the solution is more than that of pure solvent

(c) Only solute molecules solidify at the freezing point

(d) Only solvent molecules solidify at the freezing point

A. vapour pressure of pure solvent is more than that of solution

B. vapour pressure of pure solvent is less than that of solution

C. only solute molecules solidify at the freezing point

D. only solvent molecules solidify at the freezing point

**Answer: a,c**



**Watch Video Solution**

**157.** The value of  $G$  depends upon

- A. the mole mass of the solute in the solution
- B. the molar mass of the solvent in the solution
- C. the enthalpy of fusion of the solvent
- D. the freezing point of the solvent

**Answer:** b,c,d



**Watch Video Solution**

**158.** Consider 0.1 M solutions of two solutes X and Y. The solute X behaves as univalent electrolyte, while the solute

Y dimerises in solution. Select correct statement(s) regarding these solutions:

- A. The boiling point of solution of 'X' will be higher than that of 'Y'
- B. The osmotic pressure of solution of 'Y' will be lower than that of 'X'
- C. The freezing point of solution of 'X' will be lower than that of 'Y'
- D. The relative lowering of vapour pressure of both the solution will be the same

**Answer: a,b,c**



**Watch Video Solution**

**159.** Consider following solutions: (I) 1 M glucose(aq) (II) 1 M sodium chloride(aq)  
(III) 1 M acetic acid in benzene (IV) 1 M ammonium phosphate (aq)

A. all are isotonic solutions

B. III is hypotonic of I, II, IV

C. I, II, IV are hypertonic of III

D. IV is hypertonic I, II, III

**Answer: b.c.d**



**Watch Video Solution**

160. Which of the following statement is/are incorrect?

A. 0.1 M KCl solution will have the same osmotic pressure as 0.1 M glucose solution

B. 0.1 M KCl solution will have the same boiling point as 0.1 M urea solution

C. 0.1 m glucose and 0.1 m urea are ismotic

D. 0.1 m  $MgCl_2$  solution will have less relative lowering of vapour pressure than 0.1 m NaCl

**Answer: a,b,d**




**Watch Video Solution**

**161.** Consider following solution:

0.1 m  $C_6H_5NH_3^+ Cl^-$ , 0.1 m KCl, 0.1 m Glucose, 0.1 m  $Na_2C_2O_4 \cdot 10H_2O$

A. the solution with higher boiling point is 0.1



B. the solution with higher freezing point is 0.1 m

glucose

C. 0.1 m  $C_6H_5NH_3Cl$  and 0.1 m NaCl will have the

same osmotic pressure

D. 0.1 m glucose solution will have the lowest osmotic

pressure



Answer: a,b,c,d

 Watch Video Solution

162. Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of column-

**Column-I**  
(A)  $\pi_1$  : 0.1 M glucose;  $\pi_2$  : 0.1 M urea  
(B)  $\pi_1$  : 0.1 M NaCl;  $\pi_2$  : 0.1 M  $\text{Na}_2\text{SO}_4$   
(C)  $\pi_1$  : 0.1 M NaCl;  $\pi_2$  : 0.1 M KCl  
(D)  $\pi_1$  : 0.1 M  $\text{CuSO}_4$ ;  $\pi_2$  : 0.1 M sucrose

**Column-II**  
(P)  $\pi_1$  and  $\pi_2$  are isotonic  
(Q) No net migration of solvent across the membrane  
(R)  $\pi_1$  is hypertonic to  $\pi_2$   
(S)  $\pi_1$  is hypotonic to  $\pi_2$

II.

 Watch Video Solution

**163.** Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of column-

Column-I
(A) $0.1\text{ M Al}_2(\text{SO}_4)_3$
(B) $0.1\text{ M AlPO}_4$
(C) $0.1\text{ M urea}$
(D) $0.1\text{ M MgCl}_2$

Column-II
(P) Solution with highest boiling point
(Q) van't Hoff factor is greater than 1
(R) Solution with lowest osmotic pressure
(S) Solution with lowest freezing point

II.



**Watch Video Solution**

**164.** Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I

may have the matching with the same entries of column-

Column-I	Column-II
(Solute)	(Van't Hoff factor, $i$ )
(A) $\text{AlCl}_3$ if $\alpha = 0.8$	(P) $i = 3.4$
(B) $\text{BaCl}_2$ if $\alpha = 0.9$	(Q) $i = 2.8$
(C) $\text{Na}_3\text{PO}_4$ if $\alpha = 0.9$	(R) $i = 3.8$
(D) $\text{K}_4[\text{Fe}(\text{CN})_6]$ if $\alpha = 0.7$	(S) $i = 3.7$

II.

 [Watch Video Solution](#)

**165.** Column -I and Column -II contain four entries each.

Entries of column-I are to be matched with some entries

of column-II. One or more than one entries of column-I

may have the matching with the same entries of column-

Column-I	Column-II
(A) Elevation of B.P.	(P) Colligative property
(B) Osmotic pressure	(Q) Ebullioscopic constant
(C) Relative lowering in V.P.	(R) Berkeley-Heartley method
(D) Depression of F.P.	(S) Ostwald and Walker method

II.

 [Watch Video Solution](#)

**166.** (A) Increase in surface area increases rate of evaporation

(R) Stronger the intermolecular force faster rate of evaporation at a given temperature

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**



**Watch Video Solution**

**167.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instruction given below:

STATEMENT - 1 : An ideal solution obeys Raoult's law.

STATEMENT - 2 : In an ideal solution, solute-solvent as well as solvent-solvent, interactions are similar to solute - solvent interactions.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TURE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

**168.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : if a liquid solute more volatile than the solvent is added to the solvent, the vapour pressure of the solution is greater than vapour pressure of pure solvent.

STATEMENT - 2 : Vapour pressure of solution is equal to vapour pressure of solvent.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**



**Watch Video Solution**

**169.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instrution given below:

STATEMENT - 1 :  $\Delta V_{\text{mix}}$  and  $\Delta S_{\text{mix}}$ for an ideal solution is zero.

STATEMENT - 2 : A..B interaction in an ideal solution are same as between A...A and B...B.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TURE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TURE and STATEMENT-2 is FALSE



D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**



**Watch Video Solution**

**170.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Elevation in boiling point will be high if the molal elevation constant of the liquid is high.

STATEMENT - 2 : Elevation in boiling point is a colligative property.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



**Watch Video Solution**

**171.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given

below:

STATEMENT - 1 : The boiling point of 0.1 M urea solution is less than that of 0.1 M KCl solution.

STATEMENT - 2 : Elevation of boiling point is directly proportional to the number of moles of non-volatile solute particles present in the solution.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



Watch Video Solution

**172.** Statement-1 : The observed molar mass of acetic acid in benzene is more than the normal molar mass of acetic acid.

Statement-2 : Molecules of acetic acid dimerise in benzene due to hydrogen bonding.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

**173.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instrution given below:

STATEMENT - 1 : addition of ethylene glycol to water lowers the freezing point of water, therefore, used as antifreeze substance.

STATEMENT - 2 : Ethylene glycol is soluble in water.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



**Watch Video Solution**

**174.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Osmotic pressure is a colligative property.

STATEMENT - 2 : Osmotic pressure is developed in a column due to osmosis.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



**Watch Video Solution**

**175.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instrution given below:

STATEMENT - 1 : Osmotic involves movement of solvent molecules from lower concentration to higher concentration.

STATEMENT - 2 : Solutions having the same osmotic pressure are called isotonic solutions.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TURE but STATEMENT-2 is NOT the correct explanation STATEMENT-1



C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**

 [Watch Video Solution](#)

**176.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Isotonic solutions must have the same molal concentration.

STATEMENT - 2 : Solution which have the same osmotic pressure are known as isotonic solution.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**



**Watch Video Solution**

**177.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given

below:

STATEMENT - 1 : Isotonic solutions do not show phenomenon of osmosis.

STATEMENT - 2 : Isotonic solutions have same molal concentration at same temperature.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

**178.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instrution given below:

STATEMENT - 1 : When dried fruits and vegetables are placed in water, they slowly get swollen.

STATEMENT - 2 : It happens due to the phenomenon of osmosis.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TURE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

**179.** Assertion: Reverse osmosis is used in the desalination of sea water.

Reason: When pressure more than osmotic pressure is applied, pure water is squeezed out of the sea water through the membrane.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: B**



**Watch Video Solution**

**180.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : All solute becomes more soluble in water

at higher temperature.

STATEMENT - 2 : Solubility of solute depends upon temperature.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**



**Watch Video Solution**

**181.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Henry's law is always applicable for gases.

ItSTATEMENT - 2 : Raoult's law is a special case of Henry's law.

A. If both the statements are TRUE and STATEMENT-2 is

the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is

NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE



**Answer: D**

 [Watch Video Solution](#)

**182.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Increasing pressure on pure water decrease its freezing point.

STATEMENT - 2 : Density of water is maximum at 273 K.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**

 [Watch Video Solution](#)

**183.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : The molecular mass of acetic acid

determined by depression in freezing point method in benzene and water was found to be different.

STATEMENT - 2 : Water is polar and benzene is non-polar.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: A**



**Watch Video Solution**

**184.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer accoring to the instrution given below:

STATEMENT - 1 : If red blood cells were removed from the body and placed in pure water, pressure inside the cell increases.

STATEMENT - 2 : The concentration of the salt content in the cells increases.

A. If both the statements are TURE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TURE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TURE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: C**

 [Watch Video Solution](#)

**185.** Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Azeotrope is a binary mixture formed by ideal solutions.

STATEMENT - 2 : Azeotrope boils with unchanged composition.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

**Answer: D**

 [Watch Video Solution](#)

**186.** The vapour pressure of two pure liquids A and B are 5 and 10 torr respectively. Calculate the total pressure of

the solution (in torr) obtained by mixing 2 mole of A and 3 mole of B.

 [Watch Video Solution](#)

**187.** The vapour pressure of two liquids P and Q are 80 torr and 60 torr respectively. The total vapour pressure obtained by mixing 3 mole of P and 2 mole of Q would be

 [Watch Video Solution](#)

**188.** The vapour pressure of a liquid solution containing A and B is 99 torr. Calculate mole % of B in vapour phase.

(Given :  $P_{A^\circ} = 100T$  or  $r$ ,  $P_{B^\circ} = 80T$ )

 [Watch Video Solution](#)

[Watch Video Solution](#)

**189.** If 30 g a solute of molecular mass 154 is dissolved in 250 g of benzene. What will be the elevation in boiling point of the resulting solution ?

(Given :  $K_B(C_6H_6) = 2.6Kkgmol^{-1}$ )

[Watch Video Solution](#)

**190.** Calculate elevation in boiling point for 2 molal aqueous solution of glucose.

(Given  $K_b(H_2O) = 0.5kgmol^{-1}$ )

[Watch Video Solution](#)



**191.** Calculate depression of freezing point for 0.56 molal aq. Solution of KCl.

(Given :  $K_f(H_2O) = 1.8 \text{ kg mol}^{-1}$ ).



[Watch Video Solution](#)

**192.** What is the maximum value of van't Hoff factor for  $AlCl_3$  ?



[Watch Video Solution](#)

**193.** A solution containing 500 g of a protein per liter is isotonic with a solution containing 3.42 g sucrose per liter. The molecular mass of protein is  $5 \times 10^x$ , hence x is.



[Watch Video Solution](#)

 Watch Video Solution

**194.** An aqueous solution of urea has a freezing point of  $-0.515^{\circ}\text{C}$ . Predict the osmotic pressure (in atm) of the same solution at  $37^{\circ}\text{C}$ .

 Watch Video Solution

**195.** 0.2 aq. Solution of KCl is isotonic with 0.2 M  $\text{K}_2\text{SO}_4$  at same temperature. What is the van't Hoff factor of  $\text{K}_2\text{SO}_4$ ?

 Watch Video Solution

1. A saturated solution of  $XCl_3$  has a vapour pressure 17.20 mm Hg at  $20^\circ C$ , while pure water vapour pressure is 17.25 mm Hg. Solubility product ( $K_{sp}$ ) of  $XCl_3$  at  $20^\circ C$  is :

A.  $9.8 \times 10^{-2}$

B.  $10^{-5}$

C.  $2.56 \times 10^{-6}$

D.  $7 \times 10^{-5}$

**Answer: d**



**Watch Video Solution**

## Level 3 Match The Column

1. Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of column-

Column-I
(A) <i>n</i> -hexane + <i>n</i> -heptane
(B) Acetone + chloroform
(C) Chlorobenzene and bromobenzene
(D) Ethanol + water

Column-II
(P) Can be separated by fractional distillation
(Q) Maximum boiling azeotrope
(R) Cannot be separated by fractional distillation completely
(S) Minimum boiling azeotrope

II.



Watch Video Solution