



PHYSICS

BOOKS - HC VERMA

ROTATIONAL MECHANICS

Example

1. The motor of an engine is rotating about its axis with an angular velocity of 100 rev/minute. It comes to rest in 15 s, after being

switched off. Assuming constant angular deceleration, calculate the number of revolutions made by it before coming to rest.



[Watch Video Solution](#)

2. Starting from rest, a fan takes five seconds to attain the maximum speed of 400 rpm (revolutions per minute). Assuming constant acceleration find the time taken by the fan in attaining half the maximum speed.




[Watch Video Solution](#)

3. A bucket is being lowered down into a well through a rope passing over a fixed pulley of radius 10 cm. Assume that the rope does not slip on the pulley. Find the angular velocity and angular acceleration of the pulley at an instant when the bucket is going down at a speed of 20 cm/s and has an acceleration of $4.0 \frac{m}{s^2}$.



Watch Video Solution

4. Consider a pulley fixed at its centre of mass by a clamp. A light rope is wound over it and the free end is tied to block. The tension in the rope is T . a. Write the forces acting on the pulley. How are they related? B. Locate the axis of rotation. c. Find the torque of the forces about the axis of rotation.






Watch Video Solution

5. A wheel of radius 10 cm can rotate freely about its centre as shown in figure. A string is wrapped over its rim and is pulled by a force of 5.0 N. It is found that the torque produces an angular acceleration 2.0 rads^{-2} in the wheel. Calculate the moment of inertia of the

wheel.



Watch Video Solution

6. A wheel is rotating at an angular speed ω about its axis which is kept vertical. An identical wheel initially at rest is gently dropped into the same axle and the two


wheels start rotating with a common angular speed. Find this common angular speed.



[Watch Video Solution](#)

7. A wheel of moment of inertia I and radius r is free to rotate about its centre as shown in figure. A string is wrapped over its rim and a block of mass m is attached to the free end of the string. The system is released from rest. Find the speed of the block as it descends

through a height h .




Watch Video Solution

8. Consider a light rod with two heavy mass particles at its ends. Let AB be a line

perpendicular to the rod as shown in figure.


What is the moment of inertia of the system about AB?



[Watch Video Solution](#)

9. Three particles, each of mass m are situated at the vertices of an equilateral triangle ABC of side L as shown in the figure. Find the moment

of inertia of the system about the line AX
perpendicular to AB in the plane of ABC



Watch Video Solution

10. Find the moment of inertia of a uniform ring of mass M and radius R about a diameter.



Watch Video Solution

11. Find the moment of inertia of a solid cylinder of mass M and radius R about a line parallel to the axis of the cylinder and on the surface of the cylinder.



Watch Video Solution

12. A uniform sphere of mass 200 g rolls without slipping on a plane surface so that its centre moves at a speed of 2.00 cm/s. Find its kinetic energy.



[Watch Video Solution](#)

Worked Out Examples

1. A wheel rotates with a constant acceleration of $2.0 \frac{\text{rad}}{\text{s}^2}$. If the wheel starts from rest, how

many revolutions will it make in the first 10 seconds?



[Watch Video Solution](#)

2. The wheel of a motor, accelerated uniformly from rest, rotates through 2.5 radian during the first second. Find the angle rotated during the next second.



[Watch Video Solution](#)

3. A wheel having moment of inertia 2 kg m^2 about its axis, rotates at 50 rpm about this axis. Find the torque that can stop the wheel in one minute.



[Watch Video Solution](#)

4. A string is wrapped around the rim of a wheel of moment of inertia 0.20 kg-m^2 and radius 20 cm. The wheel is free to rotate about its axis. Initially, the wheel is at rest. The string


is now pulled by a force of 20 N. Find the angular velocity of the wheel after 5.0 seconds.



[Watch Video Solution](#)

5. A wheel of radius r and moment of inertia I about its axis is fixed at top of an inclined plane of inclination θ as shown in figure. A string is wrapped round the wheel and its free end supports a block of mass M which can slide on the plane. Initially, the wheel is rotating at a speed ω in direction such that


the block slides up the plane. How far will the block move before stopping?



[Watch Video Solution](#)

6. The pulley shown in figure has a moment of inertia I about its axis and its radius is R . Find the magnitude of the acceleration of the two blocks. Assume that the string is light and

does not slip on the pulley.



[Watch Video Solution](#)

7. Two small kids weighing 10 kg and 15 kg respectively are trying to balance a seesaw of

total length 5.0 with the fulcrum at the centre.

If one of the kids is sitting at an end where should the other sit?



[Watch Video Solution](#)


8. A uniform ladder of mass 10 kg leans against a smooth vertical wall making an angle of 53° with it. The other end rests on a rough horizontal floor. Find the normal force and the frictional force that the floor exerts on the ladder



Watch Video Solution

9. The ladder shown in figure has negligible mass and rests on a frictionless floor. The crossbar connects the two legs of the ladder at the middle. The angle between the two legs is 60° . The fat person sitting on the ladder has a mass of 80 kg. Find the contact force exerted by the floor on each leg and the tension in the

cross bar.



[Watch Video Solution](#)

10. Two small balls A and B each of mass m , are attached tightly to the ends of a light rod of

length d . The structure rotates about the perpendicular bisector of the rod at an angular speed ω . Calculate the angular momentum of the individual balls and of the system about the axis of rotation.



[Watch Video Solution](#)

11. Two particles of mass m each are attached to a light rod of length d , one at its centre and the other at a free end, The rod is fixed at the other end and is rotated in a plane at an

angular speed ω . Calculate the angular momentum of the particle at the end with respect to the particle at the centre



[Watch Video Solution](#)

12. A ball is thrown at a speed of 40 m/s at an angle of 60° with the horizontal. Find a. the maximum height reached and b. the range of the ball. Take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

13. A uniform circular disc of mass 200 g and radius 4.0 cm is rotated about one of its diameter at an angular speed of 10 rad/s. Find the kinetic energy of the disc and its angular momentum about the axis of rotation.



Watch Video Solution

14. A wheel rotating at an angular speed of 20 rad/s is brought to rest by a constant torque in 4.0 seconds. If the moment of inertia of the wheel about the axis of rotation is 0.20

$kg - m^2$ find the work done by the torque in the first two seconds.



[Watch Video Solution](#)

15. Two masses M and m are connect by a light string gong over a pulley of radis r . The pulley is free to rotate about its axis which is kept horizontal. The moment of inertia of the pulley about the axis is I . The system is releaed from rest. Find the angular momentum fo teh system when teh mass M has descended

through a height h . The string does not slip over the pulley.



[Watch Video Solution](#)

16. Figure shows a mass m placed on a frictionless horizontal table and attached to a string passing through a small hole in the surface. Initially, the mass moves in a circle of radius r_0 with a speed v_0 and the free end of the string is held by a person. The person pulls on the string slowly to decrease the radius of

th circle of r. a. Find the tension in the string when the mass moves in the circle of radius r.

b. Calculate the change in the kinetic energy of the mass



[Watch Video Solution](#)

17. A light rod of length has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is



[Watch Video Solution](#)

18. Four particles each of mass ' m ' are kept at the four corners of a square of edge ' a '. Find the moment of inertia of the system about a line perpendicular to the plane of the square and passing through the center of the square.



[Watch Video Solution](#)

19. Two identical spheres each of mass 1.20 kg and radius 10.0 cm are fixed at the ends of a


light rod so that the separation between the centers is 50.0 cm. Find the moment of inertia of the system about an axis perpendicular to the rod passing through its middle point.



[Watch Video Solution](#)

20. Two uniform identical rods each of mass M and length l are joined to form a cross as shown in figure. Find the moment of inertia of the cross about a bisector as shown dotted in

the figure



Watch Video Solution

21. A uniform rod of mass M and length a lies on a smooth horizontal plane. A particle of

mass m moving at a speed v perpendicular to the length of the rod strikes it at a distance $\frac{a}{4}$ from the centre and stops after the collision. Find a. the velocity of the centre of the rod and b. the angular velocity of the rod about its centre just after the collision.



[Watch Video Solution](#)

22. A wheel of perimeter 220 cm rolls on a level road at a speed of 9 km/h. How many revolutions does the wheel make per second?



[Watch Video Solution](#)

23. A cylinder is released from rest from the top of an incline of inclination θ and length 'L'. If the cylinder rolls without slipping. What will be its speed when it reaches the bottom?



[Watch Video Solution](#)


24. A sphere rolls down an inclined plane of inclination θ . What is the acceleration as the sphere reaches bottom ?



Watch Video Solution

25. Figure shows two cylinders of radii r_1 and r_2 having moments of inertia I_1 and I_2 about their respective axes. Initially the cylinders rotate about their axes with angular speed ω_1 and ω_2 as shown in the figure. The cylinders are moved closer to touch each other keeping the axes parallel. The cylinders first slip over each other at the contact but the slipping finally ceases due to the friction between them. Find the angular


speeds of the cylinders after the slipping ceases.



[Watch Video Solution](#)

26. A cylinder of mass m is suspended through two strings wrapped around it as shown in figure . Find a. the tension T in the string and b. the speed of the cylinder as it falls through a

distance h .



[Watch Video Solution](#)

27. A force F acts tangentially at the highest point of a sphere of mass m kept on a rough horizontal plane. If the sphere rolls without


slipping, find the acceleration of the centre of the sphere.



[Watch Video Solution](#)

28. A sphere of mass M and radius r shown in figure slips on a rough horizontal plane. At some instant it has translational velocity V_0 and rotational velocity about the centre $\frac{v_0}{2r}$. Find the translational velocity after the sphere

starts pure rolling.




[Watch Video Solution](#)

29. The sphere shown in figure lies on a rough plane when a particle of mass m travelling at a speed v_0 collides and sticks with it. If the line of motion of the particle is at a distance h

above the plane, find a. the linear speed of the combined system just after the collision b. the angular speed of the system about the centre of the sphere just after the collision c. the value of h for which the sphere starts pure rolling on the plane Assume that the mass M of the sphere is large compared to the mass of the particle so that the centre of mass of the combined system is not appreciably shifted from the

centre of the sphere.



[Watch Video Solution](#)

Questions For Short Answer

1. Can an object be in pure translation as well as in pure rotation?



[Watch Video Solution](#)

2. The vibrations of a simple pendulum are also known as ____.



Watch Video Solution

3. In a rotating body $a = \alpha r$ and $v = \omega r$.

Thus $\frac{a}{\alpha} = \frac{v}{\omega}$. Can you use the theorems of

ratio and proportion studied in algebra so as

to write

$$\frac{a + \alpha}{a - \alpha} = \frac{v + \omega}{v - \omega}$$




[Watch Video Solution](#)

4. A ball is whirled in a circle by attaching it to a fixed point with a string. Is there an angular rotation of the ball about its centre? If yes, is this angular velocity equal to the angular velocity of the ball about the fixed point?



[Watch Video Solution](#)

5. In the given figure, the value of x is



[Watch Video Solution](#)

6. The torque of the weight of any body about any vertical axis is zero. Is it always correct?



Watch Video Solution

7. The torque of a force \vec{F} about a point is defined as $\vec{\Gamma} = \vec{r} \times \vec{F}$. Suppose \vec{r} , \vec{F} and $\vec{\Gamma}$ are all nonzero. Is $\vec{r} \times \vec{\Gamma} \parallel \vec{F}$ always true? Is it ever true?



Watch Video Solution

8. The motion of a particle of mass m is described by $h = ut + 1/2gt^2$. Find the force

acting on particle.



[Watch Video Solution](#)

9. If several forces act on a particle, the total torque on the particle may be obtained by first finding the resultant force and then taking torque of this resultant. Prove this. Is this result valid for the forces acting on different particles of a body in such a way that their lines of action intersect at a common point?



[Watch Video Solution](#)

10. The force acting on a body it is a rest



[Watch Video Solution](#)

11. If the angular momentum of a body is found to be zero about a point is it necessary that it will also be zero about a different point?



[Watch Video Solution](#)

12. If the resultant torque of all the forces acting on a body is zero about a point is it necessary that it will be zero about any other point?



Watch Video Solution

13. A body is in translational equilibrium under the action of coplanar forces. If the torque of these forces is zero about a point is it necessary that it will also be zero about any other point?



[Watch Video Solution](#)


14. A rectangular brick is kept on a table with a part of its length projecting out. It remains at rest if the length projected is slightly less than half the total length but it falls down if the length projected is slightly more than half the total length. Give reason.



[Watch Video Solution](#)

15. When a fat person tries to touch his toes, keeping the legs straight he generally falls.

Explain with reference to figure.



Watch Video Solution

16. A ladder makes an angle of 60° with the ground, when placed against a wall. If the foot of the ladder is 2m away from the wall, then the length of the ladder (in metres) is



Watch Video Solution

17. When body is weighted on an ordinary balance we demand that the arm should be horizontal if the weights on the two pans are equal. Suppose equal weights are put on the

two pawns, the arm is kept at an angle with the horizontal and released. Is the torque of the two weights about the middle point (point of support) zero? Is the total torque zero? If so, why does the arm rotate and finally become horizontal?



[Watch Video Solution](#)

18. The density of a rod AB continuously increases from A to B. Is it easier to set it in rotation by clamping it at A and applying a

perpendicular force at B or by clamping it at B and applying the force at A?



[Watch Video Solution](#)

19. The net force acting on a body when it is at rest is



[Watch Video Solution](#)

20. If the ice at the poles melts and flows towards the equator, how will it affect the

duration of day and night?



[Watch Video Solution](#)

21. A hollow sphere and a solid sphere having same mass and same radii are rolled down a rough inclined plane.



[Watch Video Solution](#)

22. A sphere rolls on a horizontal surface. Is there any point of the sphere which has a

vertical velocity?



Watch Video Solution

Objective 1

1. Let \vec{A} be a unit vector along the axis of rotation of a purely rotating body and \vec{B} be a unit vector along the velocity of a particle P of the body away from the axis. The value of $\vec{A} \cdot \vec{B}$ is

A. 1

B. -1

C. 0

D. none of these

Answer: C



Watch Video Solution

2. Let \vec{A} be a unit vector along the axis of rotation of a purely rotating body and \vec{B} be a unit vector along the velocity of a particle P of

the body away from the axis. The value of

$\vec{A} \cdot \vec{B}$ is

A. 1

B. -1

C. 0

D. none of these

Answer: C



Watch Video Solution

3. A particle moves with a constant velocity parallel to the X-axis. Its angular momentum with respect to the origin

A. is zero

B. remains constant

C. goes on increasing

D. goes on decreasing

Answer: B



Watch Video Solution

4. A body is in pure rotation. The linear speed 'v' of a particle, the distance 'r' of the particle from the axis and the angular velocity ω of the body are related as $\omega = \frac{v}{r}$. Thus

A. $\omega \propto \frac{1}{r}$

B. $\omega \propto r$

C. $\omega = 0$


D. ω is independent of r

Answer: D



Watch Video Solution

5. From the figure find x and y .



A. $x = 2y$

B. $x = y$

C. $y = 2x$

D. none of these

Answer: C



Watch Video Solution

6. A body is rotating uniformly about a vertical axis fixed in an inertial frame. The resultant force on a particle of the body not on axis is

A. vertical

B. horizontal skew with the axis

C. horizontal and intersecting the axis

D. none of these

Answer: C



Watch Video Solution

7. A body is rotating uniformly about a vertical axis fixed in an inertial frame. The resultant force on a particle of the body not on axis is

A. vertical

B. horizontal and skew with the axis

C. horizontal and intersection

D. none of these

Answer: B



Watch Video Solution

8. If F is the force acting on a particle having position vector \vec{r} and $\vec{\tau}$ be the torque of this force about the origin, then,

A. $\vec{r} \cdot \vec{\Gamma} = 0$ and $\vec{F} \cdot \vec{\Gamma} = 0$

B. $\vec{r} \cdot \vec{\Gamma} = 0$ but $\vec{F} \cdot \vec{\Gamma} \neq 0$

C. $\vec{r} \cdot \vec{\Gamma} \neq 0$ but $\vec{F} \cdot \vec{\Gamma} = 0$

D. $\vec{r} \cdot \vec{\Gamma} \neq 0$ and $\vec{F} \cdot \vec{\Gamma} \neq 0$

Answer: A



Watch Video Solution

9. One end of a uniform rod of mass m and length l is clamped. The rod lies on a smooth horizontal surface and rotates on it about the

clamped end at a uniform angular velocity ω .

The force exerted by the clamp on the rod has a horizontal component

A. $m\omega^2 l$

B. *zero*

C. mg

D. $\frac{1}{2}m\omega^2 l$

Answer: D



Watch Video Solution

10. A uniform rod is kept vertically on a horizontally smooth surface at a point O. IF it is rotated slightly and released, it falls down on the horizontal surface. The lower end will remain

A. at O

B. at a distance less than $\frac{l}{2}$ from O

C. at a distance $\frac{l}{2}$ from O

D. at a distance larger than $\frac{l}{2}$ from O

Answer: C



11. A circular disc A of radius r is made from an iron plate of thickness t and another circular disc B of radius $4r$ is made from an iron plate of thickness $t/4$. The relation between the moments of inertia I_A and I_B is

A. $I_A > I_B$

B. $I_A = I_B$

C. $I_A < I_B$

D. depends on the actual values of t and r

Answer: C



Watch Video Solution

12. Equal torques act on the discs A and B of the previous problem, initially both being at rest. At a later instant, the linear speeds of a point on the rim of A and another point on the rim of B are V_A and V_B respectively. We have

A. $V_A < V_B$

B. $V_A = V_B$

C. $V_A < V_B$

D. the relation depends on the actual magnitude of the torques

Answer: A



Watch Video Solution

13. A closed tube partly filled with water lies in a horizontal plane. If the tube is rotated about a perpendicular bisector, the moment of inertia of the system.

A. increases

B. decreases

C. remains constant

D. increases if the rotation is clockwise and
decreases if it is anticlockwise

Answer: A



Watch Video Solution

14. The moment of inertia of a uniform semi-circular wire of mass 'M' and radius 'r' about a line perpendicular to the plane of the wire through the center is

A. a) Mr^2

B. b) $\frac{1}{2}Mr^2$

C. c) $\frac{1}{4}Mr^2$

D. d) $\frac{2}{5}Mr^2$

Answer: A



Watch Video Solution

15. Let I_1 and I_2 be the moments of inertia of two bodies of identical geometrical shape, the first made of aluminum and the second of iron.

A. $I_1 < I_2$

B. $I_1 = I_2$

C. $I_1 > I_2$

D. Relation between I_1 and I_2 depends on the actual shapes of the bodies

Answer: A



Watch Video Solution

16. A body having its centre of mass at the origin has three of its particles at $(a, 0, 0)$, $(0, a, 0)$, $(0, 0, a)$. The moments of inertia of the body about the X and Y axes are $0.20 \text{ kg} - m^2$ each. The moment of inertia about the Z-axis

A. is $0.20 \text{ kg} - m^2$

B. is $0.40 \text{ kg} - m^2$

C. is $0.20\sqrt{2} \text{ kg} - m^2$

D. cannot be deduced with this information.

Answer: D



Watch Video Solution

17. A cubical block of mass M and edge a slides down a rough inclined plane of inclination θ with a uniform velocity. The torque of the

normal force on the block about its centre has magnitude.

A. zero

B. Mga

C. $Mga \sin \theta$

D. $\frac{1}{2}Mga \sin \theta$

Answer: D



Watch Video Solution

18. A thin circular ring of mass M and radius r is rotating about its axis with an angular speed ω . Two particles having mass m each are now attached at diametrically opposite points.

The angular speed of the ring will become

A. a) $\frac{\omega M}{M + m}$

B. b) $\frac{\omega M}{M + 2m}$

C. c) $\frac{\omega(M - 2m)}{M + 2m}$

D. d) $\frac{\omega(M + 2m)}{M}$

Answer: B



Watch Video Solution

19. A man is sitting on a rotating stool with his arms outstretched. If suddenly he folds his arms the angular velocity of the man would

- A. increases
- B. decreases
- C. remains unchanged
- D. doubles

Answer: C



Watch Video Solution

20. The center of a wheel rolling on a plane surface moves with a speed v_0 . A particle on the rim of the wheel at the same level as the center will be moving at speed

A. *zero*

B. v_0

C. $\sqrt{2}v_0$

D. $2v_0$

Answer: C



Watch Video Solution

21. A wheel of radius 20 cm is pushed to move it on a rough horizontal surface. It is found to move through a distance of 60 cm on the road during the time it completes one revolution about the centre. Assume that the linear and the angular accelerations are uniform. The frictional force acting on the wheel by the surface is

A. along the velocity of the wheel

B. opposite to the velocity o the wheel

C. perpendicular to the velocity of the
wheel

D. zero

Answer: A



Watch Video Solution

22. The angular velocity of the engine (and hence of the wheel) on a scooter is proportional to the petrol input per second. The scooter is moving on a frictionless road with uniform velocity. If the petrol input is increased by 10% the linear velocity of the scooter is increased by

A. 0.5

B. 10

C. 0.2

D. 0

Answer: D



Watch Video Solution

23. A solid sphere, a hollow sphere and a disc, all having the same mass and radius, are placed at the top of an incline and released. The friction coefficients between the objects and the incline are same and not sufficient to

allow pure rolling. The least time will be taken in reaching the bottom by

- A. the solid sphere
- B. the hollow sphere
- C. the disc
- D. all will take same time

Answer: D



Watch Video Solution

24. A solid sphere, a ring and a disc all having same mass and radius are placed at the top of an incline and released. The friction coefficient between the objects and the incline are same but not sufficient to allow pure rolling. Least time will be taken in reaching the bottom by

- A. the solid sphere
- B. the hollow sphere
- C. the disc
- D. all will take same time

Answer: D



Watch Video Solution

25. In the previous question the smallest kinetic energy at the bottom of the incline will be achieved by

A. the solid sphere

B. the hollow sphere

C. the disc

D. all will achieve same kinetic energy


Answer: B



Watch Video Solution

26. A string of negligible thickness is wrapped several times around a cylinder kept on a rough horizontal surface. A man standing at a distance l from the cylinder holds one end of the string and pulls the cylinder towards him. There is no slipping anywhere. The length of the string passed through the hand of the man while the cylinder reaches his

hands is



A. 1

B. 2l

C. 3l

D. 4l

Answer: B



Watch Video Solution

Objective 2

1. The axis of rotation of a purely rotating body

A. must pass through the centre of mass

B. may pass through the centre of mass

C. must pass through a particle of the body

D. may pass through a particle of the body

Answer: B::D



Watch Video Solution

2. Consider the following two equations

$$(A). \vec{R} = \frac{1}{M} \sum_i m_i \vec{r}_i$$

$$\text{and } (B). \vec{a}_{CM} = \frac{\vec{F}}{M}$$

IN a noninertial frame

A. both A and B are true

B. A is true but B is false

C. B is true but A is false

D. both A and B are false

Answer: B



Watch Video Solution

3. A particle moves on a straight line with a uniform velocity. Its angular momentum

A. is always zero

B. is zero about a point on the straight line

C. is not zero about a point away from the straight line

D. about any given point remains constant

Answer: B::C::D



Watch Video Solution

4. If there is no external force acting on a nonrigid body, which of the following quantities must remain constant?

A. angular momentum

B. linear momentum

C. kinetic energy

D. moment of inertia

Answer: A::B



Watch Video Solution

5. Let I_A and I_B be moments of inertia of a body about two axes A and B respectively. The axis A passes through the centre of mass of the body but B does not. Then

A. $I_A < I_B$

B. If $I_A < I_B$ the axes are parallel

C. If the axes are parallel $I_A < I_B$

D. If the axes are not parallel $I_A \geq I_B$

Answer: C



Watch Video Solution

6. A sphere is rotating about a diameter

A. the particles on the surface of the sphere do not have any linear acceleration

B. the particles on the diameter mentioned above do not have any linear acceleration

C. different particles on the surface have different angular speeds.

D. All particles on the surface have same linear speed

Answer: B



Watch Video Solution

7. Find the moment of inertia of a uniform rod about an axis which is perpendicular to the rod and touches any one end of the rod.

A. angular acceleration

B. angular velocity when the rod completes
one rotation

C. angular momentum when the rod

competes one rotation


D. torque of the applied force

Answer: D



Watch Video Solution

8. Consider a wheel of a bicycle rolling on a level road at a linear speed v_0 figure.



- A. the speed of the particle A is zero
- B. the speed of B,C and D are all equal to v_0
- C. The speed of C is $2v_0$
- D. the speed of B is greater than the speed of O.

Answer: A::C::D



Watch Video Solution

9. Two uniform solid spheres having unequal radii are released from rest from the same height on a rough incline. If the spheres roll without slipping

A. the heavier sphere reaches the bottom first

B. the bigger sphere reaches the bottom first

C. the two spheres reach the bottom together

D. the information given is not sufficient to tell which sphere will reach the bottom first.

Answer: C



Watch Video Solution

10. A hollow sphere and a solid sphere having same mass and same radii are rolled down a rough inclined plane.

A. the hollow sphere reaches the bottom first

B. the solid sphere reaches the bottom with greater speed.

C. the solid sphere reaches the bottom with greater kinetic energy

D. the two spheres will reach the bottom
with same linear momentum

Answer: B



Watch Video Solution

11. A sphere cannot roll on

A. a smooth horizontal surface

B. a smooth inclined surface

C. a rough horizontal surface

D. a rough inclined surface

Answer: B



Watch Video Solution

12. In rear wheel drive cars the engine rotates the rear wheel and the front wheels rotate only because the car moves. If such a car accelerates on a horizontal road the friction

A. on the rear wheels is in the forward direction

B. on the front wheels is in the backward direction

C. on the rear wheels has larger magnitude than the friction on the front wheels

D. on the car is in the backward direction

Answer: A::B::C



Watch Video Solution

13. A sphere can roll on a surface inclined at an angle θ if the friction coefficient is more than $\frac{2}{7}g \sin \theta$. Suppose the friction coefficient is $\frac{1}{7}g \sin \theta$, and a sphere is released from rest on the incline,

- A. it will stay at rest
- B. it will make pure translation motion
- C. it will translate and rotate about the centre

D. the angular momentum of the sphere about its centre will remain constant

Answer: C



Watch Video Solution

14. A sphere is rolled on a rough horizontal surface. It gradually slows down and stops.

The force of friction tries to

A. decrease the linear velocity

B. increase the angular velocity

C. increase the linear momentum

D. decrease the angular velocity


Answer: A::B



Watch Video Solution

15. Figure shows smooth inclined plane fixed in a car accelerating on a horizontal road. The angle of incline θ is related to the acceleration a of the car as $a = g \tan \theta$. If the sphere is set

in pure rolling on the incline



- A. it will continue pure rolling
- B. it will slip down the plane
- C. its linear velocity will increase
- D. its linear velocity will slowly decrease.

Answer: A



Watch Video Solution

Exercises

1. A wheel is making revolutions about its axis with uniform angular acceleration. Starting from rest, till it reaches 100 rev/sec in 4 seconds. Find the angular acceleration.



[Watch Video Solution](#)

2. A wheel rotating with uniform angular acceleration covers 50 revolutions in the first five seconds after the start. Find the angular acceleration and the angular velocity at the end of five seconds.



[Watch Video Solution](#)

3. A wheel starting from rest is uniformly accelerated at 4 rad/s^2 for 10 seconds. It is allowed to rotate uniformly for the next 10

seconds and is finally brought to rest in the next 10 seconds. Find the total angle rotated by the wheel.



[Watch Video Solution](#)

4. A body rotates about a fixed axis with an angular acceleration of 1 rad/sec^2 . Through what angle does it rotate during the time in which its angular velocity increases from 5 rad/s to 15 rad/s.



[Watch Video Solution](#)

5. Find the angular velocity of a body rotating with an acceleration of $2re \frac{v}{s^2}$ as it completes the 5th revolution after the start



[Watch Video Solution](#)

6. A disc of radius 10 cm is rotating about its axis at an angular speed of 20 rad/s. Find the linear speed of

a. a point on the rim,

b. the middle point of the radius.



[Watch Video Solution](#)

7. A disc rotates about its axis with a constant angular acceleration of $4\pi a \frac{d}{s^2}$. Find the radial and tangential acceleration of a particle at a distance of 1 cm from the axis at the end of the first second after the disc starts rotating.



[Watch Video Solution](#)

8. A block hangs from a string wrapped on a disc of radius 20 cm free to rotate about its axis which is fixed in a horizontal position. If the angular speed of the disc is 10 rad/s at some instant, with what speed is the block going down at that instant?



Watch Video Solution

9. Three particles, each of mass 200 g are kept at the corners of an equilateral triangle of side

10 cm. Find the moment of inertia of the system about an axis joining two of the particles.



[Watch Video Solution](#)

10. Particles of masses 1g, 2g, 3g ...100g are kept at the marks 1cm, 2cm, 3cm ..., 100 cm respectively on a metre scale. Find the moment of inertia of the system of particles about a perpendicular bisector of the metre scale.



[Watch Video Solution](#)

11. Find the moment of inertia of a pair of spheres, each having a mass m and radius r kept in contact about the tangent passing through the point of contact.



[Watch Video Solution](#)

12. The moment of inertia of a uniform rod of mass 0.50 kg and length 1 m is 0.10 kg m^2 about a line perpendicular to the rod. Find the

distance of this line from the middle point of the rod.



[Watch Video Solution](#)

13. Find the radius of gyration of a circular ring of radius r about a line perpendicular to the plane of the ring and passing through one of this particles.



[Watch Video Solution](#)

14. The radius of gyration of a uniform disc about a line perpendicular to the disc equals to its radius. Find the distance of the line from the centre.



[Watch Video Solution](#)

15. Find the moment of inertia of a uniform square plate of mass M and edge a about one of its diagonals.



[Watch Video Solution](#)

16. The surface density (mass/area) of a circular disc of radius a depends on the distance from the centre as $\rho(r) = A + Br$. Find its moment of inertia about the line perpendicular to the plane of the disc through its centre.



Watch Video Solution

17. A particle of mass m is projected with speed u at an angle θ with the horizontal. Find

the torque of the weight of the particle about the point of projection when the particle is at the highest point.




[Watch Video Solution](#)

18. A simple pendulum of length l is pulled aside to make an angle θ with the vertical. Find the magnitude of the torque of the weight w of the bob about the point of suspension. When is the torque zero?




[Watch Video Solution](#)

19. When a force of 6.0 N is exerted at 30° to a wrench at a distance of 8 cm from the nut, it is just able to loosen the nut. What force F would be sufficient to loosen it if it acts perpendicularly to the wrench at 16 cm from the nut?



Watch Video Solution

20. Calculate the total torque acting on the body shown in figure about the point O.



[Watch Video Solution](#)

21. A cubical block of mass M and edge a slides down a rough inclined plane of inclination θ

with a uniform velocity. The torque of the normal force on the block about its centre has magnitude.



[Watch Video Solution](#)

22. A rod of mass m and length L , lying horizontally is free to rotate about a vertical axis through its centre. A horizontal force of constant magnitude F acts on the rod at a distance of $L/4$ from the centre. The force is always perpendicular to the rod. Find the

angle rotated by the rod during the time t after the motion starts.



[Watch Video Solution](#)

23. A square plate of mass 120g and edge 5.00 cm rotates about one of the edges. If it has a uniform angular acceleration of $0.2ra \frac{d}{s^2}$, what torque acts on the plate?



[Watch Video Solution](#)

24. Calculate the torque on the square plate of the previous problem if it rotates about a diagonal with the same angular acceleration.



Watch Video Solution

25. A flywheel of moment of inertia 5.0 kg m^2 is rotated at a speed of 60 rad/s . Because of the friction at the axle, it comes to rest in 5.0 minutes. Find a. The average torque of the friction. B. the total work done by the friction

and c. the angular momentum of the wheel 1 minute before it stops rotating.



[Watch Video Solution](#)

26. Because of the friction between the water in oceans with the earth's surface the rotational kinetic energy of the earth is continuously decreasing. If earth's angular speed decreases by 0.0016 rad/day in 100 years, find the average torque of the friction

on the earth. Radius of the earth is 6400 km and its mass is $6.0 \times 10^{24} \text{ kg}$.



[Watch Video Solution](#)

27. A flywheel rotating at a speed of 600 rpm about its axis is brought to rest by applying a constant torque for 10 seconds. Find the angular deceleration and angular velocity 5 second after the application of the torque.



[Watch Video Solution](#)

28. A wheel of mass 10 kg and radius 0.2 m is rotating at an angular speed of 100 rpm, when the motion is turned off. Neglecting the friction at the axis. Calculate the force that must be applied tangentially to the wheel to bring it to rest in 10 rev. Assumed wheel to be a disc.



Watch Video Solution

29. A cylinder rotating at an angular speed of 50 rev/s is brought in contact with an identical

stationary cylinder. Because of the kinetic friction, torques act on the two cylinders, accelerating the stationary one and decelerating the moving one. If the common magnitude of the acceleration and deceleration be one revolution per second square, how long will it take before the two cylinders have equal angular speed?




[Watch Video Solution](#)

30. The kinetic energy of a body is given by



Watch Video Solution

31. A light rod of length 1 m is pivoted at its centre and two masses of 5 kg and 2 kg are hung from the ends as shown in the figure. Find the initial angular acceleration of the rod assuming that it was horizontal in the beginning.



Watch Video Solution

32. Suppose the rod in the previous problem has a mass of 1 kg distributed uniformly over its length.

a. find the initial angular acceleration of the rod.


b. Find the tension in the supports to the blocks of mass 2 kg and 5kg`.



Watch Video Solution

33. Figure shows two blocks of masses m and M connected by a string passing over a pulley. The horizontal table over which the mass m slides is smooth. The pulley has a radius r and moment of inertia I about its axis and it can freely rotate about this axis. Find the acceleration of the mass M assuming that the


string does not slip on the pulley.



[Watch Video Solution](#)

34. A string is wrapped on a wheel of moment of inertia $0.20 \text{ kg}\cdot\text{m}^2$ and radius 10 cm and goes through a light pulley to support a block of mass 2.0 kg as shown in figure. Find the

acceleration of the block.



[Watch Video Solution](#)


35. Suppose the smaller pulley of the previous problem has its radius 5.0 cm and moment of inertia $0.10 \text{ kg}\cdot\text{m}^2$. Find the tension in the part of the string joining the pulleys.



Watch Video Solution

36. The pulleys in figure are identical, each having a radius R and moment of inertia I . Find


the acceleration of the block M .



 [Watch Video Solution](#)

37. The descending pulley shown in figure has a radius 20 cm and moment of inertia $0.20 \text{ kg}\cdot\text{m}^2$. The fixed pulley is light and the


horizontal plane frictionless. Find the acceleration of the block if its mass is 1.0 kg.



[Watch Video Solution](#)

38. The pulley shown in figure has a radius 10 cm and moment of inertia $0.5 \text{ kg}\cdot\text{m}^2$ about its axis. Assuming the inclined planes to be frictionless, calculate the acceleration of the

4.0 kg block.



[Watch Video Solution](#)

39. Solve the previous problem if the friction coefficient between the 2.0 kg block and the plane below it is 0.5 and the plane below the 4.0 kg block is frictionless.



[Watch Video Solution](#)

40. A uniform metre stick of mass 200 g is suspended from the ceiling through two vertical strings of equal lengths fixed at the ends. A small object of mass 20 g is placed on the stick at a distance of 70 cm from the left end. Find the tensions in the two strings.



Watch Video Solution

41. A uniform ladder of length 10.0 m and mass 16.0 kg is resting against a vertical smooth wall making an angle of 37° with it. An electrician weighing 60.0 kg climbs up the ladder. If he stays on the ladder at a point 8.00 m from the lower end, what will be normal force and the force of friction on the ladder by the ground? What should be the minimum coefficient of friction for the electrician to work safely?



Watch Video Solution

42. Suppose the friction coefficient between the ground and the ladder of the previous problem is 0.540. Find the maximum weight of a mechanic who could go up and do the work from the same position of the ladder.



Watch Video Solution

43. A 6.5 m long ladder rests against a vertical wall reaching a height of 6.0 m. A 60 kg man stands half way up the ladder. A. Find the

torque of the force exerted by the man on the ladder about the upper end of the ladder.

b. Assuming the weight of the ladder to be negligible as compared to the man and assuming the wall to be smooth find the force exerted by the ground on the ladder.



[Watch Video Solution](#)

44. the door of an almirah is 6 ft high, 1.5 ft wide and weights 8 kg. The door is supported by two hinges situated at a distance of 1 ft

from the ends. If the magnitude of the forces exerted by the hinges on the door are equal find this magnitude.



[Watch Video Solution](#)

45. A uniform rod of length L rests against a smooth roller as shown in figure. Find the friction coefficient between the ground and the lower end if the minimum angle that rod

can make with the horizontal is θ .




Figure 10-E9



Watch Video Solution

46. A uniform rod of mass 300 g and length 50 cm rotates at a uniform angular speed of 2 rad/s about an axis perpendicular to the rod through an end. Calculate a. the angular momentum of the rod about the axis of rotation b. its kinetic energy.



Watch Video Solution

47. A uniform square plate of mass 2.0 kg and edge 10 cm rotates about one of its diagonals

under the action of a constant torque of 0.10 Nm . Calculate the angular momentum and the kinetic energy of the plate at the end of the fifth second after the start.



[Watch Video Solution](#)

48. Calculate the ratio of the angular momentum of the earth about its axis due to its spinning motion to that about the sun due to its orbital motion. Radius of the earth

=6400 km and radius of the orbit of the earth about the sun = $1.5 \times 10^8 \text{ km}$.



Watch Video Solution

49. Two particles of masses m_1 and m_2 are joined by a light rigid rod of length r . The system rotates at an angular speed ω about an axis through the centre of mass of the system and perpendicular to the rod. Show that the angular momentum of the system is

$L = \mu r^2 \omega$ where μ is the reduced mass of the system defined as $\mu = \frac{m_1 m_2}{m_1 + m_2}$



Watch Video Solution

50. A dumb bell consists of two identical small balls of mass $1/2$ kg each connected to the ends of a 50 cm long light rod. The dumb bell is rotating about a fixed axis through the centre of the rod and perpendicular to it at an angular speed of 10 rad/s. An impulsive force of average magnitude 5.0 N acts on one of the

masses in the direction of its velocity for 0.10s.

Find the new angular velocity of the system.



[Watch Video Solution](#)

51. A wheel of moment of inertia $0.500\text{kg} - m^2$ and radius 20.0 cm is rotating about its axis at an angular speed of 20.0 rad/s. It picks up a stationary particle of mass 200 g at its edge. Find the new angular speed of the wheel.



[Watch Video Solution](#)

52. A diver having a moment of inertia of $6.0 \text{ kg} - \text{m}^2$ about an axis through its centre of mass rotates at an angular speed of 2 rad/s about this axis. If he folds his hands and feet to decrease the moment of inertia to $5.0 \text{ kg} - \text{m}^2$ what will be the new angular speed?



[Watch Video Solution](#)

53. A boy is seated in a revolving chair revolving at an angular speed of 120

revolutions per minute. Two heavy balls form part of the revolving system and the boy can pull the balls closer to himself or may push them apart. If by pulling the balls closer, the boy decreases the moment of inertia of the system from $6kg - m^2 \rightarrow 2kg - m^2$ what will be the new angular speed?



[Watch Video Solution](#)

54. A boy is standing on a platform which is free to rotate about its axis. the boys holds an

open umbrella in his hand. The axis of the umbrella coincides with that of the platform. The moment of inertia of the platform plus the boy system is $3.0 \times 10^{-3} \text{ kg} \cdot \text{m}^2$ and that of the umbrella is $2.0 \times 10^{-3} \text{ kg} \cdot \text{m}^2$. The boy starts spinning the umbrella about the axis at an angular speed of 2 rev/s with respect to himself. Find the angular velocity imparted to the platform.



[Watch Video Solution](#)

55. A wheel of moment of inertia $0.10 \text{ kg} \cdot \text{m}^2$ is rotating about a shaft at an angular speed of $160 \frac{\text{rev}}{\text{min}}$. A second wheel is set into rotation at 300 rev/minute and is coupled to the same shaft so that both the wheels finally rotate with a common angular speed of 200 rev/minute . Find the moment of inertia of the second wheel.



[Watch Video Solution](#)

56. A kid of mass M stands at the edge of a platform of radius R which can be freely rotated about its axis. The moment of inertia of the platform is I . The system is at rest when a friend throws a ball of mass m and the kid catches it. If the velocity of the ball is v horizontally along the tangent to the edge of the platform when it was caught by the kid find the angular speed of the platform after the event.



[Watch Video Solution](#)

57. Suppose the platform of the previous problem is brought to rest with the ball in the hand of the kid standing on the rim. The kid throws the ball horizontally to his friend in a direction tangential to the rim with a speed v as seen by his friend. Find the angular velocity with which the platform will start rotating.



Watch Video Solution

58. Suppose the platform with the kid in the previous problem is rotating in anticlockwise direction at an angular speed ω . The kid starts walking along the rim with a speed v relative to the platform also in the anticlockwise direction. Find the new angular speed of the platform.



Watch Video Solution

59. A uniform rod of mass m and length l is struck at an end by a force F perpendicular to the rod for a short time interval t . Calculate

- the speed of the centre of mass ,
- the angular speed of the rod about centre of mass,
- the kinetic energy of the rod and
- the angular momentum of the rod about the centre of mass after the force has stopped to act. Assume that t is so small that the rod does not appreciably change its direction while the force acts.



Watch Video Solution

60. A uniform rod of length L lies on a smooth horizontal table. A particle moving on the table strikes the rod perpendicularly at an end and stops. Find the distance travelled by the centre of the rod by the time it turns through a right angle. Show that if the mass of the rod is four times that of the particle, the collision is elastic



Watch Video Solution

61. Suppose the particle of the previous problem has a mass m and a speed v before the collision and it sticks to the rod after the collision. The rod has a mass M . a. Find the velocity of the particle with respect to C of the system consisting of the rod plus the particle. b. Find the velocity of the particle with respect to C before the collision. c. Find the velocity of the rod with respect to C before the collision. d. Find the moment of inertia of the system about the vertical axis through the centre of mass C after the collision. e. Find the velocity of the rod with respect to C after the collision. f. Find the velocity of the particle with respect to C after the collision.

of the centre of mass C and the angular velocity of the system about the centre of mass after the collision.



[Watch Video Solution](#)

62. Two small balls A and B , each of mass m , are joined rigidly by a light horizontal rod of length L . The rod is clamped at the centre in such a way that it can rotate freely about a vertical axis through its centre. The system is rotated with an angular speed ω about the axis. A

particle P of mass m kept at rest sticks to the ball A as the ball collides with it. Find the new angular speed of the rod.




[Watch Video Solution](#)

63. Two small balls A and B each of mass m , are joined rigidly to the ends of a light rod of length L figure. The system translates on a frictionless horizontal surface with a velocity v_0 in a direction perpendicular to the rod. A particle P of mass kept at rest on the surface

sticks to the ball A as the ball collides with it .

Find

- the linear speeds of the balls A and B after the collision,
- the velocity of the centre of mass C of the system A+B+P and
- the angular speed of the system about C after the collision.



Watch Video Solution

64. Suppose the rod with the balls A and B of the previous problem is clamped at the centre in such a way that it can rotate freely about a horizontal axis through the clamp. The system is kept at rest in the horizontal position. A particle P of the same mass m is dropped from a height h above the ball B. The particle collides with B and sticks to it. a. Find the angular momentum and the angular speed of the system just after the collision. b. What should

be the minimum value of h so that the system makes a full rotation after the collision.



[Watch Video Solution](#)

65. Two blocks of masses 400 g and 200 g are connected through a light string going over a pulley which is free to rotate about its axis.

The pulley has a moment of inertia $1.6 \times 10^{-4} \text{ kg} \cdot \text{m}^2$ and a radius 2.0 cm. Find

a. the kinetic energy of the system as the 400

block falls through 50 cm. b. the speed of the block at this instant.




[Watch Video Solution](#)

66. The pulley shown in figure has a radius of 20 cm and moment of inertia $0.2 \text{ kg}\cdot\text{m}^2$. The string going over it is attached at one end to a vertical spring of spring constant 50 N/m fixed from below and supports a 1 kg mass at other end. The system is released from rest with the spring at its natural length. Find the speed of

the block when it has descended through 10 cm.

Take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

67. A metre stick is held vertically with one end on a rough horizontal floor. It is gently

allowed to fall on the floor. Assuming that the end at the floor does not slip find the angular speed of the rod when it hits the floor.



[Watch Video Solution](#)

68. A metre stick weighing 240 g is pivoted at its upper end in such a way that it can freely rotate in a vertical plane through this end figure. A particle of mass 100 g is attached to the upper end of the stick through a light string of length 1 m. Initially the rod is kept

vertical and the string horizontal when the system is released from rest. The particle collides with the lower end of the stick and sticks there. Find the maximum angle through which the stick will rise.




Figure 10-E9



Watch Video Solution

69. A uniform rod pivoted at its upper end hangs vertically. It is displaced through an angle of 60° and then released. Find the magnitude of the force acting on a particle of mass dm at the tip of the rod when the rod makes an angle of 37° with the vertical.



Watch Video Solution

70. A cylinder rolls on a horizontal plane surface. If the speed of the centre is 25 m/s, what is the speed of the highest point?



Watch Video Solution

71. A sphere of mass m rolls on a plane surface. Find its kinetic energy at an instant when its centre moves with speed v .



Watch Video Solution

72. A string is wrapped over the edge a uniform disc and the free end is fixed with the ceiling. The disc moves down, unwinding the string. Find the downward acceleration of the disc.



Watch Video Solution

73. A small spherical ball is released from a point at a height h on a rough track shown in figure. Assuming that it does not slip anywhere, find its linear speed when it rolls on

the horizontal part of the track.



[Watch Video Solution](#)

74. A small disc is set rolling with a speed v on the horizontal part of the track of the previous problem from right to left. To what height will it climb up the curved part?



[Watch Video Solution](#)

75. A sphere starts rolling down an incline of inclination θ . Find the speed of its centre when it has covered a distance l .



Watch Video Solution

76. A hollow sphere is released from the top of an inclined plane of inclination θ . A. What should be the minimum coefficient of friction between the sphere and the plane to prevent sliding? B. Find the kinetic energy of the ball


as it moves down a length l on the incline if the friction coefficient is half the value calculated in part a.



[Watch Video Solution](#)

77. Consider the situation as shown in the figure. A solid sphere of mass m is released from rest from the rim of a hemispherical cup so that it rolls along the surface. Find the normal contact force between the solid sphere

and the cup at the bottom most point.



[Watch Video Solution](#)

78. Figure shows a rough track a portion of which is in the form of a cylinder of radius R . With what minimum linear speed should a sphere of radius r be set rolling on the

horizontal part so that it completely goes round the circle on the cylindrical part.



[Watch Video Solution](#)

79. Figure shows a small spherical ball of mass m rolling down the loop track. The ball is released on the linear portion at a vertical height H from the lowest point. The circular


part show has a radius R .

a. find the kinetic energy of the ball when it is at a point where the radius makes an angle θ with the horizontal.

Find the radial and the tangential accelerations of the center when the ball is at

A. c. find the normal force and the frictional force acting on the ball if $H=60$ cm, $R=10$ cm

$\theta = 0$ and $m=70$ fg.



Watch Video Solution

80. A hollow sphere of radius R lies on a smooth horizontal surface. It is pulled by a horizontal force acting tangentially from the highest point. Find the distance travelled by

the sphere during the time it makes one full rotation.



[Watch Video Solution](#)

81. A uniform wheel of radius R is set lying on a rough horizontal surface. It is hit by a cue in such a way that the line of action passes through the centre of the wheel. As a result the wheel starts moving with a linear speed v without any initial angular velocity. Find the

linear speed of the shell after it starts pure rolling on the surface.



[Watch Video Solution](#)

82. A uniform wheel of radius R is set lying on a rough horizontal surface is hit by a cue in such a way that the line of action passes through the centre of the shell. As a result the shell starts moving with a linear speed v without any initial angular velocity. Find the

linear speed of the shell after it starts pure rolling on the surface.



[Watch Video Solution](#)

83. _____ is the distance travelled by a body per unit time.



[Watch Video Solution](#)

84. A solid sphere of mass 0.50 kg is kept on a horizontal surface. The coefficient of static

friction between the surfaces in contact is $\frac{2}{7}$.


What maximum force can be applied at the highest point in the horizontal direction so that the sphere does not slip on the surface?



[Watch Video Solution](#)

85. A solid sphere is set into motion on a rough horizontal surface with a linear speed v in the forward direction and an angular speed $\frac{v}{R}$ in the anticlockwise direction as shown in figure . Find the linear speed of the sphere a.

where it stops rotating and b. when slipping finally ceases and pure rolling starts.



[Watch Video Solution](#)

86. A solid sphere rolling on a rough horizontal surface with a linear speed v collides elastically with a fixed, smooth, vertical wall. Find the

speed of the sphere after it has started pure rolling in the backward direction.



Watch Video Solution