



CHEMISTRY

BOOKS - NARENDRA AWASTHI

CHEMICAL EQUILIBRIUM

Exercise

1. A reversible reaction is one which

- A. proceeds in one direction
- B. proceeds in both directions
- C. proceeds spontaneously
- D. all the statements are wrong

Answer: b



Watch Video Solution

2. The equilibrium constant K_c for the reaction



is 1.4 at $400^\circ C$. Suppose that 3 moles of $P_4(g)$ and 2 moles of $P_2(g)$ are mixed in 2 litre container at $400^\circ C$. What is the value of reaction quotient (Q_c)?

A. $\frac{3}{2}$

B. $\frac{2}{3}$

C. 1

D. none of these

Answer: b

[Watch Video Solution](#)

3. In a chemical reaction, equilibrium is said to have been established when the

- A. opposing reaction ceases
- B. concentrations of reactants and product are equal
- C. velocity of opposing reaction is the same as that of forward reaction
- D. reaction ceases to generate heat

Answer: bc

 [Watch Video Solution](#)

4. The equilibrium constant for a reaction is K , and the reaction quotient is Q . For a particular reaction mixture, the ratio $\frac{K}{Q}$ is 0.33. this means that:

- A. the reaction mixture will equilibrate to form more reactant species

- B. the reaction mixture will equilibrate to form more product species
- C. the equilibrium ratio of reactant to product concentration will be 3
- D. the equilibrium ratio of reactant to product concentrations will be 0.33

Answer: b

 [Watch Video Solution](#)

5. Consider the reaction $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ for which $K_c = 278M^{-1}$. 0.001 mole of each of the reagents $SO_2(g)$, $O_2(g)$ and $SO_3(g)$ are mixed in a 1.0 L flask. Determine the reaction quotient of the system and the spontaneous direction of the system:

- A. $Q_c = 1000$, the equilibrium shifts to the right
- B. $Q_c = 1000$, the equilibrium shifts to the left
- C. $Q_c = 0.001$, the equilibrium shifts to the left

D. $Q_c = 0.001$, the equilibrium shifts to the right

Answer: a

 [Watch Video Solution](#)

6. Calculate the wavelength and frequency of a light wave whose period is 2×10^{-10}

 [Watch Video Solution](#)

7. For the reaction $A(g) + 3B(g) \rightleftharpoons 2C(g)$ at $27^\circ C$, 2 moles of A, 4 moles of B and 6 moles of C are present in 2 litre vessel. If K_c for the reaction is 1.2, the reaction will proceed in :

- A. Forward direction
- B. backward direction
- C. neither direction

D. none of these

Answer: a

 [Watch Video Solution](#)

8. For a reversible gaseous reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$ at equilibrium , if some moles of H_2 are replaced by same number of moles of T_2 (T is tritium , isotope of H and assume isotopes do not have different chemical properties) without affecting other parameters , then:

A. the sample of ammonia obtained after something will be radioactive .

B. moles of N_2 after the change will be different as compared to moles of N_2 present before the change

C. the value of K_p or K_c will change


D.

Answer: a




Watch Video Solution


9. For the synthesis of ammonia by the reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$ in the Haber's process, the attainment of equilibrium is correctly predicted by the curve




A.



B.



C.




D.

Answer: a

 Watch Video Solution

10. The figure shows the change in concentration of species A and B as a function of time.

The equilibrium constant K_c for the reaction $A(g) \rightleftharpoons 2B(g)$ is :



A. $K_c > 1$

B. $K < 1$

C. $K = 1$


D. data insufficient

Answer: a

 Watch Video Solution

11. Attainment of the equilibrium $A(g) \rightleftharpoons 2C(g) + B(g)$ gave the following graph . Find the correct option .

(% dissociation = $\frac{\text{Fraction dissociated}}{\text{Initial concentration}} \times 100$)



A. At $t = 5 \text{ sec}$ equilibrium has been reached and $K_c = 40(\text{mol / litre})^2$


- B. At $t=5$ sec equilibrium has been reached and % dissociation of A is 20%
- C. At $t=5$ sec equilibrium has been reached and % dissociation of A is 30%
- D. none of these

Answer: b



Watch Video Solution

12. Using molar concentrations, what is the unit of K_c for the reaction ?




- A. M^{-2}
- B. M^2
- C. M^{-1}
- D. M

Answer: b



Watch Video Solution

13. What is the unit of K_p for the reaction ?



A. atm

B. atm^{-2}

C. atm^2


D. atm^{-1}

Answer: b



Watch Video Solution

14. What is the equilibrium expression for the reaction



A. $K_c = [O_2]^5$

B. $K_c = [P_4O_{10}] / 5[P_4][O_2]$


C. $K_c = [P_4O_{10}] / [P_4][O_2]^5$

D. $K_c = 1 / [O_2]^5$


Answer: d

 Watch Video Solution

15. At $527^\circ C$, the reaction given below has $K_c = 4$



what is the K_p for the reaction ?



A. $16 \times (800R)^2$

B. $\left(\frac{800R}{4}\right)^{-2}$

C. $\left(\frac{1}{4 \times 800R}\right)^2$

D. none of these

Answer: c



Watch Video Solution

16. The equilibrium constant for the reaction

$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ at temperature T is 4×10^{-4} . The value of K_c

for the reaction

$NO(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{1}{2}O_2(g)$ at the same temperature is

A. 4×10^{-4}

B. 50

C. 2.5×10^2


D. 0.02

Answer: b



Watch Video Solution

17. The equilibrium constant K_c for the following reaction at 842°C is 7.90×10^{-3} . What is K_p at same temperature?



A. 8.64×10^{-5}

B. 8.26×10^{-4}

C. 7.90×10^{-2}

D. 7.56×10^{-2}


Answer: d



Watch Video Solution

18. The equilibrium constant K_p for the following reaction at 191°C is 1.24.

What is K_c ?



A. 6.7

B. 0.61

C. 8.30

D. 7.6

Answer: d

 [Watch Video Solution](#)

19. For the equilibrium $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$, what is the temperature at which $\frac{K_p(atm)}{K_c(M)} = 3$?

A. $0.027K$

B. $0.36K$


C. $36.54K$

D. $273K$

Answer: c

 [Watch Video Solution](#)

20. For the reversible reaction



at $500^\circ C$, the value of K_p is 1.44×10^{-5} when the partial pressure is measured in atmosphere. The corresponding value of K_c with concentration in $\text{mol } L^{-1}$ is

A. $1.44 \times 10^{-5} / (0.082 \times 500)^{-2}$

B. $1.44 \times 10^{-5} / (8.314 \times 773)^{-2}$

C. $1.44 \times 10^{-5} / (0.082 \times 773)^2$

D. $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$

Answer: d

 [Watch Video Solution](#)

21. For the reaction $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ the value of $\left(\frac{K_c}{K_P} \right)$

is equal to :

A. \sqrt{RT}

B. RT

C. $\frac{1}{RT}$

D. 1.0

Answer: b



Watch Video Solution

22. The concentration of a pure solid or liquid phase is not include in the expression of equilibrium constant because :

A. density of solid and liquid are independent of their quantities .

B. solids and liquids react slowly.

C. solids and liquids at equilibrium do not interact with gaseous phase.

D. the molecules of solids and liquids cannot migrate to the gaseous phase.

Answer: a



Watch Video Solution

23. A catalyst is a substance which

- A. increase the equilibrium concentration of the product.
- B. change the equilibrium constant of the reaction.
- C. shortens the time to reach equilibrium.
- D. supplies energy to the reaction.

Answer: c



Watch Video Solution

24. What will be the effect of the equilibrium constant on increasing temperature. If the reaction neither absorbs heat nor releases heat?

- A. Equilibrium constant will remain constant.
- B. Equilibrium constant will decrease .
- C. Equilibrium constant will increase.
- D. Can not be predicted.

Answer: a



[Watch Video Solution](#)

25. The equilibrium constant for a reaction

$N_2(g) + O_2(g) = 2NO(g)$ is 4×10^{-4} at $2000K$. In the presence of catalyst, the equilibrium constant is attained 10 times faster. The equilibrium constant in the presence of catalyst, at $2000K$ is

- A. 40×10^{-4}

B. 4×10^{-4}

C. 4×10^{-3}

D. difficult to compute without more data

Answer: a



Watch Video Solution

26. For the reaction $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$

the equilibrium constant K_p changes with

A. total pressure

B. catalyst

C. concentration of H_2 and I_2




D. temperature

Answer: d



Watch Video Solution

27. Consider the reaction :-



Which of the following relation is correct ?

A. $K_3 = \frac{K_1}{K_2}$

B. $K_3 = \frac{K_1^2}{K_2^2}$

C. $K_3 = K_1 K_2$

D. $K_3 = \sqrt{K_1} \cdot K_2$

Answer: d



Watch Video Solution

28. For the reaction $2NO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons N_2O_5(g)$ if the equilibrium constant is K_p , then the equilibrium constant for the reaction

$2N_2O_5(g) \rightleftharpoons 4NO_2(g) + O_2(g)$ would be :

A. K_P^2

B. $\frac{2}{K_P}$

C. $\frac{1}{K_p^2}$


D. $\frac{1}{\sqrt{K_p}}$

Answer: c




Watch Video Solution

29. The equilibrium constant (K_c) for the reaction



is 4×10^{-34} at $25^\circ C$. what is the equilibrium constant for the reaction ?



A. 2×10^{-17}

B. 2.5×10^{33}



C. 5×10^{16}

D. none of these


Answer: d

 [Watch Video Solution](#)

30. At a certain temperature , the following reactions have the equilibrium constants as shown below:



what is the equilibrium constant K_c for the reaction at the same temperature?



A. 2.5×10^{76}

B. 4×10^{23}

C. 4×10^{-77}




D. none of these

Answer: c




Watch Video Solution

31. Given



The equilibrium constant for



will be

A. $K_1K_2K_3$

B. $\frac{K_1K_2}{K_3}$

C. $\frac{K_1K_3^2}{K_2}$

D. $\frac{K_1K_3^2}{K_3}$

Answer: d



[Watch Video Solution](#)

32. In the reaction $X(g) + Y(g) \rightleftharpoons 2Z(g)$, 2 mole of X, 1 mole of Y and 1 mole of Z are placed in a 10 litre vessel and allowed to reach equilibrium. If final concentration of Z is 0.2 M, then K_c for the given reaction is :

A. 1.60

B. $\frac{80}{3}$

C. $\frac{16}{3}$

D. none of these

Answer: c

[Watch Video Solution](#)

33. An equilibrium mixture of the reaction $2H_2S(g) \rightleftharpoons 2H_2(g) + S_2(g)$ had 0.5 mole H_2S , 0.10 mole H_2 and 0.4 mole S_2 in one litre vessel. The value of equilibrium constants (K) in mole $litre^{-1}$ is

A. 0.0004

B. 0.008

C. 0.016

D. 0.160

Answer: c




Watch Video Solution

34.

Given

$[CS_2] = 0.120M$, $[H_2] = 0.10$, $[H_2S] = 0.20$ and $[CH_4] = 7.40 \times 10^{-5}M$

for the following reaction at $900^\circ C$ at eq. Calculate the equilibrium constant (K_c).



A. 0.0120

B. 0.0980

C. 0.280


D. 0.120

Answer: c

 [Watch Video Solution](#)

35. The equilibrium constant for the following reaction is 10.5 at 500 K .A system at equilibrium has

$[CO] = 0.250M$ and $[H_2] = 0.120M$ what is the $[CH_3OH]$?



A. 0.0378

B. 0.435

C. 0.546

D. 0.0499

Answer: a

 [Watch Video Solution](#)

36. When sulphur (in the form of S_8) is heated at temperature T , at equilibrium , the pressure of S_8 falls by 30 % from 1.0atm , because $S_8(g)$ is partially converted into $S_2(g)$.

Find the value of K_P for this reaction.

A. 2.96

B. 6.14

C. 204.8

D. none of these

Answer: a

 [Watch Video Solution](#)

37. 9.2 grams of $N_2O_4(g)$ is taken in a closed one litre vessel and heated till the following equilibrium is reached $N_2O_4(g) \rightleftharpoons 2NO_2(g)$. At equilibrium, 50 % $N_2O_4(g)$ is dissociated. What is the equilibrium constant (in mol litre^{-1}) (Molecular weight of $N_2O_4 = 92$) ?

A. 0.1

B. 0.4

C. 0.2

D. 2

Answer: c



Watch Video Solution

38. Two moles of NH_3 when put into a previously evacuated vessel (one litre) partially dissociate into N_2 and H_2 . If at equilibrium one mole of NH_3 is present, the equilibrium constant is

A. $3/4 \text{ mol}^2 \text{ litre}^{-2}$

B. $27/64 \text{ mol}^2 \text{ litre}^{-2}$


C. $27/32 \text{ mol}^2 \text{ litre}^{-2}$

D. $27/16 \text{ mol}^2 \text{ litre}^{-2}$

Answer: d

 [Watch Video Solution](#)

39. In the presence of excess of anhydrous (in torr) of water taken up is governed by $K_p = 10^{12} atm^{-4}$ for the following reaction at $273K$



What is equilibrium vapour pressure (in torr) of water in a closed vessel that contains $SrCl_2 \cdot 2H_2O(s)$?

A. 0.001 torr

B. 10^3 torr


C. 0.76 torr

D. 1.31 torr

Answer: c

 [Watch Video Solution](#)

40.




If the vapour pressure of water is 38 toor then percentage of relative humidity is :(Assume all data at constant temperture)

- A. 4
- B. 10
- C. 40
- D. none of these

Answer: c

 [Watch Video Solution](#)



The equilibrium pressure at 25°C is 0.660 atm . What is K_p for the reaction?

- A. 0.109

B. 0.218

C. 1.89

D. 2.18

Answer: a



Watch Video Solution

42. For the reaction $2A(g) \rightleftharpoons B(g) + 3C(g)$, at a given temperature , $K_c = 16$. What must be the volume of the flask , if a mixture of 2 mole each of A,B and C exist in equilibrium ?

A. $\frac{1}{4}$

B. $\frac{1}{2}$

C. 1

D. none of these

Answer: b



Watch Video Solution

43. How many neutrons and protons are there in the following nuclei?

A. 1. C atomic mass 13

B.


C.

D.

Answer: a



Watch Video Solution



This reaction is set-up in aqueous medium. We start with 1 mol of I_2 and 0.5 mol of I^\ominus in 1L flask. After equilibrium reached, excess of $AgNO_3$ gave 0.25 mol of yellow precipitate. Equilibrium constant is

A. 1.33

B. 2.66

C. 2.0

D. 3.0

Answer: a



[Watch Video Solution](#)

45. How many neutrons and protons are there in the following nuclei?

A. ${}^2_0\text{O}$ atomic mass 16

B.

C.

D.

Answer: a



[Watch Video Solution](#)

46. In the equilibrium $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$, the partial pressure of SO_2 , O_2 and SO_3 are 0.662, 0.10 and 0.331 atm respectively . What should be the partial pressure of Oxygen so that the equilibrium concentrations of SO_3 are equal ?

A. 0.4atm

B. 1.0atm

C. 0.8atm

D. 0.25atm

Answer: a



[Watch Video Solution](#)

47. How many neutrons and protons are there in the following nuclei?

A. 3. Mg atomic mass 24

B.

C.

D.

Answer: c



Watch Video Solution

48. In the system $A_{(s)} \rightleftharpoons 2B_{(g)} + 3C_{(g)}$, if the concentration of C at equilibrium is increased by a factor of 2, it will cause the equilibrium concentration of B to change to:

A. Two times original value

B. One half of its original value

C. $2\sqrt{2}$ times to the original value

D. $\frac{1}{2\sqrt{2}}$ times the original value

Answer: d



Watch Video Solution

49. $A + B \rightleftharpoons C + D$. If finally the concentrations of A and B are both equal but at equilibrium concentration of D will be twice of that of A then what will be the equilibrium constant of reaction.

A. $\frac{4}{9}$

B. $\frac{9}{4}$

C. $\frac{1}{9}$


D. 4

Answer: d



Watch Video Solution

50. The equilibrium constant K_c for the reaction



1 mole of each of all the four gases is taken in 1dm^3 vessel , the equilibrium concentration of NO would be:

A. $0.4M$

B. $0.6M$

C. $1.4M$

D. $1.6M$

Answer: d



Watch Video Solution

51. On increasing the temperature , the rate of a reaction:

A. always increases

B. always decreases

C. first increases and then decreases

D. may increase or decrease depending upon the nature of the reaction

Answer: a

 [Watch Video Solution](#)


52. A catalyst increases the rate of a reaction by:

- A. increasing the activation energy of a reaction
- B. increasing the value of rate constant (K_f and K_b)
- C. increasing the enthalpy change of the reaction
- D. decreasing the enthalpy change of the reaction

Answer: b

 [Watch Video Solution](#)

53. At a certain temperature, only 50% HI is dissociated at equilibrium in the following reaction:



the equilibrium constant for this reaction is:

A. 0.25

B. 1.0

C. 3.0


D. 0.5

Answer: a



[Watch Video Solution](#)

54. The equilibrium constant K_p for the reaction



is 4.0 at $1660^\circ C$. Initially, 0.80 mol H_2 and 0.80 mol CO_2 are injected into a 5.0 litre flask. What is the equilibrium concentration of $CO_2(g)$?

A. 0.533

B. 0.0534

C. 0.535

D. none of these

Answer: b

 [Watch Video Solution](#)

55. A sample of gas occupies 1.50L at 25 o C. If the temperature is raised to 60 o C, what is the new volume of the gas if pressure remains constant?

 [Watch Video Solution](#)

56. The equilibrium constant for the reaction $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$ is 5. How many moles of CO_2 must

be added to 1 litre container already containing 3 moles each of CO and H_2O to make 2 M equilibrium concentration of CO?

A. 15

B. 19

C. 5

D. 20

Answer: b



[Watch Video Solution](#)

57. A nitrogen-hydrogen mixture initially in the molar ratio of 1 : 3 reached equilibrium to form ammonia when 25% of the N_2 and H_2 had reacted. If the pressure of the system was 21 atm, the partial pressure of ammonia at the equilibrium was :

A. 4.5 atm

B. 3.0 atm

C. 2.0atm

D. 1.5atm

Answer: b

 [Watch Video Solution](#)

58. NH_3 is heated at 15 at, from 25°C to 347°C assuming volume constant. The new pressure becomes 50 atm at equilibrium of the reaction $2\text{NH}_3 \rightleftharpoons \text{N}_2 + 3\text{H}_2$. Calculate % moles of NH_3 actually decomposed.

A. 65 %

B. 61.3 %

C. 62.5 %

D. 64 %

Answer: b

 [Watch Video Solution](#)

59. 0.1 mole of $N_2O_4(g)$ was sealed in a tube under one atmospheric conditions at $25^\circ C$ Calculate the number of moles of $NO_2(g)$ present , if the equilibrium $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ ($K_P = 0.14$) is reached after some time :

A. 1.8×10^2

B. 2.8×10^2

C. 0.034

D. 2.8×10^{-2}

Answer: c

[Watch Video Solution](#)

60. 5 moles of SO_2 and 5 moles of O_2 are allowed to react .At equilibrium , it was found that 60 % of SO_2 is used up .If the pressure of the

equilibrium mixture is one atmosphere, the partial pressure of O_2 is :

A. 0.52atm

B. 0.21tm


C. 0.41atm

D. 0.82atm

Answer: c



[Watch Video Solution](#)



For the reaction initially the mole ratio was 1:3 of $N_2 : H_2$. At equilibrium 50% of each has reacted. If the equilibrium pressure is P, the partial pressure of NH_3 at equilibrium is :

A. $\frac{p}{3}$

B. $\frac{P}{4}$


C. $\frac{P}{6}$

D. $\frac{p}{8}$

Answer: a

 [Watch Video Solution](#)

62. 2.0 mole of PCl_5 were introduced in a vessel of 5.0 L capacity of a particular temperature. At equilibrium, PCl_5 was found to be 35 % dissociated into PCl_3 and Cl_2 . The value of K_c for the reaction



A. 0.075

B. 0.377


C. 1.33

D. 13.3

Answer: d

 [Watch Video Solution](#)

63. At certain temperature compound $AB_2(g)$ dissociates according to the reaction



With degree of dissociation α which is small compared with unity, the expression of K_p in terms of α and initial pressure P is :

A. $p \frac{\alpha^3}{2}$

B. $\frac{P\alpha^2}{3}$

C. $P \frac{\alpha^3}{3}$

D. $\frac{P\alpha^2}{2}$

Answer: a



Watch Video Solution

64. For the reaction

$H_2(g) + CO(g) \rightleftharpoons CO(g) + H_2O(g)$, if the initial concentration of

$[H_2] = [CO_2]$ and x moles /litres of hydrogen is consumed at equilibrium, the correct expression of K_p is :

A. $\frac{x^2}{(1-x)^2}$

B. $\frac{(1-x)^2}{(1-x)^2}$

C. $\frac{x^2}{(2+x)^2}$

D. $\frac{x^2}{(1-x)^2}$

Answer: a



Watch Video Solution

65. If D_T and D_o are the theoretical and observed vapour densities at a definite temperature and α be the degree of dissociation of a substance, then α in the terms of D_o , D_T and n (number of moles of products formed from 1 mole reactant) is calculated by the formula :

A. $\alpha = \frac{D_o - D_T}{(1-n)D_T}$

B. $\alpha = \frac{D_T - D_o}{(n-1)D_T}$

$$C. \alpha = \frac{D_T - D_o}{(n - 1)D_o}$$



$$D. \alpha = \frac{D - D_T}{(n - 1)D_T}$$


Answer: c



Watch Video Solution

66. For the dissociation of PCl_5 into PCl_3 and Cl_2 in gaseous phase reaction, if d is the observed vapour density and D the theoretical vapour density with ' α ' as degree of dissociation, variation of D/d with ' α ' is given by ?





C.

D. none of these

Answer: a

 [Watch Video Solution](#)

67. At $27^{\circ}C$ and 1 atm pressure, N_2O_4 is 20% dissociation into NO_2 . What is the density of equilibrium mixture of N_2O_4 and NO_2 at $27^{\circ}C$ and 1 atm?

A. $3.11g/litre$

B. $2.11g/litre$

C. $4.5g/litre$

D. none of these

Answer: a



Watch Video Solution

68. $COCl_2$ gas dissociates according to the equation, $COCl_2 \rightleftharpoons CO(g) + Cl_2(g)$. When heated to 700 K the density of the gas mixture at 1.16 atm and at equilibrium is $1.16g/litre$ The degree of dissociation of $COCl_2$ at 700K is :

A. 0.28

B. 0.50

C. 0.72

D. 0.42

Answer: c



Watch Video Solution

69. A gas occupies a volume of 250 ml at 745 mm Hg and 25 ° C. What additional pressure is required to reduce the gas volume to 200 ml at the same temperature?

A. 1.57atm

B. 2.57atm

C. 3.57atm


D. 4.57atm

Answer: d



Watch Video Solution

70. Determine the value of equilibrium constant (K_C) for the reaction



if 10 moles of A_2 , 15 moles of B_2 and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M:

A. 4.5

B. 1.5


C. 0.6

D. none of these

Answer: a

 [Watch Video Solution](#)

71. At $87^\circ C$, the following equilibrium is established



If 0.50 mole of hydrogen and 1.0 mole of sulphur are heated to $87^\circ C$

calculate partial pressure of H_2S at eq.

A. 0.966 atm

B. 1.38n atm

C. 0.0327 atm

D. 1atm


Answer: a



Watch Video Solution

72. Pure PCl_5 is introduced into an evacuated chamber and to equilibrium at $247^\circ C$ and 2.0 atm .The equilibrium gases mixture contains 40% chlorine by volume .

Calculate K_p at $247^\circ C$ for the reaction



- A. 0.625 atm
- B. 4atm
- C. 1.6atm
- D. none of these

Answer: c



Watch Video Solution

73. For the reaction

Calculate K_p at 900K where the equilibrium steam-hydrogen mixture was 45% H_2 by volume :

A. 1.49

B. 1.22

C. 0.67

D. none of these

Answer: a



Watch Video Solution

74. For the reaction

$XCO_3 \rightleftharpoons XO(s) + CO_2(g)$, $K_p = 1.642 \text{ atm}$ at $727^\circ C$ If 4 moles of XCO_3

was put into a 50 litre container and heated to $727^\circ C$

What mole percent of the XCO_3 remains unreacted at equilibrium ?

A. 20

B. 25

C. 50

D. none of these

Answer: d



Watch Video Solution

75. $Fe_2O_3(s)$ may be converted to Fe by the reaction

$Fe_2O_3(s) + 3H_2(g) \rightleftharpoons 2Fe(s) + 3H_2O(g)$ for which $K_c = 8$ at temp .
 $720^\circ C$.

What percentage of the H_2 remains unreacted after the reaction has come to equilibrium ?

A. ~22 %

B. ~34 %

C. ~66 %

D. ~78 %

Answer: b

 [Watch Video Solution](#)

76. $AB_3(g)$ dissociates as $AB_3(g) \rightleftharpoons AB(g) + \frac{1}{2}B_2(g)$

When the initial pressure of AB_3 is 800 torr and the pressure developed at equilibrium is 900 torr, what fraction of $AB_3(g)$ is dissociated?

A. 10 %

B. 20 %


C. 25 %

D. 30 %

Answer: c

 [Watch Video Solution](#)

77. At 1000 K , a sample of pure NO_2 gases decomposes as :



The equilibrium constant K_P is 156.25 atm .Analysis shows that the partial pressure of O_2 is 0.25 atm at equilibrium .The parital pressure of NO_2 at equilibrium is :

- A. 0.01
- B. 0.02
- C. 0.04
- D. none of these

Answer: b



[Watch Video Solution](#)

78. 34.05 mL of phosphorus vapour weighs 0.0625 g at $546^\circ C$ and 0.1 bar pressure. What is the molar mass of phosphorus?



[Watch Video Solution](#)

79. At a certain temperature the equilibrium constant K_c is 0.25 for the reaction



If we take 1 mole of each of the four gases in a 10 litre container, what would be equilibrium concentration of A_2 (g)?

A. 0.331 M

B. 0.033M

C. 0.133M

D. 1.33M

Answer: c



Watch Video Solution

80. At 200° PCl_5 dissociates as follows :



It was found that the equilibrium vapours are 62 times as heavy as hydrogen. The degree of dissociation of PCl_5 at $200^\circ C$ is nearly :

A. 10 %

B. 42 %

C. 50 %

D. 68 %

Answer: d



Watch Video Solution

81. For the dissociation reaction $N_2O_4(g) \rightleftharpoons 2NO_2(g)$, the degree of dissociation (α) in terms of K_p and total equilibrium pressure P is:

A. $\alpha = \sqrt{\frac{4P + K_p}{K_p}}$

B. $\alpha = \sqrt{\frac{K_p}{4P + K_p}}$

C. $\alpha = \sqrt{\frac{K_p}{4P}}$

D. none of these


Answer: b

 Watch Video Solution


82. Consider the following equilibrium




Then select the correct graph, which shows the variation in concentrations of N_2O_4 Against concentrations of NO_2 :




A.



B.



C.



D.

Answer: b

 **Watch Video Solution**

83. The vapour pressure of mercury is 0.002 mm Hg at 27°C . K_c for the process $\text{Hg}(l) \rightleftharpoons \text{Hg}(g)$ is :

A. 0.002

B. 8.12×10^{-5}


C. 6.48×10^{-5}

D. 1.068×10^{-7}

Answer: d

 [Watch Video Solution](#)

84. Determine the value of equilibrium constant (K_C) for the reaction



if 10 moles of A_2 , 15 moles of B_2 and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M:

- A. 1.333
- B. 2.66
- C. 20
- D. none of these

Answer: b

 [Watch Video Solution](#)

85. A sample of helium has volume of 520mL at 100°C. calculate the temperature at which the volume becomes 260ml assume that the pressure is constant.

 [Watch Video Solution](#)




86. Given the following reaction at equilibrium $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$. Some inert gas at constant pressure is added to the system. Predict which of the following facts:

- A. more NH_3 is produced
- B. Less $NH_3(g)$ is produced
- C. No effect on the equilibrium
- D. K_p of the reaction is decreased

Answer: b

 [Watch Video Solution](#)

87. In which of the following equilibrium, change in volume of the system does not alter the number of moles:




Answer: a



Watch Video Solution

88. For the reaction



The number of moles of H_2 at equilibrium will increase if :

A. volume is increased

B. volume is decreased

C. argon gas is added at constant volume

D. NH_3 is removed

Answer: a

 [Watch Video Solution](#)

89. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction



When equilibrium is re-established :

A. The amount of $Cl_2(g)$ remains unchanged

B. the amount of $Cl_2(g)$ increases

C. The amount of $SO_2Cl_2(g)$ decreases

D. The amount of $SO_2(g)$ decreases

Answer: b

 [Watch Video Solution](#)

90. Some inert gas is added at constant volume to the following reaction at equilibrium

$N_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$ Itbgt predict the effec of adding the inert gas:

- A. The equilibrium shifts in the forward dircetion
- B. The equilibrium shifts in the backward direction
- C. The equilibrium remins unaffected
- D. The value of K_p is increased

Answer: c

[Watch Video Solution](#)

91. Consider thr reaction where $K_p = 0.497$ at 500K



If the three gases are mixed in a rigid container so that the partial pressure of each gas is initially 1 atm, then which is the correct observation?

- A. More PCl_5 will be produced
- B. More PCl_3 will be produced
- C. Equilibrium will be established when 50% reaction is complete
- D. none of these


Answer: a



Watch Video Solution

92. The preparation of $SO_3(g)$ by reaction $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$ is an exothermic reaction. If the preparation follows the following temperature-pressure relationship for % yield, then for temperatures

T_1 , T_2 and T_3 the correct option is:



A. $T_3 > T_2 > T_1$

B. $T_1 > T_2 > T_3$

C. $T_1 = T_2 = T_3$

D. Nothing could be predicated about temperature though given information




Answer: b

 [Watch Video Solution](#)

93. A balloon is filled with hydrogen at room temperature, it will burst if pressure exceeds 0.2 bar. If at 1 bar pressure the gas occupies 2.27 L volume, upto what volume can the balloon be expanded?




 [Watch Video Solution](#)

94. Le - Chatelier principle is not applicable to :



Answer: b

95. Consider the following reactions .In which case the formation of product is favoured by decrease in pressure?



A. 2, 3




B. 3, 4

C. 2, 4

D. 1, 4

Answer: d

96. In which of the following reactions, the formation of product is favoured by decrease in temperature ?



A. 1, 2

B. 2 only

C. 1,2,3



D. 3,4

Answer: d



Watch Video Solution




97. For which of the following reaction is product formation favoured by low pressure and high temperature?



Answer: b



 [Watch Video Solution](#)

98. For which of the following reaction is product formation favoured by low pressure and high temperature?



Answer: c

99. Consider the following reaction at equilibrium and determine which of the indicated changes will cause the reaction to proceed to right.



A. 2, 3


B. 1, 4

C. 2, 4

D. 2, 3, 4

Answer: a

100. If the pressure in a reaction vessel for the following reaction is increased by decreasing the volume ,what will happen to the concentrations of Coand CO_2 ?



- A. both the $[CO]$ and $[CO_2]$ will decrease
- B. neither the $[Co]$ nor the $[CO_2]$ will change
- C. the $[CO]$ will decrease and the $[CO_2]$ will increase
- D. both the $[CO]$ and $[CO_2]$ will increase

Answer: d



Watch Video Solution

101. A System at equilibrium is described by the equation of fixed temperature T.



What effect will be the effect on equilibrium, if total pressure is increased by reducing volume?

- A. Concentration of $SO_2Cl_2(g)$ increases
- B. Concentrations of $SO_2(g)$ increases
- C. Concentration of $Cl_2(g)$ increases
- D. Concentration of all gases increases

Answer: d

 [Watch Video Solution](#)

102. For the reaction $2NO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons N_2O_5(g)$ if the equilibrium constant is K_p , then the equilibrium constant for the reaction $2N_2O_5(g) \rightleftharpoons 4NO_2(g) + O_2(g)$ would be :

- A. equilibration of this gas mixture will be slower at high temperature
- B. A mole of N_2O_4 will occupy twice the volume of a mole of NO_2 at the same?

C. the equilibrium will move to the right if an equilibrium mixture is cooled

D. the position of equilibrium will move to the left with increasing gas pressure

Answer: c

 [Watch Video Solution](#)

103. give the symbol of the atom whose ground state corresponds to each of the following configuration : (1) $1s^2 2s^2 2p_x^2 2p_y^2 2p_z^1$

 [Watch Video Solution](#)

104. For an equilibrium $H_2O(s) \rightleftharpoons H_2O(l)$, which of the following statements is true ?

A. The pressure changes do not affect the equilibrium

- B. More of ice melts if pressure on the system is increased
- C. More of liquid freezes if pressure on the system is increased
- D. The pressure changes may increase may increase or decrease the degree of advancement of the process

Answer: b

 [Watch Video Solution](#)

105. A pressure cooker reduces cooking time because :

- A. the higher pressure inside the cooker crushes the food material
- B. cooking involves chemical change helped by a rise in temperature
- C. heat is more evenly distributed in the cooking space
- D. boiling point of water involved in cooking is increased

Answer: d

 [Watch Video Solution](#)

106. The vapour pressure of a liquid in a closed container depends upon

- A. 1 only
- B. 2 only
- C. 1 and 3 only
- D. 1,2,and3


Answer: a

 [Watch Video Solution](#)


107. Draw lewis structure of Al^{3+}

 [Watch Video Solution](#)


108. An exothermic reaction is represented by the graph :




A.



B.



C.




Answer: c




Watch Video Solution


109. An endothermic reaction is represented by the graph :




A.



B.



C.



D.

Answer: b



Watch Video Solution

110. The enthalpy change for the reaction $\text{N}_2(\text{g}) + 3 \text{H}_2(\text{g}) \rightarrow 2 \text{NH}_3(\text{g})$ is -91.8 kJ at 298 K . Calculate the enthalpy of formation of ammonia?

 [Watch Video Solution](#)

111. The correct relationship between free energy change in a reaction and the corresponding equilibrium constant K_c is:

A. $\Delta G^\circ = RT \ln K$

B. $\Delta G^\circ = -RT \ln K$


C. $\Delta G = RT \ln K$

D. $\Delta G = -RT \ln K$

Answer: b

 [Watch Video Solution](#)

112. K_p has the value of 10^{-6} atm^3 and 10^{-4} atm^3 at 298 K and 323 K respectively for the reaction



$\Delta_r H^\circ$ for the reaction is :

A. 7.7 KJ/mol

B. -147.41 KJ/mol

C. 147.41 KJ/mol

D. none of these

Answer: c



Watch Video Solution

113. Van's Hoff's equation shows the effect of temperature on equilibrium constants K_c and K_p . the K_P varies with tempertaure according to the realation:

A. $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta H^\circ}{2.303R} \left(\frac{T_1 - T_2}{T_1 T_2} \right)$

B. $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta H^\circ}{2.303R} \left(\frac{T_2 - T_1}{T_1 T_2} \right)$

C.

D.

Answer: b



Watch Video Solution

114. For a reaction, the value of K_p increases with increase in temperature.

The ΔH for the reaction would be :

A. positive

B. negative





C. zero

D. cannot be predicted

Answer: A

 [Watch Video Solution](#)


115. The most stable oxides of nitrogen will be :



Answer: A

 [Watch Video Solution](#)

116. When 1 mole of pure ethyl alcohol (C_2H_5OH) is mixed with 1 mole of acetic acid at $25^\circ C$. the equilibrium mixture contains $2/3$ mole each of ester and water



The ΔG° for the reaction at $298K$ is :

A. 3435 J

B. 4 J

C. -3435 J

D. zero

Answer: C



Watch Video Solution

117. The value of ΔG° for a reaction in aqueous phase having $K_c = 1$, would be :

A. $-RT$

B. -1

C. 0

D. $+RT$

Answer: C

 [Watch Video Solution](#)


118. A plot of Gibbs energy of a reaction mixture against the extent of the reaction is :

- A. minimum at equilibrium
- B. zero at equilibrium
- C. maximum at equilibrium
- D. None of these

Answer: A

 [Watch Video Solution](#)

119. For the reaction at $300K$



$$\Delta_r H^\circ = -30 \text{ kJ/mol}, \Delta_r S^\circ = -0.1 \text{ K}^{-1} \cdot \text{mol}^{-1}$$

What is the value of equilibrium constant ?

A. 0

B. 1


C. 10

D. None of these

Answer: B

 [Watch Video Solution](#)

120. Solid $Ca(HCO_3)_2$ decomposes as



If the total pressure is 0.2 bar at 420K, what is the standard free energy change for the given reaction ($\Delta_r G^\circ$) ?

A. 840kJ/mol

B. 3.86kJ/mol

C. 6.98kJ/mol

D. 16.083kJ/mol

Answer: D

 [Watch Video Solution](#)

121. The standard free energy change of a reaction is $\Delta G^\circ = -115 \text{ kJ/mol}^{-1}$ at 298 K . Calculate the value of $\log_{10} K_p$ ($R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$)

A. 20.16

B. 2.303

C. 2.016

D. 13.83

Answer: A

 [Watch Video Solution](#)

122. On a ship sailing in Pacific Ocean where temp. is 23.4°C A balloon is filled with 2L air, what will be the volume of balloon where the ship reaches Indian Ocean where temp is 26.1°C

 [Watch Video Solution](#)

123. What will be the pH of a soft drink if hydronium ion concentration is sample is $3.8 \times 10^{-3}\text{M}$?

 [Watch Video Solution](#)

124. In three moles of ethane. calculate the number of moles of carbon atoms.

 [Watch Video Solution](#)

125. In three moles of ethane. calculate the number of moles of hydrogen atoms.

 [Watch Video Solution](#)

126. For the reaction $C_2H_6(g) \rightleftharpoons C_2H_4(g) + H_2(g)$

K_p is 5×10^{-2} atm. Calculate the mole per cent of $C_2H_6(g)$ at equilibrium if pure C_2H_6 at 1 atm is passed over a suitable catalyst at $900K$:

A. 20

B. 33.33

C. 66.66

D. None of these

Answer: C

 [Watch Video Solution](#)

127. $2NOBr(g) \rightleftharpoons 2NO(g) + Br_2(g)$. If nitrosyl bromide (NOBr) 40 % dissociated at certain temp. and a total pressure of 0.30 atm K_p for the reaction $2NO(g) + Br_2(g) \rightleftharpoons 2NOBr(g)$ is

A. 45

B. 25

C. 0.022

D. 0.25

Answer: A



[Watch Video Solution](#)

128. Consider the partial decomposition of A as

$2A(g) \rightleftharpoons 2B(g) + C(g)$ At equilibrium 700mL gaseous mixture contains 100mL of gas C at 10 atm and 300K what is the value of K_p for the reaction ?

A. $\frac{40}{7}$

B. $\frac{1}{28}$

C. $\frac{10}{28}$

D. $\frac{28}{10}$


Answer: C

 [Watch Video Solution](#)

129. In the three moles of ethane. Calculate the number of molecules of ethane.

 [Watch Video Solution](#)

130. 0.020 g of selenium vapour at equilibrium occupying a volume of 2.463 mL at 1 atm and $27^{\circ}C$. The selenium is in a state of equilibrium according to reaction



What is the degree of association of selenium ?

(At.mass of se = 79)

A. 0.205

B. 0.315

C. 0.14

D. None of these

Answer: B



[Watch Video Solution](#)

131. Determine the degree of association (polymerization) for the reaction in aqueous solution, if observed (mean) molar mass of HCHO and $C_6H_{12}O_6$ is 150:

A. 0.50

B. 0.833

C. 0.90

D. 0.96

Answer: D



Watch Video Solution

132. A reaction system in equilibrium according to reaction $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ in one litre vessel at a given temperature was found to be 0.12 mole each of SO_2 and SO_3 and 5 mole of O_2 . In another vessel of one litre contains 32 g of SO_2 at the same temperature. What mass of O_2 must be added to this vessel in order that at equilibrium 20% of SO_2 is oxidized to SO_3 ?

A. 0.4125

B. 11.6 g

C. 1.6 g

D. None of these

Answer: B



Watch Video Solution

133. The equilibrium constant K_p for the following reaction is 4.5

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$ What would be the average molar mass (in g/mol) of an equilibrium mixture of N_2O_4 and NO_2 formed by the dissociation of pure N_2O_4 at a total pressure of 2 atm ?

A. 69

B. 57.5

C. 80.5

D. 85.5

Answer: B



Watch Video Solution

134. Express the following in scientific notation : 8008



Watch Video Solution

135. Express the following in scientific notation : 500.0



Watch Video Solution

136. For the reaction $2A(g) + B(g) \rightleftharpoons C(g) + D(g)$, $K_c = 10^{12}$.if initially 4,2,6,2 moles of A,B,C,D respectively are taken in a 1 litre vessel, then the equilibrium concentration of A is :

A. 4×10^{-4}

B. 2×10^{-4}

C. 10^{-4}


D. 8×10^{-4}

Answer: A



Watch Video Solution

137. The equilibrium constant for the following reaction in aqueous solution is 0.90.



How many mole of glycerin should be added per litre of $0.10M H_3BO_3$ so that 80 % of the H_3BO_3 is converted to the boric-acid glycerin complex ?

A. 4.44

B. 4.52

C. 3.6

D. 0.08

Answer: B



[Watch Video Solution](#)

138. Rate of diffusion of ozonized oxygen is $0.4\sqrt{5}$ times that of pure oxygen what is the per cent degree of association of oxygen assuming

pure O_2 in the sample initially ?

A. 20

B. 40

C. 60

D. None of these

Answer: C



[Watch Video Solution](#)

139. Express the following in scientific notation : 6.0012



[Watch Video Solution](#)

140. At $800^\circ C$, the following equilibrium is established as



The composition of equilibrium may be determined by measuring the

rate of effusion of the mixture through a pin hole. It is found that at 800°C and 1 atm mixture effuses 1.6 times as fast as SO_2 effuses under the similar conditions. (At. mass of F = 19) what is the value of K_p (in atm) ?

A. 0.315

B. 0.685

C. 0.46


D. 1.49

Answer: D



[Watch Video Solution](#)

141. The equilibrium constant K_p for the reaction



is 4.0 at 1660°C . Initially 0.80 mol H_2 and 0.80 mol CO_2 are injected into a 5.0 litre flask. What is the equilibrium concentration of $\text{CO}_2(\text{g})$?

A. ≈ 12.3

B. ≈ 11.3

C. ≈ 11.45

D. None

Answer: B

 [Watch Video Solution](#)

142. When N_2O_5 is heated at certain temperature, it dissociates as $N_2O_5(g) \rightleftharpoons N_2O_3(g) + O_2(g)$, $K_c = 2.5$ At the same time N_2O_3 also decomposes as :

$N_2O_3(g) \rightleftharpoons N_2O(g) + O_2(g)$. "If initially" 4.0 moles of N_2O_5 "are taken in" 1.0 litre flask and allowed to dissociate. Concentration of O_2 at equilibrium is 2.5 M. "Equilibrium concentration of " N_2O_5 is :

A. 1.0 M

B. 1.5M

C. $2.166M$



D. $1.846 M$

Answer: D



Watch Video Solution

143. Two solid compounds X and Y dissociates at a certain temperature as follows



The total pressure of gases over a mixture of X and Y is :

A. 4.5 atm

B. 0.45 atm

C. 0.6 atm

D. None of these

Answer: B



Watch Video Solution

144. Calculate the mass % of different elements present in sodium sulphate (Na_2SO_4).



Watch Video Solution

145. For a gaseous reaction



equilibrium constants K_c , K_p and K_x are represented by the following reaction

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, K_p = \frac{P^c \cdot P_D^d}{P_A^a} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where $[A]$ represents molar concentration of A , p_A represents partial pressure of A and P represents total pressure, x_A represents mole fraction of A

For the reaction $\text{SO}_2\text{Cl}_2(g) \rightleftharpoons \text{SO}_2(g) + \text{Cl}_2(g)$, $K_p > K_x$ is obtained at :

A. 0.5 atm

B. 0.8 atm

C. 1 atm

D. 2atm

Answer:

 [Watch Video Solution](#)

146. For a gaseous reaction




equilibrium constants K_c , K_p and K_x are represented by the following relation

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, K_p = \frac{P_C^c \cdot P_D^d}{P_A^a} \text{ and } K_x = \frac{x_C^c \cdot x_D^d}{x_A^a \cdot x_B^b}$$

where $[A]$ represents molar concentration of A , p_A represents partial pressure of A and P represents total pressure, x_A represents mole fraction of A . For the following equilibrium relation between K_c and K_c (in

terms of mole fraction) is



A. $K_c = K_x(RT)^{-1}$

B. $K_c = K_x(RT)$

C. $K_c = K_x\left(\frac{RT}{P}\right)$

D. $K_c = K_x\left(\frac{P}{RT}\right)$

Answer:

 [Watch Video Solution](#)



147. Calculate the amount of carbon dioxide that would be produced when 1 mole of carbon is burnt in air.

 [Watch Video Solution](#)

148. Calculate the amount of carbon dioxide that would be produced when 1 mole of carbon is burnt in 16 g dioxygen.

 [Watch Video Solution](#)

149. N_2O_3 is an unstable oxide of nitrogen and it decomposes into NO (g) and $NO_2(g)$ where $NO_2(g)$ is further dimerise into N_2O_4 as



A flask is initially filled with pure $N_2O_3(g)$ having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 bar.

The equilibrium partial pressure of $N_2O_3(g)$ is :



- A. 0.5bar
- B. 1.0 bar
- C. 1.5 bar
- D. 0.1 bar

Answer:



Watch Video Solution

150. N_2O_3 is an unstable oxide of nitrogen and it decomposes into NO (g) and $NO_2(g)$ where $NO_2(g)$ is further dimerise into N_2O_4 as



A flask is initially filled with pure $N_2O_3(g)$ having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 ber.

The equilibrium partial presure of $NO_2(g)$ is:

A. 0.066bar

B. 0.133 bar

C. 0.423 bar



D. 0.83 bar

Answer:



Watch Video Solution

151. N_2O_3 is an unstable oxide of nitrogen and it decomposes into NO (g) and $NO_2(g)$ where $NO_2(g)$ is further dimerise into N_2O_4 as



A flask is initially filled with pure $N_2O_3(g)$ having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of NO (g) was found to be 1.5 ber.

The value of K_{P_2} is

A. 0.16bar^{-1}

B. 0.32bar^{-1}

C. 0.48bar^{-1}

D. 0.64bar^{-1}

Answer:



[Watch Video Solution](#)

152. AX'(g) solute when dissolved in water heat is evolved. Then solubility of 'X' will increase :

- A. high temperature, low pressure
- B. low temperature, high pressure
- C. high temperature, high pressure
- D. low temperature, high pressure

Answer:



[Watch Video Solution](#)

153. If a system at equilibrium is subjected to a change of any one of the factors such as concentration , pressure or temperature, the system

adjusts itself in such a way so as to minimise the effect of that change.



Above equilibrium is favoured at :

- A. high pressure, low temperature
- B. high pressure, high temperature
- C. low pressure, high temperature
- D. low pressure, low temperature


Answer:



[Watch Video Solution](#)

154. If a system at equilibrium is subjected to a change of any one of the factors such as concentration , pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

For the reaction



If pressure is increased by reducing the volume of the container then :

- A. total pressure at equilibrium will remain same
- B. concentration of all the component at equilibrium will change
- C. concentration of all the component at equilibrium will ramin same
- D. equilibrium will shift in the beckward direction

Answer:

 [Watch Video Solution](#)

155. A catalyst :

- A. increase the average kinetic energy of reactiong molecules
- B. decreases the activation energy
- C. can alters the reaction mechanism
- D. Can change pre-exponential factor

Answer:

 [Watch Video Solution](#)

156. Which of the following is correct about the chemical equilibrium ?

A. $(\Delta G)_{T,p} = 0$

B. Equilibrium constant is independent of initial concentration of reactants

C. Catalyst has no effect on equilibrium state

D. Reaction stops at equilibrium

Answer:

 [Watch Video Solution](#)

157. For the reaction



If α is negligible w.r.t 1 then degree of dissociation (α) of AB_2 is proportional to :

A. $\frac{1}{P}$

B. $\frac{1}{V}$

C. $\frac{1}{\sqrt{P}}$





D. \sqrt{V}

Answer:



Watch Video Solution


158. Consider the reaction given below. In which cases will the reaction proceed toward right by increasing the pressure ?



Answer:

 [Watch Video Solution](#)

159. Ammonia is a weak base that reacts with water according to the equation



Select the correct option (s) that can increase the moles of ammonium ion in water:

- A. Addition of HCl
- B. Addition of NaOH
- C. Addition of NH_4Cl
- D. Addition of H_2O

Answer:

 [Watch Video Solution](#)

160. Consider the reaction $2CO(g) + O_2(g) \rightleftharpoons 2CO_2(g) + \text{Heat}$

Under what conditions shift is undetminable ?

- A. Addition of O_2 and decrease in volume
- B. Addition of CO and removal of CO_2 at constant volume
- C. Increase in temperature and decrease in volume
- D. Addition of CO and increase in temperature at constant volume

Answer:



[Watch Video Solution](#)

161. What will be the effect of addition of catalyst at constant temperature ?

- A. The equilibrium constant will remain constant
- B. ΔH of the reaction will remain constant
- C. K_f and K_b wil increase upto same extent

D. equilibrium composition will change

Answer:

 [Watch Video Solution](#)

162. For the reaction $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$, the forward reaction at constant temperature favorrd by :

- A. introducing an inert gas at constant volume
- B. introducing chlorine gas at constant volume
- C. introducing an inert gas at constant pressure
- D. increasing the volume of the container

Answer:

 [Watch Video Solution](#)

163. For the reaction :
 $Cl_2(g) + 3F_2(g) \rightleftharpoons 2ClF_3(g)$, $\Delta H = -329kJ$, dissociation of $ClF_3(g)$ will be favoured by :

- A. increasing the temperature
- B. increasing the volume of the container
- C. adding of F_2 gas
- D. adding of inert gas at constant pressure

Answer:

 [Watch Video Solution](#)

164. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction



When equilibrium is re-established :

A. formation of more $H_2O(l)$

B. formation of more $H_2O(g)$


C. increase in b.p of $H_2O(l)$

D. decrease in b.p. of $H_2O(l)$

Answer:

 [Watch Video Solution](#)

165. Heating a II group metal carbonate leads to decomposition as :



Equilibrium will shift left

A. by addition of $BaO(s)$

B. by addition of $CO_2(g)$

C. by decreasing the temperature

D. by decreasing the volume of the vessel

Answer:

 [Watch Video Solution](#)

166. $N_2(g)$ and $H_2(g)$ are allowed to react in a closed vessel at given temp. and pressure for the formation of $NH_3(g)$, $[N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g) + 22.4kcal]$ If he (g) is added at equilibrium at constant pressure than which is/are correct ?

- A. Concentration of $N_2(g)$, $H_2(g)$ and $NH_3(g)$ decrease.
- B. Moles of $NH_3(g)$ decreases.
- C. The extent of cooling depends on amount of he (g) added.
- D. Concentration of N_2 and H_2 increases and concentration of NH_3 decreases.

Answer:

 [Watch Video Solution](#)

167. Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with, some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II

Column-I	Column-II
(A) $\text{CaCO}_3(s) \rightleftharpoons \text{CaO}(s) + \text{CO}_2(g)$	(P) $K_p > K_c$ above room temperature
(B) $\text{CO}(g) + \text{Cl}_2(g) \rightleftharpoons \text{COCl}_2(g)$	(Q) $K_p = K_c$ above room temperature
(C) $\text{H}_2(g) + \text{I}_2(g) \rightleftharpoons 2\text{HI}(g)$	(R) $K_p < K_c$ above room temperature
(D) $\text{HCl}(g) \rightleftharpoons \text{H}^+(aq) + \text{Cl}^-(aq)$	(S) K_p and K_c not defined

 [Watch Video Solution](#)

168. Calculate the mass % of calcium, phosphorous and oxygen in calcium phosphate.

 [Watch Video Solution](#)

169. Calculate the energy of a photon of light having frequency of $6 \times 10^{15} \text{ s}^{-1}$

 [Watch Video Solution](#)

 [Watch Video Solution](#)

170. Explain Le Chatelier's principle.

 [Watch Video Solution](#)

171. Calculate the mass % of carbon in ethanol.

 [Watch Video Solution](#)

172. Which of the following is isoelectronic : Na^+ , Ar , Mg^{2+} , S^{2-}

 [Watch Video Solution](#)

173. Assertion (A): The endothermic reactions are favoured at lower temperature and the exothermic reactions are favoured at higher temperature.

Reason (R) : when a system in equilibrium is disturbed by changing the

temperature, it will tend to adjust itself so as to overcome the effect of the change.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

 [Watch Video Solution](#)

174. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The melting point of ice decreases with increase of pressure.

STATEMENT-2: Ice contracts on melting .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

175. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according

to the instruction given below:

STATEMENT-1: The equilibrium of $A(g) \rightleftharpoons B(g) + c(g)$ is not affected by changing the volume.

STATEMENT-2: K_c for the reaction does not depend on volume of the container.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



Watch Video Solution

176. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1:For a chemical reaction at initial stage rate of forward reaction (r_f) is greater than rate of reversed reaction(r_b)

STATEMENT-2: When $r_f = r_b$,chemical reaction is at equilibrium.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

177. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: If $Q_p < K_p$ reaction moves in direction of reactants.

STATEMENT-2: Reaction quotient is defined in the same way as equilibrium constant at any stage of the reaction.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

178. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The gas phase reaction $PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$ shifts to the right on increasing pressure.

STATEMENT-2: When pressure increase, equilibrium shifts towards more number of moles.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



Watch Video Solution

179. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For a reaction at equilibrium, the Gibb's free energy of reaction is minimum at constant temp. and pressure.

STATEMENT-2: The Gibb's free energy of both reactants and products increases and become equal at equilibrium.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



[Watch Video Solution](#)

180. STATEMENT-1: The physical equilibrium is not static but dynamic in nature.

STATEMENT-2: The physical equilibrium is a state in which two opposing processes are proceeding at the same rate.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

181. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.

STATEMENT-2: Equilibrium constant depends upon the way in which the reaction is written.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

182. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The equilibrium of $A(g) \rightleftharpoons B(g) + C(g)$ is not affected by changing the volume.

STATEMENT-2: K_c for the reaction does not depend on volume of the container.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

183. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the reaction $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ if the volume of vessel is reduced to half of its original volume, equilibrium concentration of all gases will be doubled.

STATEMENT-2: According to Le- Chatelier's principle, reaction shifts in a direction that tends to minimized the effect of the stess.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

184. STATEMENT-1: The equilibrium constant of the exothermic reaction at high temperature decreases.

STATEMENT-2: Since $\ln \frac{K_2}{K_1} = \frac{\Delta H^\circ}{R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$ and for exothermic reaction ,

$$\Delta H^\circ = -ve \text{ and thereby, } \frac{K_2}{K_1} < 1$$

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE


Answer: A



185. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: For the reaction at certainn temperature



there will be no effect by addition of inert gas at constant volume.

STATEMENT-2: Molar concentration of all gases remains constant.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

186. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The melting point of ice decreases with increase of pressure.

STATEMENT-2: Ice contracts on melting .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



Watch Video Solution

187. Each question contains STATEMENT-1 (Assertion) and STATEMENT-2(Reason).

Examine the statements carefully and mark the correct answer according to the instruction given below:

STATEMENT-1: The catalyst does not alter the equilibrium constant.

STATEMENT-2: Because for the catalysed reaction and uncatalysed reaction ΔH remains same and equilibrium constant depends of ΔH .

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE


Answer: A

 [Watch Video Solution](#)

188. In the reaction $C(s) + CO_2(g) \rightleftharpoons 2CO(g)$, the equilibrium pressure is 12 atm. If 50 % of CO_2 reacts, calculate K_p .

 [Watch Video Solution](#)

189. Calculate partial pressure of B at equilibrium in the following equilibrium



 [Watch Video Solution](#)

190. In a gaseous reaction $A + 2B \rightleftharpoons 2C + D$ the initial concentration of B was 1.5 times that of A. At equilibrium the concentration of A and D were equal. Calculate the equilibrium constant K_C .



Watch Video Solution

191. For the reaction $A(g) \rightleftharpoons B(g)$, $K_C = 10$

$B(g) \rightleftharpoons C(g)$, $K_C = 2$

$C(g) \rightleftharpoons D(g)$, $K_C = 0.01$

Calculate K_C for the reaction $D(g) \rightleftharpoons A(g)$.



Watch Video Solution

192. 5 litre vessel contains 2 moles of each of gases A and B at equilibrium. If 1 mole each of A and B are removed. Calculate K_C for the reaction $A(g) \rightleftharpoons B(g)$



Watch Video Solution

193. Calculate K_P for the reaction $A(g) \rightleftharpoons B(s) + 2C(g)$, $K_C = 0.2$ at 305 K.



Watch Video Solution

[Watch Video Solution](#)

194. A mixture of 3 moles of SO_2 , 4 moles of NO_2 , 1 mole of SO_3 and 4 moles of NO is placed in a 2.0L vessel.



At equilibrium, the vessel is found to contain 1 mole of SO_2 . Calculate the value of K_C .

[Watch Video Solution](#)


195. Write electronic configuration of elements with atomic number 17 , 29

[Watch Video Solution](#)

196. If chemical equilibrium is attained t standard states then what is the value of ΔG° ?

[Watch Video Solution](#)

197. Calculate the equilibrium concentration ratio of C to A if equimolar ratio of A and B were allowed to come to equilibrium at 300K.




 [Watch Video Solution](#)

198. A definite amount of solid NH_4HS is placed in a flask already containing ammonia gas at a certain temperature and 0.1 atm pressure. NH_4HS decomposes to give NH_3 and H_2S and at equilibrium total pressure in flask is 1.1 atm. If the equilibrium constant K_P for the reaction $NH_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$ is represented as $z \times 10^{-1}$ then find the value of z.

 [Watch Video Solution](#)

199. The gaseous reaction : $A(g) + nB(g) \rightleftharpoons mC(g)$ is represented by following curves



What is the value of $n+m$?

[Watch Video Solution](#)

Level 1 Q 93 To Q 122

1. A sample of nitrogen gas occupies a volume of 1.0L at a pressure of 0.5 bar and at 40°C. Calculate the pressure the gas if compressed to 0.225ml at -6°C

[Watch Video Solution](#)

1. Calculate $\Delta_r G$ for the reaction at $27^\circ C$



Given : $P_{H_2} = 0.5 \text{ bar}$, $[Ag^+] = 10^{-5} M$,

$[H^+] = 10^{-3} M$, $\Delta_r G^\circ [Ag^+(aq)] = 77.1 kJ/mol$

A. $-154.2 kJ/mol$

B. $-178.9 kJ/mol$

C. $-129.5 kJ/mol$

D. None of these

Answer: C



[Watch Video Solution](#)

1. Calculate the mass of sodium acetate (CH_3COONa) required to make 500 mL of 0.375 molar aqueous solution. Molar mass of sodium acetate is $82.0245 \text{ g mol}^{-1}$

 [Watch Video Solution](#)

2. Define Entropy

 [Watch Video Solution](#)

3. Calculate energy of one mole of photons of radiation whose frequency is $5 \times 10^{14} \text{ hz}$

 [Watch Video Solution](#)