



CHEMISTRY

BOOKS - NARENDRA AWASTHI

DILUTE SOLUTION

Exercise

1. The vapour pressure of a give liquid will decrease if :

- A. surface area of liquid decreased
- B. the volume of liquid in the container is decreased
- C. the volume of the vapour phase is increased
- D. the temperature is dexreased

Answer: d





Watch Video Solution

2. The normal boiling point of water is 373 K. Vapour of water at temperature T is 19 mm Hg. If enthalpy of vapourisation is 40.67 kJ/mol, then temperature T would be

(use : $\log 2 = 0.3$, $R : 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$):

- A. 250 K
- B. 291.4 K
- C. 230 K
- D. 290 K

Answer: b



Watch Video Solution

3. A sample of liquid H_2O at 18.0 g is injected into an evacuated 7.6 L flask maintained at 27.0°C . If vapour pressure of H_2O at 27.0°C is 24.63

mm Hg, what weight of the water will be vaporized when the system comes to equilibrium? Assume water vapour behaves as an ideal gas. The volume occupied by the liquid water is negligible compared to the volume of the container:

- A. 0.01
- B. 0.1
- C. 0.18
- D. 0.2

Answer: a

 [Watch Video Solution](#)

4. Raoult's law is obeyed by each constituent of binary liquid solution when :

- A. the forces of attraction between like molecules are greater than those between unlike molecules

- B. the forces of attraction between like molecules are smaller than those between unlike molecules
- C. the forces of attraction between like molecules are identical with those between unlike molecules
- D. the volume occupied by unlike molecules are different

Answer: c



Watch Video Solution

5. For a binary ideal liquid solution, the total vapour of the solution is given as:

A. $P_{\text{total}} = P_A^\circ + (P_A^\circ - P_B^\circ)X_B$

B. $P_{\text{total}} = P_B^\circ + (P_A^\circ - P_B^\circ)X_A$

C. $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ)X_A$

D. $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ)X_B$

Answer: b



Watch Video Solution

6. Two liquids A and B have P_A° and P_B° in the ratio of 1 : 3 and the ratio of number of moles of A and B in liquid phase are 1 : 3 then mole fraction of 'A' in vapour phase in equilibrium with the solution is equal to :

A. $Y_a < Y_b$

B. $X_A > X_B$

C. $\frac{Y_A}{Y_B} > \frac{X_A}{X_B}$

D. $\frac{Y_A}{Y_B} < \frac{X_A}{X_B}$

Answer: c



Watch Video Solution

7. X_A and X_B are the mole fraction of A and B respectively in liquid phase y_A and y_B are the mole fraction of A and B respective in vapour phase. Find out the slope of straight line if a graph is plotted $\frac{1}{y_A}$ along Y-axis against $\frac{1}{x_A}$ along X-axis gives straight line [p_A° and p_B° are vapour pressure of pure components A and B].

A. $\frac{P_B^\circ}{P_A^\circ}$

B. $\frac{P_A^\circ}{P_B^\circ}$

C. $P_B^\circ - P_A^\circ$

D. $P_A^\circ - P_B^\circ$

Answer: a



Watch Video Solution

8. For a dilute solution, Raoult's law states that :

- A. the lowering of vapour pressure is equal to the mole fraction of solute
- B. the relative lowering of vapour pressure is equal to the mole fraction of solute
- C. the relative lowering of vapour pressure is proportional to the amount of solute in solution
- D. the vapour pressure of the solution is equal to the mole fraction of solvent

Answer: b

 [Watch Video Solution](#)

9. A solution containing 30 g of non-volatile solute exactly in 90 g of water has a vapour pressure of 2.8 kPa at 298 K. Further, 18 g of water is then added to the solution and the new vapour pressure becomes 2.9 kPa at 298 K. Calculate: vapour pressure of water at 298 K.

A. $X < Y < Z$

B. $X > Y > Z$

C. $Z > X = Y$

D. $X > Y = Z$

Answer: d







Watch Video Solution

10. The boiling point of

C_6H_6 , CH_3OH , $C_6H_5NH_2$ and $C_6H_5NO_2$ are $80^\circ C$, $65^\circ C$, $184^\circ C$ and $212^\circ C$

respectively. Which will show highest vapour pressure at room temperature :



Answer: B

 [Watch Video Solution](#)

11. 6.0 g of urea (molecules mass = 60) was dissolved in 9.9 moles of water.

If the vapour pressure of pure water is P° , the vapour pressure of solution is :

A. $0.10 P^\circ$

B. $1.10 P^\circ$

C. $0.90 P^\circ$

D. $0.99 P^\circ$

Answer: d

 [Watch Video Solution](#)

12. An ideal solution was found to have a vapour pressure of 80 torr when the mole fraction of a non-volatile solute was 0.2. What would be the vapour pressure of the pure solvent at the same temperature?

- A. 64 torr
- B. 80 torr
- C. 100 torr
- D. 400 torr

Answer: c



[Watch Video Solution](#)

13. If the vapor pressure of a dilute aqueous solution of glucose is 750mm of Hg at 373K, then molality of solute is

- A. 0.26
- B. 0.73

C. 0.74

D. 0.039

Answer: c

 [Watch Video Solution](#)

14. Name the solute and solvent in an aqueous solution of sugar.

A. 10 torr

B. 18 torr

C. 13.45 torr

D. 24 torr

Answer: c

 [Watch Video Solution](#)

15. Calculate the weight of non – volatile solute having molecular weight 40, which should be dissolved in 57gm octane to reduce its vapour pressure to 80 % :

A. 47.2 g

B. 5 g

C. 106.2 g

D. None of these

Answer: b



[Watch Video Solution](#)

16. Equal mass of a solute are dissolved in equal mass of two solvents A and B and formed very dilute solution. The relative lowering of vapour pressure for the solution B has twice the relative lowering of vapour pressure for the solution A. If m_A and M_B are the molecules mass of solvents A and B respectively, then :

A. $M_A = M_B$

B. $M_B = 2 \times M_A$

C. $M_A = 4M_B$

D. $M_A = 2M_B$

Answer: b

 [Watch Video Solution](#)

17. An ideal solution has two components A is more volatile than B, i.e.

$P_A^\circ > P_B^\circ$ and also $p_A^\circ > P_{\text{total}}$. If X_A and Y_A are mole fraction of

components A in liquid and vapour phases, than :

A. $X_A = Y_A$

B. $X_A > Y_A$

C. $X_A < Y_A$

D. Data insufficient

Answer: c

 [Watch Video Solution](#)

18. At $25^{\circ}C$, the vapour pressure of pure liquid A (mol. Mas = 40) is 100 torr, and pure liquid B is 40 torr (mol. = 80). The vapour pressure at $25^{\circ}C$ of a solution containing 20 g of each A and B is :

- A. 80 torr
- B. 59.8 torr
- C. 68 torr
- D. 48 torr

Answer: a

 [Watch Video Solution](#)

19. The liquid A and B form ideal solutions. At 300 K, the vapour pressure of solution containing 1 mole of A and 3 mole of B is 550 mm Hg. At the same temperature, if one more mole of B is added to this solution, the vapour pressure of the solution increases by 10 mm Hg. Determine the vapour pressure of A and B in their pure states (in mm Hg):

A. 400, 600

B. 500, 500

C. 600, 400

D. none of these

Answer: a

 [Watch Video Solution](#)

20. Two liquid A and B have vapour pressure in the ratio $P_A^\circ : P_B^\circ = 1.3$ at a certain temperature. Assume A and B form an ideal solution and the ratio

of mole fractions of A to B in the vapour phase is 4 : 3, then the mole fraction of B in the solution at the same temperature is :

A. $\frac{1}{5}$

B. $\frac{2}{3}$

C. $\frac{4}{5}$

D. $\frac{1}{4}$

Answer: a



Watch Video Solution

21. Two liquids A and B have P_A° and P_B° in the ratio of 1 : 3 and the ratio of number of moles of A and B in liquid phase are 1 : 3 then mole fraction of 'A' in vapour phase in equilibrium with the solution is equal to :

A. 0.1

B. 0.2

C. 0.5

D. 1


Answer: a

 Watch Video Solution

22. Calculate the oxidation number of nitrogen (N) in N_2H_4

 Watch Video Solution

23. Which represents correct difference when non-volatile solute is present in an ideal solution?



A. I, II, III

B. I, III

C. II, III

D. I, II

Answer: a



Watch Video Solution

24. Select correct statement :

A. Solution has more molecules randomness than a pure solvent. The

entropy change between solution and solid is larger than the

entropy change between pure solvent and solid

B. Heat of fusion of solution are positive

C. Solution containing sugar freezes at a lower temperature than

pure water

D. All are correct statements

Answer: d

 [Watch Video Solution](#)

25. Select correct statement :


- A. Heats of vaporisation for a pure solvent and for a solution are similar because similar intermolecular forces between solvent molecules must be overcome in both cases
- B. Entropy change between solution and vapour is smaller than the entropy change between pure solvent and vapour
- C. Boiling point of the solution is larger than that of the pure solvent
- D. All are correct statements

Answer: d

 [Watch Video Solution](#)

26. The vapour pressure curves of the same solute in the same solvent are shown below. The curves are parallel to each other and do not intersect.

The concentrations of solutions are in order of :



A. $I < II < III$

B. $I = II = III$

C. $I > II > III$

D. $I > III > II$

Answer: a



Watch Video Solution

27. What is the importance of (a) melting point of solid and (b) boiling point of a liquid ?

- A. Composition of the still (residus) will approach pure liquid B only
- B. composition of the distillate will approach pure A only
- C. Composition of distillate and residus will approach pure A and B respectively
- D. Neither of the componement can be obtained in pure state

Answer: c



Watch Video Solution

28. Dettol is a mixture of :

- A. the mixture will show negative deviation from Raoult's law
- B. the mixture will show positive deviation from Raoult's law
- C. the mixture can be considered as pure solution
- D. this mixture can be considered as pure solution

Answer: b

 [Watch Video Solution](#)

29. Formation of a solution from two components can be considered as

:

- (i) pure solvent \rightarrow separated solvent molecules, ΔH_1
- (ii) Pure solute \rightarrow separated molecules, ΔH_2
- (iii) separated solvent and solute molecules \rightarrow solution, ΔH_3

solution so formed will be ideal if :

A. $\Delta H_{\text{soln}} = \Delta H_1 + \Delta H_2 + \Delta H_3$

B. $\Delta H_{\text{soln}} = \Delta H_1 + \Delta H_2 - \Delta H_3$

$$C. \Delta H_{\text{soln}} = \Delta H_1 - \Delta H_2 - \Delta H_3$$

$$D. \Delta H_{\text{soln}} = \Delta H_3 - \Delta H_1 - \Delta H_2$$

Answer: a

 [Watch Video Solution](#)

30. Total vapour pressure of mixture of 1 mole of volatile components A ($P_{a\%}$)=100 mm Hg) and 3 mole of volatile component B ($P_B^\circ = 80\text{mmHg}$) is 90 mm Hg. For such case:

- A. there is a negative deviation from Raoult's law
- B. there is a positive deviation from Raoult's law
- C. there is no deviation from Raoult's law
- D. can not be decided

Answer: a

 [Watch Video Solution](#)

31. In a mixture of A and B, components show positive deviation when:

A. A-B interaction is stronger than A-A and B-B interaction

B. A-B interaction is weaker than A-A and B-B interaction

C. $\Delta V_{\text{mix}} < 0$, $\Delta S_{\text{mix}} > 0$

D. $\Delta V_{\text{mix}} = 0$, $\Delta S_{\text{mix}} > 0$


Answer: b



Watch Video Solution

32. A liquid mixture having composition corresponding to point Z in the figure shown is subjected to distillation at constant pressure. Which of

the following statements is correct about the process?



- A. The composition of distillate differs from the mixture
- B. The boiling point goes on changing
- C. The mixture has highest vapour pressure than for any other composition
- D. Composition of an azeotrope alters on changing the external pressure

Answer: d



Watch Video Solution

33. Which will form maximum boiling azeotrope ?

A. $C_6H_6 + C_6H_5CH_3$ solution

B. $HNO_3 + H_2O$ solution

C. $C_2H_5OH + H_2O$ solution

D. n-hexane and n-heptane

Answer: b



Watch Video Solution

34. Total vapour pressure of mixture of 1 mole of volatile components A (

P_A°) = 100 mm Hg) and 3 mole of volatile component B ($P_B^\circ = 80 \text{ mm.Hg}$)

is 90 mm Hg. For such case:

A. There is positive deviation from Raoult's law

B. boiling point has been lowered

- C. force of attraction between A and B is weaker than that between A and A or between B and B
- D. All the above statements are correct

Answer: d

 [Watch Video Solution](#)

35. The azeotropic mixture of water ($B. P. = 100^{\circ}C$) and HCl ($B. P. = 86^{\circ}C$) boils at about $120^{\circ}C$. During fractional distillation of this mixture it is possible to obtain :

- A. pure HCl
- B. pure H_2O
- C. pure H_2O as well as pure HCl
- D. Neither C_2H_5OH nor HCl

Answer: d



Watch Video Solution

36. The azeotropic mixture of water ($B. P. = 100^{\circ}C$) and HCl ($B. P. = 86^{\circ}C$) boils at about $120^{\circ}C$. During fractional distillation of this mixture it is possible to obtain :

- A. pure C_2H_5OH only
- B. Pure water only
- C. Neither C_2H_5OH nor water
- D. Both water and C_2H_5OH in pure state

Answer: c



Watch Video Solution

37. Which of the following has highest boiling point ?

- A. shows positive deviation from Raoult's law

B. shows negative deviation from Raoult's law

C. shows ideal behaviour

D. is saturated

Answer: b

 [Watch Video Solution](#)

38. If two liquids A ($P_A^\circ = 100\text{torr}$) and ($P_B^\circ = 200\text{ torr}$) which are completely immiscible with each other (each one will behave independently of the other) are present in a closed vessel, the total vapour pressure of the system will be :

A. less than 100 torr

B. greater than 200 torr

C. between 100 to 200 torr

D. 300 torr

Answer: d



Watch Video Solution

39. A solution contain 8 g of a carbohydrate in 100 g of water has a density 1.025 g/mL and an osmotic pressure of 5 atm at $27^{\circ}C$. What is the molar mass of the carbohydrate?

- A. 7.975 g/mol
- B. 166 g/ mol
- C. 145.8 g/mol
- D. None of these

Answer: c



Watch Video Solution

40. If the density of some lake water is 1.25 g mL^{-1} and contains 92 g of Na^+ ions per kg of water, calculate the molality of Na^+ ions in the lake.

- A. 50
- B. 60
- C. 78.3
- D. 38.46

Answer: d

 [Watch Video Solution](#)

41. Which of the following is a colligative property?

- A. Vapour pressure
- B. Depression in f.pt.
- C. Elevation in b.pt.

D. Osmotic pressure

Answer: a

 [Watch Video Solution](#)

42. The degree of an electrolyte is a and its van't Hoff factor is i . The number of ions obtained by complete dissociation of 1 molecule of electrolyte as :

A. $\frac{i + a - 1}{a}$

B. $i - a - 1$

C. $\frac{i - 1}{a}$

D. $i + 1 + \frac{a}{1 - a}$

Answer: a

 [Watch Video Solution](#)

43. One mole of a solute A is dissolved in a given volume of a solvent. The association of the solute take place as follows: $nA \rightarrow A_n$. If a is the degree of association of A, the van't hoff factor i is expressed as:

A. $i = 1 - a$

B. $i = 1 + \frac{a}{n}$

C. $i = \frac{1 - a + \frac{a}{n}}{1}$

D. $i = 1$

Answer: C



Watch Video Solution

44. The van't Hoff factor i for an electrolyte which undergoes dissociation and association in solvents respectively are:

A. greater than one and less than one

B. less than one and greater than one

C. less than one and less than one

D. greater than one and greater than one

Answer: A

 [Watch Video Solution](#)

45. Which solution has the highest vapour pressure ?

A. 0.02 M NaCl at 50°C

B. 0.03 M sucrose at 15°C

C. 0.005 M CaCl_2 at 50°C

D. 0.005 M CaCl_2 at 25°C

Answer: c

 [Watch Video Solution](#)

46. An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase ?

- a) addition of water
- b) addition of NaCl
- c) addition of Na_2SO_4
- d) Addition of 1.0 molal KI

A. addition of water

B. addition of NaCl

C. addition of Na_2SO_4

D. Addition of 1.0 molal KI

Answer: a



[Watch Video Solution](#)

47. Calculate the oxidation number of U in $(UO_2)_2^{+}$



[Watch Video Solution](#)

48. Calculate the oxidation number of S in K_2S

 [Watch Video Solution](#)

49. The vapour pressure of an aqueous solution of sucrose at 373 K is found to be 750 mm Hg. The molality of the solution at the same temperature will be :

A. 6.3×10^{-4}

B. 3.5×10^{-2}

C. 5×10^{-3}

D. None of these

Answer: c

 [Watch Video Solution](#)

50. When 1 mole of a solute is dissolved in 1 kg of H_2O , boiling point of solution was found to be $100.5^\circ C$. K_b for H_2O is :

- A. 0.5
- B. 100
- C. 100.5
- D. 95.5

Answer: a



Watch Video Solution

51. Chloroform has $\Delta H_{\text{vaporization}} = 29.2 \text{ kJ/mol}$ and boils at $61.2^\circ C$.

What is the value of $\Delta S_{\text{vaporization}}$ for chloroform ?





- A. 61.9
- B. 62
- C. 52.2

D. 62.67

Answer: d

 [Watch Video Solution](#)

52. When 36.0 g of a solute having the empirical formula CH_2O is dissolved in 1.20 kg of water, the solution freezes at $-0.93^\circ C$. What is the molecular formula of the solute? ($K_f = 1.86^\circ Ckgmol^{-1}$)



Answer: d

 [Watch Video Solution](#)

53. A solution of 0.640 g of azulene in 100.0 g of benzene is $80.23^\circ C$. The boilingpoint of benzene is $80.10^\circ C$, and K_b is $2.53^\circ C/\text{molal}$. What is the molecular mass of azulene?

- a) 108
- b) 99
- c) 125
- d) 134

A. 108

B. 99

C. 125

D. 134

Answer: c



Watch Video Solution

54. One molal solution of a carboxylic acid in benzene shows the elevation of boiling point of 1.518 K. The degree of association for dimerization of the acid in benzene is (K_b for benzene = $2.53 \text{ K kg mol}^{-1}$):

A. 0.6

B. 0.7

C. 0.75

D. 0.8

Answer: d



Watch Video Solution

55. The normal boiling point of toluene is 110.7°C and its boiling point elevation constant $3.32 \text{ K kg mol}^{-1}$. The enthalpy of vaporization of toluene is nearly :

A. 17.0 kJ mol^{-1}

B. 34.0 kJ mol^{-1}

C. 51.0 kJ mol^{-1}

D. 68.0 kJ mol^{-1}

Answer: b



Watch Video Solution

56. Which one of the following aqueous solutions will exhibit highest boiling point?

A. 0.015 M urea

B. 0.01 M KNO_3

C. 0.10 M Na_2SO_4

D. 0.015 m glucose

Answer: c



Watch Video Solution

57. Calculate the percentage degree of dissociation of an electrolyte XY_2 (Normal molar mass = 164) in water if the water if the observed molar mass by measuring elevation in boiling point is 65.6

A. 75 %

B. 25 %

C. 65 %

D. None of these

Answer: a



Watch Video Solution

58. if the elevation in boiling point of a solution of non-volatile, non-electrolytic and non-associating solute in solvent ($K_b = x K kg mol^{-1}$) is yK , then the depression in freezing point of solution of same

concentration would be

(K_f) of the solvent = $zk. \text{kgmol}^{-1}$)

A. $2x \frac{y}{y}$

B. $y \frac{z}{x}$

C. $x \frac{z}{y}$

D. $y \frac{z}{2x}$

Answer: b



Watch Video Solution

59. When a solution containing non-volatile solute freezes, which equilibrium would exist?

A. solid solvent \Leftrightarrow liquid solvent

B. solid solute \Leftrightarrow liquid solution

C. solid solute \Leftrightarrow liquid solvent

D. solid solvent \Leftrightarrow liquid solution

Answer: d

 [Watch Video Solution](#)

60. Bromoform has a normal freezing point of 7.734°C and $K_f = 14.4^{\circ}\text{C}/m$. A solution of 2.60 g of an unknown substance in 100 g of bromoform freezes at 5.43°C . What is the molecular mass of the unknown substance?

A. 16.25

B. 162.5

C. 100

D. none of these

Answer: b

 [Watch Video Solution](#)

61. C_6H_6 freezes at $5.5^\circ C$. At what temperature will a solution of 10.44 g of C_4H_{10} in 200 g of C_6H_6 freeze? $K_f(C_6H_6) = 5.12^\circ C/m$

A. $4.608^\circ C$

B. $0.892^\circ C$

C. $5.5^\circ C$

D. none of these

Answer: b



Watch Video Solution

62. How much ethyl alcohol must be added to 1.00L of water so that the solution will not freeze at $-4^\circ F$?

A. 20 g

B. 10.75 g

C. 492.2 g

D. 495 g

Answer: d



Watch Video Solution

63. The freezing point of a solution of 2.40 g of biphenyl($C_{12}H_{10}$) in 75.0 g of benzene (C_6H_6) is $4.40^\circ C$. The normal freezing point of benzene is $5.50^\circ C$. What is the molal freezing point constant ($^\circ C/m$) for benzene ?

A. 5.3

B. 5.1

C. 4.6




D. 4.8

Answer: a



Watch Video Solution

64. A solution containing 1.8 g of a compound (empirical formula CH_2O) in 40 g of water is observed to freeze at -0.465°C . The molecules formula of the compound is (K_f of water $=1.86\text{kg Kmol}^{-1}$):



Answer: d



Watch Video Solution

65. Freezing point of the following equilibrium, liquid solvent \rightleftharpoons solid solvent is :

a) $\frac{\Delta H - \Delta G}{T \Delta S}$

b) $\frac{\Delta H}{\Delta S}$

$$\text{c) } \frac{\Delta G}{\Delta S}$$

$$\text{d) } \frac{\Delta S}{\Delta H}$$

$$\text{A. } \frac{\Delta H - \Delta G}{T \Delta S}$$

$$\text{B. } \frac{\Delta H}{\Delta S}$$

$$\text{C. } \frac{\Delta G}{\Delta S}$$

$$\text{D. } \frac{\Delta S}{\Delta H}$$

Answer: b

 [Watch Video Solution](#)

66. Freezing point of a solvent containing a non volatile solute

A. ΔH of solution and solvent is almost identical since intermolecular force between solvent molecules are involved

B. ΔS solution (between solution and solid) is larger than that of the ΔS of solvent (between solvent and solid)

C. ΔS of the solution is smaller than that of the solvent

D. ΔH of the solution is much higher than of solvent but ΔS of solvent than that of the solvent

Answer: b

 [Watch Video Solution](#)

67. When 36.0 g of a solute having the empirical formula CH_2O is dissolved in 1.20 kg of water, the solution freezes at $-0.93^\circ C$. What is the molecular formula of the solute? ($K_f = 1.86^\circ Ckgmol^{-1}$)

A. C_2H_4O

B. $C_2H_2O_2$

C. $C_2H_4O_3$

D. $C_2H_4O_2$

Answer: d



Watch Video Solution

68. Calculate the molar mass of a substance whose 7.0% by mass solution in water freezes at -0.93°C . The cryoscopic constant of water is $1.86^{\circ}\text{C kg mol}^{-1}$:

A. 140 g mol^{-1}

B. 150.5 g mol^{-1}

C. 160 g mol^{-1}

D. 155 g mol^{-1}

Answer: b



Watch Video Solution

69. Camphor is often used in molecular mass determination because

A. it is readily available

B. it has a very high cryoscopic constant

C. it is volatile

D. it is solvent for organic substances




Answer: b


 [Watch Video Solution](#)

70. Calculate the oxidation number of nitrogen (N) in N_2O_5

 [Watch Video Solution](#)

71. Which of the following 0.1 M aqueous solution will have lowest freezing point?









Answer: d



Watch Video Solution

72. Calculate depression of freezing point for 0.56 molal aq. Solution of KCl.

(Given : $K_{f(H_2O)} = 1.8 \text{ kg mol}^{-1}$).






Answer: c



Watch Video Solution

73. A complex is represented as $CoCl_3 \cdot xNH_3$. Its 0.1 molal solution in aqueous solution shows $\Delta T_f = 0.558^\circ$. (K_f for H_2O is $1.86 K molality^{-1}$) Assuming 100% ionisation of complex and coordination number of Co as six, calculate formula of complex.







D. none of these

Answer: b

 Watch Video Solution

74. The freezing point of equimolar aqueous solutions will be highest for





Answer: d



Watch Video Solution

75. The freezing point of 4% aqueous solution of 'A' is equal to the freezing point 10% aqueous solution of 'B'. If the molecules mass of 'A' is 60, then the molecules mass of 'B' will be:

A. 160

B. 90

C. 45

D. 180

Answer: a



Watch Video Solution

76. The depression in freezing point of $0.01m$ aqueous CH_3COOH solution is 0.02046° , $1m$ urea solution freezes at $-1.86^\circ C$. Assuming molality equal to molarity, pH of CH_3COOH solution is

- A. 2
- B. 3
- C. 4
- D. 5

Answer: b

 [Watch Video Solution](#)

77. When mercuric iodide is added to the aqueous solution of potassium iodide, then:

- A. freezing point is raised

- B. Freezing point is lowered
- C. freezing point does not change
- D. boilingpoint does not change

Answer: a

 [Watch Video Solution](#)

78. Draw structure of acetone.

 [Watch Video Solution](#)

79. The temperature of a city was found to be -9.3°C . A car used, whose rasiator was filled with 5 L of water . What minimum quantity of antifreezing agent ethylene glycol were added to water of raiator in order to use the car for teavelling? (K_f of water 1.86 k mol^{-1})

A. 3200 g

B. 1670 g

C. 1550 g

D. 2100 g

Answer: c



Watch Video Solution

80. The cryoscopic constant of water is $1.86 \text{ K kg mol}^{-1}$. A 0.01 molal acetic acid solution produces a depression of 0.0194°C in the freezing point.

The degree of dissociation of acetic acid is :

A. zero

B. 0.043

C. 0.43

D. 1

Answer: b



Watch Video Solution

81. In a 0.5 molal solution KCl, KCl is 50% dissociated. The freezing point of solution will be ($K_f = 1.86 \text{ K kg mol}^{-1}$):

- A. 274.674 K
- B. 271.60 K
- C. 273 K
- D. none of these

Answer: b



Watch Video Solution

82. A 1.0 g sample of $\text{co}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_3\text{Cl}_3$ is dissolved in 25.0 g of water and the freezing point of the solution is -0.87°C . How many ions are produced per mole of compound? The K_f of water is 1.86°C/molal

A. 2

B. 3

C. 4

D. 5

Answer: c

 [Watch Video Solution](#)

83. Draw structure of metoxymethane.

 [Watch Video Solution](#)

84. phenol associates in benzene to a certain extent in dimerisation reaction. A solution containing 0.02 kg of phenol in 1.0 kg of benzene has its freezing point depressed 0.69 k. Hence, the degree of association of phenol dimerized will be, [$K_f(C_6H_6) = 5.12\text{ k mol}^{-1}$]

a) 0.63

b) 0.73

c) 0.83

d) 0.93

A. 0.63

B. 0.73

C. 0.83

D. 0.93

Answer: b



[Watch Video Solution](#)

85. Draw structure of methanal



[Watch Video Solution](#)

86. In a 0.2 molal aqueous solution of a weak acid HX the degree of dissociation is 0.25. The freezing point of the solution will be nearest to: (

$$K_f = 1.86 \text{ K kg mol}^{-1})$$

- a) -0.26°C
- b) 0.465°C
- c) -0.48°C
- d) -0.465°C

A. -0.26°C

B. 0.465°C

C. -0.48°C

D. -0.465°C

Answer: d



Watch Video Solution

87. An aqueous solution of 0.01 M KCl cause the same elevation in boiling point as an aqueous solution of urea. The concetration of urea solution

is :

a) 0.01 m

b) 0.005 M

c) 0.02 M

d) 0.04 M

A. 0.01 m

B. 0.005 M

C. 0.02 M

D. 0.04 M

Answer: c



Watch Video Solution

88. Write structure o 1,1- dichloropropane.



[Watch Video Solution](#)

89. Write structural formula of but-2-en-1-oic acid



[Watch Video Solution](#)

90. Write structural formula of hexa-1,3,5-triene



[Watch Video Solution](#)

91. Write bond line formula for tert- butylcyclopentane.



[Watch Video Solution](#)

92. Write bond line formula for isopropylalcohol.



[Watch Video Solution](#)

93. Write bond line formula for 2,3-dimethylbutanal.

 [Watch Video Solution](#)

94. Which one of the following pairs of solutions can be expected to be isotonic at the same temperature ?

A. 0.1 M urea and 0.1 M NaCl

B. 0.1 M urea and 0.2 M $MgCl_2$




C. 0.1 M NaCl and 0.1 M Na_2SO_4

D. 0.1 M $Ca(NO_3)_2$ and 0.1 M Na_2SO_4

Answer: d

 [Watch Video Solution](#)

95. The empirical formula of a non-electrolyte is CH_2O . A solution containing 3 g L^{-1} of the compound exerts the same osmotic pressure as that of 0.05 M glucose solution. The molecules formula of the compound is :



Answer: b

 [Watch Video Solution](#)

96. A semipermeable membrane used in the measurement of osmotic pressure of a solution allows the passage of

A. solute molecular through it

- B. solvent molecules though it
- C. both solvent and solute molecules
- D. either solvent or solute

Answer: b

 [Watch Video Solution](#)

97. In the case of osmosis, solvent molecules move from :

- A. higher vapour pressure to lower vapour pressure
- B. higher concentration to lower concentration
- C. lower vapour pressure to higher vapour pressure
- D. higher osmotic pressure to lower osmotic pressure

Answer: a

 [Watch Video Solution](#)

98. The osmotic pressures of equimolar solutions of urea, $BaCl_2$ and $AlCl_3$ will be in the order :

A. $AlCl_3 > BaCl_2 > urea$

B. $BaCl_2 > AlCl_3 > urea$

C. $urea > BaCl_2 > AlCl_3$

D. $BaCl_2 > urea > AlCl_3$

Answer: a



Watch Video Solution

99. Assuming each salt to be 90 % dissociated which of the following will have the highest osmotic pressure?

A. decimolar aluminium sulphate

B. decimolar barium chloride solution

C. decimolar sodium sulphate solution

D. solution of volume of decimolar barium chloride and decimolar sodium sulphate solutions

Answer: a



Watch Video Solution

100. consider 0.1 M solutions of two solutes X and Y. The X behaves as a univalent electrolyte while the solute Y dimerises in solution. Which of the following statements are correct regarding these solutions?

- (1) The boiling point of the solution of X will be higher than that of Y
 - (2) The osmotic pressure of the solution of Y will be lower than that of X
 - (3) The freezing point of the solution of X will be lower than that of Y
 - (4) The relative lowering of vapour pressure of both the solutions will be the same
- It is to be noted that Select the correct answer from the option given below

A. 1, 2 and 3

B. 2, 3 and 4

C. 1, 2 and 4

D. 1, 3 and 4

Answer: a

 [Watch Video Solution](#)

101. If M_{normal} is the normal molecular mass and α is the degree of ionization of $K_3[Fe(CN)_6]$, then the abnormal molecular mass of the complex in the solution will be :

A. $M_{\text{normal}}(1 + 2\alpha)^{-1}$

B. $M_{\text{normal}}(1 + 3\alpha)^{-1}$

C. $M_{\text{normal}}(1 + \alpha)^{-1}$

D. equal to M_{normal}

Answer: b

 [Watch Video Solution](#)

102. Equal volumes of 0.1 M urea and 0.1 M glucose are mixed. The mixture will have :

- A. lower osmotic pressure
- B. same osmotic pressure
- C. higher osmotic pressure
- D. none of these

Answer: b

 [Watch Video Solution](#)

103. A 5% (w/V) solution of cane sugar (molecular mass = 342) is isotonic with 1% (w/V) solution of a substance X. The molecular mass of X is :

- A. 34.2
- B. 171.2

C. 68.4

D. 136.8

Answer: c



[Watch Video Solution](#)

104. A 1% (mass/vol) KCl solution is ionised to the extent of 80%. The osmotic pressure at 27° C of the solution will be :

A. 3×10^5

B. 9×10^5

C. 4.5×10^5

D. 5.16×10^6

Answer: d



[Watch Video Solution](#)

105. An aqueous solution of sucrose ($C_{12}H_{22}O_{11}$) having a concentration of 34.2 gram/ litre has an osmotic pressure of 2.38 atmospheres at $17^{\circ}C$. For an aqueous solution of glucose ($C_6H_{12}O_6$) to be isotonic with this solution, its concentration should be :

- A. 34.2 gram per liter
- B. 17.1 gram per liter
- C. 18.0 gram per liter
- D. 36.0 gram per liter

Answer: c

 [Watch Video Solution](#)

106. Which of the following experimental methods is adopted to determine osmotic pressure?

- A. Berkley- Hartely's method

B. Beckmann's method

C. Landsberger's method

D. Differential method

Answer: a

 [Watch Video Solution](#)

107. Estimste the lowering of vapour pressure due to the solute (glucose) in a 1.0 m aqueous solution at $100^{\circ}C$:

A. 10.5 atm

B. 21 atm

C. 12.2 atm

D. 6.09 atm

Answer: c

 [Watch Video Solution](#)

108. A 1% (mass/vol) KCl solution is ionised to the extent of 80%. The osmotic pressure at 27° C of the solution will be :

A. 6.95 atm

B. 5.94 atm

C. 2.71 atm

D. 3.30 atm

Answer: b

 [Watch Video Solution](#)

109. Osmotic pressure of blood is 7.40 atm, at 27° C. Number of moles of glucose to be used per liter for an intravenous injection that is to have same osmotic pressure of blood is :

A. 0.3

B. 0.2

C. 0.1

D. 0.4

Answer: a

 [Watch Video Solution](#)

110. The relationship between osmotic pressure (π_1, π_2 and π_3) at a definite temperature when 1 g glucose, 1 g urea and 1 g sucrose are dissolved in 1 litre of water is (assume $i = 1$ for all):

A. $\pi_1 > \pi_2 > \pi_3$

B. $\pi_3 > \pi_1 > \pi_2$

C. $\pi_2 > \pi_1 > \pi_3$

D. $\pi_2 > \pi_3 > \pi_1$





Answer: c

111. van't Hoff proved that osmotic pressure (π) is a colligative property.

For an ideal solution, osmotic pressure(π) is helpful to determine that

molecular mass of solute using $M_B = \frac{W_B RT}{\pi \cdot V}$ Relation can expressed

by the curve (C = concentration) :



Answer: a



Watch Video Solution

112. A solution containing 4.0 g of PVC in 2 liter of dioxane (industrial solvent) was found to have an osmotic pressure 3.0×10^{-4} atm at $27^\circ C$.

The molar mass of the polymer (g/mol) will be :

A. 1.6×10^4

B. 1.6×10^5

C. 1.6×10^3

D. 1.6×10^2

Answer: b



Watch Video Solution

113. The osmotic pressures of 0.010 M solutions of KI and sucrose ($C_{12}H_{22}O_{11}$) are 0.432 atm and 0.24 atm respectively. The van't Hoff factor for KI is :

A. 1.8

B. 0.8

C. 1.2

D. 1

Answer: a



[Watch Video Solution](#)

114. What is the correct sequence of osmotic pressure of 0.01M aq. solution of :

(a) $Al_2(SO_4)_3$ (b) Na_3PO_4 (c) $BaCl_2$ (d) *Glucose*

A. $\pi_4 > \pi_2 > \pi_3 > \pi_1$

B. $\pi_3 > \pi_4 > \pi_2 > \pi_1$

C. $\pi_3 > \pi_4 > \pi_1 > \pi_2$

D. $\pi_1 > \pi_2 > \pi_3 > \pi_4$

Answer: d

 [Watch Video Solution](#)

115. Osmotic pressure of a solution is 0.0821 atm at temperature of 300 K.

The concentration of solution in *mol / litre* will be

A. 0.5 mole of AgCl

B. 1.0 mole of AgCl

C. 1.5 mole of AgCl

D. 3.0 mole of AgCl

Answer: b

 [Watch Video Solution](#)

116. A 1.0 g sample of $\text{Co}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_3\text{Cl}_3$ is dissolved in 25.0 g of water and the freezing point of the solution is -0.87°C . How many ions are produced per mole of compound? The K_f of water is $1.86^\circ\text{C}/\text{molal}$

A. 1

B. 4

C. 2

D. 3

Answer: c



Watch Video Solution

117. Which of the following aqueous solutions should have the highest osmotic pressure?

A. 0.011 M AlCl_3 at 50°C

B. 0.03 m NaCl at 25° C

C. 0.012 m (NH₄)₂SO₄ at 25°

D. 0.03 m NaCl at 50° C


Answer: d

 Watch Video Solution


118. Write iupac name of CH₂Cl₂

 Watch Video Solution


119. Which of the following curves represents the Henry's law ?




A.



B.



C.



D.

Answer: a



Watch Video Solution

120. According to Henry's law, the solubility of a gas in a given volume of liquid increases with increases in :

- A. trmperature
- B. pressure
- C. Both (a) and (b)
- D. none of these

Answer: b

 [Watch Video Solution](#)

121. Determine the amount of $CaCl_2$ ($i = 2.47$) dissolved in 2.5 litre of water such that its osmotic pressure is 0.75 atm at $27^\circ C$.

- A. 0.1 g
- B. 1.24 g
- C. 0.48 g
- D. 4.8 g

Answer: b

 [Watch Video Solution](#)

122. 1 kg of water under a nitrogen pressure of 1 atmosphere dissolves 0.02 gm of nitrogen at 293 k. Calculate Henry's law constant :

- A. 7.2×10^{-4} L/atm

B. 7.7×10^3 atm

C. 2×10^{-5} atm

D. 2×10^{-2} atm

Answer: a

 [Watch Video Solution](#)

123. According to Henry's law, the partial pressure of gas (p'_g) is directly proportional to mole fraction of gas in dissolved state, i.e.,

$P_{\text{gas}}' = K_H, X_{\text{gas}}$ where K_H is Henry's constant. Which are correct?

A. K_H is characteristic constant for a given gas-solvent system

B. Higher is the value of K_H , lower is solubility of gas for a given partial pressure of gas

C. K_H has temperature dependence

D. K_H decreases with increase of temperature

Answer: d



Watch Video Solution

124. At 760 torr pressure and $20^{\circ}C$ temperature, 1 L of water dissolves 0.04 gm of pure oxygen or 0.02 gm of pure nitrogen. Assuming that dry air is compound of 20% oxygen and 80% nitrogen (by volume), the masses (in g/L) of oxygen and nitrogen dissolved by 1 L of water at $20^{\circ}C$ exposed to air at a total pressure of 706 torr are respectively :

A. 0.008, 0.016

B. 0.016, 0.008

C. 0.16, 0.08

D. 0.04, 0.02

Answer: a



Watch Video Solution

125. Write bond line formula for Heptan-4-one.

 [Watch Video Solution](#)

126. At $48^{\circ}C$, the vapour pressure of pure CS_2 is 850 torr . A solution of 2.0 g of sulphur in 100g of CS_2 has a vapour pressure 844.9 torr.

Determine the atomicity of sulphur molecule :

A. 1

B. 2

C. 4

D. 8

Answer: d

 [Watch Video Solution](#)

127. For a binary ideal liquid solution, the total vapour of the solution is given as:

- A. 150
- B. 180
- C. 188.88
- D. 198.88

Answer: c



[Watch Video Solution](#)

128. Draw structure of 3-chlorobutan-1-ol



[Watch Video Solution](#)

129. The functional group present in butanal is carboxylic acid.



[Watch Video Solution](#)

130. A certain non-volatile electrolyte contains 40% carbon, 6.7% hydrogen and 53.3% oxygen. An aqueous solution containing 5% by mass of the solute boils at 100.15°C . Determine the molecular formula of the compound ($K_b = 0.51^\circ\text{C}/m$):

- A. HCHO
- B. CH_3OH
- C. $\text{C}_2\text{H}_5\text{OH}$
- D. $\text{C}_6\text{H}_{12}\text{O}_6$

Answer: d

131. Draw the structure of 2-methylpropane.

132. An aqueous solution boils at 101°C . What is the freezing point of the same solution?

(Gives : $K_f = 1.86^{\circ}\text{C}/m$ and $K_b = 0.51^{\circ}\text{C}/m$)

A. 3.647°C

B. -3.647°C

C. -0.199°C

D. none of these

Answer: b

 [Watch Video Solution](#)

133. Draw the structure of simplest alkene.

 [Watch Video Solution](#)

134. Ratio of $\frac{\Delta T_b}{K_b}$ of 10 g AB_2 and 14 g A_2B per 100 g of solvent in their respective, solution (AB_2 and A_2B both are non-electrolytes) is 1 mole/ kg in both cases. Hence, atomic wt. of A and B are respectively :

- A. 100, 40
- B. 60, 20
- C. 20, 60
- D. None of these

Answer: b

 [Watch Video Solution](#)

135. The freezing point of solution containing 0.2g of acetic acid in 20.0g of benzene is lowered by $0.45^\circ C$. Calculate the degree of association of acetic acid in benzene.

$$(K_f = 5.12K^\circ mol^{-1}kg^{-1})$$

A. 0.527

B. 0.8

C. 0.945

D. None of these

Answer: c



Watch Video Solution

136. An aqueous solution boils at $101^{\circ}C$. What is the freezing point of the same solution?

(Gives : $K_f = 1.86^{\circ}C/m$ and $K_b = 0.51^{\circ}C/m$)

A. $0.361^{\circ}C$

B. $-0.361^{\circ}C$

C. $-3.61^{\circ}C$

D. None of these

Answer: b

 [Watch Video Solution](#)

137. When 36.0 g of a solute having the empirical formula CH_2O is dissolved in 1.20 kg of water, the solution freezes at $-0.93^\circ C$. What is the molecular formula of the solute? ($K_f = 1.86^\circ Ckgmol^{-1}$)

A. 95.5 g

B. 4.5 g

C. 45.5 g

D. 47.8 g

Answer: d

 [Watch Video Solution](#)

138. In a 0.5 molal solution KCl, KCl is 50% dissociated. The freezing point of solution will be ($K_f = 1.86 \text{ K kg mol}^{-1}$):

A. 0.2

B. 0.25

C. 0.4

D. 0.5

Answer: b

 [Watch Video Solution](#)

139. 0.1 M KI and 0.2 M $AgNO_3$ are mixed in 3 : 1 volume ratio. The depression of freezing point of the resulting solution will be [$K_b(H_2O) = 1.86 \text{ K kg mol}^{-1}$]:

A. 3.72 K

B. 1.86 K

C. 0.93 K

D. 0.279 K

Answer: d

 [Watch Video Solution](#)

140. If 0.1 M $H_2SO_4(aq.)$ solution shows freezing point $0.3906^\circ C$ then what is the K_{a2} for H_2SO_4 ? (Assume $m = M$ and $K_f(H_2O) = 1.86 Kkgmol^{-1}$)

A. 0.122

B. 0.0122

C. 1.11×10^{-3}

D. None of these

Answer: b

 [Watch Video Solution](#)

141. A living cell contains a solution which is isotonic with 0.2 M glucose solution. What osmotic pressure develops when the cell is placed in 0.05 M $BaCl_2$ solution at 300 K ?

- A. 1.23 atm
- B. 3.69 atm
- C. 6.15 atm
- D. None of these

Answer: a



[Watch Video Solution](#)

142. What is the correct sequence of osmotic pressure of 0.01 M aq. Solution of :

- (1) $Al_2(SO_4)_3$
- (2) Na_3PO_4
- (3) $BaCl_2$
- (4) Glucose

A. 4.926 atm

B. 0.5024 atm

C. 5.024 atm

D. None of these

Answer: c

 [Watch Video Solution](#)

143. A solution contain 8 g of a carbohydrate in 100 g of water has a density 1.025 g/mL and an osmotic pressure of 5 atm at 27°C . What is the molar mass of the carbohydrate?

A. 387

B. 374

C. 3740


D. None of these

Answer: b



Watch Video Solution

144. Study the following figure and choose the correct options. Assuming complete dissociation of electrolyte:



- A. There will be net movement of any substance across the membrane
- B. $MgCl_2$ will flow towards the $Al_2(SO_4)_3$ solution
- C. $Al_2(SO_4)_3$ will flow towards the $MgCl_2$ solution
- D. The π (osmotic pressure) of 0.1 M $MgCl_2$ is higher than the π of 0.05 M $Al_2(SO_4)_3$

Answer: d



Watch Video Solution

145. Write iupac name of $\text{CH}_3\text{COCH}(\text{Cl})\text{CH}_3$


 [Watch Video Solution](#)

146. Write iupac name of $\text{CH}_3\text{-C}(\text{CH}_3)_2\text{-CH}_2\text{-CH}_3$

 [Watch Video Solution](#)

147. Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.



Loss in mass of solvent (w_{II}) will be proportional to :

A. $P^\circ - P$

B. $P - P^\circ$

C. $\frac{P}{P^\circ}$

D. $P \times P^\circ$

Answer: a

[Watch Video Solution](#)


148. What do you mean by alkanes.

[Watch Video Solution](#)

149. Write structure of benzoic acid.

150. Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighed separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.




Loss in mass of solvent (w_{II}) will be proportional to :

- A. 50
- B. 62
- C. 70
- D. 80

Answer: b

 [Watch Video Solution](#)

151. A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant K_b . The solute dimerises in the solution according to the following equation. The degree of association is (a) :



The van't Hoff factor will be:

A. $i = 1 - 2a$

B. $i = 1 - \frac{a}{2}$


C. $i = 1 + \frac{a}{2}$

D. $i = 1 + a$

Answer: b

 [Watch Video Solution](#)

152. A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant K_b . The solute dimerises in the solution according to the following equation. The degree of association is (a) :



The molecular mass observed will be:

- a) greater than actual molecular mass
- b) lesser than actual molecular mass
- c) equal to the actual molecular mass
- d) cannot be predicted by the data given

A. greater than actual molecular mass

B. lesser than actual molecular mass

C. equal to the actual molecular mass

D. cannot be predicted by the data given

Answer: a

 [Watch Video Solution](#)

153. A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant K_b . The solute dimerises in the solution according to the following equation. The degree of association is (a) :



The degree of association is equal to :

$$\text{a) } a = \frac{(K_b x - \Delta T_b)}{\Delta T_b 2}$$

$$\text{b) } a = \frac{2(K_b x - \Delta T_b)}{K_b x}$$

$$\text{c) } a = 2 + \frac{2 \Delta T_b}{K_b x}$$

$$\text{d) } a = \frac{\Delta T_b}{2K_b x}$$

$$\text{A. } a = \frac{(K_b x - \Delta T_b)}{\Delta T_b 2}$$

$$\text{B. } a = \frac{2(K_b x - \Delta T_b)}{K_b x}$$

$$\text{C. } a = 2 + \frac{2 \Delta T_b}{K_b x}$$

$$\text{D. } a = \frac{\Delta T_b}{2K_b x}$$

Answer: b



Watch Video Solution

154. Which of the following statement(s) is/are correct, if intermolecular forces in liquids A, B and C are in the order of A It B It C ?

- A. B evaporates more readily than A
- B. B evaporates more readily than C
- C. A evaporate more readily than C
- D. all evaporates at same rate at same temperature.

Answer: b,c



[Watch Video Solution](#)

155. When non-volatile solute is added to a pure solvent, the:

- A. vapour pressure of the solution becomes lower than the vapour pressure of the pure solvent
- B. rate of evaporation of solvent is reduced
- C. solute does not affect the rate of condensation

D. none of these

Answer: a,b,c

 [Watch Video Solution](#)

156. The total vapour pressure of a binary solution is given by

$$P = (100X_A + 260X_B)\text{mm Hg}$$

where, X_A and X_B are the mole fractions of components A and B. This indicates that the:

- A. vapour pressure of solution is less than the pure B component
- B. vapour pressure of solution is less than the pure A component
- C. vapour pressure of pure A is 100 mm Hg and that of pure B is 260 mm Hg
- D. the vapour pressure of pure A and B are 260 mm Hg and 100 mm hg respectively

Answer: a,b,c



Watch Video Solution

157. Which of the following is correct for an ideal solution?

a) $\Delta H_{\text{mix}} = 0$ and $\Delta V_{\text{mix}} = 0$

b) $\Delta V_{\text{mix}} = 0$ and $\Delta S_{\text{mix}} > 0$

c) $\Delta H_{\text{mix}} > 0$ and $\Delta S_{\text{mix}} > 0$

d) $\Delta G_{\text{mix}} < 0$ and $\Delta S_{\text{mix}} > 0$

A. $\Delta H_{\text{mix}} = 0$ and $\Delta V_{\text{mix}} = 0$

B. $\Delta V_{\text{mix}} = 0$ and $\Delta S_{\text{mix}} > 0$

C. $\Delta H_{\text{mix}} > 0$ and $\Delta S_{\text{mix}} > 0$

D. $\Delta G_{\text{mix}} < 0$ and $\Delta S_{\text{mix}} > 0$

Answer: a,b,d



Watch Video Solution

158. Which of the following is correct for a non-ideal solution of liquids A and B showing negative deviation?

A. $\Delta H_{\text{mix}} = -ve$

B. $\Delta V_{\text{mix}} = -ve$

C. $\Delta S_{\text{mix}} = -ve$

D. $\Delta G_{\text{mix}} = -ve$

Answer: a,b,d



Watch Video Solution

159. A binary solution of liquids A and B will show positive deviation from Raoult's law if it fulfils the following condition:

A. $P_A > X_A P_{A^\circ}$ and $P_B > X_B P_{B^\circ}$

B. The intermolecular forces of $A - B < A - A, B - B$

C. ΔH_{mixing} is positive

D. ΔV mixing is negative

Answer: a,b,c

 [Watch Video Solution](#)

160. Which of the following statement is/are correct about acetone and trichloromethane mixture?

- a) Mixtures of acetone and trichloromethane show positive deviation from Raoult's law
- b) The forces of attraction acting between molecules of acetone and trichloromethane in a mixture are stronger than those acting between the molecules in pure acetone
- c) Pure acetone can be obtained by the careful fractional distillation of any mixture of acetone and trichloromethane
- d) When acetone and trichloromethane are mixed, the enthalpy change is negative

- A. Mixtures of acetone and trichloromethane show positive deviation from Raoult's law
- B. The forces of attraction acting between molecules of acetone and trichloromethane in a mixture are stronger than those acting between the molecules in pure acetone
- C. Pure acetone can be obtained by the careful fractional distillation of any mixture of acetone and trichloromethane
- D. When acetone and trichloromethane are mixed, the enthalpy change is negative

Answer: b,d

 [Watch Video Solution](#)

161. The azeotropic solution of two miscible liquids:

- A. can be separated by simple distillation

B. may show positive or negative deviation from Raoult's law

C. are supersaturated solution

D. behave like a single component and boil at a constant temperature

Answer: b,d



Watch Video Solution

162. For exact determination of molecular mass through colligative properties measurement :

A. solute must be volatile

B. solution must be very dilute

C. solution must be formed by similar nature of substances

D. solute must not be dissociated

Answer: b,d



Watch Video Solution

163. Formula of toluene is



Watch Video Solution

164. The rate constant depends on

- A. the mole mass of the solute in the solution
- B. the molar mass of the solvent in the solution
- C. the enthalpy of fusion of the solvent
- D. the freezing point of the solvent

Answer: b,c,d



Watch Video Solution

165. consider 0.1 M solutions of two solutes X and Y. The X behaves as a univalent electrolyte while the solute Y dimerises in solution. Which of

the following statement are correct regarding these solutions?

- (1) The boiling point of the solution of X will be higher than that of Y
 - (2) The osmotic pressure of the solution of Y will be lower than that of X
 - (3) The freezing point of the solution of X will be lower than that of Y
 - (4) The relative lowering of vapour pressure of both the solutions will be the same
- It brgt Select the correct answer from the option given below

- A. The boiling point of solution of 'X' will be higher than that of 'Y'
- B. The osmotic pressure of solution of 'Y' will be lower than that of 'X'
- C. The freezing point of solution of 'X' will be lower than that of 'Y'
- D. The relative lowering of vapour pressure of both the solution will be the same

Answer: a,b,c



Watch Video Solution

166. Consider following solutions: (I) 1 M glucose(aq) (II) 1 M sodium chloride(aq)

(III) 1 M acetic acid in benzene (IV) 1 M ammonium phosphate (aq)

a) all are isotonic solutions

b) III is hypotonic of I, II, IV

c) I, II, IV are hypertonic of III

d) IV is hypertonic I, II, III

A. all are isotonic solutions

B. III is hypotonic of I, II, IV

C. I, II, IV are hypertonic of III

D. IV is hypertonic I, II, III

Answer: b.c.d



Watch Video Solution

167. Which of the following statement is (are) incorrect?

- A. 0.1 M KCl solution will have the same osmotic pressure as 0.1 M glucose solution
- B. 0.1 M KCl solution will have the same boiling point as 0.1 M urea solution
- C. 0.1 m glucose and 0.1 m urea are ismotic
- D. 0.1 m $MgCl_2$ solution will have less relative lowering of vapour pressure than 0.1 m NaCl

Answer: a,b,d

 [Watch Video Solution](#)

168. Consider following solution:

0.1 m $C_6H_5NH_3^+ Cl^-$, 0.1 m KCl, 0.1 m Glucose, 0.1 m $Na_2C_2O_4 \cdot 10H_2O$

- A. the solution with higher boiling point is 0.1 $Na_2C_2O_4 \cdot 10H_2O$
- B. the solution with higher freezing point is 0.1 m glucose

C. 0.1 m $C_6H_5NH_3Cl$ and 0.1 m NaCl will have the same osmotic pressure

D. 0.1 m glucose solution will have the lowest osmotic pressure

Answer: a,b,c,d

 [Watch Video Solution](#)

169. Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of

Column-I	Column-II
(A) <i>n</i> -hexane + <i>n</i> -heptane	(P) Can be separated by fractional distillation
(B) Acetone + chloroform	(Q) Maximum boiling azeotrope
(C) Chlorobenzene and bromobenzene	(R) Cannot be separated by fractional distillation completely
(D) Ethanol + water	(S) Minimum boiling azeotrope

column-II.

 [Watch Video Solution](#)

170. Write structure of 4-oxo -2-hydroxypentanal



[Watch Video Solution](#)

171. what is iodoform.



[Watch Video Solution](#)

172. What is bromoform.



[Watch Video Solution](#)

173. What is alkanols?



[Watch Video Solution](#)

174. What is the formula of butanoic acid



[Watch Video Solution](#)

175. Write the IUPAC names of (i) $\text{CH}_3\text{CH}(\text{CH}_3)\text{CHO}$

 [Watch Video Solution](#)

176. Write the IUPAC names of (ii) $\text{CH}_3\text{CH}(\text{Br})\text{CH}_2\text{CH}_3$.

 [Watch Video Solution](#)

177. Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Elevation in boiling point will be high if the molal elevation constant of the liquid is high.

STATEMENT - 2 : Elevation in boiling point is a colligative property.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

178. Write the IUPAC names of (III) CH_3CHCl_2

 [Watch Video Solution](#)

179. Statement-1 : The observed molar mass of acetic acid in benzene is more than the normal molar mass of acetic acid.

Statement-2 : Molecules of acetic acid dimerise in benzene due to hydrogen bonding.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

180. STATEMENT - 1 : addition of ethylene glycol to water lowers the freezing point of water, therefore, used as antifreeze substance.

STATEMENT - 2 : Ethylene glycol is soluble in water.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

181. Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Osmotic pressure is a colligative property.

STATEMENT - 2 : Osmotic pressure is developed in a column due to osmosis.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

182. Each question contains STATEMENT-I(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Osmosis involves movement of solvent molecules from lower concentration to higher concentration.

STATEMENT - 2 : Solutions having the same osmotic pressure are called isotonic solutions.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

183. Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Isotonic solutions must have the same molal concentration.

STATEMENT - 2 : Solution which have the same osmotic pressure are known as isotonic solution.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

 [Watch Video Solution](#)

184. STATEMENT - 1 : Isotonic solutions do not show phenomenon of osmosis.

STATEMENT - 2 : Isotonic solutions have same molal concentration at same temperature.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

185. STATEMENT - 1 : When dried fruits and vegetables are placed in water, they slowly get swollen.

STATEMENT - 2 : It happens due to the phenomenon of osmosis.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A

 [Watch Video Solution](#)

186. Statement- Reverse osmosis is used to purify saline water.

Explanation- Solvent molecules pass from concentrate solution to dilute solution through semipermeable membrane if high pressure is applied on solution side.

A. If both the statements are TRUE and STATEMENT-2 is the correct

explanation STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the

correct explanation STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

187. Each question contains STATEMENT-I(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : All solute becomes more soluble in water at higher temperature.

STATEMENT - 2 : Solubility of solute depends upon temperature.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



Watch Video Solution

188. Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Henry's law is always applicable for gases. It
STATEMENT - 2 : Raoult's law is a special case of Henry's law.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



Watch Video Solution

189. Each question contains STATEMENT-1(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Osmotic pressure is a colligative property.

STATEMENT - 2 : Osmotic pressure is developed in a column due to osmosis.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



Watch Video Solution

190. Each question contains STATEMENT-I(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : The molecular mass of acetic acid determined by depression in freezing point method in benzene and water was found to be different.

STATEMENT - 2 : Water is polar and benzene is non-polar.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



191. Each question contains STATEMENT-I(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : If red blood cells were removed from the body and placed in pure water, pressure inside the cell increases.

STATEMENT - 2 : The concentration of the salt content in the cells increases.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



Watch Video Solution

192. Each question contains STATEMENT-I(Assertion) and STATEMENT-2(Reason).the statement carefully and mark the correct answer according to the instruction given below:

STATEMENT - 1 : Osmotic pressure is a colligative property.

STATEMENT - 2 : Osmotic pressure is developed in a column due to osmosis.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation STATEMENT-1
- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



Watch Video Solution

193. The vapour pressure of two pure liquids A and B are 5 and 10 torr respectively. Calculate the total pressure of the solution (in torr) obtained by mixing 2 mole of A and 3 mole of B.

- A. 2 torr
- B. 4 torr
- C. 8 torr
- D. 6 torr

Answer: 8

[Watch Video Solution](#)

194. The vapour pressure of two pure liquids A and B are 50 and 40 torr respectively. If 8 molrs of A is mixed with x moles of B , then vapour pressure of solution obtained is 48 torr. What is the value of x.



 [Watch Video Solution](#)

195. The vapour pressure of a liquid solution containing A and B is 99 torr.

Calculate mole % of B in vapour phase. (Given :

$$P_{A^\circ} = 100 \rightarrow rr, P_{B^\circ} = 80 \rightarrow rr)$$

 [Watch Video Solution](#)

196. If 30 g a solute of molecular mass 154 is dissolved in 250 g of benzene. What will be the elevation in boiling point of the resulting solution ?

$$\text{(Given : } K_B(C_6H_6) = 2.6 K kg mol^{-1}\text{)}$$

 [Watch Video Solution](#)

197. Calculate elevation in boiling point for 2 molal aqueous solution of glucose.

$$\text{(Given } K_b(H_2O) = 0.5 kg mol^{-1}\text{)}$$



Watch Video Solution

198. Calculate depression of freezing point for 0.56 molal aq. Solution of KCl.

(Given : $K_f(H_2O) = 1.8 \text{ kg mol}^{-1}$).



Watch Video Solution

199. for an aqueous solution of $K_4[Fe(CN)_6]$, the value of van't Hoff factor, i is 5 (approx).



Watch Video Solution

200. A solution containing 500 g of a protein per liter is isotonic with a solution containing 3.42 g sucrose per liter. The molecular mass of protein is 5×10^x , hence x is.



Watch Video Solution

201. An aqueous solution of urea has a freezing point of -0.515°C .

Predict the osmotic pressure (in atm) of the same solution at 37°C .

 [Watch Video Solution](#)

202. for an aqueous solution of $\text{K}^4[\text{Fe}(\text{CN})_6]$, the value of van't Hoff factor, i is 5 (approx).

 [Watch Video Solution](#)

Level 1 Q 62 To Q 91

1. If 1m solution of benzoic acid in benzene has a freezing point depression of 3.84°C . ($K_f = 5.12^{\circ}\text{Cmol}^{-1}\text{kg}$) and boiling point elevation of 2.53°C ($K_b = 2.53^{\circ}\text{Cmol}^{-1}\text{kg}$), then select the correct statement/s : It brgtstatement I : there are dimer formation when under =going freezing

Statement II : there are no change when undergoing boiling

Statement III : reverse of I and II Itbr. Statement IV : dimer formation in freezing and boiling state

A. I, II

B. II, III

C. III, I

D. only I

Answer: a



[Watch Video Solution](#)

Level 2

1. Write bond line formula for cyclohexanone.



[Watch Video Solution](#)

2. Write structural formula for 5-oxohexanoic acid.

 [Watch Video Solution](#)

3. The total vapour pressure of a 4 mole % solution of NH_3 in water at 293 K is 50.0 torr. The vapour pressure of pure water is 17.0 torr at this temperature . Applying Henry's and Raoult's laws, calculate the total vapour pressure for a 5 mole % solution:

A. 58.25 torr

B. 33 torr

C. 42.1 torr

D. 52.25 torr

Answer: a

 [Watch Video Solution](#)

Level 3 Match The Column

1. Column -I and Column -II contain four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of

Column-I
(A) <i>n</i> -hexane + <i>n</i> -heptane
(B) Acetone + chloroform
(C) Chlorobenzene and bromobenzene
(D) Ethanol + water

Column-II
(P) Can be separated by fractional distillation
(Q) Maximum boiling azeotrope
(R) Cannot be separated by fractional distillation completely
(S) Minimum boiling azeotrope

column-II.



[Watch Video Solution](#)