



CHEMISTRY

BOOKS - NARENDRA AWASTHI

GASEOUS STATE

Exercise

1. Which of the following statements is not correct about the three states of matter, i.e., solid, liquids and gas?


- A. Molecules of a solid possess least energy whereas those of a gas possess highest energy
- B. The density of solids is highest whereas that of gases is lowest
- C. Gases and liquids possess definite volumes
- D. Molecules of a solid possess vibratory motion

Answer: c




Watch Video Solution


2. Which of the following plots does not represent Boyle's law?




A.



B.



C.



D.

Answer: c



Watch Video Solution

3. A certain sample of gas has a volume of 0.2 litre measured at 1 atm pressure and 0°C . At the same pressure but at 273°C , its volume will be

A. 0.4 litre

B. 0.8 litre

C. 27.8 litres


D. 55.6 litres

Answer: a




Watch Video Solution


4. Among the following curves, which is not according to Charle's law ?




A.



B.



C.



D.

Answer: d

5. Initial temperature of an ideal gas is $75^{\circ}C$. At what temperature, the sample of neon gas would be heated to double its pressure, if the initial volume of gas is reduced by 15%?

A. $319^{\circ}C$


B. $592^{\circ}C$

C. $128^{\circ}C$


D. $60^{\circ}C$

Answer: a


6. Which is correct curve for Charle's law, when the curve is plotted at 0.821 atm pressure for 10 mole ideal gas?




A.



B.



C.



D.

Answer: b



Watch Video Solution

7. The temperature of an ideal gas increases in an:


- A. increase in the average molecular speed
- B. decrease in rate of collision amongst molecules
- C. increase in molecular attraction
- D. decrease in mean free path

Answer: a




Watch Video Solution


8. Which of the following curve does not represent Gay-lusacc's law?




A.



B.



C.



D.

Answer: d



Watch Video Solution

9. Three flasks of equal volumes contain CH_4 , CO_2 , and Cl_2 gases respectively. They will contain equal number of molecules if :

A. the mass of all the gasses is same

B. the mass of all the gas is same but temperature is different

C. temperature and pressure of all the flasks are same

D. temperature, pressure and masses same in the flasks

Answer: c

 [Watch Video Solution](#)

10. Write the formula and identify the functional group , if any : N-methyl ethanamine

 [Watch Video Solution](#)

11. Equal volumes of gases at the same temperature and pressure contain equal number of particles. This statement is a direct consequence of

A. Ideal gas equation

B. Charle's law

C. Dalton's law of partial pressure

D. Avogadro's law

Answer: a



Watch Video Solution

12. A 2.24L cylinder of oxygen at 1 atm and 273 K is found to develop a leakage. When the leakage was plugged the pressure dropped to 570 mm of Hg. The number of moles of gas that escaped will be :

A. 0.025

B. 0.05


C. 0.075


D. 0.09

Answer: a


 [Watch Video Solution](#)

13. Which of the following curve is correct for an ideal gas ?






B.



C.



D.

Answer: c



Watch Video Solution

14. The value of universal gas constant R depends on :

A. $8.314 \text{ JK}^{-1} \text{ mol}^{-1}$

B. $0.0821 \text{ atm L mol}^{-1} \text{K}^{-1}$

C. $0.8314 \text{ bar L mol}^{-1} \text{K}^{-1}$

D. $2 \text{ cal mol}^{-1} \text{K}^{-1}$

Answer: c



Watch Video Solution

15. At 0°C and one atm pressure, a gas occupies 100 cc. If the pressure is increased to one and a half-time and temperature is increased by one-third of absolute temperature, then final volume of the gas will be:

A. 80 cc

B. 88.9 cc

C. 66.7 cc

D. 100 cc

Answer: b



Watch Video Solution

16. 10 g of a gas at 1 atm and 273 K occupies 5 litres.

The temperature at which the volume becomes double

for the same mass of gas at the same pressure is:

A. 273 K

B. $-273^{\circ}C$



C. $273^{\circ}C$


D. $546^{\circ}C$

Answer: c


 Watch Video Solution

17. Which of the following curve does not represent Gay-lusacc's law?





C.



D.

Answer: a



Watch Video Solution

18. Densities of two gases are in the ratio 1:2 and their temperatures are in the ratio 2:1, then the ratio of their respective pressure is

A. 1:1

B. 1 : 2

C. 2 : 1

D. 4 : 1

Answer: a



Watch Video Solution

19. Two separate bulbs contain ideal gas A and B . The density of a gas A is twice that of a gas B . The molecular mass of A is half that of gas B . The two gases are at the same temperature. The ratio of the pressure of A to that gas B is

A. 2

B. $1/2$

C. 4

D. $1/4$

Answer: c



Watch Video Solution

20. Volume of the air that will be expelled from a vessel of 300cm^3 when it is heated from 27°C to 37°C at the same pressure will be

A. 310 cm^3

B. 290 cm^3

C. 10 cm^3

D. 37 cm^3


Answer: c



Watch Video Solution

21. For an ideal gas $V - T$ curves at constant pressure

P_1 & P_2 are shown in figure, from the figure



A. $P_1 > P_2$

B. $P_1 < P_2$

C. $P_1 = P_2$

D. All of these

Answer: b



Watch Video Solution

22. Two flasks A and B have equal volumes. A is maintained at 300 K and B at 600 K. A contains H_2 gas, B has an equal mass of CO_2 gas. Find the ratio of total K.E. of gases in flask A to that of B.

A. the same number of atoms

B. the same number of molecules

C. more number of moles of molecules in flask A as compared to flask B


D. the same amount of gases

Answer: b



Watch Video Solution

23. 2.8 g of a gas at 1atm and 273K occupies a volume of 2.24 litres. The gas can not be:



Answer: a



Watch Video Solution

24. Five grams each of the following gases at 87°C and 750 mm pressure are taken. Which of them will have the least volume ?

A. HF

B. HCL

C. HBr

D. HI

Answer: d



Watch Video Solution

25. At what pressure a quantity of gas will occupy a volume of 60 mL, if it occupies a volume of 100mL at a pressure of 720 mm (while temperature is constant) :

A. 700 mm

B. 800 mm

C. 100 mm

D. 1200 mm

Answer: d



Watch Video Solution

26. 20 g of ideal gas contains only atoms of S and O occupies 5.6 L at 1 atm and 273 K. What is the molecular mass of gas?

A. 44.8 g/L

B. 11.4 g/L

C. 2 g/L

D. 3 g/L

Answer: c



Watch Video Solution

27. A small bubble rises from the bottom of a lake, where the temperature and pressure are 8°C and 6.0atm , to the water's surface, where the temperature is 25°C and pressure is 1.0atm . Calculate the final volume of the bubble if its initial volume was 2mL .

- A. 14 mL
- B. 12.72 mL
- C. 11.31 mL
- D. 15 mL

Answer: b



Watch Video Solution

28. Argon is an inert gas used in light bulbs to retard the vaporization of the filament. A certain light-bulb containing argon at 1.25 atm and 18°C is heated to 85°C at constant volume. Calculate its final pressure.

A. 1.53 atm

B. 1.25 atm

C. 1.35 atm


D. 2 atm

Answer: a



Watch Video Solution

29. Calculate the value of O_2 at 1 atm and 273 K required for the complete combustion of 2.64 L of acetylene (C_2H_2) at 1 atm and 273 K.



A. 3.6 L

B. 1.056 L

C. 6.6 L

D. 10 L

Answer: c



Watch Video Solution

30. The density of $O_2(g)$ is maximum at :

A. STP

B. 273 K and 2 atm

C. 546 K and 1 atm

D. 546 K and 2 atm

Answer: b



Watch Video Solution

31. At $27^\circ C$ a sample of ammonia gas exerts a pressure of 5.3 atm. What is the pressure when the

volume of the gas is reduced to one-tenth of the original value at the same temperature ?

A. 0.53 atm

B. 5.3atm

C. 53 atm

D. None of these

Answer: c



Watch Video Solution

32. A certain amount of gas at $2.5^{\circ}C$ and at a pressure of 0.80 atm is kept in a glass vessel. Suppose that the

vessel can withstand a pressure of 2.0 atm. How high can you raise the temperature of the gas without bursting the vessel?

A. $745^{\circ}C$

B. $472^{\circ}C$

C. $500^{\circ}C$


D. None of these

Answer: b




Watch Video Solution


33. Which one of these graphs for an ideal gas having a fixed amount, the arrow indication, is incorrectly marked ?




A.



B.



C.



D.

Answer: b



Watch Video Solution

34. The pressure of sodium vapour in a 1.0 L container is 10 torr at 1000°C . How many atoms are in the container?

A. 9.7×10^{17}

B. 7.6×10^{19}

C. 4.2×10^{17}

D. 9.7×10^{19}

Answer: b



Watch Video Solution

35. An ideal gaseous mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 28 litre at 1 atm and 273 K. The mixture reacts completely with 128 g O_2 to produce CO_2 and H_2O . Mole fraction of C_2H_6 in the mixture is ,

A. 0.6

B. 0.4

C. 0.5

D. 0.8

Answer: a



Watch Video Solution

36. Give reason : a gas exerts pressure on the walls of the container.

A. 2

B. 5

C. 6

D. 7

Answer: d



Watch Video Solution

37. Air entering the lungs ends up in tiny sacs called alveoli. From the alveoli, the oxygen diffuses into the blood. The average radius of the alveoli is 0.0050 cm and the air inside contains 14 per cent oxygen. Assuming that the pressure in the alveoli is 1.0 atm and the temperature is 37°C , calculate the number of oxygen molecules in one of the alveoli.

A. 6×10^{13}

B. 10^{24}

C. 1.7×10^{22}

D. 1.7×10^{12}

Answer: d

38. Starting out on a trip into the mountains, you inflate the tires on your automobile to a recommended pressure of 3.21×10^5 Pa on a day when the temperature is -5.0°C . You drive to the beach, where the temperature is 28.0°C . Assume that the volume of the tire has increased by 3%. What is the final pressure in the tyres?

A. 350 Pa

B. 3500 Pa

C. 3.5×10^5 Pa

D. None of these

Answer: c



Watch Video Solution

39. Temperature of 1 mole of a gas is increased by $2^{\circ}C$ at constant pressure. Work done is :

A. 1264 L

B. 126 L

C. 12600 L

D. 45 L

Answer: a



Watch Video Solution

40. A high-altitude balloon is filled with $1.41 \times 10^4 L$ of hydrogen at a temperature of $21^\circ C$ and a pressure of 745 torr. What is the volume of the balloon at a height of 20 km, where the temperature is $-48^\circ C$ and the pressure is 63.1 torr?

A. $1.274 \times 10^5 L$

B. $1.66 \times 10^5 L$

C. $1.66 \times 10^4 L$

D. None of these

Answer: a



Watch Video Solution

41. The atmospheric pressure on Mars is 0.61 kPa. What is the pressure in mm Hg?

A. 0.63

B. 4.6

C. 6.3

D. 3.2

Answer: b



Watch Video Solution

42. Mercury has an angle of contact equal to 140° with soda lime glass. A narrow tube of radius 1.00 mm made of this glass is dipped in a trough containing mercury. By what amount does the mercury dip down in the tube relative to the liquid surface outside ? Surface tension of mercury at the temperature of the experiment is 0.465 Nm^{-1} . Density of mercury = $13.6 \times 10^3 \text{ kgm}^{-3}$.

A. 322

B. 285

C. 165

D. 210

Answer: c



[Watch Video Solution](#)

43. Write the structural formulae for : 3-ethylpent-2-ene



[Watch Video Solution](#)

44. A balloon contains 14.0 L of air at 760 torr. What will be the volume of the balloon when it is taken to a depth of 10 ft. in a swimming pool? Assume that the

temperature of the air and water are equal. (density :

Hg=13.6g/mL.)

A. 11

B. 11.3

C. 10

D. 10.8

Answer: d

 [Watch Video Solution](#)

45. Write the structural formulae for : cyclohexene

 [Watch Video Solution](#)

46. Equal volumes of oxygen gas and a second gas weigh 1.00 and 2.375 grams respectively under the same experimental conditions. Which of the following is the unknown gas?

A. NO

B. SO_2

C. CS_2

D. CO

Answer: c



Watch Video Solution

47. A high altitude balloon contains 6.81 g of helium in 1.16×10^4 L at -23°C . Assuming ideal gas behaviour, how many grams of helium would have to be added to increase the pressure to 4.0×10^{-3} atm?

A. 1.27 g

B. 1.58 g

C. 2.68 g

D. 2.23 g

Answer: d



Watch Video Solution

48. A 4.40 g piece of solid CO_2 (dry ice) is allowed to sublime in a balloon. The final volume of the balloon is 1.00 L at 300 K. What is the pressure (atm) of the gas?

A. 0.122

B. 2.46

C. 122

D. 24.6

Answer: b



Watch Video Solution

49. Give the iupac name of : $CH_2=CHCH=CHCH=CH_2$



[Watch Video Solution](#)

50. Give the iupac name of : $\text{CH}_3\text{-C}(\text{CH}_3)=\text{CH-C}(\text{CH}_3)_3$



[Watch Video Solution](#)

51. Give the iupac name of : $(\text{C}_2\text{H}_5)_2\text{C}=\text{C}(\text{C}_2\text{H}_5)_2$



[Watch Video Solution](#)

52. Give the iupac name of : $\text{CH}_3\text{-C}(\text{CH}_3)=\text{C}(\text{CH}_3)\text{-CH}_3$



[Watch Video Solution](#)

53. Give the iupac name of $\text{CH}_3\text{-CH}(\text{CH}_3)\text{-C}(\text{CH}_3)_2\text{-C}=\text{CH}$

 [Watch Video Solution](#)

54. Write the structural formula of the following compound : but-2-en-1-oic acid

 [Watch Video Solution](#)

55. An open flask containing air is heated from 300K to 500K . What percentage of air will be escaped to the atmosphere, if the pressure is kept constant ?

A. 80

B. 40

C. 60

D. 20

Answer:



Watch Video Solution

56. The value of universal gas constant R depends on :

A. temperature of gas

B. volume of gas

C. number of moles of gas


D. units of volume and pressure

Answer: d

 [Watch Video Solution](#)


57. A open ended mercury manometer is used to measure the pressure exerted by a trapped gas as shown in the figure. Initially manometer shows no difference in mercury level in both columns as shown in diagram.

After sparking 'A' dissociates according to following reaction



If pressure of Gas "A" dissociates to 0.8 atm, then

(Assume temperature to be constant and is 300 K)



A. 9 cm Hg

B. 18 cm Hg

C. 27 cm Hg

D. None of these

Answer: a



Watch Video Solution

[Watch Video Solution](#)

58. Write the structural formula of the following compound : 2,4,4- trimethyl-3-isopropylpent-1-ene

 [Watch Video Solution](#)

59. Give the iupac name of : $\text{CH}_3\text{-CH}_2\text{-CH(OH)-CH}_2\text{-CH}_2\text{-CH(CH}_3\text{)-CH}_2\text{-CH}_3$

 [Watch Video Solution](#)

60. Give the iupac name of : $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-C(CH}_2\text{-CH}_3\text{)=CH}_2$



[Watch Video Solution](#)

61. Give structure of cyclohexyl benzene



[Watch Video Solution](#)

62. The following organic compound are popularly known by their common names. write their structural formula and iupac name : neopentane



[Watch Video Solution](#)

63. The following organic compound are popularly known by their common names. write their structural formula and iupac name : acetone

 [Watch Video Solution](#)

64. The following organic compound are popularly known by their common names. write their structural formula and iupac name : glycerol

 [Watch Video Solution](#)

65. The following organic compound are popularly known by their common names. write their structural formula and iupac name : acetylene

 [Watch Video Solution](#)

66. The following organic compound are popularly known by their common names. write their structural formula and iupac name : vinyl chloride

 [Watch Video Solution](#)

67. The following organic compound are popularly known by their common names. write their structural formula and iupac name : tert-amyl alcohol

 [Watch Video Solution](#)

68. The following organic compound are popularly known by their common names. write their structural formula and iupac name : acetonitrile

 [Watch Video Solution](#)

69. The vapour pressure of water at $80^{\circ}C$ is 355 mm of Hg. 1 L vessel contains O_2 at $80^{\circ}C$, which is saturated with water and the total pressure being 760 mm of Hg. The contents of the vessel were pumped into 0.3 L vessel at the same temperature. What is the partial pressure of O_2 ?

A. 1350 Hg

B. 2178.3 Hg

C. 121.5 Hg

D. 355 Hg

Answer: a



Watch Video Solution

70. Which of the following gaseous mixture does not follow Dalton's law of partial pressure?

A. SO_2 and Cl_2

B. CO_2 and N_2

C. CO and CO_2

D. CO and N_2

Answer: a



Watch Video Solution

71. Equal masses of methane and oxygen are mixed in an empty container at $25^{\circ}C$. The fraction of the total pressure exerted by oxygen is:

A. $\frac{2}{3}$

B. $\frac{1}{3} \times \frac{273}{298}$

C. $\frac{1}{3}$

D. $\frac{1}{2}$

Answer: c



Watch Video Solution

72. A box of 1 L capacity is divided into two equal compartments by a thin partition which are filled with 2g H_2 and 16 g CH_4 respectively. The pressure in each compartment is recorded as P atm. The total pressure when partition is removed will be:

A. P

B. 2P

C. $P/2$

D. $P/4$

Answer: a



Watch Video Solution

73. If $10^{-4} dm^3$ of water is introduced into a $1.0 dm^3$ flask to $300K$ how many moles of water are in the vapour phase when equilibrium is established ? (Given vapour pressure of H_2O at $300K$ is $3170 PaR = 8.314 JK^{-1} mol^{-1}$) .

A. 1.27×10^{-3} mole

B. 5.56×10^{-3} mole

C. 1.53×10^{-2} mole

D. 4.46×10^{-2} mole

Answer: a



Watch Video Solution

74. Dalton's law of partial pressures is not applicable to

- A. H_2 and N_2 mixture
- B. H_2 and Cl_2 mixture
- C. H_2 and CO_2 mixture
- D. none of these

Answer: b



Watch Video Solution

75. 56 g of nitrogen and 96 g of oxygen are mixed isothermally and at a total pressure of 10 atm. The partial pressures of oxygen and nitrogen (in atm) are respectively :

A. 4,6

B. 5,5

C. 2,8

D. 6,4

Answer: d



Watch Video Solution

76. The closed containers of the same capacity and at the same temperature are filled with 44 g of H_2 in one and 44 g of CO_2 in the other . If the pressure of carbon dioxide in the second container is 1 atm , then pressure of hydrogen in the first container would be :

A. 1 atm

B. 10 atm

C. 22 atm

D. 44 atm

Answer: c



Watch Video Solution

77. A jar contains a gas and a few drops of water at T_K . The pressure in the jar is 830mm of Hg. The temperature of the jar is reduced by 1% . The vapour pressure of water at two temperatures are 30 and 25 mm of Hg. Calculate the new pressure in the jar.

A. 792 mm of Hg

B. 817 mm of Hg

C. 800 mm of Hg

D. 840 mm of Hg

Answer: b



Watch Video Solution

78. In the equilibrium $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$, the partial pressure of SO_2 , O_2 and SO_3 are 0.662, 0.10 and 0.331 atm respectively . What should be the partial pressure of Oxygen so that the equilibrium concentrations of SO_3 are equal ?

A. 0.36

B. 0.036

C. 3.6

D. 36

Answer: b



Watch Video Solution

79. A gaseous mixture contains three gaseous A , B and C with a total number of moles of 10 and total pressure of 10atm . The partial pressure of A and B are 3atm and 1atm respectively and if C has molecular weight of 2g/mol . Then, the weight of C present in the mixture will be :

A. 8 g

B. 12 g

C. 3 g

D. 6 g

Answer: b



Watch Video Solution

80. A rigid container contains 5 mole H_2 gas at some pressure and temperature. The gas has been allowed to escape by simple process from the container due to which pressure of the gas becomes half of its initial pressure and temperature become $(2/3)^{rd}$ of its initial. The mass of gas remaining is :

A. 7.5 g

B. 1.5 g

C. 2.5 g

D. 3.5 g

Answer: a



Watch Video Solution

81. Pressure of 1 g ideal gas X at 300 K is 2 atm. When 2 g of another gas Y is introduced in the same vessel at same temperature, the pressure become 3 atm. The correct relationship between molar mass of X and Y is :

A. $M_Y = 2M_X$

B. $M_Y = 4M_X$

C. $M_X = 4M_Y$

D. None of these

Answer: b



Watch Video Solution

82. Dry ice is solid carbon dioxide. A 0.050 g sample of dry ice is placed in an evacuated 4.6 L vessel at 30°C . Calculate the pressure inside the vessel after all the dry has been converted to CO_2 gas.

A. 6.14 atm

B. 0.614 atm

C. 0.0614 atm

D. 6.14×10^{-3} atm

Answer: d



Watch Video Solution

83. A mixture of helium of neon gases is collected over water at 28.0°C and 745 mmHg. If the partial pressure of helium is 368 mmHg, what is the partial pressure of neon?

A. 348.7 mmHg


B. 377 mmHg

C. 384.7 mmHg

D. none of these

Answer: a

 Watch Video Solution



Consider the following apparatus. Calculate the partial pressure of He after opening the valve. The temperature is remain constant at $16^{\circ}C$

A. 0.164 atm

B. 1.64 atm

C. 0.328 atm

D. 1 atm

Answer: a



Watch Video Solution

85. Oxygen gas generated by the decomposition of potassium chlorate is collected. The volume of oxygen collected at $24^{\circ}C$ and atmospheric pressure of 760mmHg is 128mL . Calculate the mass (in grams) of oxygen gas obtained. The pressure of water vapour at $24^{\circ}C$ is 22.4mmHg .

A. 1.36 g

B. 1.52 g

C. 0.163 g

D. 1.63 g

Answer: c



Watch Video Solution

86. The quantity $\frac{PV}{k_B T}$ represents the (k_B : Boltzmann constant)

A. number of particles of the gas

B. mass of the gas

C. number of moles of the gas

D. translation energy of the gas

Answer: a



Watch Video Solution

87. Which of the following statements about kinetic energy (K.E.) is true?

A. All objects moving with the same velocity have the same K.E.

B. The K.E. of a body will quadruple if its velocity doubles

C. As the velocity of a body increases, its K.E. decreases

D. The K.E. of a body is independent of its mass

Answer: b

 [Watch Video Solution](#)

88. At what temperature most probable speed of SO_2 molecule have the same value as root mean square speed of O_2 molecules at 300 K?

A. II, IV, V

B. I, III, V

C. I, II, III

D. I, II

Answer: b



Watch Video Solution

89. Which of the following is NOT a postulate of the kinetic molecular theory of gases?

- A. The gas molecules possess a volume that is negligibly small compared to the container
- B. The pressure and volume of a gas are inversely related

C. Gases consist of discrete particles that are in constant chaotic motion

D. The average kinetic energy of the molecules is directly proportional to the absolute temperature

Answer: B

 [Watch Video Solution](#)

90. Which one of the following relationships when graphed does not give a straight line for helium gas?

I. K.E. and T at constant pressure and volume

II. P v/s V at constant temperature for a constant mass

III. V v/s $1/T$ at constant pressure for a constant mass

A. II

B. II and III

C. III

D. I

Answer: b



Watch Video Solution

91. Consider Three one -litre flasks labeled A,B and C filled with the gases NO , NO_2 , and N_2O ,

respectively, each at 1 atm and 273 K. In which flask do the molecules have the highest average kinetic energy?

A. Flask C

B. All are the same

C. Flask A

D. None

Answer: b



Watch Video Solution

92. Which of the following statements is false?

- A. The product of pressure and volume of fixed amount of a gas is independent of temperature.
- B. Molecules of different gasses have the same kinetic energy at a given temperature.
- C. The gas equation is not valid at high pressure and low temperature.
- D. The gas constant per molecule is known as Boltzmann's constant.

Answer: a



Watch Video Solution

93. Which is not correct in terms of kinetic theory of gases?

- A. Gaseous particles are considered as point mass.
- B. The gaseous molecules are in random motion.
- C. When gaseous molecules collide, they lose energy.
- D. When the gas is heated, the molecules moves faster.

Answer: c



Watch Video Solution

94. Two flasks A and B have equal volumes. A is maintained at 300 K and B at 600 K. A contains H_2 gas, B has an equal mass of CO_2 gas. Find the ratio of total K.E. of gases in flask A to that of B.

A. 1 : 2

B. 11 : 1

C. 33 : 2

D. 55 : 7

Answer: b



Watch Video Solution

95. Kinetic energy and pressure of a gas of unit volume are related as:

A. $P = \frac{2}{3}E$

B. $P = \frac{3}{2}E$

C. $P = \frac{E}{2}$

D. $P=2E$

Answer: a



Watch Video Solution

96. Two flask A and B of equal volumes maintained at temperature $300K$ and $700K$ contain equal mass of

$He(g)$ and $N_2(g)$ respectively. What is the ratio of total translational kinetic energy of gas in flask A to that of flask B ?

A. 1 : 3

B. 3 : 1

C. 3 : 49

D. None of these

Answer: b



Watch Video Solution

97. Which of the following change is observed occurs when a substance X is converted from liquid to vapour phase at the standard boiling point?

I. Potential energy of the system decreases

II. The distance between molecules increases

III. The average kinetic energy of the molecules in both phases are equal

A. I only

B. II only

C. III only

D. II and III only

Answer: d



Watch Video Solution

98. A mixture of Ne and Ar kept in a closed vessel at 250 K has a total K.E.=3 kJ. The total mass of Ne and Ar is 30 g. Find mass % of Ne in gaseous mixture at 250 K.

A. 61.63

B. 38.37

C. 0.5

D. 28.3

Answer: d



Watch Video Solution

99. In two vessels of 1 litre each at the same temperature 1g of H_2 and 1g of CH_4 are taken. For these gases:

- A. V_{rms} values will be same
- B. Kinetic energy per mol will be same
- C. Total kinetic energy will same
- D. Pressure will be same

Answer: b



Watch Video Solution

100. Four particles have speed 2, 3, 4 and 5 cm/s respectively Their *RMS* speed is .

A. 3.5 cm/s

B. $(27/2)$ cm/s

C. $\sqrt{54}$ cm/s

D. $\sqrt{54}/2$ cm/s

Answer: d



Watch Video Solution

101. A gaseous mixture contains 4 molecules with a velocity of 6 cm sec^{-1} , 5 molecules with a velocity of 2 cm sec^{-1} and 10 molecules with a velocity of 3 cm sec^{-1} . What is the rms speed of the gas:

A. 2.5 cm sec^{-1}

B. 1.9 cm sec^{-1}

C. 3.6 cm sec^{-1}

D. 4.6 cm sec^{-1}

Answer: c



Watch Video Solution

102. The ratio between the root mean square speed of H_2 at $50K$ and that of O_2 at $800K$ is

A. 4

B. 2

C. 1

D. $1/4$

Answer: c



Watch Video Solution

103. If C_1, C_2, C_3, \dots represent the speeds on n_1, n_2, n_3, \dots molecules, then the root mean square

speed is

$$\text{A. } \sqrt{\frac{n_1 C_1^2 + n_2 C_2^2 + n_3 C_3^2 + \dots}{n_1 + n_2 + n_3 + \dots}}$$

$$\text{B. } \sqrt{\frac{(n_1 + n_2 + n_3 + \dots)^2}{n_1 C_1^2 + n_2 C_2^2 + n_3 C_3^2 + \dots}}$$

$$\text{C. } \sqrt{\frac{(n_1 C_1)}{n_1} + \frac{(n_2 C_2)}{n_2} + \frac{(n_3 C_3)}{n_3}}$$

$$\text{D. } \sqrt{\frac{(n_1 C_1 + n_2 C_2 + n_2 C_3 + \dots)^2}{n_1 + n_2 + n_3 + \dots}}$$

Answer: a



Watch Video Solution

104. The root mean square speed of hydrogen is $\sqrt{5}$ times than that of nitrogen. If T is the temperature of

the gas, then :

A. $T_{H_2} = T_{N_2}$

B. $T_{H_2} > T_{N_2}$

C. $T_{H_2} < T_{N_2}$

D. $T_{H_2} = \sqrt{7}T_{N_2}$


Answer: c



Watch Video Solution

105. At a definite temperature (T), the distribution of speeds is given by the curve. In the curve points A, B

and C indicates the speeds corresponding to :



A. most probable, average and root mean square speeds

B. average, root mean square and most probable speeds

C. root mean square, average and most probable speeds

D. most probable, root mean square and average speeds

Answer: a



Watch Video Solution

106. Find an expression for the rotational kinetic energy of a body.

A. 480 J

B. 240 J

C. 120 J

D. none of these

Answer: b



Watch Video Solution

107. At what temperature will most probable speed of the molecules of the second member of homologous series C_nH_{2n-2} be the same as that of SO_2 at $527^\circ C$.

A. 259.4 K

B. 400 K

C. 532.4 K

D. None of these

Answer: c



Watch Video Solution

108. Addition of 0.643g of a compound to 43.95g of benzene lowers the freezing point from 5.51°C to 5.03°C . If K_f for benzene is 5.12K kg mol^{-1} , calculate the molar mass of the compound.

- A. 0.02 kg/mol
- B. 0.001 kg/mol
- C. 0.003 kg/mol
- D. 1 kg/mol

Answer: d



Watch Video Solution

109. 6×10^{22} gas molecules each of mass 10^{-24} kg are taken in a vessel of 10 litre. What is the pressure exerted by gas molecules ? The root mean square speed of gas molecules is 100 m/s .

A. 20 Pa

B. $2 \times 10^4 \text{ Pa}$

C. $2 \times 10^5 \text{ Pa}$

D. $2 \times 10^7 \text{ Pa}$

Answer: b



110. At what temperature will the speed of sound in air become double of its value of $0^{\circ} C$?

A. $500^{\circ} C$

B. $727^{\circ} C$

C. $227^{\circ} C$

D. None of these

Answer: c



Watch Video Solution

111. The most probable speed of 8 g of H_2 is 200ms^{-1} .

Average kinetic energy (neglect rotational and vibrational energy) of H_2 gas is :

A. 120 J

B. 240 J

C. 360 J

D. None of these

Answer: c



Watch Video Solution

112. Consider the following statement regarding Maxwell's distribution of speeds. The correct statement(s) is/are :

A. $T_1 > T_2 > T_3$

B. $T_1 < T_2 < T_3$

C. $T_1 = T_2 = T_3$

D. None of these

Answer: b



Watch Video Solution

113. The density of a gas filled in electric lamp is 0.75 kg/m^3 . After the lamp has been switched on, the pressure in it increases from $4 \times 10^4 \text{ Pa}$ to $9 \times 10^4 \text{ Pa}$.

What is increases in u_{rms} ?

A. 100

B. 200

C. 300

D. None of these

Answer: b



Watch Video Solution

114. Define root mean square value of an alternating current.

A. d^2

B. d

C. \sqrt{d}

D. $1/\sqrt{d}$

Answer: d



Watch Video Solution

115. The ratio among most probable velocity, mean velocity and root mean velocity is given by

A. $1 : 2 : 3$

B. $1 : \sqrt{2} : \sqrt{3}$

C. $\sqrt{2} : \sqrt{3} : \sqrt{8/\pi}$

D. $\sqrt{2} : \sqrt{8/\pi} : \sqrt{3}$

Answer: d



Watch Video Solution

116. The average speed at temperature $T^\circ C$ of

$CH_4(g)$ is $\sqrt{\frac{28}{88}} \times 10^3 ms^{-1}$. What is the value of T ?

A. $240.55^\circ C$

B. $-32.45^{\circ}C$

C. $3000^{\circ}C$

D. $-24.055^{\circ}C$

Answer: b



Watch Video Solution

117. At what temperature will most probable speed of the molecules of the second member of homologous series C_nH_{2n-2} be the same as that of SO_2 at $527^{\circ}C$.

A. 150 K

B. 600 K

C. 750 K

D. 900 K

Answer: d



Watch Video Solution

118. The *rms* speed of N_2 molecules in a gas is u . If the temperature is doubled and the nitrogen molecules dissociate into nitrogen atoms, the *rms* speed becomes

A. $u/2$

B. $2u$

C. $4u$

D. $14u$

Answer: b



Watch Video Solution

119. The rate of diffusion of a gas is proportional to

A. $\frac{P}{\sqrt{d}}$

B. $\frac{P}{d}$

C. $\sqrt{\frac{P}{d}}$

D. $\frac{\sqrt{P}}{d}$

Answer: a



Watch Video Solution

120. At constant volume and temperature conditions, the rates of diffusion r_A and r_B of gases A and B having densities P_A and P_B are related by the expression :

A. $r_A = r_B \cdot (p_A/p_B)^2$

B. $r_A = r_B(p_A/p_B)^{1/2}$

C. $r_A = (r_B \cdot p_A/p_B)^{1/2}$

D. $r_A = r_B(p_B/p_A)^{1/2}$

Answer: b



Watch Video Solution

121. What is the ratio of diffusion rate of oxygen to hydrogen?

A. 1 : 4

B. 4 : 1

C. 1 : 8

D. 8 : 1

Answer: a



Watch Video Solution

122. The molecular weight of a gas which diffuses through a porous plug at $1/6^{th}$ of the speed of hydrogen under identical condition is:

A. 27

B. 72

C. 36

D. 48

Answer: b



Watch Video Solution

123. $X\text{ mL}$ of H_2 gas effuses through a hole in a container in 5 s . The time taken for the effusion of the same volume of the gas specified below, under identical conditions, is

A. 10 sec : He

B. 20 sec : O_2

C. 25 sec : CO_2

D. 55 sec : CO_2

Answer: b



Watch Video Solution

124. At identical temperature and pressure the rate of diffusion of hydrogen gas is $3\sqrt{3}$ times that of a hydrocarbon having molecular formula C_nH_{2n-2} . What is the value of n ? .

A. 1

B. 4

C. 3

D. 8

Answer: b



Watch Video Solution

125. Calculate relative rate of effusion of O_2 to CH_4 from a container containing O_2 and CH_4 in 3 :2 mass ratio.

A. $\frac{3\sqrt{2}}{4}$

B. $\frac{3}{4\sqrt{2}}$

C. $\frac{3}{2\sqrt{2}}$

D. none of these

Answer: b



Watch Video Solution

126. The following organic compound are popularly known by their common names. write their structural formula and iupac name : allyl alcohol



Watch Video Solution

127. The vapour density of a mixture containing NO_2 and N_2O_4 is 27.6 . The mole fraction of N_2O_4 in the mixture is :

A. 2 : 2 : 3

B. 6 : 6 : 1

C. $\sqrt{2} : \sqrt{2} : 3$

D. 4: 4: 3

Answer: d



Watch Video Solution

128. 80 mL of O_2 takes 2 minutes to pass through the hole. What volume of SO_2 will pass through the hole in 3 minute?

A. $\frac{120}{\sqrt{2}}$ mL

B. $120 \times \sqrt{2}$ mL

C. $\frac{12}{\sqrt{2}}$ mL


D. None of these

Answer: a



Watch Video Solution

129. When two cotton plugs, one moistened with ammonia and the other with hydrochloric acid, are simultaneously inserted into opposite ends of a glass tube 87.0 cm long, a white ring of NH_4Cl forms where gaseous NH_3 and gaseous HCl first come into contact.



At what distance from the ammonia-moistened plug does this occur?

A. 51.7 cm from NH_3 end

B. 51.7 cm from HCl end

C. 43.5 at mid point

D. None of these

Answer: a

 [Watch Video Solution](#)

130. 4 gm of sulphur dioxide gas diffuses from a container in 8 min. Mass of helium gas diffusing from the same container over the same time interval is :

A. 0.5 g

B. 1 g

C. 2 g

D. None of these

Answer: b



Watch Video Solution

131. Under identical conditions of pressure and temperature, 4 L of gaseous mixture (H_2 and CH_4) effuses through a hole in 5 min whereas 4 L of a gas X of molecular mass 36 takes to 10 min to effuse through the same hole. The mole ratio of $H_2 : CH_4$ in the mixture is:

A. 1 : 2

B. 2 : 1

C. 2 : 3

D. 1 : 1

Answer: d



Watch Video Solution

132. A ballon weighing $50kg$ is filled with $685kg$ of helium at $1atm$ pressure and $25^{\circ}C$. What will be its pay load if it displaced $5108kg$ of air ?

A. 4373 kg

B. 4423 kg

C. 5793 kg

D. none of these

Answer: a



Watch Video Solution

133. According to the law of equipartition of energy, the energy associated with each degree of freedom is :

A. $\frac{1}{3}K_B T$

B. $\frac{1}{2}K_B T$

C. $K_B T$

D. $\frac{3}{2}K_B T$

Answer: b



Watch Video Solution

134. Calculate γ (ratio of C_p and C_v) for triatomic linear gas at high temperature. Assume that the contribution of vibrational degree of freedom is 75% :

A. 1.222

B. 1.121

C. 1.18

D. 1.33

Answer: c



Watch Video Solution

135. If one mole of a mono-atomic gas ($\gamma = 5/3$) is mixed with one mole of a diatomic gas ($\gamma = 7/5$), the value of γ for the mixture is :

- A. 1.4
- B. 1.428
- C. 1.5
- D. 1.33

Answer: c



Watch Video Solution

136. If one mole of a monatomic gas $\left(\gamma = \frac{5}{3}\right)$ is mixed with one mole of a diatomic gas $\left(\gamma = \frac{7}{5}\right)$, the value of gamma for mixture is

- A. 1.4
- B. 1.5
- C. 1.53
- D. 3.07

Answer: b



Watch Video Solution

137. Select the correct statement :

- A. Internal energy of a real gas at a given temperature increases as the volume increases
- B. Internal energy of an ideal gas at given temperature increase as the volume increases
- C. Internal energy of an ideal gas molecules is not a function of temperature
- D. The internal energy of a real gas at a constant temperature is independent of change in volume

Answer: a



Watch Video Solution

138. Which gas shows real behaviour?

A. 16 g O_2 at 1 atm & 273 K occupies 11.2 L

B. 1 g H_2 in 0.5 L flask exerts pressure of 24.63 atm
at 300 K

C. 1 mole NH_3 at 300 K and 1 atm occupies volume
22.4 L

D. 5.6 L of CO_2 at 1 atm & 273 K is equal to 11 g

Answer: c



Watch Video Solution

139. Consider an ideal gas contained in a vessel. If the intermolecular interaction suddenly begins to act, which of the following will happen ?

- A. Pressure decreases
- B. Pressure increases
- C. Pressure remains unchanged
- D. Gas collapses

Answer: b



Watch Video Solution

140. The pressure of real gas is less than the pressure of an ideal gas because of

- A. increase in number of collisions
- B. finite size of molecule
- C. increase in KE of molecules
- D. intermolecular forces of attraction

Answer: d



Watch Video Solution

141. A real gas behaves like an ideal gas if its

- A. high pressure and low temperature
- B. low pressure and high temperature
- C. high pressure and high temperature
- D. low pressure and low temperature

Answer: b



Watch Video Solution

142. If temperature and volume are same, the pressure of a gas obeying van der Waal's equation is :

- A. less than that of an ideal gas
- B. more than that of an ideal gas

C. same as that of an ideal gas

D. none of these

Answer: a



Watch Video Solution

143. A and B appear for interview. The probability of their selection is : $\frac{1}{2}$ and $\frac{1}{3}$ respectively. Find the probability that only one of them is selected.

A. $P_1 < P_2 < P_3$

B. $P_2 < P_1 < P_3$

C. $P_2 < P_3 < P_1$

$$D. P_1 = P_2 = P_3$$

Answer: c



Watch Video Solution

144. A gas obeys the equation of state $P(V - b) = RT$ (The parameter b is a constant). The slope for an isochore will be .

A. negative

B. zero

C. $R/(V - b)$

D. R/P

Answer: c

 **Watch Video Solution**

145. van der Waals constant b of helium is 24 mL mol^{-1} . Find molecular diameter of helium.

A. $1.335 \times 10^{-10} \text{ cm}$

B. $1.335 \times 10^{-8} \text{ cm}$

C. $2.67 \times 10^{-8} \text{ cm}$

D. $4.34 \times 10^{-8} \text{ cm}$

Answer: c

 **Watch Video Solution**

146. Which of the following can be must readily liquefied? The given value of a for NH_3 is 4.17, CO_2 is 3.59, SO_2 is 6.71, and Cl_2 is 6.49.

A. NH_3

B. Cl_2

C. SO_2

D. CO_2

Answer: c



Watch Video Solution

147. For which of the following gasses should the correction for the molecular volume be largest :

CO , CO_2 , NH_3 or SF_6 ?

A. CO

B. CO_2

C. NH_3

D. SF_6

Answer: d



Watch Video Solution

148. Under which of the following sets of conditions is a real gas expected to deviate from ideal behaviour?

(I) High pressure, small volume

(II) High temperature , low pressure

(III) Low temperature, high pressure

A. only I

B. only II

C. only III

D. I and III both

Answer: d



Watch Video Solution

149. For a certain gas which deviates a little from ideal behaviour, a plot between P/p vs P was found to be non-linear . The intercept on y-axis will be :

A. $\frac{RT}{M}$

B. $\frac{M}{RT}$

C. $\frac{MZ}{RT}$

D. $\frac{R}{TM}$

Answer: a



Watch Video Solution

150. At low pressure, the van der Waal's equation become :

A. $PV = RT$

B. $P(V - b) = RT$

C. $\left(P + \frac{a}{V^2}\right)V = RT$

D. $P = \frac{RT}{V} + \frac{a}{V^2}$

Answer: c



Watch Video Solution

151. At low pressure, if $RT = 2\sqrt{a.P}$, then the volume occupied by a real gas is :

A. $\frac{2RT}{P}$

B. $\frac{2P}{RT}$

C. $\frac{RT}{2P}$

D. $\frac{2RT}{P}$

Answer: c



Watch Video Solution

152. For a gas deviation from ideal behaviour is maximum at :

A. $0^{\circ}C$ and 1.0 atm

B. $100^{\circ}C$ and 2.0 atm

C. $-13^{\circ}C$ and 1.0 atm

D. $-13^{\circ}C$ and 2.0 atm

Answer: d



Watch Video Solution

153. At low pressures, the van der Waals equation is

$$\text{written as } \left[P + \frac{a}{V^2} \right] V = RT$$

The compressibility factor is then equal to

A. $\left(1 - \frac{a}{RTV} \right)$

B. $\left(1 - \frac{RTV}{a} \right)$

C. $\left(1 + \frac{a}{RTV} \right)$

D. $\left(1 + \frac{RTV}{a} \right)$

Answer: a



Watch Video Solution

154. The compressibility factor for a real gas at high pressure is .

A. 1

B. $1 + \frac{Pb}{RT}$

C. $1 - \frac{Pb}{RT}$

D. $1 + \frac{RT}{Pb}$

Answer: b



Watch Video Solution

155. The compressibility of a gas is greater than unity at 1 atm and 273 K. Therefore :

A. $V_m > 22.4L$

B. $V_m < 22.4L$

C. $V_m = 22.4L$

D. $V_m = 44.8L$

Answer: a



Watch Video Solution

156. At $273K$ temp, and 9 atm pressure, the compressibility fog a gas is 0.9. The volume of 1 mill-moles of gas at this temperature and pressure is :

A. 2.24 litre

B. 0.020 mL

C. 2.24 mL

D. 22.4 mL

Answer: c



Watch Video Solution

157. The compressibility factor for nitrogen at $330K$ and $800atm$ is 1.90 and at $570K$ and $200atm$ is 1.10 . A certain mass of N_2 occupies a volume of $1dm^3$ at $330K$ and $800atm$ calculate volume occupied by same quantity of N_2 gas at $570K$ and $200atm$

A. 1 L

B. 2 L

C. 3 L

D. 4 L

Answer: d



Watch Video Solution

158. Consider the equation $Z = \frac{PV}{RT}$. Which of the following statements is correct?

A. When $Z > 1$, real gasses are easier to compress than the ideal gas

B. When $Z = 1$, real gases get compressed easily

C. When $Z > 1$, real gases are difficult to compress

D. When $Z = 1$, real gases are difficult to compress

Answer: c



Watch Video Solution

159. What is the compressibility factor (Z) for 0.02 mole of a van der Waals's gas at pressure of 0.1 atm.

Assume the size of gas molecules is negligible.

Given : $RT=20 \text{ L atm mol}^{-1}$ and $a=1000 \text{ atm L}^2\text{mol}^{-2}$

A. 2

B. 1

C. 0.02

D. 0.5

Answer: d



Watch Video Solution

160. Consider the following statements If th van der

Waals' parameters of two gases are given as

$$a / dm^6 barmol^{-2} \quad b / dm^3 mol^{-1}$$

Gas A 6.5 0.056

Gas B 18.0 0.011

then which of the following statements is//are correct

?.

A. 1 alone

B. 1 and 2

C. 1, 2 and 3

D. 2 and 3

Answer: c



[View Text Solution](#)

161. The van der Waals parameters for gases W, X, Y and Z are

Gas	$a(\text{atmL}^2\text{mol}^{-2})$	$b(\text{Lmol}^{-1})$
<i>W</i>	4.0	0.027
<i>X</i>	8.0	0.030
<i>Y</i>	6.0	0.032
<i>Z</i>	12.0	0.027

Which one of these gases has the highest critical temperature?

- A. W
- B. X
- C. Y
- D. Z

Answer: d



Watch Video Solution

162. Pressure remaining the constant, the volume of a given mass of an ideal gas increases for every degree centigrade rise in temperature by definite fraction of its volume at:

A. $0^{\circ} C$

B. its critical temperature

C. absolute zero

D. its Boyle temperature

Answer: a



Watch Video Solution

163. Define critical temperature.

- A. the temperature above which the substance undergoes decomposition
- B. the temperature above which a substance can exist only in gaseous state
- C. boiling point of the substance
- D. all are wrong

Answer: b



Watch Video Solution

164. The excluded volume of a gas will be larger, if $\frac{T_C}{P_C}$ is :

A. small

B. large

C. equal to 1

D. less than unity

Answer: b



Watch Video Solution

165. Select incorrect statement :

A. we can condense vapour simply by applying pressure

B. to liquefy a gas one must lower the temperature below T_C and also apply pressure

C. at T_C there is no distinction between liquid and vapour state hence density of the liquid is nearly equal to density of the vapour

D. However great the pressure applied, a gas cannot be liquified below it's critical temp.

Answer: d



Watch Video Solution

166. The correct order of temperature of a real gas is :

(I) Boyle's temperature

(II) Critical temperature

(III) Inversion temperature

A. $III > I > II$

B. $I > II > III$

C. $II > I > III$

D. $I > III > II$

Answer: a



Watch Video Solution

167. The temperature at which the second virial coefficient of a real gas is zero is called .

A. Critical temperature

B. Triple point

C. Boiling point

D. Boyle's temperature

Answer: d



Watch Video Solution

168. The following organic compound are popularly known by their common names. write their structural formula and iupac name : formaldehyde



Watch Video Solution

169. van der Waal equation is true for :

A. both $\left(\frac{\partial P}{\partial V}\right)_T$ and $\left(\frac{\partial^2 P}{\partial V^2}\right)_T$ are zero

B. only $\left(\frac{\partial^2 P}{\partial V^2}\right)_T$ is not zero

C. $\left(\frac{\partial P}{\partial V}\right)_T$ is zero but $\left(\frac{\partial^2 P}{\partial V^2}\right)_T$ is non-zero

D. $\left(\frac{\partial P}{\partial V}\right)_T$ is non-zero but $\left(\frac{\partial^2 P}{\partial V^2}\right)_T$ is zero

Answer: a



Watch Video Solution

170. However great the pressure , a gas cannot be liquified above its :

- A. Boyle temperature
- B. Inversion temperature
- C. Critical temperature
- D. Room temperature

Answer: c



Watch Video Solution

171. Inversion temperature $\left(T_i = \frac{2a}{Rb}\right)$ is defined as the temperature above which if gas is expanded adiabatically it gets warm up but if temperature of gas is lower than T_i then it will cool down. What will happen to gas if it is adiabatically expanded at $50^\circ C$ if its Boyle's temperature is $20^\circ C$

- A. Heating
- B. Cooling
- C. Constant
- D. None

Answer: a

 **Watch Video Solution**

172. The van der Waals' equation of law of corresponding states for 1 mole of gas is :

A. $\left(P_r + \frac{3}{V_r^2}\right)(3V_r - 1) = 8T_r$

B. $\left(P_r - \frac{3}{V_r^2}\right)(3V_r - 1) = 8T_r$

C. $\left(P_r + \frac{3}{V_r^2}\right)(3V_r + 1) = 8\pi T_r$

D. $\left(P_r + \frac{3}{V_r^2}\right)(3V_r + 1) = 8$

Answer: a

 **Watch Video Solution**

173. Calculate the volume occupied by 16gram O_2 at $300K$ and $8.31Mpa$ if $\frac{P_C V_C}{RT_C} = 3/8$ and $\frac{P_r V_r}{T_r} = 2.21$ (Given : $R = 8.314MPa / K - mol$)

A. 125.31 mL

B. 124.25 mL

C. 248.62 mL

D. none of these


Answer: b



Watch Video Solution

174. Consider the composite system, which is held at 300 K, shown in the following figure. Assuming ideal gas behavior, calculate the total pressure if the barriers separating the compartments are removed. Assume that the volume of the barriers is negligible.

(Given : $R = 0.082 \text{ atm L/mol. K}$)



- A. 1 atm
- B. 2 atm
- C. 2.3 atm
- D. 3.2 atm


Answer: b



Watch Video Solution

175. 11 moles of N_2 and 12 moles of H_2 mixture reacted in 20 litre vessel at 800 K. After equilibrium was reached, 6 mole of H_2 was present. 3.58 litre of liquid water is injected in equilibrium mixture and resultant gaseous mixture suddenly cooled to 300K. What is the final pressure of gaseous mixture? Neglect vapour pressure of liquid solution. Assume (i) all NH_3 dissolved in water (ii) no change in volume of liquid (iii) At 300 K no reaction takes place between N_2 and

H_2



A. 18.47 atm

B. 60 atm

C. 22.5 atm

D. 45 atm

Answer: c



Watch Video Solution

176. Two vessels connected by a valve of negligible volume. One container (I) has 2.8 g of N_2 at temperature $T_1(K)$. The other container (II) is completely evacuated. The container (I) is heated to $T_2(K)$ while container (II) is maintained at $T_2/3(K)$. volume of vessel (I) is half that of vessel (II). If the valve is opened then what is the mass ratio of N_2 in both vessel (W_I/W_{II})?

A. 1:2

B. 1:3

C. 1:6



D. 3:1

Answer: c



Watch Video Solution

177. A mixture of $NH_3(g)$ and $N_2H_4(g)$ is placed in a sealed container at $300K$. The total pressure is $0.5atm$. The container is heated to $1200K$, at which time both substances decompose completely according to the equations:



After decomposition is complete, the total pressure at $1200K$ is found to be $4.5atm$. Find the amount (mole) per cent of $N_2H_4(g)$ in the original mixture.

A. 0.2

B. 0.25

C. 0.5

D. 0.75

Answer: b



Watch Video Solution

178. Correct expression for density of an ideal gas mixture of two gases 1 and 2, where m_1 and m_2 are masses and n_1 and n_2 are moles and M_1 and M_2 are molar masses.

$$\text{A. } d = \frac{(m_1 + m_2)}{(M_1 + M_2)}$$

$$\text{B. } d = \frac{(m_1 + m_2)}{(n_1 + n_2)} \frac{P}{RT}$$

$$\text{C. } d = \frac{(n_1 + n_2)}{(m_1 + m_2)} \times \frac{P}{RT}$$

D. None of these

Answer: b



Watch Video Solution

179. Two closed vessels of equal volume containing air at pressure P_1 and temperature T_1 are connected to each other through a narrow tube. If the temperature in one of the vessels is now maintained at T_1 and that

in the other at T_2 , what will be the pressure in the vessels?

A. $\frac{T_1}{2P_1T_2}$

B. $\frac{2P_1T_2}{T_1 + T_2}$

C. $\frac{2P_1T_2}{T_1 - T_2}$

D. $\frac{2P_1}{T_1 + T_2}$

Answer: b



Watch Video Solution

180. The following organic compound are popularly known by their common names. write their structural

formula and iupac name : acetic acid

 [Watch Video Solution](#)

181. The following organic compound are popularly known by their common names. write their structural formula and iupac name : salicylic acid

 [Watch Video Solution](#)

182. What is the density of wet air with 75% relative humidity at 1 atm and 300 K? Given : vapour pressure of H_2O is 30 torr and average molar mass of air is 29 g mol^{-1} .

A. 1.614 g/L

B. 0.96 g/L

C. 1.06 g/L

D. 1.164 g/L

Answer: d



Watch Video Solution

183. 7 moles of a tetra-atomic non-linear gas 'A' at 10 atm and T K are mixed with 6 moles of another gas B at $\frac{T}{3}$ K and 5 atm in a closed, rigid vessel without energy transfer with surroundings. If final temperature

of mixture was $\frac{5T}{6}K$, then gas B is ? (Assuming all modes of energy are active)

A. monoatomic

B. diatomic

C. triatomic

D. tetra atomic

Answer: b




Watch Video Solution

184. Three closed rigid vessels, A, B and C, which initially contain three different gases at different

temperatures are connected by tube of negligible volume, without any energy transfer with surroundings. The vessel A contain 2 mole Ne gas, at 300 K, vessel 'B' contain 2 mole SO_2 gas at 400 K and vessel 'C' contain 3 mole CO_2 gas at temperature 500 K. What is the final pressure (in atm) attained by gases when all valves of connecting three vessels are opened and additional 15.6 kcal heat supplied to vessels through valve. The volume of A, B and C vessel is 2, 2 and 3 litre respectively

Given $R = 2$ calorie/mol-K,

$$C_v(Ne) = 3/2R, C_v(CO) = 5/2R \text{ and } C_v(SO_2) = 3R$$



A. 73.89 atm

B. 67.31 atm

C. 80 atm

D. none of these

Answer: a



Watch Video Solution

185.6×10^{22} gas molecules each of mass 10^{-24} kg are taken in a vessel of 10 litre. What is the pressure exerted by gas molecules ? The root mean square speed of gas molecules is 100 m/s .

A. 298 K

B. 25 K

C. 250 K

D. 2500 K

Answer: c



Watch Video Solution

186. A balloon of diameter 21 meter weight 100 kg.

Calculate its pay-load, if it is filled with He at 1.0 atm

and $27^{\circ}C$. Density of air is $1.2 \text{ kg } m^{-3}$. (Given :

$R=0.082 \text{ L atm } K^{-1}mol^{-1}$)

A. 4952.42 kg

B. 4932.42 kg

C. 493.242 kg

D. none of these

Answer: b

 [Watch Video Solution](#)

187. A given volume of ozonised oxygen (containing 60% oxygen by volume) required 220 sec to effuse while an equal volume of oxygen took 200 sec only under identical conditions. If density of O_2 is 1.6 g/L then find density of O_3 .

A. 1.963 g/L

B. 2.16 g/L

C. 3.28 g/L

D. 2.24 g/L

Answer: d

 [Watch Video Solution](#)

188. The following organic compound are popularly known by their common names. write their structural formula and iupac name : styrene

 [Watch Video Solution](#)

189. The following organic compound are popularly known by their common names. write their structural formula and iupac name : resorcinol



Watch Video Solution

190. At low pressure, if $RT = 2\sqrt{a.P}$, then the volume occupied by a real gas is :

A. 443.34 L

B. 246.3 L

C. 12.315 L


D. 24.63 L

Answer: d




Watch Video Solution

191. At room temperature following reaction goes to completion



AB_2 is solid with negligible vapour pressure below $0^\circ C$. At 300 K, the AB in the smaller flask exerts a pressure of 3 atm and in the larger flask B_2 exerts a pressure of 1 atm at 400 K when they are separated out by a close valve, The gases are mixed by opening

the stop cock and after the end of the reaction the flask are cooled to 250 K



The final pressure is :

- A. 0.156 atm
- B. 0.3125 atm
- C. 0.625 atm
- D. 3.2 atm

Answer: c




Watch Video Solution

192. A vessel of uniform cross-section of length 500 cm as shown in figure is divided in two parts by a weightless and frictionless piston one part contains 5 moles of $\text{He}(g)$ and other part 2 moles of $\text{H}_2(g)$ and 4 moles of $\text{O}_2(g)$ added at the same temperature and pressure in which reaction takes place finally vessel cooled to 300 K and 1 atm. What is the length of He compartment?

(Assume volume of piston and vol. of $\text{H}_2\text{O}(l)$ formed

are

negligible)



At initial stage

A. 187.5 cm

B. 300 cm

C. 312.5 cm

D. none of these

Answer: c



Watch Video Solution

193. For a real gas (mol.mass =60) if density at critical point is $0.80g/cm^{-3}$ and its $T_c = \frac{4 \times 10^5}{821}K$, then van der Waals' constant a (in atm L^2mol^{-2}) is

A. 0.3375

B. 3.375

C. 1.68

D. 0.025

Answer: b



Watch Video Solution

194. The van der Waals' constant 'b' of a gas is $4\pi \times 10^{-4} L/mol$. How near can the centres of the two molecules approach each other? [Use : $N_A = 6 \times 10^{23}$]

A. 10^{-7} m

B. 10^{-10} m

C. $5 \times 10^{-11} \text{ m}$

D. $5 \times 10^{-9} \text{ m}$

Answer: b



Watch Video Solution

195. The following organic compound are popularly known by their common names. write their structural formula and iupac name : o-cresol

 [Watch Video Solution](#)

196. The following organic compound are popularly known by their common names. write their structural formula and iupac name : catechol

 [Watch Video Solution](#)

197. The following organic compound are popularly known by their common names. write their structural

formula and iupac name : salicylaldehyde

 [Watch Video Solution](#)

198. The following organic compound are popularly known by their common names. write their structural formula and iupac name : pyrogallol

 [Watch Video Solution](#)

199. At a constant temperature what should be the percentage increase in pressure for a 10% decrease in the volume of gas ? .

A. 11.1 %

B. 20 %

C. 10 %

D. 50 %

Answer: a

 [Watch Video Solution](#)

200. The following organic compound are popularly known by their common names. write their structural formula and iupac name : quinol

 [Watch Video Solution](#)

201. Match the items of columns I and II.

Column-I		Column-II	
(P)	Z for ideal gas behaviour	(1)	$3/8$
(Q)	Z for real gas at low pressure	(2)	$\left(1 + \frac{Pb}{RT}\right)$
(R)	Z for real gas at high pressure	(3)	1
(S)	Z for critical state	(4)	$\left(1 - \frac{a}{RTV}\right)$

A. $P \quad Q \quad R \quad S$
1 2 4 3

B. $P \quad Q \quad R \quad S$
3 4 2 1

C. $P \quad Q \quad R \quad S$
2 1 4 3

D. $P \quad Q \quad R \quad S$
1 2 3 4

Answer: b



Watch Video Solution

202. The following organic compound are popularly known by their common names. write their structural formula and iupac name : benzyl alcohol

 **Watch Video Solution**

203. van der Waal's equation for calculating the pressure of a non ideal gas is

$$\left(P + \frac{an^2}{V^2}\right)(V - nb) = nRT$$

van der Waal's suggested that the pressure exerted by an ideal gas , P_{ideal} , is related to the experiventally measured pressure, P_{ideal} by the equation

$$P_{\text{ideal}} = \underset{\substack{\uparrow \\ \text{observed pressure}}}{P_{\text{real}}} + \underset{\substack{\uparrow \\ \text{currection term}}}{\frac{an^2}{V^2}}$$

Constant 'a' is measure of intermolecular interaction between gaseous molecules that gives rise to nonideal behavior. It depends upon how frequently any two molecules approach each other closely. Another correction concerns the volume occupied by the gas molecules. In the ideal gas equation, V represents the volume of the container. However, each molecule does occupy a finite, although small, intrinsic volume, so the effective volume of the gas becomes $(V-nb)$, where n is the number of moles of the gas and b is a constant. The term nb represents the volume occupied by gas particles present in n moles of the gas .

Having taken into account the corrections for pressure and volume, we can rewrite the ideal gas equation as follows :

$$\left(P + \frac{an^2}{V^2} \right) (V - nb) = nRT$$

corrected pressure corrected volume

Using van der Waals' equation, find the constant 'a' (in $\text{atm L}^2\text{mol}^{-2}$) when two moles of a gas confined in 4 L flask exerts a pressure of 11.0 atmospheres at a temperature of 300 K. The value of b is 0.05 L mol^{-1} .
($R = 0.082 \text{ atm.L/K mol}$)

A. $PV = RT - a/V_m$

B. $PV = aRT/V_m^2$

C. $P = RT - a/V_m^2$

D. $PV_m = RT + Pb$

Answer: d



Watch Video Solution

204. For non-zero value of force of attraction between gas molecules at large volume, gas equation will be :

A. $PV = nRT - \frac{n^2a}{V}$

B. $PV = nRT + nbP$

C. $P = \frac{nRT}{V - b}$

D. $PV = nRT$

Answer: a



Watch Video Solution

205. The following organic compound are popularly known by their common names. write their structural formula and iupac name : phenyl acetaldehyde

 **Watch Video Solution**

206. The van der Waals' constant 'b' of a gas is $4\pi \times 10^{-4} L/mol$. How near can the centres of the two molecules approach each other? [Use : $N_A = 6 \times 10^{23}$]

A. 2.62

B. 2.64

C. 6.24

D. 6.46

Answer: d

 [Watch Video Solution](#)


207. Write the formula and identify the functional group, if any : 4-hydroxy-3-methoxybenzaldehyde

 [Watch Video Solution](#)

208. Write the structure of benzyl chloride

 [Watch Video Solution](#)

209. A monometer contains a liquid of density 5.44g/cm^3 is attached to a flask containing gas 'A' as follows



If the same liquid is used in barometer to measure the atmospheric pressure, then what will be the length of

the liquid column, which exerts pressure equal to 1 atm ? (density of Hg = 13.6 g / cm^3)

A. 190 cm

B. 76 cm

C. 30.4 cm

D. 266 cm

Answer:




[Watch Video Solution](#)

210. Write the structure of benzal chloride



[Watch Video Solution](#)

211. A monometer contains a liquid of density 5.44g/cm^3 is attached to a flask containing gas 'A' as follows



If the same liquid is used in barometer to measure the atmospheric pressure, then what will be the length of the liquid column which exerts pressure equal to 1 atm ? (density of $Hg = 13.6\text{g/cm}^3$)

A. 190 cm

B. 76 cm

C. 30.4 cm

D. 266 cm

Answer:

 [Watch Video Solution](#)

212. Write the structure of benzochloride


 [Watch Video Solution](#)

213. Write structure of phthalic acid




Watch Video Solution


214. Which of the following curves represent(s) Boyle's law?




A.



B.



C.



D.

Answer: a,b,c,d



Watch Video Solution

215. If a gas expands at constant temperature

A. the pressure decreases

B. the kinetic energy of the molecules remains the same

C. the K.E. of the molecules decrease

D. the number of molecules of the gas increase

Answer: a,b

 [Watch Video Solution](#)

216. Which of the following statements are correct ?

A. It is not possible to compress a gas at a temperature below T_C

B. At a temperature below T_C the molecules are close enough for the attractive forces to act, and condensation occurs

C. No condensation takes place above T_C



D. Due to higher kinetic energy of the gas molecules above T_C , it is considered as supercritical fluid

Answer: b,c,d



Watch Video Solution

217. What conclusion would you draw from the following graphs for an ideal gas?



- A. As the temperature is reduced, the volume as well as the pressure increase
- B. As the temperature is reduced, the volume becomes zero and the pressure reaches infinity

C. As the temperature is reduced, the pressure decrease

D. A point is reached where, theoretically, the volume become zero

Answer: c,d

 [Watch Video Solution](#)

218. Write the formula and identify the functional group , if any cyclo hex-1-en-1-ol

 [Watch Video Solution](#)

219. Indicate the correct statement for equal volumes of $N_2(g)$ and $CO_2(g)$ at $25^\circ C$ and 1 atm.

- A. The average translational K.E. per molecule is the same for N_2 and CO_2
- B. The rms speed is same for both N_2 and CO_2
- C. The density of N_2 is less than that of CO_2
- D. The total translational K.E. of both N_2 and CO_2 is the same

Answer: a,c,d



Watch Video Solution

220. Which of the following is correct for critical temperature ?

A. It is the highest temperature at which liquid and vapour can coexist

B. Beyond this temperature, there is no distinction between the two phases and a gas cannot be liquefied by compression.

C. At this temperature, the gas the liquid phases have different critical densities

D. All are correct

Answer: a,b

221. Consider the following statement regarding Maxwell's distribution of speeds. The correct statement(s) is/are :

A. as temperature increases, the peak (maxima) of a curve is shifted towards right side

B. as temperature increases, the most probable speed of molecules increases but fraction of molecules of maximum speed decreases

C. the area under the curve at all the temperatures is the same because it represents the number of

gaseous molecules

D. the fractions of molecules having different speeds are different at a given temperature

Answer: a,b,c,d

 [Watch Video Solution](#)


222. Write structure of 2-amino phenol

 [Watch Video Solution](#)

223. Write structure of isophthalic acid



224. Observe the figure given below and name the gas produced also give the chemical equation.



A. Total area under the two curves is independent to moles of gas

B. u_{mp} decrease as temperature decreases

C. $T_1 > T_2$ and hence higher the temperature,
sharper the curve


D. The fraction of molecules having speed u_{mp}
decreases as temperature increases

Answer: a,b,d



Watch Video Solution

225. Two container each containing liquid water are
connected as shown in diagram.



Given that vapour pressure of $H_2O(l)$ at 300 K and 350 K are 20 torr and 40 torr, select correct statement(s) :

- A. The final pressure in each container if valve is opened while keeping the containers at the given temperature is 22 torr
- B. The final pressure in each container if valve is opened while keeping the containers at the given temperature is 40 torr

C. Mass of $H_2O(l)$ is decreased in container X

D. Mass of $H_2O(l)$ is decreased in container Y

Answer: a,d



Watch Video Solution

226. Select the correct statement(s) :

A. At Boyle's temperature a real gas behaves like an

ideal gas at low pressure

B. Above critical conditions, a real gas behave like

an ideal gas

C. For hydrogen gas 'b' dominates over 'a' at all temperatures

D. At high pressure van der Waals' constant 'b' dominates over 'a'

Answer: a,b,d


 [Watch Video Solution](#)

227. Write structure of benzoic anhydride.

 [Watch Video Solution](#)


228. A open ended mercury manometer is used to measure the pressure exerted by a trapped gas as shown in the figure. Initially manometer shows no difference in mercury level in both columns as shown in diagram.

After sparking 'A' dissociates according to following reaction



If pressure of Gas "A" dissociates to 0.8 atm, then

(Assume temperature to be constant and is 300 K)



- A. total pressure increases to 1.3 atm
- B. total pressure increases by 0.3 atm
- C. total pressure increases by 22.3 cm of Hg
- D. difference in mercury level is 228 mm.

Answer: a,b,d

 [Watch Video Solution](#)

229. Write the formula and identify the functional group , if any : ethane nitrile

 [Watch Video Solution](#)

230. Write the formula and identify the functional group, if any : methyl methanoate



Watch Video Solution

231. Which is/are correct for real gases?

A. $\lim_{P \rightarrow 0} (PV_m) = \text{constant}$ at constant high temperature

B. $\lim_{V_m \rightarrow 0} (PV_m) = \text{constant}$ at constant low temperature

C. $\lim_{P \rightarrow 0} \left(\frac{PV_m}{RT} \right) = 1$ at high temperature

D. $\lim_{V \rightarrow 0} \left(\frac{PV_m}{RT} \right) = R$

Answer: a,c



Watch Video Solution

232. Write the formula and identify the functional group , if any : 2,2,4-trimethyl pentane



Watch Video Solution

233. Write the formula and identify the functional group , if any : cyclocta-1,5-diene



Watch Video Solution

234. Write the formula and identify the functional group , if any : hexanedial



Watch Video Solution

235. Write the formula and identify the functional group , if any : 3-nitrocyclohexene



Watch Video Solution

236. Write the formula and identify the functional group , if any : 4-nitropent-1-yne



Watch Video Solution

237. Write the formula and identify the functional group , if any : 4-ethyl-aniline



Watch Video Solution

238. What do you mean by chain isomerism?




Watch Video Solution

239. Write the formula and identify the functional group , if any : butan-2-one



Watch Video Solution

240. Match the following columns



Column-I

- (A) Speed at point 'A' is
 (B) Speed at point 'B' is
 (C) Speed at point 'C' is
 (D) The speed possessed by maximum fraction of the gas particles

Column-II

- (P) $\sqrt{\frac{8RT}{\pi M}}$
 (Q) $\sqrt{\frac{3RT}{M}}$
 (R) U_{rms}
 (S) $\sqrt{\frac{2RT}{M}}$



Watch Video Solution

241. STATEMENT-1 : The Heat absorbed during the isothermal expansion of an ideal gas against vacuum is zero.

STATEMENT-2 : The volume occupied by the molecules of an ideal gas is zero.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



242. STATEMENT-1 : A lighter gas diffuses mor rapidly than a heavier gas.

STATEMENT-2 : At a given temperature, the rate of diffusion of a gas is inversely proportional to density.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C

 [Watch Video Solution](#)

243. Assertion: The value of van der Waals constant a is larger for ammonia than for nitrogen.

Reason: Hydrogen bonding is present in ammonia.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

244. Assertion: Helium shows only positive deviations from ideal behaviour.

Reason: Helium is an inert gas.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

245. Statement-1: CH_4, CO_2 has value of Z (compressibility factor) less than one, generally.
Statement-2: $Z < 1$ is due to repulsive forces among the molecules.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

246. Assertion (A): The Joules -Thomon coefficient for an ideal gas is zero.

Reason (R) : There are no intermolecular attractive forces in an ideal gas.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A

 [Watch Video Solution](#)

247. STATEMENT-1 : The average translational kinetic energy per molecule of the gas per degree of freedom

is $\frac{1}{2} kT$.

STATEMENT-2 : For every molecule there are three rotational degree of freedom.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



Watch Video Solution

248. STATEMENT-1 : On increasing the temperature, the height of the peak of the Maxwell distribution curve decreases.

STATEMENT-2 : The fraction of molecules is very less at the higher speeds.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is

FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is

TRUE

Answer: B



Watch Video Solution

249. STATEMENT-1 : The gases He and H_2 are very different in their behaviour at any temperature and pressure but their compressibility factors are nearly the same at the critical point.

STATEMENT-2 : They have nearly the same critical constant.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



250. Assertion: Most probable velocity is the velocity possessed by maximum fraction of molecules at the same temperature.

Reason: On collision, more and more molecules acquire higher speed at the same temperature.

A. If both the statement are TRUE and STATEMENT-

2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2

is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C

 [Watch Video Solution](#)

251. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: For each $10^{\circ}C$ rise of temperature the k

is nearly double.

STATEMENT -2: Energy wise distribution of molecules in a gas sample is an exponential function of temperature so $e^{-E_a/RT}$ is doubled.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C

 [Watch Video Solution](#)

252. STATEMENT-1 : 1 mol of H_2 and O_2 each occupy 22.7 L of volume at $0^\circ C$ and 1 bar pressure, considering ideal behaviour.

STATEMENT-2 : Molar volume for all ideal gases at the same temperature and pressure are equal.

A. If both the statement are TRUE and STATEMENT-

2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2

is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A

 [Watch Video Solution](#)

253. STATEMENT-1 : The gases He and H_2 are very different in their behaviour at any temperature and pressure but their compressibility factors are nearly the same at the critical point.

STATEMENT-2 : They have nearly the same critical constant.

A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



254. STATEMENT-1 : $1/4^{th}$ of the initial mole of the air is expelled, if air present in an open vessel is heated from $27^{\circ}C$ to $127^{\circ}C$.

STATEMENT-2 : Rate of diffusion of a gas is inversely proportional to the square root of its molecular mass.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

255. STATEMENT-1: All strong monoprotic acid with same concentration in dilute solution show same pH.

STATEMENT-2: Water shows levelling effect.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

256. STATEMENT-1 : Wet air is heavier than dry air.

STATEMENT-2 : The density of dry air is less than the density of water at 1 atm and 273K.

- A. If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



[Watch Video Solution](#)

257. A bubble of gas released at the bottom of a lake increases to four times its original volume when it reaches the surface. Assuming that atmospheric pressure is equivalent to the pressure exerted by a column of water 10 m high, what is the depth of the lake?




[Watch Video Solution](#)

258. A gaseous mixture containing equal mole of H_2 , O_2 and He is subjected to series of effusion steps.

The composition (by moles) of effused mixture after 4 effusion steps is $x : 1 : y$ respectively. Then find the value of $\left(\frac{x}{y}\right)$.

[Watch Video Solution](#)

259. Find $x+y+z$



[Watch Video Solution](#)

260. 1 mole of a diatomic gas present in 10 L vessel at certain temperature exert a pressure of 0.96 atm. Under similar conditions an ideal gas exerted 1.0 atm pressure. If volume of gas molecule is negligible, then find the value of van der Waals' constant "a" (in $\text{atm L}^2 / \text{mol}^2$).


 [Watch Video Solution](#)

261. The graph of compressibility factor (Z) vs. P for one mole of a real gas is shown in following diagram. The graph is plotted at constant temperature 273K. If the slope of graph at very high pressure $\left(\frac{dZ}{dP}\right)$ is

$\left(\frac{1}{2.8}\right) atm^{-1}$, then calculate volume of one mole of real gas molecules (in L/mol)

Given : $N_A = 6 \times 10^{23}$ and

$$R = \frac{22.4}{273} LatmK^{-1}mol^{-1}$$



[Watch Video Solution](#)

262. Under the identical conditions of temperature, the density of a gas X is two times to that of gas Y while molecular mass of gas Y is three times that of X. Calculate the ratio of pressure of X and Y.



Watch Video Solution

263. The time for a certain volume of a gas A to diffuse through a small hole is 2 minute If takes 5.65 minute for oxygen to diffuse under similar conditions Find the molecular weight of A .



Watch Video Solution

264. Excess $F_2(g)$ reacts at $150^\circ C$ and 1.0 atm pressure with $Br_2(g)$ to give a compound BrF_n . If 423 mL of $Br_2(g)$ at the same temperature and


pressure produced 4.2 g of BrF_n , what is n? [Atomic mass Br =80, F =19]

 [Watch Video Solution](#)

265. Initially bulb "a" contained oxygen gas at $27^\circ C$ and 950 mm of Hg and bulb "B" contained neon gas at $27^\circ C$ and 900 mm of Hg. These bulbs are connected by a narrow tube of negligible volume equipped with a stopcock and gases were allowed to mix-up freely. The pressure obtained in the combined system was found to be 910 mm of Hg.


If volume of bulb B was measured to be 10 L, then find

the volume of oxygen gas present initially in bulb "A" .



[Watch Video Solution](#)

266. Which phenomenon of light is shown in fig below:



[Watch Video Solution](#)

267. A flask containing air at $107^{\circ} C$ and 722 mm of Hg is cooled to 100 K and 760 mm of Hg. If density in the initial condition $1g/cm^3$, then what is the final density (g/cm^3)?

 [Watch Video Solution](#)

268. If an ideal gas at 100 K is heated to 109 K in a rigid container, the pressure increases by X%. What is the value of X?

 [Watch Video Solution](#)

269. The van der Waals' constantes for a gas are

$$a = 3.6 \text{ atm L}^2 \text{ mol}^{-2}, b = 0.6 \text{ L mol}^{-1} \quad .\text{if}$$

$$R = 0.08 \text{ Latm K}^{-1} \text{ mol}^{-1} \quad \text{and} \quad \text{the Boyle's}$$

temperature (K) is T_B of this gas, then what is the

value of $\frac{T_B}{15}$?

 [Watch Video Solution](#)

270. A flask has 10 molecules out of which four

molecules are moving at 7 ms^{-1} and the remaining

ones are moving at same speed of $X \text{ ms}^{-1}$. If rms of

the gas is 5 ms^{-1} , what is X ?

 [Watch Video Solution](#)

Level 1 Q 31 To Q 60

1. Nitrogen forms several gaseous oxides. One of them has a density of 1.33 g/L measured 764 mmHg and $150^{\circ}C$. Write the formula of the compound.

A. NO

B. N_2O

C. NO_2

D. N_2O_5


Answer: c



Watch Video Solution

Level 1 Q 121 To Q 150

1. Dimethyl ether decomposes as



When $CH_3OCH_3(g)$ decomposes to 20% extent under certain conditions, what is the ratio of diffusion of pure $CH_3OCH_3(g)$ with methane?

A. 0.59: 1

B. 1.18: 1

C. 2.36: 1

D. 1.77: 1

Answer: c



Watch Video Solution

Level 1 Q 151 To Q 176

1. The temperature at which real gases obey the ideal gas laws over a wide range of low pressure is called:

- A. Critical temperature
- B. Inversion temperature
- C. Boyle temperature
- D. Reduced temperature

Answer: c

 [Watch Video Solution](#)


Level 2

1. The following organic compound are popularly known by their common names. write their structural formula and iupac name : hydroquinol

 [Watch Video Solution](#)

Level 3 Passage

1. Find $x+y+z$



- A. independent to temperature
- B. decreases with rise in temperature
- C. increases with rise in temperature
- D. directly proportional to T^2

Answer: c

 [Watch Video Solution](#)

Level 3 Subjective Problems

1. One mole of an ideal gas undergoes a change of state (2.0 atm, 3.0 L) to (2.0 atm, 7.0 L) with a change in internal energy (ΔU) = 30 L-atm. The change in enthalpy (ΔH) of the process in L-atm :



[Watch Video Solution](#)