



MATHS

BOOKS - OBJECTIVE RD SHARMA ENGLISH

DISCRETE PROBABILITY DISTRIBUTIONS

Illustration

1. If the probability of a random variable X is a given

below :

$$X = x: \quad -2 \quad -1 \quad 0 \quad 1 \quad 2 \quad 3$$

$$P(X = x): \quad \frac{1}{10} \quad k \quad \frac{1}{5} \quad 2k \quad \frac{3}{10} \quad k$$

Then the value of k , is :

A. $\frac{1}{10}$

B. $\frac{2}{10}$

C. $\frac{3}{10}$

D. $\frac{7}{10}$

Answer: A



Watch Video Solution

2. A random variable X takes values $0, 1, 2, 3, \dots$ with probability proportional to $(x + 1) \left(\frac{1}{5}\right)^x$.

$P(X = 0)$ equals

A. $\frac{7}{25}$

B. $\frac{16}{25}$

C. $\frac{18}{25}$

D. $\frac{19}{25}$

Answer: B



Watch Video Solution

3. A random variable 'X' has the following probability distribution :

X	0	1	2	3	4	5	6	7
P(X)	k	k	2k	3k	k ²	2k ²	7k ²	2k

The values of k is

A. $\frac{7}{81}$

B. $\frac{5}{81}$

C. $\frac{2}{81}$

D. $\frac{1}{81}$

Answer: D



Watch Video Solution

4. A random variable X has the following probability distribution :

$X:$	1	2	3	4	5	6	7	8
$P(X):$	0.15	0.23	0.12	0.10	0.20	0.08	0.07	0.08

for the events $E = [X \text{ is a prime number}]$

$F = \{X < 4\}$, the probability $P(E \cup F)$, is.



Watch Video Solution

5. A random, variable X has the following probability distribution:

$$X: 0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7$$

$$P(X): 0 \ k \ 2k \ 2k \ 3k \ k^2 \ 2k^2 \ 7k^2 + k$$

Find each of the following: k ii. $P(X < 6)$ iii. $P(X \geq 6)$ iv. $P(0$

A. $\frac{19}{100}$

B. $\frac{81}{100}$

C. $\frac{9}{100}$

D. $\frac{91}{100}$

Answer: A



Watch Video Solution

6. about to only mathematics

A. 0.1

B. 0.2

C. 0.3

D. 0.4

Answer: D



Watch Video Solution

7. A random variable X has the following probability distribution.

$X:$	1	2	3	4	5
$P(X):$	0	0.1	0.2	0.3	0.4

Find the mean.



[Watch Video Solution](#)

8. Five coins are tossed simultaneously. The probability that at least one head is turning up, is

A. $1/32$

B. $5/32$

C. $7/16$

Answer: D**Watch Video Solution**

9. The probability that a marksman will hit a target is given is $1/5$. Then the probability that at least once hit in 10 shots is $1 - (4/5)^{10}$ b. $1/5^{10}$ c. $1 - (1/5)^{10}$ d. $(4/5)^{10}$

A. $1 - \left(\frac{4}{5}\right)^{10}$

B. $\frac{1}{5^{10}}$

C. $1 - \frac{1}{5^{10}}$

D. none of these

Answer: A



Watch Video Solution

10. The probability that an event A happens in one trial of an experiment, is 0.4. Three independent trials of the experiments are performed. The probability that the event A happens atleast once, is :

A. 0.936

B. 0.784

C. 0.904

D. none of these

Answer: B



Watch Video Solution

11. एक प्रयोग के सफल होने का संयोग उसके असफल होने से दो गुना है। प्रायिकता ज्ञात कीजिए कि अगले छः परीक्षणों में कम से कम 4 सफल होंगे।

A. $\frac{496}{729}$

B. $\frac{233}{729}$

C. $\frac{432}{729}$

D. $\frac{256}{729}$

Answer: A



View Text Solution

12. One hundred identical coins, each with probability 'p' of showing heads are tossed once. If $0 < p < 1$ and the probability of heads showing on 50 coins is equal to that of heads showing on 51 coins, then the value of p is

A. $\frac{1}{2}$

B. $\frac{49}{101}$

C. $\frac{50}{101}$

D. $\frac{51}{101}$

Answer: D



Watch Video Solution

13. A man takes a step forward with probability 0.4 and backward with probability 0.6. Then find the probability that at the end of eleven steps he is one step away from the starting point.

A. ${}^{11}C_5 (0.4)^6 (0.6)^5$

B. ${}^{11}C_6 (0.4)^5 (0.6)^6$

C. ${}^{11}C_5 (0.4)^5 (0.6)^6$

D. ${}^n C_5 (0.4)^5 (0.6)^5$

Answer: C



Watch Video Solution

14. India play two matches each with West Indies and Australia. In any match the probabilities of India getting 0,1 and 2 points are 0.45, 0.05 and 0.50 respectively. Assuming that the outcomes are independent, the probability of India getting at least 7 points is 0. 0875 b. $1/16$ c. 0. 1125 d. none of these

A. $\frac{1}{80}$

B. $\frac{7}{80}$

C. $\frac{7}{8}$

D. $\frac{1}{8}$

Answer: B



Watch Video Solution

15. The probability of India winning a test match against England is $\frac{2}{3}$. Assuming independence from match to match, the probability that in a 7 match series India's third win occurs at the fifth match, is

A. $\frac{8}{27}$

B. $\frac{16}{81}$

C. $\frac{8}{81}$

D. $\frac{32}{81}$

Answer: B



Watch Video Solution

16. about to only mathematics

A. $1/4$

B. $1/32$

C. $1/16$

D. $1/8$

Answer: B

 [Watch Video Solution](#)

17. A dice is thrown 100 times . If getting an even number is considered a success , then the variance of the number of successes is

A. 50

B. 25

C. 10

D. 100

Answer: B

 [Watch Video Solution](#)

18. In eight throws of a die 1 or 3 is considered a success. Then, the standard deviation of success, is

A. $\frac{16}{9}$

B. $\frac{8}{3}$

C. $\frac{4}{3}$

D. $\frac{2}{3}$

Answer: C



Watch Video Solution

19. A die is tossed 5 times. Getting an odd number is considered a success. Then, the variance of distribution of success, is

A. $\frac{8}{3}$

B. $\frac{3}{8}$

C. $\frac{4}{5}$

D. $\frac{5}{4}$

Answer: D



Watch Video Solution

20. A box contains 15 green and 10 yellow balls. If 10 balls are randomly drawn, one -by - one with replacement, then the variance of the number of green balls drawn, is

A. $\frac{12}{5}$

B. 6

C. 4

D. $\frac{6}{25}$

Answer: A



Watch Video Solution

Section I Solved Mcqs

1. In a box containing 100 bulbs, 10 are defective. The probability that out of a sample of 5 bulbs, none is defective is (A) 10^{-1} (B) $\left(\frac{1}{2}\right)^5$ (C) $\left(\frac{9}{10}\right)^5$ (D) $\frac{9}{10}$

A. $\left(\frac{9}{10}\right)^5$

B. $\frac{9}{10}$

C. 10^{-5}

D. $\left(\frac{1}{2}\right)^2$

Answer: A



Watch Video Solution

2. For a binomial variate X , if $n = 4$ and $P(X=0) = \frac{16}{81}$

then $P(X=4)$ is

A. $\frac{1}{16}$

B. $\frac{1}{81}$

C. $\frac{1}{27}$

D. $\frac{1}{8}$

Answer: B



Watch Video Solution

3. A rifleman is firing at a distance target and hence has only 10% chance of hitting it. Find the number of rounds; he must fire in order to have more than 50% chance of hitting it at least once.

A. 11

B. 9

C. 7

D. 5

Answer: C



Watch Video Solution

4. A fair coin is tossed n times . If the probability of getting seven heads is equal to the probability of getting 9 heads , then the probability of getting two heads is

A. $15/2^8$

B. $2/15$

C. $15/2^{13}$

D. none of these

Answer: C



Watch Video Solution

5. A fair coin is tossed 100 times. The probability of getting tails an odd number of times is $\frac{1}{2}$ b. $\frac{1}{8}$ c. $\frac{3}{8}$ d. none of these

A. $\frac{1}{2}$

B. $\frac{1}{8}$

C. $\frac{3}{8}$

D. none of these

Answer: A



Watch Video Solution

6. A fair die is thrown 20 times. The probability that on the 10th throw, the fourth six appears is

A. $\frac{{}^{20}C_{10} \times 5^5}{6^{20}}$

B. $\frac{120 \times 5^7}{6^{10}}$

C. $\frac{84 \times 5^6}{6^{10}}$

D. none of these

Answer: C



Watch Video Solution

7. If X has binomial distribution with mean np and variance npq , then $\frac{P(X = r)}{P(X = r - 1)}$ is equal to

A. $1/2$

B. $1/3$

C. $1/4$

D. none of these

Answer: A



Watch Video Solution

8. A fair coin is tossed n times. Let X denote the number of times head occurs. If $P(X = 4)$, $P(X = 5)$ and $P(X = 6)$ are in arithmetic progression, then the value of n can be

A. 7,14

B. 10,14

C. 12,7

D. 14,12

Answer: A



Watch Video Solution

9. One hundred identical coins, each with probability 'p' of showing heads are tossed once. If $0 < p < 1$ and the probability of heads showing on 50 coins is equal to that of heads showing on 51 coins, then the value of p is

A. $1/2$

B. $51/101$

C. $49/101$

D. none of these

Answer: B



Watch Video Solution

10. A fair coin is tossed 99 times. Let X be the number of times heads occurs. Then $(P(X=r))$ is maximum when r is

A. 49,50

B. 50,51

C. 51,52

D. none of these

Answer: A



Watch Video Solution

11. A coin is tossed n times. The probability of getting head atleast once is greater than 0.8. Then the least value of such n is

A. 7

B. 6

C. 5

D. 3

Answer: D



Watch Video Solution

12. If the mean and the variance of a binomial variate X are 2 and 1 respectively, then the probability that X takes a value greater than one is equal to

A. $\frac{2}{3}$

B. $\frac{4}{5}$

C. $\frac{7}{8}$

D. $\frac{15}{16}$

Answer: D



Watch Video Solution

13. A biased coin with probability p , $0 < p < 1$, of heads is tossed until a head appears for the first time. If the probability that the number of tosses required is even is $2/5$, then p is equal to

A. $1/3$

B. $2/3$

C. $2/5$

D. $3/5$

Answer: A



Watch Video Solution

14. If X follows a binomial distribution with parameters

$n = 8$ and $p = 1/2$, then $p(|x - 4| \leq 2)$ is equal to :

A. $\frac{118}{128}$

B. $\frac{119}{128}$

C. $\frac{117}{128}$

D. none of these

Answer: B



[Watch Video Solution](#)

15. If X follows a binomial distribution with parameters

$n = 100$ and $p = \frac{1}{3}$, then $P(X=r)$ is maximum, when r

equals

A. 32

B. 34

C. 33

D. 31

Answer: C



Watch Video Solution

16. A fair die is tossed eight times. The probability that

a third six is observed in eight throw is $\frac{{}^7C_{10} \times 5^7}{6^7}$ b.

$\frac{{}^7C_2 \times 5^2}{6^8}$ c. $\frac{{}^7C_2 \times 5^5}{6^6}$ d. none of these

A. $\frac{{}^7C_2 \times 5^5}{6^7}$

B. $\frac{{}^7C_2 \times 5^5}{6^8}$

C. $\frac{{}^7C_2 \times 5^5}{6^6}$

D. none of these

Answer: B



Watch Video Solution

17. Four digit numbers with different digits are formed using the digits 1, 2, 3, 4, 5, 6, 7, 8. One number from them is picked up at random. The chance that the selected number contains the digit '1' is

A. $1/8$

B. $1/4$

C. $1/2$

D. none of these

Answer: C



Watch Video Solution

18. A die is thrown $2n + 1$ times, $n \in \mathbb{N}$. The probability that faces with even numbers show odd number of times is :

A. $\frac{1}{2}$

B. $< \frac{1}{2}$

C. $> \frac{1}{2}$

D. none of these

Answer: A



Watch Video Solution

19. A fair coin is tossed repeatedly. The probability of getting a result in fifth toss different from those obtained in the first four tosses is

A. $1/2$

B. $1/32$

C. $\frac{31}{32}$

D. $\frac{1}{16}$

Answer: D



Watch Video Solution

20. An ordinary dice is rolled a certain number of times. If the probability of getting an odd number 2 times is equal to the probability of getting an even number 3 times. Then the probability of getting an odd number and odd number of times, is

A. $\frac{1}{32}$

B. $5/16$

C. $1/2$

D. none of these

Answer: C



Watch Video Solution

21. A coin is tossed $2n$ times. The chance that the number of times one gets head is not equal to the

number of times one gets tails is $\frac{(2n!)}{(n!)^2} \left(\frac{1}{2}\right)^{2n}$ b.

$1 - \frac{(2n!)}{(n!)^2}$ c. $1 - \frac{(2n!)}{(n!)^2} \frac{1}{(4^n)}$ d. none of these

A. ${}^{2n}C_n \left(\frac{1}{2}\right)^{2n}$

B. $1 - {}^{2n}C_n$

C. $1 - {}^{2n}C_n \left(\frac{1}{4^n} \right)$

D. none of these

Answer: C



Watch Video Solution

22. A card is drawn from a pack of 52 playing cards. The card is replaced and the pack is reshuffled. If this is done six times. The probability that 2 hearts, 2 diamond and 2 black cards are drawn is

A. $90 \times \left(\frac{1}{4} \right)^6$

B. $\frac{45}{2} \times \left(\frac{3}{4}\right)^4$

C. $90 \times \left(\frac{1}{2}\right)^{10}$

D. none of these

Answer: C



Watch Video Solution

23. From a box containing 20 tickets marked with numbers 1 to 20, four tickets are drawn one by one. After each draw, the ticket is replaced. The probability that the largest value of tickets drawn is 15 is.

A. $\left(\frac{3}{4}\right)^4$

B. $\frac{27}{320}$

C. $\frac{27}{1280}$

D. none of these

Answer: B



Watch Video Solution

24. A coin is tossed 3 times by 2 persons. The probability that both get equal number of heads, is

A. $3/8$

B. $1/9$

C. $5/16$

D. none of these

Answer: C



Watch Video Solution

25. The mean and the variance of a binomial distribution are 4 and 2 respectively. then, the probability of 2, successes is

A. $28 / 256$

B. $219 / 256$

C. $128 / 256$

D. $37 / 256$

Answer: A



Watch Video Solution

26. If two coins are tossed five times, then the probability of getting 5 heads and 5 tails is

A. $\frac{63}{256}$

B. $\frac{1}{1024}$

C. $\frac{2}{205}$

D. $\frac{9}{64}$

Answer: A



Watch Video Solution

27. If X and Y are independent binomial variates $A\left(5, \frac{1}{2}\right)$ and $B\left(7, \frac{1}{2}\right)$. Find the value of $P(X+Y=3)$.

A. $35/47$

B. $55/1024$

C. $220/512$

D. $11/204$

Answer: B



Watch Video Solution

28. In a binomial distribution $B\left(n, p = \frac{1}{4}\right)$, if the probability of at least one success is greater than or equal to $\frac{9}{10}$, then n is greater than (1)

$$(1) \frac{(\log)_{10}^4 - (\log)_{10}^3}{1}$$

$$(2) \frac{(\log)_{10}^4 + (\log)_{10}^3}{9}$$

$$(3) \frac{(\log)_{10}^4 - (\log)_{10}^3}{4}$$

$$(4) \frac{(\log)_{10}^4 - (\log)_{10}^3}{1}$$

$$A. \frac{1}{\log_{10} 4 - \log_{10} 3}$$

$$B. \frac{1}{\log_{10} 4 + \log_{10} 3}$$

$$C. \frac{9}{\log_{10} 4 - \log_{10} 3}$$

$$D. \frac{4}{\log_{10} 4 - \log_{10} 3}$$

Answer: A



Watch Video Solution

29. Two cards are drawn successively with replacement from a well shuffled deck of 52 cards, then the mean of the number of aces is

A. $1/13$

B. $3/13$

C. $2/13$

D. none of these

Answer: C



Watch Video Solution

30. A dice is thrown 100 times . If getting an even number is considered a success , then the variance of the number of successes is

A. 10

B. 25

C. 18

D. 12

Answer: B



Watch Video Solution

31. There are 12 white and 12 red ball in a bag. Balls are drawn one by one with replacement from the bag. The probability that 7th drawn ball is 4th white, is

A. $1/4$

B. $5/32$

C. $3/16$

D. $5/16$

Answer: B



Watch Video Solution

32. A die is rolled thrice . If the event of getting an even number is a success , then the probability of getting atleast two sucessess is

A. $\frac{7}{8}$

B. $\frac{1}{4}$

C. $\frac{2}{3}$

D. $\frac{1}{2}$

Answer: D



Watch Video Solution

33. IF the mean and the variance of a binomial distribution are 4 and 3 respectively ,then the probability of six successes is

A. ${}^{16}C_6 \left(\frac{1}{4}\right)^6 \left(\frac{3}{4}\right)^{10}$

B. ${}^{16}C_6 \left(\frac{1}{4}\right)^{16} \left(\frac{3}{4}\right)^{20}$

C. ${}^{16}C_6 \left(\frac{1}{4}\right)^8 \left(\frac{3}{4}\right)^{12}$

D. ${}^{16}C_9 \left(\frac{1}{4}\right)^{16} \left(\frac{3}{4}\right)^{20}$

Answer: A



Watch Video Solution

34. The sum of mean and variance of a binomial distribution is 15 and the sum of their squares is 117. Determine the distribution.

A. 6

B. 9

C. 3

D. 12

Answer: B



Watch Video Solution

35. Consider 5 independent Bernoulli's trials each with probability of at least one failure is greater than or equal to $\frac{31}{32}$, then p lies in the interval

A. $\left(\frac{1}{2}, \frac{3}{4}\right]$

B. $\left(\frac{3}{4}, \frac{11}{12}\right]$

C. $\left[0, \frac{1}{2}\right]$

D. $\left(\frac{11}{12}, 1\right]$

Answer: C



Watch Video Solution

36. A random variable X takes values $-1, 0, 1, 2$ with probabilities $\frac{1+3p}{4}, \frac{1-p}{4}, \frac{1+2p}{4}, \frac{-14p}{4}$ respectively, where p varies over R . Then the minimum and maximum values of the mean of X are respectively

A. $-\frac{7}{4}$ and $\frac{1}{2}$

B. $-\frac{1}{16}$ and $\frac{5}{16}$

C. $-\frac{7}{4}$ and $\frac{5}{16}$

D. $-\frac{1}{16}$ and $\frac{5}{4}$

Answer: D



View Text Solution

37. A multiple choice examination has 5 questions. Each question has three alternative answers of which exactly one is correct. The probability that a student will get 4 or more correct answers just guessing is

A. $\frac{17}{3^5}$

B. $\frac{13}{3^5}$

C. $\frac{11}{3^5}$

D. $\frac{1}{3^5}$

Answer: C



Watch Video Solution

38. about to only mathematics

A. 7

B. 9

C. 8

D. 5

Answer: C



Watch Video Solution

39. एक प्रयोग के सफल होने का संयोग उसके असफल होने से दो गुना है। प्रायिकता ज्ञात कीजिए कि अगले छः परीक्षणों में कम से कम 4 सफल होंगे।

A. $\frac{192}{729}$

B. $\frac{256}{729}$

C. $\frac{240}{729}$

D. $\frac{4962}{729}$

Answer: B



View Text Solution

Section II Assertion Reason Type

1. Statement -1 : The probability of getting a tail most of the times in 10 tosses of a unbiased coin is

$$\frac{1}{2} \left\{ 1 - \frac{10!}{2^{10} 5! 5!} \right\}.$$

Statement

-2:

$${}^2 nC_0 + {}^2 nC_1 + {}^2 nC_2 + \dots + {}^2 nC_n = 2^2 n - 1, n \in N$$

.

A. 1

B. 2

C. 3

D. 4

Answer: C



[View Text Solution](#)

Exercise

1. A random variable has the following probability distribution:

$$X = x_i: 0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7$$

$$P(X = x_i): 0 \ 2p \ 2p \ 3p \ p^2 \ 2p^2 \ 7p^2 \ 2p$$

The value of p is

a. $\frac{1}{10}$

b. -1

c. $-\frac{1}{10}$

d. $\frac{1}{5}$

A. $1/10$

B. -1

C. $-1/10$

D. none of these

Answer: A



Watch Video Solution

2. If X is a random variable with probability distribution as given below: $X = x_i: 0 \ 1 \ 2 \ 3$

$P(X = x_i): k \ 3k \ 3k \ k$ The value of k and its

variance are $1/8, 22/27$ b. $1/8, 23/27$ c. $1/8, 24/27$

d. $1/8, 3/4$

A. $1/8, 22/27$

B. $1/8, 23/27$

C. $1/8, 24/27$

D. $1/8, 3/4$

Answer: D



Watch Video Solution

3. If in a distribution each x is replaced by corresponding value of $f(x)$, then the probability of getting $f(x)$, when the probability of getting x_i is p_i , is.

A. p_i

B. $f(p_i)$

C. $f\left(\frac{1}{p_i}\right)$

D. none of these

Answer: A



Watch Video Solution

4. A man takes a step forward with probability 0.4 and backward with probability 0.6. Then find the probability that at the end of eleven steps he is one step away from the starting point.

A. ${}^C_{(6)} (0.24)^5$

B. ${}^{11}C_6(0.4)^6(0.6)^5$

C. ${}^{11}C_6(0.6)^6(0.4)^5$

D. none of these

Answer: A



Watch Video Solution

5. The probability that a man can hit a target is $\frac{3}{4}$.

He tries 5 times. The probability that he will hit the target at least three times is

A. $\frac{291}{364}$

B. $\frac{371}{464}$

C. 471/502

D. 459/512

Answer: D



Watch Video Solution

6. Six ordinary dice are rolled. The probability that at least half of them will show at least 3 is

A. $41 \times \frac{2^4}{3^6}$

B. $\frac{2^4}{3^6}$

C. $20 \times \frac{2^4}{3^6}$

D. none of these

Answer: A



Watch Video Solution

7. Two persons each makes a single throw with a pair of dice. Find the probability that the throws are unequal.

A. $\frac{1}{6^3}$

B. $\frac{73}{6^3}$

C. $\frac{51}{6^3}$

D. none of these

Answer: C



Watch Video Solution

8. If the range of a random variable X is 0, 1, 2, 3, at

$P(X = K) = \left(\frac{K + 1}{3^k}\right) a$ for $k \geq 0$, then a equals

A. $\frac{2}{3}$

B. $\frac{4}{9}$

C. $\frac{8}{27}$

D. $(16)/(81)$

Answer: B



Watch Video Solution

9. An experiment succeeds twice as often as it fails.

Find the probability that in the next six trials there will be at least 4 successes.

A. $\frac{64}{729}$

B. $\frac{192}{729}$

C. $\frac{240}{79}$

D. $\frac{496}{729}$

Answer: D



Watch Video Solution

10. The probability that a candidate secure a seat in Engineering through EAMCET is $\frac{1}{10}$ Seven candidate are selected at random from a centre. The probability that exactly two will get seats is

A. $15(0.1)^2(0.9)^5$

B. $20(0.1)^2(0.9)^5$

C. $21(0.1)^2(0.9)^5$

D. $23(0.1)^2(0.9)^5$

Answer: C



Watch Video Solution

11. Six coins are tossed simultaneously. The probability of getting at least 4 heads, is

A. $\frac{11}{64}$

B. $\frac{11}{32}$

C. $\frac{15}{44}$

D. $\frac{21}{32}$

Answer: B



Watch Video Solution

12. about to only mathematics

A. $\left(\frac{9}{16}\right)^6$

B. $\left(\frac{8}{15}\right)^7$

C. $\left(\frac{3}{5}\right)^5$

D. none of these

Answer: D



Watch Video Solution

13. Two players toss 4 coins each. The probability that they both obtain the same number of heads is a. $5/256$ b. $1/16$ c. $35/128$ d. none of these

A. $5/256$

B. $1/16$

C. $35/128$

D. none of these

Answer: C



Watch Video Solution

14. A box contains 24 identical balls of which 12 are white and 12 are black. The balls are drawn at random from the box one at a time with replacement. The probability that a white ball is drawn for the 4th time on the 7th draw is

A. $\frac{5}{64}$

B. $\frac{27}{32}$

C. $\frac{5}{32}$

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

15. Two dice are tossed 6 times. Then the probability that 7 will show an exactly four of the tosses , is

A. $\frac{225}{18442}$

B. $\frac{116}{20003}$

C. $\frac{125}{15552}$

D. none of these

Answer: C



Watch Video Solution

16. If X follows a binomial distribution with parameters $n = 6$ and p . If $4P(X = 4) = P(X = 2)$, then $P =$

A. $1/2$

B. $1/4$

C. $1/6$

D. $1/3$

Answer: D



Watch Video Solution

17. The number of times a die must be tossed to obtain a 6 at least one with probability exceeding 0.9 is at least

A. 13

B. 19

C. 25

D. none of these

Answer: A



Watch Video Solution

18. Seven chits are numbered 1 to 7. Four chits are drawn one by one with replacment. The probability that the least number appearing on any selected chit is 5 is :

A. $\left(\frac{3}{7}\right)^4$

B. $\left(\frac{6}{7}\right)^3$

C. $\frac{5 \times 4 \times 3}{7^3}$

D. $\left(\frac{3}{4}\right)^4$

Answer: A





Watch Video Solution

19. If the mean of a binomial distribution is 25, then its standard deviation lies in the interval

A. $[0, 5]$

B. $(0, 5]$

C. $[0, 25)$

D. $(0, 25]$

Answer: A



Watch Video Solution

20. The value of C for which $P(X = k) = Ck^2$ can serve as the probability function of a random variable X that takes value 0, 1, 2, 3, 4, is

A. $\frac{1}{30}$

B. $\frac{1}{10}$

C. $\frac{1}{3}$

D. $\frac{1}{15}$

Answer: A



Watch Video Solution

21. In order to get a head at least once with probability ≥ 0.9 , the minimum number of times a unbiased coin needs to be tossed is

A. 3

B. 4

C. 5

D. none of these

Answer: B



Watch Video Solution

22. The probability that a man will hit a target in shooting practise is 0.3. If he shoots 10 times, the probability that he hits the target , is

A. 1

B. $1 - (0.7)^{10}$

C. $(0.7)^{10}$

D. $(0.3)^{10}$

Answer: B



Watch Video Solution

23. If A and B each toss three coins. The probability that both get the same number of heads is

A. $1/9$

B. $3/16$

C. $5/16$

D. $3/8$

Answer: C



Watch Video Solution

24. In a box containing 100 bulbs, 10 bulbs are defective. Probability that out of a sample of 5 bulbs, none is defective, is

A. $\left(\frac{1}{10}\right)^5$

B. $\left(\frac{1}{2}\right)^5$

C. $\left(\frac{9}{10}\right)^5$

D. $\frac{9}{10}$

Answer: C



Watch Video Solution

25. The box contains tickets numbered from 1 to 20. Three tickets are drawn from the box with replacement. The probability that the largest number on the tickets is 7 is $\frac{2}{19}$ b. $\frac{7}{20}$ c. $1 - \left(\frac{7}{20}\right)^3$ d. none of these

A. $\frac{2}{19}$

B. $\frac{7}{20}$

C. $1 - \left(\frac{7}{20}\right)^3$

D. none of these

Answer: D



Watch Video Solution

26. An unbiased coin is tossed a fixed number of times. If the probability of getting 4 heads equals the probability of getting 7 heads, then the probability of getting 2 heads, is

A. $\frac{55}{2048}$

B. $\frac{3}{4096}$

C. $\frac{1}{1024}$

D. none of these

Answer: A



Watch Video Solution

27. A coin is tossed n times. The probability that head will turn up an odd number of times, is

A. $\frac{1}{2}$

B. $\frac{n+1}{2n}$

C. $\frac{n-1}{2n}$

D. $\frac{2^n - 1 - 1}{2^n}$

Answer: A



Watch Video Solution

28. Two coins are tossed five times. The probability that an odd number of heads are obtained, is

A. $\left(\frac{1}{2}\right)^5$

B. $\frac{3}{5}$

C. $\frac{2}{5}$

D. none of these

Answer: D



Watch Video Solution

29. A six-faced dice is so biased that it is twice as likely to show an even number as an odd number when thrown. It is thrown twice, the probability that the sum of two numbers thrown is even is

A. $\frac{4}{9}$

B. $\frac{5}{9}$

C. $\frac{1}{9}$

D. none of these

Answer: B



Watch Video Solution

30. A fair coin is tossed n times. if the probability that head occurs 6 times is equal to the probability that head occurs 8 times, then find the value of n .

A. 7

B. 14

C. 16

D. 19

Answer: B



Watch Video Solution

31. An unbiased coin is tossed n times. Let X denote the number of times head occurs. If $P(X = 4)$, $P(X = 5)$ and $P(X = 6)$ are in A.P, then the value of n can be

A. 7,14

B. 10,14

C. 12,7

D. none of these

Answer: A



Watch Video Solution

32. A fair coin is tossed is fixed number of times. If the probability of getting 4 heads equals the probability of getting 7 heads, then the probability of getting 2 heads, is

A. $55/2048$

B. $1/1024$

C. $3/4096$

D. none of these

Answer: A



Watch Video Solution

33. A carton contains 20 bulbs ,5 of which are defective. The probability that,if a sample of 3 bulbs in chosen at random from the carton, 2 will be defective, is

A. $1/16$

B. $\frac{3}{64}$

C. $\frac{9}{64}$

D. $\frac{2}{3}$

Answer: C



Watch Video Solution

34. In a precision bombing attack, there is a 50% chance that any one bomb will strike the target. Two direct hits are required to destroy the target completely. The number of bombs which should be dropped to give a 99% chance or better of completely destroying the target can be 12 b. 11 c. 10 d. 13

A. 10

B. 11

C. 12

D. none of these

Answer: A



Watch Video Solution

35. The probability of a man hitting a target is $\frac{1}{4}$. How many times must he fire so that the probability of his hitting the target at least once is greater than $\frac{2}{3}$?

A. 3

B. 4

C. 5

D. none of these

Answer: B



Watch Video Solution

36. IF the mean and S.D. of binomial distribution are 20 and 4 respectively. Than the number of trials , is

A. 50

B. 25

C. 100

D. 80

Answer: B



Watch Video Solution

37. The probability of India winning a test match against West Indies is $1/2$. Assuming independence from match to match, find the probability that in a match series India's second win occurs at the third test.

A. $1/8$

B. $1/4$

C. $\frac{1}{2}$

D. $\frac{2}{3}$

Answer: B



Watch Video Solution

38. The mean and standard deviation of a binomial variate X are 4 and $\sqrt{3}$ respectively. Then ,
 $P(X \geq 1) =$

A. $1 - \left(\frac{1}{4}\right)^{16}$

B. $1 - \left(\frac{3}{4}\right)^{16}$

C. $1 - \left(\frac{2}{3}\right)^{16}$

$$D. 1 - \left(\frac{1}{3}\right)^{16}$$

Answer: B



Watch Video Solution

39. The probability distribution of a random variable X is given by.

$X = x:$	0	1	2	3	4
$P(X = x):$	0.4	0.3	0.1	0.1	0.1

The variance of X , is

A. 1.76

B. 2.45

C. 3.2

D. 4.8

Answer: A



Watch Video Solution

40. For a binomial variate X with $n=6$, if

$P(X = 4) = \frac{135}{2^{12}}$ then its variance is

A. $8/9$

B. $1/4$

C. $9/8$

D. 4

Answer: C



Watch Video Solution