


## CHEMISTRY

### BOOKS - CENGAGE CHEMISTRY (ENGLISH)


### CHEMICAL EQUILIBRIUM

#### Solved Example


1. Which graph will show equilibrium condition?



A.



B.



C.

D. None of these

**Answer: C**

 [Watch Video Solution](#)

2. For  $A + B \rightleftharpoons C + D$  , the equilibrium constant is  $K_1$  and for  $C + D \rightleftharpoons A + B$ , the equilibrium constant is  $K_2$ . The correct relation between  $K_1$  and  $K_2$  is

A.  $K_1 \times K_2 = 1$

B.  $K_1 \times (K_2 - 1) = 0$

C.  $K_1/K_2 = 1$

D. All of these

**Answer: A**

 [Watch Video Solution](#)

**3.** For the reactions,

$A \rightleftharpoons B, K_c = 1, B \rightleftharpoons C, K_c = 3, C \rightleftharpoons D, K_c = 5.$   $K_c$  for the reaction  $A \rightleftharpoons D$  is

A. 15

B. 5

C. 3

D. 1

**Answer: A**

 [Watch Video Solution](#)

**4.** The law of mass action was enunciated by

A. Guldberg and Waage

B. Le Chatelier and Braun



C. Kossel and Lewis

D. vant Hoff


**Answer: A**

 [Watch Video Solution](#)

5. At a certain temperature, the following reactions have the equilibrium constants as shown below:



what is the equilibrium constant  $K_c$  for the reaction at the same temperature?



A.  $K_1 = K_2$



B.  $K_2^2 = K_1$

C.  $K_1^2 = K_2$

D.  $K_2 = \sqrt{K_1}$

**Answer: C**

 [Watch Video Solution](#)

6. When 4mol of A is mixed with 4mol of B, 2mol of C and D are formed at equilibrium, according to the reaction



the equilibrium constant is

A. (a)  $\sqrt{2}$

B. (b) 2

C. (c) 1

D. (d) 4

**Answer: C**



**Watch Video Solution**

7. The rate at which a substance reacts, depends on its:

- A. Active mass
- B. molecular mass
- C. Equivalent mass
- D. Atomic mass

**Answer: A**



**Watch Video Solution**

8. The state of equilibrium refers to

- A. State of rest

B. Dynamic state

C. Stationary state

D. State of inertness

**Answer: B**

 [Watch Video Solution](#)

9. For the reaction,  $A + 2B \rightleftharpoons C$ , the expression for equilibrium constant is

A.  $\frac{[A][B]^2}{[C]}$

B.  $\frac{[A][B]}{[C]}$

C.  $\frac{[C]}{[A][B]^2}$

D.  $\frac{[C]}{[2B][A]}$

**Answer: C**

 [Watch Video Solution](#)

10. For the reaction:



Two moles each of  $A$  and  $B$  were taken into a flask. The following must always be true when the system attained equilibrium

A.  $[A] = [B]$

B.  $[A] < [B]$

C.  $[B] = [C]$

D.  $[A] > [B]$

**Answer: B**



[Watch Video Solution](#)

11. In a chemical reaction equilibrium is established when :

A. Concentrations of reactants and products are equal

B. Opposing reactions cease

C. Speeds of opposing reactions become equal

D. Temperature of opposing reactions are equal

**Answer: C**

 [Watch Video Solution](#)

**12.** Active mass is defined as

A. Number of g equivalent per unit volume

B. Number of g mol per L



C. Amount of substance in g per unit volume

D. Number of g mol in 100L

**Answer: B**

 [Watch Video Solution](#)


13. For the reaction





write down the expression for equilibrium constants  $K_c$  and  $K'_c$ . How is  $K_c$  related to  $K'_c$ ?

 [Watch Video Solution](#)

14. The equilibrium constant for the reaction




at a particular temperature is 100. Write down the equilibrium law equations for the following reaction and determine the values of equilibrium constants.





 [Watch Video Solution](#)

15. Determine  $K_c$  for the reaction



from the following data at 298K.

The equilibrium constants for the following reaction





are  $2.4 \times 10^{30}$  and 1.4, respectively.




[Watch Video Solution](#)

16. For the hypothetical reactions, the equilibrium constant ( $K$ ) value are given



The equilibrium constant ( $K$ ) for the reaction



A. 48

B. 6

C. 27




D. 24

**Answer: D**




**Watch Video Solution**

17. Consider the following gases equilibrium given below:



The equilibrium constant for the reaction ,



A.  $K_1K_2K_3$

B.  $\frac{K_1K_2}{K_3}$

C.  $\frac{K_1K_3^2}{K_2}$



$$D. \frac{K_2 K_3^3}{K_1}$$

**Answer: D**

 [Watch Video Solution](#)

18. In a reversible reaction, study of its mechanism says that both the forward and reverse reaction follows first-order kinetics. If the half-life of forward reaction  $(t_{1/2})_f$  is 400s and that of reverse reaction  $(t_{1/2})_b$  is 250s, the equilibrium of the reaction is

A. 1.6

B. 0.433

C. 0.625

D. 1.109

**Answer: C**

 [Watch Video Solution](#)

**19.** A vessel at  $1000\text{K}$  contains carbon dioxide with a pressure of  $0.5\text{atm}$ . Some of the carbon dioxide is converted to carbon monoxide on addition of graphite. Calculate the value of  $K_p$  if total pressure at equilibrium is  $0.8\text{atm}$ .

 [Watch Video Solution](#)



**20.** A sample of  $\text{CaCO}_3(\text{s})$  is introduced into a sealed container of volume  $0.654\text{L}$  and heated to  $1000\text{K}$  until equilibrium is reached. The equilibrium constant for the reaction




is  $3.9 \times 10^{-2}$  atm at this temperature. Calculate the mass of  $\text{CaO}$  present at equilibrium.

 [Watch Video Solution](#)

21. Form the given data of equilibrium constants of the following reactions:




Calculate the equilibrium constant of the reaction,






 [Watch Video Solution](#)

22. Given that at 1000K



Calculate K for the following equations:



 [Watch Video Solution](#)

23. If  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  then  $2N_2 + 6H_2 \rightleftharpoons 4NH_3$ ;  $K'$  is equal to

- A.  $K^2$
- B.  $(K)^{1/3}$
- C.  $1/\sqrt{K}$
- D.  $1/K^2$

**Answer: A**

 [Watch Video Solution](#)

24. Equilibrium constants for four different reaction are given as  $K_1 = 10^6$ ,  $K_2 = 10^{-4}$ ,  $K_3 = 10$ , and  $K_4 = 1$ . Which reaction will take maximum time to attain equilibrium?

- A.  $K_1 = 10^6$
- B.  $K_2 = 10^{-4}$
- C.  $K_3 = 10$

D.  $K_4 = 1$

**Answer: B**



**Watch Video Solution**

25. In the equilibrium constant for  $A \rightleftharpoons B + C$  is  $K_{eq}^{(1)}$  and that of  $B + C = P$  is  $K_{eq}^{(2)}$ , the equilibrium constant for  $A \rightleftharpoons P$  is :

A.  $K_1 + K_2 + K_3$

B.  $K_1 \times K_2 \times K_3$





C.  $K_1 K_2 / 3$

D. None

**Answer: B**



**Watch Video Solution**



26.

Which oxide is most stable?

A. AO

B. BO

C. CO

D. DO

**Answer: A**



**Watch Video Solution**

27. Theory of 'active mass' indicates that the rate of a chemical reaction is directly proportional to the

- A. Equilibrium constant
- B. Properties of reactants
- C. Volume of apparents
- D. Concentration of reactants

**Answer: D**



[Watch Video Solution](#)

28. In the system  $X + 2Y \rightleftharpoons Z$ , the equilibrium concentration are,

$$[X] = 0.06 \text{ mol L}^{-1}, [Y] = 0.12 \text{ mol L}^{-1},$$

$[Z] = 0.216 \text{ mol L}^{-1}$ . Find the equilibrium constant of the reaction.

- A. 120
- B. 400

C.  $4 \times 10^{-3}$

D. 250

**Answer: D**

 [Watch Video Solution](#)

29. Equilibrium constants ( $K$ ) for the reaction

$2NO(g) + Cl_2(g) \rightleftharpoons 2NOCl(g)$  is correctly given by the expression

A. (a)  $\frac{[NOCl]^2}{[NO]^2 [Cl_2]}$

B. (b)  $\frac{[2NOCl]}{[2NO] [Cl_2]}$

C. (c)  $\frac{[NO]^2 + [Cl_2]}{[NOCl]}$



D. (d)  $\frac{[NO]^2 [Cl_2]}{[NOCl]^2}$

**Answer: A**

 [Watch Video Solution](#)



30. Consider the following equilibrium:



What is the relation between  $K_1$  and  $K_2$ ?

A. (a)  $K_1 = \frac{1}{K_2}$

B. (b)  $K_1 = \frac{1}{\sqrt{K_2}}$

C. (c)  $K_1 = K_2$


D. (d)  $K_1 = \frac{1}{K_2^2}$

**Answer: B**








**Watch Video Solution**

31. Consider the following reaction:



In the figure below, identify the curves  $X$ ,  $Y$ , and  $Z$  associated with the three species in the reaction



**Answer: A**



**Watch Video Solution**

32. Two equilibrium  $AB \rightleftharpoons A^+ + B^-$  and  $AB + B^- \rightleftharpoons AB_2^-$  are simultaneously maintained in a solution with equilibrium constants  $K_1$  and  $k_2$ , respectively. Ratio of  $[A^+]$  to  $[AB_2^-]$  in the solution is


- A. Directly proportional to  $[B^-]$
- B. Inversely proportional to  $[B^-]$
- C. Directly proportional to  $[B^-]^2$
- D. Inversely proportional to  $[B^-]^2$

Answer: D



Watch Video Solution

33. At 1400K,  $K_c = 2.5 \times 10^{-3}$  for the reaction



A 10L reaction vessel at 1400K contains 2.0mol of  $CH_4$ , 4.0mol of  $H_2S$ , 3.0mol of  $CS_2$ , 3.0mol of  $H_2$ . In which direction does the reaction proceed to reach equilibrium?

A. Forward

B. Backward


C. May be forward or backward

D. Reaction is in equilibrium

**Answer: B**

 [Watch Video Solution](#)


**34.** For the reaction




Which of the following sketches may represent above equilibrium?

Assume equilibrium can be achieved from either side and by taking any


one or more components initially. Give  $K_c$  for the reaction  $< 2$ ?




**A.**



B.



C.






D.



Watch Video Solution

35. a. For which of the following reactions,  $K_p$  is equal to  $K_c$  ?





b. For which of the following cases does the reaction go farthest to completion:

$K = 1$ ,  $K = 10^{10}$ ,  $K = 10^{-10}$ ?



Watch Video Solution

**36.** Both metals Mg and Fe can reduce copper metal from a solution having copper ions ( $\text{Cu}^{2+}$ ). According to the equilibria:




Which metal will remove cupric ion from the solution to a greater extent?




Watch Video Solution

**37.** The equilibrium constant of the reaction




at  $426^\circ\text{C}$  is 55.3, what will be the value of equilibrium constant

- if the reaction is reversed and
- if the given reaction is represented as



Watch Video Solution

38. What will be the effect on the equilibrium constant for the reaction





$\Delta H = -22.4$  kcal, when

- Pressure is increased
- Concentration of  $N_2$  is increased and
- Temperature is raised at equilibrium ?




Watch Video Solution

39. At a certain temperature, the following reactions have the equilibrium constants as shown below:



what is the equilibrium constant  $K_c$  for the reaction at the same temperature?



A. 0.78

B. 2.0

C. 16.2

D. 1.28

**Answer: B**



**Watch Video Solution**

**40.** For a reversible reaction



$\left(\frac{dx}{dt}\right) = 2.0 \times 10^3 \text{Lmol}^{-1}\text{s}^{-1}[A][B] - 1.0 \times 10^2 \text{s}^{-1}[C]$  where  $x$  is the amount of 'A' dissociated. The value of equilibrium constant ( $K_{eq}$ ) is

A. 10

B. 0.05

C. 20

D. Cannot be calculated





Answer: C



Watch Video Solution

41. The formation of ammonia from nitrogen and hydrogen gases can be written by the following two equations:



The two equations have equilibrium constants  $K_1$  and  $K_2$  respectively.

The relationship between the equilibrium constant is

A.  $K_1 = K_2^2$

B.  $K_1^3 = K_2^2$

C.  $K_1^{2/3} = K_2$

D.  $K_1 = K_2^{3/2}$

Answer: C::D




Watch Video Solution

**42.** The following concentrations were obtained for the formation of  $NH_3$  from  $N_2$  and  $H_2$  at equilibrium at 500K.  $[N_2] = 1.5 \times 10^{-2}M$ .  $[H_2] = 3.0 \times 10^{-2}M$  and  $[NH_3] = 1.2 \times 10^{-2}M$ . Calculate equilibrium constant.


 [Watch Video Solution](#)

**43.** At equilibrium the concentrations of  $N_2 = 3.0 \times 10^{-3}M$ ,  $O_2 = 4.2 \times 10^{-3}M$  and  $NO = 2.8 \times 10^{-3}M$  in a sealed vessel at 800K. What will be  $K_c$  for the reaction



 [Watch Video Solution](#)

**44.** In the reaction,



The concentration of  $H_2$ ,  $I_2$ , and  $HI$  at equilibrium are 8.0, 3.0 and 28.0 mol per  $L$  respectively. Determine the equilibrium constant.

 [Watch Video Solution](#)






**45.** 40% of a mixture of 0.2 mol of  $N_2$  and 0.6 mol of  $H_2$  react to give  $NH_3$  according to the equation :  $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$  at constant temperature and pressure. Then the ratio of the final volume to the initial volume of gases are :

- A. 4:5
- B. 5:4
- C. 7:10
- D. 8:5

**Answer: A**





 [Watch Video Solution](#)

46. Arrange the following in order of increasing tendency of the forward reactions to proceed towards completion at 298K and one atmospheric pressure :



 [Watch Video Solution](#)


47. The equilibrium constant of the dissociation of various of an element A are given at constant temperature:



Write the stability of these oxides in increasing order.

 [Watch Video Solution](#)

48. At



Assuming complete decomposition of  $\text{NH}_3$  and  $\text{N}_2\text{H}_4$

$$P = 0.3 \text{ atm}, P = 2.7 \text{ atm}$$

$$T = 300 \text{ K}, T = 200 \text{ K}$$

$V_L, V_L$





mole % of  $\text{NH}_3$  in original mixture is (assume both concentration same volume)

- A. 25 %
- B. 20 %
- C. 75 %
- D. 37.5 %

Answer: C



49. Write the expression for equilibrium constant for the following reactions. If the concentrations are expressed in  $\text{molL}^{-1}$ , give the units in each case.




a.



Watch Video Solution


50. At a certain temperature, the equilibrium constant ( $K_c$ ) is 16 for the reaction:



If we take one mole of each of the equilibrium concentration of  $NO$  and  $NO_2$ ?

 [Watch Video Solution](#)


**51.** A mixture of  $SO_3$ ,  $SO_2$  and  $O_2$  gases is maintained in a  $10L$  flask at a temperature at which the equilibrium constant for the reaction is 100:



- If the number of moles of  $SO_2$  and  $SO_3$  in the flask are equal. How many moles of  $O_2$  are present?
- If the number of moles of  $SO_3$  in flask is twice the number of moles of  $SO_2$ , how many moles of oxygen are present?

 [Watch Video Solution](#)


**52.** The value of  $K_c$  for the reaction



is 0.50 at 400 ° C. Find the value of  $K_p$  at 400 ° C when concentrations are expressed in  $\text{mol L}^{-1}$  and pressure in atm.

 [Watch Video Solution](#)

53. For an ideal gas reaction



the value of  $K_p$  will be:

A.  $K_p = \frac{n_C n_D}{n_A^2 n_B} \cdot \frac{V}{RT^2}$

B.  $K_p = \frac{n_C n_D}{n_A^2 n_B} \cdot \frac{V}{RT}$

C.  $K_p = \frac{n_C n_D}{n_A^2 n_B} \cdot \frac{RT}{V}$

D.  $K_p = \frac{n_C n_D}{4n_A^2 n_B} \cdot \frac{V}{RT}$

**Answer: B**

 [Watch Video Solution](#)



54. For a reaction



at equilibrium, the heat of reaction at constant volume is 1500 cal more than at constant pressure. If the temperature is  $27^\circ\text{C}$  then

A.  $K_p = K_c$

B.  $K_p > K_c$

C.  $K_p < K_c$

D. None of these





**Answer: B**



[Watch Video Solution](#)

55. Given that  $K_c$  for equation (i) given below has a value of 256 at 1000K.

Calculate the numerical values of  $K_c$  for other reactions (ii), (iii), and (iv).



i.



[Watch Video Solution](#)

56. When 3.06g of solid  $NH_4HS$  is introduced into a two-litre evacuated flask at  $27^\circ C$ , 30% of the solid decomposes into gaseous ammonia and hydrogen sulphide. (i) Calculate  $K_c$  and  $K_p$  for the reaction at  $27^\circ C$ . (ii) What would happen to the equilibrium when more solid  $NH_4HS$  is introduced into the flask?



[Watch Video Solution](#)

57. At 540K, 0.10mol of  $PCl_5$  is heated in a 8L flask. The pressure of equilibrium mixture is found to be 1.0atm. Calculate  $K_p$  and  $K_c$  for the


reaction.

 [Watch Video Solution](#)

**58.** Prove that the pressure necessary to obtain 50 % dissociation of  $PCl_5$  at  $250^\circ C$  is numerically three times of  $K_p$ .

 [Watch Video Solution](#)

**59.** For the reaction




Hydrogen gas is introduced into a five-litre flask at  $327^\circ C$ , containing 0.2 mol of  $CO(g)$  and a catalyst, until the pressure is  $4.92 atm$ . At this point, 0.1 mol of  $CH_3OH(g)$  is formed. Calculate the equilibrium constants  $K_p$  and  $K_c$ .

 [Watch Video Solution](#)

60. When sulphur in the form of  $S_8$  is heated at  $900K$ , the initial pressure of  $1 \text{ atm}$  falls by  $10\%$  at equilibrium. This is because of conversion of some  $S_8$  to  $S_2$ . Find the value of equilibrium constant for this reaction.

 [Watch Video Solution](#)

61. Two solid  $X$  and  $Y$  dissociate into gaseous products at a certain temperature as follows:




At a given temperature, the pressure over excess solid  $X$  is  $40\text{mm}$  and total pressure over solid  $Y$  is  $80\text{mm}$ . Calculate

- The value of  $K_p$  for two reactions.
- The ratio of moles of  $A$  and  $B$  in the vapour state over a mixture of  $X$  and  $Y$ .
- The total pressure of gases over a mixture of  $X$  and  $Y$ .

 [Watch Video Solution](#)

**62.** For a homogenous gaseous reaction



at 473K, the value of  $K_c = 0.35$  concentration units. When 2 moles of Y are mixed with 1 mole of X, at what pressure 60 % of X is converted to Z?

 [Watch Video Solution](#)

**63.** Solid  $NH_4HS(s)$  (ammonium hydrogen sulphate) dissociates to give  $NH_3(g)$  and  $H_2S(g)$  and is allowed to attain equilibrium at  $100^\circ C$ . If the value of  $K_p$  for its dissociation is found to be 0.34, find the total pressure at equilibrium at  $100^\circ C$ . If the value of  $K_p$  for its dissociation is found to be 0.34, find the total pressure at equilibrium and partial pressure of each component.

 [Watch Video Solution](#)


**64.** At 700K, the equilibrium constant  $K_p$  for the reaction



is  $1.80 \times 10^{-3} \text{ kPa}$ . What is the numerical value of  $K_c$  in moles per litre for this reaction at the same temperature?

 [Watch Video Solution](#)

65. The equilibrium of formation of phosgene is represented as :



The reaction is carried out in a  $500 \text{ mL}$  flask. At equilibrium,  $0.3 \text{ mol}$  of phosgene,  $0.1 \text{ mol}$  of  $\text{CO}$ , and  $0.1 \text{ mol}$  of  $\text{Cl}_2$  are present.

The equilibrium constant of the reaction is

- A. 30
- B. 15
- C. 5
- D. 25

**Answer: B**

 [Watch Video Solution](#)

66. Which of the following relation(s) holds good for gaseous and reversible reactions?

A.  $\frac{K_p}{K_c} = (RT)^{(\Delta n)_g}$

B.  $\frac{K_p}{K_c} = (P)^{(\Delta n)_g}$

C.  $\frac{K_c}{K_p} = \left(\frac{P}{RT}\right)^{(\Delta n)_g}$



D.  $\frac{K_c}{K_p} = (P)^{- (\Delta n)_g}$

**Answer: A:B**



**Watch Video Solution**

67. If two gases  $AB_2$  and  $B_2C$  are mixed, following equilibria are readily established:



If the reaction started only with  $AB_2$  with  $B_2C$ , then which of the following is necessarily true at equilibrium?

A.  $[AB_3]_{eq} = [BC]_{eq}$

B.  $[AB_2]_{eq} = [B_2C]_{eq}$

C.  $[AB_3]_{eq} > [B_3C_2]_{eq}$

D.  $[AB_3]_{eq} > [BC]_{eq}$

**Answer: C::D**



[Watch Video Solution](#)


**68.** The degree of dissociation of  $HI$  at a particular temperature is 0.8. Calculate the volume of  $2MNa_2S_2O_3$  solution required to neutralise the iodine present in an equilibrium mixture of a reaction when 2 mol each of  $H_2$  and  $I_2$  are heated in a closed vessel of 2L capacity and the equilibrium mixture is freezed.



[Watch Video Solution](#)




69. At 1000K, the pressure of iodine gas is found to be  $0.1\text{atm}$  due to partial dissociation of  $I_2(g)$  into  $I(g)$ . Had there been no dissociation, the pressure would have been  $0.07\text{atm}$ . Calculate the value of  $K_p$  for the reaction:



 [Watch Video Solution](#)


70. Calculate the percentage dissociation of  $H_2S(g)$  if 0.1 mol of  $H_2S$  is kept in a 0.5L vessel at 1000K. The value of  $K_c$  for the reaction



is  $1.0 \times 10^{-7}$ .

 [Watch Video Solution](#)

71. For the reaction



The degree of dissociation ( $\alpha$ ) of  $HI(g)$  is related to equilibrium constant

$K_p$  by the expression

a.  $\frac{1 + 2\sqrt{K_p}}{2}$ , b.  $\sqrt{\frac{1 + 2K_p}{2}}$

c.  $\sqrt{\frac{2K_p}{1 + 2K_p}}$ , d.  $\frac{2\sqrt{K_p}}{1 + 2\sqrt{K_p}}$

 [Watch Video Solution](#)

**72.** At certain temperature compound  $AB_2(g)$  dissociates according to the reaction



With degree of dissociation  $\alpha$  Which is small compared with unity, the expression of  $K_p$  in terms of  $\alpha$  and initial pressure P is :

 [Watch Video Solution](#)


**73.** For the dissociation reaction



the equilibrium constant  $K_p$  is 0.120 atm at 298K and total pressure of system is 2 atm. Calculate the degree of dissociation of  $N_2O_4$ .

 [Watch Video Solution](#)


**74.** A sample of air consisting of  $N_2$  and  $O_2$  was heated to 2500K until the equilibrium



was established with an equilibrium constant,  $K_c = 2.1 \times 10^{-3}$ . At equilibrium, the mole % of  $NO$  was 1.8. Estimate the initial composition of air in mole fraction of  $N_2$  and  $O_2$ .

 [Watch Video Solution](#)

**75.**  $PCl_5$  dissociates into  $PCl_3$  and  $Cl_2$ , thus




If the total pressure of the system in equilibrium is  $P$  at a density  $\rho$  and temperature  $T$ , show that the degree of dissociation  $\alpha = \frac{PM}{\rho RT} - 1$ , where

$M$  is the relative molar mass of  $PCl_5$ . If the vapour density of the gas mixture at equilibrium has the value of 62 when the temperature is  $230^\circ C$ , what is the value of  $P/\rho$ ?

 [Watch Video Solution](#)


**76.** The equilibrium constant  $K_p$  for the reaction



at  $497^\circ C$  is found to be  $636\text{mmHg}$ . If the pressure of the gas mixture is  $182\text{mm}$ , calculate the percentage dissociation of  $N_2O_4$ . At what pressure will it be dissociated?

 [Watch Video Solution](#)

**77.** For the reaction



The degree of dissociation ( $\alpha$ ) of  $HI(g)$  is related to equilibrium constant  $K_p$  by the expression

$$\text{a. } \frac{1 + 2\sqrt{K_p}}{2}, \text{ b. } \sqrt{\frac{1 + 2K_p}{2}}$$

$$\text{c. } \sqrt{\frac{2K_p}{1 + 2K_p}}, \text{ d. } \frac{2\sqrt{K_p}}{1 + 2\sqrt{K_p}}$$

$$\text{A. } \frac{1 + 2\sqrt{K_p}}{2\sqrt{K_p}}$$

$$\text{B. } \sqrt{\frac{1 + 2K_p}{2}}$$

$$\text{C. } \sqrt{\frac{2K_p}{1 + 2K_p}}$$

$$\text{D. } \frac{2\sqrt{K_p}}{1 + 2\sqrt{K_p}}$$



Watch Video Solution

**78.** At a given temperature and a total pressure of 1.0 atm for the homogenous gaseous reaction




the partial pressure of  $NO_2$  is 0.5atm.

a. Find the value of  $K_p$ .

b. If volume of the vessel is decreased to half of its original volume, at constant temperature, what are the partial pressures of the components of the equilibrium mixture?


 [Watch Video Solution](#)

79. In the following reactions, the system will shift towards the forward reaction by adding inert gas at constant pressure?



 [Watch Video Solution](#)

80.  $N_2O_4$  dissociates as




At  $40^\circ C$  and one atmosphere % decomposition of  $N_2O_4$  is 50.3%. At what pressure and same temperature, the equilibrium mixture has the ratio of  $N_2O_4: NO_2$  as 1:8?



 Watch Video Solution

**81.** At  $627^{\circ}\text{C}$  and  $1\text{ atm}$   $\text{SO}_3$  is partially dissociated into  $\text{SO}_2$  and  $\text{O}_2$  by the reaction




The density of the equilibrium mixture is  $0.925\text{gL}^{-1}$ . What is the degree of dissociation?




Watch Video Solution

**82.** Density of equilibrium mixture of  $\text{N}_2\text{O}_4$  and  $\text{NO}_2$  at  $1\text{ atm}$  and  $384\text{K}$  is  $1.84\text{gdm}^{-3}$ . Calculate the equilibrium constant of the reaction.



Watch Video Solution

**83.** For the reaction



Show that the degree of dissociation of  $NH_3$  is given as

$$\alpha = \left[ 1 + \frac{3\sqrt{3}}{4} \frac{p}{K_p} \right]^{-1/2}$$

where  $p$  is equilibrium pressure. If  $K_p$  of the above reaction is  $78.1 \text{ atm}$  at  $400^\circ \text{ C}$ , calculate  $K_c$ .

 [Watch Video Solution](#)

**84.** The vapour density of  $N_2O_4$  at a certain temperature is 30. Calculate the percentage dissociation of  $N_2O_4$  this temperature.

 [Watch Video Solution](#)

**85.**  $3g$  mol of phosphorus is heated in a flask of  $4L$  volume. At equilibrium, it dissociates to give  $40\%$  of phosphorus trichloride and chlorine. Calculate the equilibrium constant.

 [Watch Video Solution](#)



86.  $N_2O_4$  is 25% dissociated at  $37^\circ C$  and one atmosphere pressure.

Calculate (i)  $K_p$  and (ii) the percentage dissociation at 0.1 atm and  $37^\circ C$ .

 [Watch Video Solution](#)

87. The equation  $\alpha = \frac{D - d}{(n - 1)d}$  is correctly matched for: ( $\alpha$  is the degree of dissociation,  $D$  and  $d$  are the vapour densities before and after dissociation, respectively).

A. 

B. 

C. 

D. 

**Answer: B**

 [Watch Video Solution](#)

88. The vapour density of the equilibrium mixture of the reaction:



is 50. The percent dissociation of  $SO_2Cl_2$  is

A. 33.00

B. 35.0

C. 30.0


D. 66.00

**Answer: B**



[Watch Video Solution](#)

89. Consider the following equilibrium in a closed container:



At a fixed temperature, the volume of the reaction container is halved.

For this change which of the following statements holds true regarding

the equilibrium constant ( $K_p$ ) and the degree of dissociation ( $\alpha$ )?

- A. Neither  $K_p$  nor alpha changes
- B. Both  $K_p$  and alpha change
- C.  $K_p$  does not change but alpha changes
- D.  $K_p$  changes, but alpha does not change

**Answer: D**

 [Watch Video Solution](#)

**90.** At certain temperature compound  $AB_2(g)$  dissociates according to the reaction



With degree of dissociation  $\alpha$  Which is small compared with unity, the expression of  $K_p$  in terms of  $\alpha$  and initial pressure P is :


- A.  $256P^3\alpha^5$
- B.  $4P\alpha^2$
- C.  $8P^3\alpha^5$

D. None of these

Answer: C

 Watch Video Solution

91. The following reaction occurs at 700K. Arrange them in the order of increasing tendency to proceed to completion.



I.

A.  $II < I < IV < III$

B.  $III < IV < I < II$

C.  $I < III < IV < II$


D.  $IV < III < I < II$

Answer: B



Watch Video Solution

92. At  $727^\circ\text{C}$  and  $1.2\text{atm}$  of total equilibrium pressure,  $\text{SO}_3$  is partially dissociated into  $\text{SO}_2$  and  $\text{O}_2$  as:



The density of equilibrium mixture is  $0.9\text{g/L}$ . The degree of dissociation is:,  $\left[ \text{Use } R = 0.08\text{atmLmol}^{-1}\text{K}^{-1} \right]$

A.  $\frac{1}{3}$

B.  $\frac{2}{3}$

C.  $\frac{1}{4}$

D.  $\frac{1}{4}$

Answer: B



Watch Video Solution

93.  $K_p$  for the reaction



at  $250^\circ C$  is 0.82. Calculate the degree of dissociation at given temperature under a total pressure of  $5\text{atm}$ . What will be the degree of dissociation if the equilibrium pressure is  $10\text{atm}$ , at same temperature.

 [Watch Video Solution](#)

94. In reaction:



if the initial pressure of  $CH_3COCH_3(g)$  is  $150\text{mm}$  and at equilibrium the mole fraction of  $CO(g)$  is  $\frac{1}{3}$ , then the value  $K_p$  is

- A.  $50\text{mm}$
- B.  $100\text{mm}$
- C.  $33.3\text{mm}$
- D.  $75\text{mm}$


**Answer: A**

 [Watch Video Solution](#)

**95.** When  $PCl_5$  is heated, it dissociates into  $PCl_3$  and  $Cl_2$ . The vapour density of the gas mixture at  $200^\circ C$  and at  $250^\circ C$  is 70 and 58, respectively. Find the degree dissociation at two temperatures.

 [Watch Video Solution](#)

**96.** 0.25 mol of  $CO$  taken in a 1.5L flask is maintained at 500K along with a catalyst so that the following reaction can take place:



Hydrogen is introduced until the total pressure of system is  $8.2\text{atm}$ , at equilibrium, and  $0.1\text{mol}$  of methanol is formed. Calculate

a.  $K_p$  and  $K_c$


b. The final pressure if the same amount of  $CO$  and  $H_2$  as before are used but no catalyst so that the reaction does take place.

 [Watch Video Solution](#)

97. Ammonia under a pressure of  $15\text{atm}$ , at  $27^\circ\text{C}$  is heated to  $327^\circ\text{C}$  in a vessel in the presence of catalyst. Under these conditions,  $\text{NH}_3$  partially decomposes to  $\text{H}_2$  and  $\text{N}_2$ . The vessel is such that the volume remains effectively constant, whereas the pressure increases to  $50\text{atm}$ . Calculate the percentage of  $\text{NH}_3$  actually decomposed.

 [Watch Video Solution](#)

98. Solid Ammonium carbamate dissociates as:



In a closed vessel, solid ammonium carbonate is in equilibrium with its dissociation products. At equilibrium, ammonia is added such that the partial pressure of  $\text{NH}_3$  at new equilibrium now equals the original total pressure. Calculate the ratio of total pressure at new equilibrium to that of original total pressure. Also find the partial pressure of ammonia gas added.



 [Watch Video Solution](#)

99. The degree of dissociation of  $I_2$  molecule of  $1000^\circ C$  and under atmospheric pressure is 40% by volume. If the dissociation is reduced to 20% at the same temperature, total equilibrium pressure on the gas is:

A.  $1.57\text{atm}$

B.  $2.57\text{atm}$

C.  $3.57\text{atm}$

D.  $4.57\text{atm}$

**Answer: D**

 [Watch Video Solution](#)

100.  $I_2(aq) + I^-(aq) \rightleftharpoons (aq)$ . We started with 1 mole of  $I_2$  and 0.5 mole of  $I^-$  in one litre flask. After equilibrium is reached, excess of  $AgNO_3$  gave 0.25 mole of yellow precipitate. Equilibrium constant is :

A. 1.33

B. 2.66


C. 2.00

D. 3.00

**Answer: A**

 [Watch Video Solution](#)


**101.** At  $25^{\circ}\text{C}$  and 1 atm,  $\text{N}_2\text{O}_4$  dissociates the reaction



If it is 35% dissociated at given condition, find the volume of above mixture will difuse if 20mL of pure  $\text{O}_2$  diffuse 10 minutes at same yemperature and pressure.

 [Watch Video Solution](#)

**102.** For the reaction



Show that the degree of dissociation of  $NH_3$  is given as

$$\alpha = \left[ 1 + \frac{3\sqrt{3}}{4} \frac{p}{K_p} \right]^{-1/2}$$

where  $p$  is equilibrium pressure. If  $K_p$  of the above reaction is  $78.1 \text{ atm}$  at  $400^\circ \text{C}$ , calculate  $K_c$ .

 [Watch Video Solution](#)

**103.** For the formation of ammonia the equilibrium constant data at  $673\text{K}$  and  $773\text{K}$ , respectively, are  $1.64 \times 10^{-4}$  and  $1.44 \times 10^{-5}$  respectively.

Calculate heat of reaction ( $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )

 [Watch Video Solution](#)

**104.** For the reaction




K is 0.63 at 700 ° C and 1.66 at 1000 ° C.

a. What is the average  $\Delta H^\ominus$  for the temperature range considered?

b. What is the value of K at 800 ° C?

 [Watch Video Solution](#)




**105.** The value of K for the reaction



Changed from 0.096 at 298K to 1.4 at 373K. Above what temperature will the reaction become thermodynamically spontaneous in the forward direction assuming that  $\Delta H^\ominus$  and  $\Delta S^\ominus$  values for the reaction do not change with change in temperature? Given that  $\Delta S_{298}^\ominus = 10.296JK^{-1}$ .

 [Watch Video Solution](#)

106. Which of the following graph represents exothermic reaction ?



A. 

B. 

C. 

D. 

Answer: D

 Watch Video Solution

107. A reversible reaction is endothermic in forward direction. Then which of the following is (are) correct?

A.  $\ln K$  vs  $1/T$  will be a straight line with negative slope

B.  $\frac{d}{dT} \ln K > 0$

C. A plot of  $d \ln K$  against  $1/T^2$  will have positive slope

D. An increase in temperature will shift the reaction in the forward direction.

Answer: A::B::C



Watch Video Solution

108. The activation energy of

$H_2 + I_2 \rightleftharpoons 2HI(g)$  in equilibrium for the forward reaction is  $167 \text{kJmol}^{-1}$

whereas for the reverse reaction is  $180 \text{kJmol}^{-1}$ . The presence of catalyst

lowers the activation energy by  $80 \text{kJmol}^{-1}$ . Assuming that the reactions

are made at  $27^\circ\text{C}$  and the frequency factor for forward and backward reactions are  $4 \times 10^{-4}$  and  $2 \times 10^{-3}$  respectively, calculate  $K_c$ .

 [Watch Video Solution](#)

**109.** Variation of equilibrium constant  $K$  with temperature is given by van't Hoff equation

$$\ln K = \frac{\Delta_r S^\circ}{R} - \frac{\Delta_r H^\circ}{RT}$$

for this equation,  $(\Delta_r H^\circ)$  can be evaluated if equilibrium constants  $K_1$  and  $K_2$  at two temperatures  $T_1$  and  $T_2$  are known.

$$\log\left(\frac{K_2}{K_1}\right) = \frac{\Delta_r H^\circ}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

Select the correct statement :

A.  $\log \frac{K_2}{K_1} = - \frac{\Delta H}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$

B.  $\log \frac{K_2}{K_1} = \frac{\Delta H}{2.303R} \left[ \frac{1}{T_2} - \frac{1}{T_1} \right]$

C.  $\log \frac{K_2}{K_1} = - \frac{\Delta H}{2.303R} \left[ \frac{1}{T_2} - \frac{1}{T_1} \right]$

D. None of the above

 [Watch Video Solution](#)

110. It is known that the heat is needed to dissociate ammonia into  $N_2$  and  $H_2$ . For the reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$ ,  $K_f$  is the velocity constant for forward reaction and  $K_b$  is velocity constant for backward reaction,  $K_c$  is equilibrium constant for the reaction shown. Then  $\frac{dk_f}{dT}$  (where T is symbol for absolute temp.):





- A. Is greater than  $dk_b/dT$
- B. Is less than  $dk_b/dT$
- C. Is equal to  $dk_b/dT$
- D. Cannot be compared with  $dk_b/dT$

**Answer: B**

 [Watch Video Solution](#)



111. For the chemical equilibrium  $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ ,  $\Delta_r H^\circ$  can be determined from which one of the following plots?



Answer: A



**112.** Solubility of a solute in water is dependent on temperature as given by



$$S = Ae^{-\Delta H/RT}, \text{ where } \Delta H = \text{heat of solution}$$



For given solution, variation of  $\log S$  with temperature is shown

graphically. Hence, solution is

$\log s$  with temperature is



**Answer: D**

113. In the preparation of CaO from  $\text{CaCO}_3$  using the equilibrium,



$K_p$  is expressed as

$$\log K_p = 7.282 - \frac{8500}{T}$$

For complete decomposition of  $\text{CaCO}_3$ , the temperature in celsius to be used is:

A. 1167


B. 894

C. 8500

D. 850

**Answer: B**

**114.** The partial pressure of  $CO_2$  in the reaction





is 0.773 mm at  $500^\circ C$ . Calculate  $K_p$  at  $600^\circ C$  for the above reaction,  $\Delta H$  of the reaction is 43.2 kcal per mole and does not change in the given range of temperature.


 [Watch Video Solution](#)

**115.** For the reaction  $Br_2 \rightleftharpoons 2Br$ , the equilibrium constants at  $327^\circ C$  and  $527^\circ C$  are, respectively,  $6.1 \times 10^{-12}$  and  $1.0 \times 10^{-7}$ . What is the nature of the reaction?

 [Watch Video Solution](#)

**116.** From the following data






Show whether reaction (iii) is exothermic or endothermic.

 [Watch Video Solution](#)

**117.** The equilibrium constant  $K_p$  for the reaction,




is  $1.64 \times 10^{-4}$  at  $400^\circ\text{C}$  and  $0.144 \times 10^{-4}$  at  $500^\circ\text{C}$ . Calculate the mean heat of formation of 1 mol of  $\text{NH}_3$  from its elements in this temperature range.

 [Watch Video Solution](#)

**118.** For the reaction  $2\text{NOCl}(g) \rightleftharpoons 2\text{NO}(g) + \text{Cl}_2(g)$ , the equilibrium constant is  $2.8 \times 10^{-5}$  at  $300\text{K}$  and  $7.0 \times 10^{-1}$  at  $400\text{K}$ . What is the activation energy for the reaction?

 [Watch Video Solution](#)

119. A schematic plot of  $\log K_{eq}$  vs inverse of temperature for a reaction is shown in the figure. The reaction must be:



- A. Exothermic
- B. Endothermic
- C. One with negligible enthalpy change
- D. Highly spontaneous at ordinary temperature



Watch Video Solution

**120.** For the reaction



K is 0.63 at 727 °C and 1.26 at 927 °C.

a. What is the average  $\Delta H$  for the temperature range considered? [Use

$\log 2=0.3$ ]


b. What is the value of K at 1227 °C?

 [Watch Video Solution](#)

**121.** The equilibrium constant  $K_p$ , for the reaction  $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g)$  is  $1.6 \times 10^{-4}$  at 400 °C. What will be the equilibrium constant at 500 °C if the heat of reaction in this temperature range is -25.14 kcal?

 [Watch Video Solution](#)

**122.** The equilibrium constant for the reaction






is 18.5 at 925K and 9.25 at 1000K, respectively. Calculate the enthalpy of the reaction.

 [Watch Video Solution](#)


**123.** Consider the reaction



at 375 ° C, the value of equilibrium constant for the reaction is 0.0032. It was observed that the concentration of the three species is  $0.050\text{molL}^{-1}$  each at a certain instant. Discuss what will happen in the reaction vessel?

 [Watch Video Solution](#)

**124.** Consider the reaction



When the system is at equilibrium at 100 °, the concentrations are found to be  $[X] = 0.2M$ ,  $[Y] = [Z] = 0.4M$

a. If the pressure of the container is suddenly halved at 100 ° C, find

equilibrium concentration.

b. If the pressure of the container is suddenly doubled at  $100^\circ\text{C}$ , find the equilibrium concentration


 [Watch Video Solution](#)

**125.** The value of  $K_c$  for the reaction

$2A \rightleftharpoons B + C$  is  $2 \times 10^{-3}$ . At a given time, the composition of reaction is  $[A] = [B] = [C] = 3 \times 10^{-4}\text{M}$ . In which direction the reaction will proceed?

 [Watch Video Solution](#)


**126.** The value of  $K_c$  for the reaction:



at  $100^\circ\text{C}$  is 49. If 1.0L flask containing one mole of  $A_2$  is connected with a 2.0L flask containing one mole of  $B_2$ , how many moles of AB will be formed at  $100^\circ\text{C}$ ?

 [Watch Video Solution](#)

**127.** The value of  $K_c$  for the reaction



is 64 at 773K. If one "mole" of  $H_2$ , one mole of  $I_2$ , and three moles of HI are taken in a 1L flask, find the concentrations of  $I_2$  and HI at equilibrium at 773K.

 [Watch Video Solution](#)

**128.** In a 1.0L aqueous solution when the reaction



reaches equilibrium,  $[Cu^{2+}] = xM$  and  $[Ag^{\oplus}] = yM$ .

reaches equilibrium,  $[Cu^{2+}] = xM$  and  $[Ag^{\oplus}] = yM$ .

If the volume of solution is doubled by adding water, then at equilibrium:

A.  $[Cu^{2+}] = \frac{x}{2}M$ ,  $[Ag^{\oplus}] = \frac{y}{2}M$

B.  $[Cu^{2+}] > \frac{x}{2}M$ ,  $[Ag^{\oplus}] > \frac{y}{2}M$

C.  $[Cu^{2+}] < \frac{x}{2}M$ ,  $[Ag^{\oplus}] > \frac{y}{2}M$

$$D. [Cu^{2+}] < \frac{x}{2}M, [Ag^{\oplus}] < \frac{y}{2}M$$




Watch Video Solution

**129.**  $H_2$  and  $I_2$  are mixed at  $400^\circ C$  in a  $1.0L$  container, and when equilibrium is established, the following concentrations are present:  $[HI] = 0.8M$ ,  $[H_2] = 0.08M$ , and  $[I_2] = 0.08M$ . If now an additional  $0.4$  mol of  $HI$  is added, what are the new equilibrium concentrations, when the new equilibrium  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$  is re-established?



Watch Video Solution

**130.** At  $448^\circ C$ , the equilibrium constant ( $K_c$ ) for the reaction



is  $50.5$ . Predict the direction in which the reaction will proceed to reach equilibrium at  $448^\circ C$ , if we start with  $2.0 \times 10^{-2}$  mol of  $HI$ ,  $1.0 \times 10^{-2}$  mol of  $H_2$  and  $3.0 \times 10^{-2}$  mol of  $I_2$  in a  $2.0L$  container.




[Watch Video Solution](#)

**131.** The value of  $\Delta G^\ominus$  for the phosphorylation of glucose in glycolysis is  $13.8\text{kJ/mol}$ . Find the value of  $K_c$  at  $298\text{K}$ .



[Watch Video Solution](#)

**132.** Hydrolysis of sucrose gives.



Equilibrium constant  $K_c$  for the reaction is  $2 \times 10^{13}$  at  $300\text{K}$ . Calculate  $\Delta G^\ominus$  at  $300\text{K}$ .



[Watch Video Solution](#)


**133.** If  $K_c$  is not numerically equal to  $K_p$ , how can both of the following equations be valid?

$$\Delta G^\ominus = -2.303RT \log K_c, \Delta G^\ominus = -2.303RT \log K_p$$



[Watch Video Solution](#)

**134.** The value of  $K_p$  at 298K for the reaction



is found to be 826.0, partial pressure being measured atmospheric units.

Calculate  $\Delta G^\ominus$  at 298K.

 [Watch Video Solution](#)

**135.** For the reaction,




Calculate the standard equilibrium constant at 298K. Given that the value of  $\Delta H^\ominus$  and  $\Delta S^\ominus$  of the reaction at 298K are  $77.2kJmol^{-1}$  and  $122JK^{-1}mol^{-1}$ .

 [Watch Video Solution](#)

136.  $\Delta G^\ominus$  for  $\frac{1}{2}N_2(g) + \frac{3}{2}H_2(g) \rightleftharpoons NH_3(g)$  is  $-16.5kJmol^{-1}$ . Find out  $K_p$  for the reaction at  $25^\circ C$ .

 [Watch Video Solution](#)

137. In the reaction equilibrium



When 5 mol of each is taken and the temperature is kept at  $298K$ , the total pressure was found to be 20 bar.

Given :  $\Delta_f G_{N_2O_4}^\ominus = 100kJ$ ,  $\Delta_f G_{NO_2}^\ominus = 50KJ$

a. Find  $\Delta G^\ominus$  of the reaction at  $298K$ .

b. Find the direction of the reaction.

 [Watch Video Solution](#)

138. A large positive value of  $\Delta G^\ominus$  corresponds to which of these?

A. Small positive  $K$

B. Small negative  $K$

C. Large positive  $K$

D. Large negative  $K$

**Answer: A**



**Watch Video Solution**

**139.** For the reaction



in a closed flask, the equilibrium pressure is  $P$  atm. The standard free energy of the reaction would be:

A.  $-RT \ln p$

B.  $-RT(\ln p - \ln 2)$

C.  $-2RT \ln p$

D.  $-2RT(\ln p - \ln 2)$



**Answer: D**

 [Watch Video Solution](#)

**140.**  $\Delta G^\ominus$  for the reaction  $X + Y \rightleftharpoons C$  is  $-4.606\text{kcal}$  at  $1000\text{K}$ . The equilibrium constant for the reverse mode of the reaction will be:

A. 100

B. 10


C. 0.01

D. 0.1

**Answer: D**

 [Watch Video Solution](#)

**141.** For the following reaction:  $K = 1.7 \times 10^7$  at  $25^\circ\text{C}$



What is the value of  $\Delta G^\ominus$  in kJ ?

 [Watch Video Solution](#)

142. In an equilibrium reaction for which  $\Delta G^\ominus = 0$ , the equilibrium constant  $K$  should be equal to :

A. Zero

B. 10


C. 1

D. 2

**Answer: C**

 [Watch Video Solution](#)

143. What is  $\Delta G^\ominus$  for the following reaction?



A.  $-26.5\text{kJmol}^{-1}$

B.  $11.5\text{kJmol}^{-1}$

C.  $-2.2\text{kJmol}^{-1}$

D.  $-0.97\text{kJmol}^{-1}$

**Answer: A**



**Watch Video Solution**

**144.** If the  $E_{\text{cell}}^{\circ}$  for a given reaction has a negative value, which of the following gives correct relationships for the value of  $\Delta G^{\circ}$  and  $K_{\text{eq}}$ ?

A.  $\Delta G^{\circ} > 0, K_{\text{eq}} < 1$

B.  $\Delta G^{\circ} > 0, K_{\text{eq}} > 1$





C.  $\Delta G^{\circ} < 0, K_{\text{eq}} > 1$

D.  $\Delta G^{\circ} < 0, K_{\text{eq}} < 1$

**Answer: A**



145. Which of the following graph correctly represent for equilibrium reaction whose  $K_p > 1$ ?



**Answer: A**

 [Watch Video Solution](#)

**146.** The equilibrium constant  $K_p$  for the homogeneous reaction is  $10^{-3}$ .

The standard Gibbs free energy change  $\Delta G^\ominus$  for the reaction at

$27^\circ\text{C}$  (using  $R = 2\text{calK}^{-1}\text{mol}^{-1}$ ) is

A. Zero

B.  $-1.8\text{kcal}$


C.  $-4.145\text{kcal}$

D.  $+4.145\text{kcal}$

**Answer: D**

 [Watch Video Solution](#)

147. The free energy of formation of NO is  $78\text{kJmol}^{-1}$  at the temperature of an automobile engine (1000K). What is the equilibrium constant for this reaction at 1000K?



A.  $8.4 \times 10^{-5}$

B.  $7.1 \times 10^{-9}$

C.  $4.2 \times 10^{-10}$

D.  $1.7 \times 10^{-19}$

**Answer: A**



[Watch Video Solution](#)

148. The densities of graphite and diamond are  $22.5$  and  $3.51\text{ gm cm}^{-3}$ . The  $\Delta_f G^\ominus$  values are  $0\text{Jmol}^{-1}$  and  $2900\text{Jmol}^{-1}$  for graphite and diamond, respectively. Calculate the equilibrium pressure for the conversion of graphite into diamond at 298K.




Watch Video Solution

**149.** Calculate the pressure of  $CO_2$  gas at  $700K$  in the heterogenous equilibrium reaction  $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ , if  $\Delta G^\ominus$  for this reaction is  $130.2kJmol^{-1}$ .



Watch Video Solution

**150.** For the equilibrium



$\Delta G^\ominus (Jmol^{-1}) = -20700 - 11.97T$ . Calculate the temperature at which the product gases at equilibrium at 1 atm will contain 400 ppm of carbon monoxide.



Watch Video Solution

151.  $K_c$  for the reaction  $N_2O_4 \rightleftharpoons 2NO_2$  in chloroform at 291K is 1.14. Calculate the free energy change of the reaction when the concentration of the two gases are  $0.5 \text{ mol dm}^{-3}$  each at the same temperature.

$$\left( R = 0.082 \text{ LatmK}^{-1} \text{ mol}^{-1} \right)$$



Watch Video Solution

152. A reaction mixture containing  $H_2$ ,  $N_2$  and  $NH_3$  has partial pressures 1 atm, 2 atm, and 3 atm. Respectively, at 725K. If the value of  $K_p$  for the reaction,  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$  is  $4.28 \times 10^{-5} \text{ atm}^{-2}$  at 725K, in which direction the net reaction will go?

- A. Forward
- B. Backward
- C. No net reaction
- D. Direction of reaction cannot be predicted.



Watch Video Solution



153. i. The initial pressure of  $PCl_5$  present in one litre vessel at  $200K$  is 2 atm. At equilibrium the pressure increases to 3 atm with temperature increasing to 250. The percentage dissociation of  $PCl_5$  at equilibrium is

A. 30 %

B. 60 %

C. 0.2 %

D. 20 %

**Answer: D**

 Watch Video Solution

154. One mole of  $N_2O_4(g)$  at  $100K$  is kept in a closed container at 1.0 atm pressure. It is heated to  $300K$ , where 30 % by mass of  $N_2O_4(g)$  decomposes to  $NO_2(g)$ . The resultant pressure will be

A.  $3.9\text{atm}$

B.  $1.95\text{atm}$

C.  $1.0\text{atm}$

D.  $3.0\text{atm}$

 [Watch Video Solution](#)

**155.** The density of an equilibrium mixture of  $N_2O_4$  and  $NO_2$  at 1 atm is  $3.62\text{gL}^{-1}$  at 288K and  $1.84\text{gL}^{-1}$  at 348K. Calculate the entropy change during the reaction at 348K.

 [Watch Video Solution](#)

**156.** Which of the following conditions help melting of ice?

A. High pressure, temperature below  $0^\circ\text{C}$

- B. High pressure, temperature above  $0^{\circ}\text{C}$
- C. Low pressure, temperature above  $0^{\circ}\text{C}$
- D. Low pressure, temperature below  $0^{\circ}\text{C}$

**Answer: B**

 [Watch Video Solution](#)

**157.** Densities of diamond and graphite are  $3.5$  and  $2.3\text{gmL}^{-1}$ , respectively.

The increase of pressure on the equilibrium  $C_{\text{diamond}} \rightleftharpoons C_{\text{graphite}}$

- A. Favours backward reaction
- B. Fovours forward reaction
- C. Have no effect
- D. Increases the reaction rate

**Answer: C**

 [Watch Video Solution](#)

158.  $K_p$  for an endothermic chemical reaction is 10 atm. Then backward reaction is favoured at

- A. High pressure, high temperature
- B. High pressure, low temperature
- C. Low pressure, high temperature
- D. Low pressure, low temperature

**Answer: C**



[Watch Video Solution](#)

159. For the following reaction, the value of  $K$  change with



- A. Change in pressure
- B. Change in concentration of oxygen





C. concentration of  $NO(g)$

D. Change in temperature

**Answer: D**

 [Watch Video Solution](#)

**160.** Which among the following reactions is favoured in forward direction by increase of temperature?



**Answer: B**

 [Watch Video Solution](#)

161. The exothermic formation of  $ClF_3$  is represented by the equation:



Which of the following will increase the quantity of  $ClF_3$  in an equilibrium mixture of  $Cl_2$ ,  $F_2$ , and  $ClF_3$ ?


- A. Increasing the temperature
- B. Removing  $Cl_2$
- C. Increasing the volume of the container
- D. Adding  $F_2$

Answer: D




Watch Video Solution

162. For the following reaction through stages I, II and III



quantity of the product formed (x) varies with temperature (T) as given.

Select the correct statement.






- A. Stages I and III are endothermic but II is exothermic.
- B. Stages I and III are exothermic but II is endothermic
- C. Stages II and III are exothermic but I is endothermic
- D. Stage I is exothermic but stages II and III are endothermic.

**Answer: A**



[Watch Video Solution](#)

**163.** Which among the following reactions will be favoured at low pressure?




**Answer: C**



[Watch Video Solution](#)

**164.** Consider the following reversible reaction at equilibrium:



Which one of the following changes in conditions will lead to maximum decomposition of  $H_2O(g)$ ?

A. Increasing both temperature and pressure



- B. Decreasing temperature and increasing pressure
- C. Increasing temperature and decreasing pressure
- D. Decreasing temperature and decreasing pressure

**Answer: C**


 [Watch Video Solution](#)

**165.** A gas  $X(g)$  is when dissolved in water heat is evolved. Then solubility of  $X$  will increase :

- A. Low pressure, high temperature
- B. Low pressure, low temperature
- C. High pressure, high temperature
- D. High pressure, low temperature

**Answer: D**

 [Watch Video Solution](#)



above mentioned equilibrium is favoured at

- A. High pressure, low temperature
- B. High pressure, high temperature
- C. Low pressure, high temperature
- D. Low pressure, low temperature

**Answer: C**



[Watch Video Solution](#)

167. What is the direction of a reversible reaction when one of the products of the reaction is removed?

- A. The reaction moves towards right hand side.
- B. The reaction moves towards left hand side

C. The reaction moves towards both hand side

D. The reaction stops.

**Answer: A**

 [Watch Video Solution](#)

**168.** According to Le- Chatelier's principle. Adding heat to a solid  $\rightleftharpoons$  liquid equilibrium will cause the.

A. Amount of solid to decrease

B. Amount of liuid to decrease

C. Temperature to rise

D. Temperature to fall

**Answer: A**

 [Watch Video Solution](#)

**169.** The equilibrium constant for the reaction,  $A + B \rightleftharpoons C + D$  is 2.85 at room temperature and  $1.4 \times 10^{-2}$  at 698K. This shows that the forward reaction is

- A. Exothermic
- B. Endothermic
- C. Unpredictable
- D. There is no relationship between  $\Delta H$  and  $K$ .

**Answer: A**

 [Watch Video Solution](#)

**170.** Le - Chatelier principle is not applicable to :

- A. Only homogeneous chemical reversible reactions
- B. Only heterogeneous chemical reversible reactions
- C. Only physical equilibria

D. All system, chemical or physical, in equilibrium

**Answer: D**

 [Watch Video Solution](#)

**171.** Solubility of a gas in liquid increases on

- A. Addition of a catalyst
- B. Decreaseing of pressure
- C. Increasing of pressure
- D. Increasing of temperature

**Answer: C**

 [Watch Video Solution](#)

172. When any system in equilibrium is subjected to a change in pressure, concentration, or temperature, the equilibrium is shifted in the direction which tends to undo the effect of the change. This statement is known as

- A. First law of thermodynamics
- B. Le Chatelier's principle
- C. Hess's law
- D. Ostwald's law

**Answer: B**



[Watch Video Solution](#)

173. The equilibrium constant for the reaction  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$  is  $4.0 \times 10^{-4}$  at  $2000K$ . In the presence of a catalyst, the equilibrium is attained 10 times faster. Therefore, the equilibrium constant in presence of the catalyst at  $2000K$  is

A.  $4 \times 10^{-4}$

B.  $40 \times 10^{-4}$

C.  $4 \times 10^{-2}$

D. Difficult to compute without more data

**Answer: A**

 [Watch Video Solution](#)

**174.** When KOH is dissolved in water, heat is evolved. If the temperature is raised, the solubility of KOH

A. Increases

B. Decreases

C. Remains the same





D. Cannot be predicted

**Answer: B**



Watch Video Solution

175. Le - Chatelier principle is not applicable to :




Answer: A



Watch Video Solution

176. Consider the reaction



in closed container at equilibrium. What would be the effect of addition of  $CaCO_3$  on the equilibrium concentration of  $CO_2$ ?



- A. Increase
- B. Decreases
- C. Remains unaffected
- D. Data is not sufficient to predict it

**Answer: C**

 [Watch Video Solution](#)

**177.** The equilibrium constant for a reaction  $A + B \rightleftharpoons C + D$  is  $1 \times 10^{-2}$  at  $298K$  and is  $2$  at  $273K$ . The chemical process resulting in the formation of  $C$  and  $D$  is

- A. Exothermic
- B. Endothermic
- C. Unpredictable
- D. There is no relationship between  $\Delta H$  and  $K$ .

**Answer: A**



[Watch Video Solution](#)

**178.** In a flask colourless  $N_2O_4$  is in equilibrium with brown coloured  $NO_2$ . At equilibrium, when the flask is heated at  $100^\circ C$  the brown colour deepens and on cooling it becomes less coloured. The change in enthalpy,  $\Delta H$  for this system is



- A. Negative
- B. Positive
- C. Zero
- D. Not defined

**Answer: A**



[Watch Video Solution](#)

179. Consider the following equilibria:



Which of the above will be disturbed by an increase in pressure?

A. II

B. I, II

C. I, II, III,


D. None of these

**Answer: D**



**Watch Video Solution**

180. The following two reactions:



are simultaneously in equilibrium in a container at constant volume. A

few moles of  $CO(g)$  are later introduced into the vessel. After some time, the new equilibrium concentration of

A.  $PCl_5$  will remain unchanged

B.  $Cl_2$  will be greater

C.  $PCl_5$  will become greater

D.  $PCl_5$  will become less

**Answer: D**

 [Watch Video Solution](#)

**181.** The oxidation of  $SO_2$  by  $O_2$  to  $SO_3$  is an exothermic process. The yield of  $SO_3$  is maximum if

i. Temperature is increased and pressure is kept constant

ii. Temperature is reduced and pressure is kept constant

iii. Pressure is increased

iv. Temperature and pressure both are increased

The correct option is:

A. I, ii

B. i, iii

C. ii, iii

D. ii, iii, ive

**Answer: C**

 [Watch Video Solution](#)

**182.** The position of equilibrium will shift, by the addition of inert gas at constant pressure condition, in the following case(s):

a.  $N_2(g) + 3F_2(g) \rightleftharpoons 2NF_2(g)$ , forward direction

b.  $COCl_2(g) \rightleftharpoons CO(g) + Cl_2(g)$ , forward direction

c.  $CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$ , backward direction

d.  $2C(s) + O_2(g) \rightleftharpoons 2CO(g)$ , forward direction

 [Watch Video Solution](#)

183.  $\text{COCl}_2$  gas decomposes as:



If one mole of He gas is added in the vessel at equilibrium at constant pressure then

A.  $[\text{COCl}_2]$  increases.

B. "moles" of CO will increase.

C. The reaction goes in forward direction.


D.  $K_c = 1$



**Answer: B::C**



[Watch Video Solution](#)


184. What would be the effect of increasing the volume of each of the following system at equilibrium?





Watch Video Solution

**185.** What happens when an inert gas is added to






at equilibrium at : (a) constant pressure and temperature and temperature, and (b) at constant volume and temperature.



Watch Video Solution

**186.** What is the effect of temperature and pressure on the yields of products?





 [Watch Video Solution](#)

**187.** What would happen to a reversible reaction at equilibrium, when

- a. The temperature is raised, given that its  $\Delta H$  is positive.
- b. The temperature is lowered, given that its  $\Delta H$  is positive.
- c. The temperature is lowered, given that its  $\Delta H$  is negative.
- d. The pressure is lowered, given that  $\Delta n$  is negative.
- e. The pressure is increased, given  $\Delta n$  is negative.

 [Watch Video Solution](#)

**188.** Which of the following factors will increase the solubility of  $\text{NH}_3$  gas in  $\text{H}_2\text{O}$ ?

- a. Increase in pressure
- b. Addition of water




c. Increase in temperature

d. Decrease in pressure

 [Watch Video Solution](#)

**189.** An aqueous solution of hydrogen sulphide shows the equilibrium:



If dilute hydrochloric acid is added to an aqueous solution of  $H_2S$ , without any change in temperature, the

a. The equilibrium constant will change.


b. The concentration  $HS^{\ominus}$  will increase.

c. The concentration of un-dissociated hydrogen sulphide will decrease.

d. The concentration of  $HS^{\ominus}$  will decrease.

 [Watch Video Solution](#)

**190.** Consider the equilibrium



How would the following affect the position of equilibrium?

- a. Addition of  $PCl_3$
- b. Addition of  $Cl_2$
- c. Removal of  $PCl_5$
- d. Addition of He without a change in volume








 [Watch Video Solution](#)

**191.** The reaction between  $H_2$  and  $CO_2$  to form  $CO$  and  $H_2O$  in the gas phase is exothermic. Predict the changes that take place when the system originally at equilibrium is stressed in each of the following ways

- a.  $CO_2$  is removed.
- b.  $CO$  is removed.
- c. The temperature is decreased.
- d. The pressure of the system is increased.
- e. The volume of the system is increased.

 [Watch Video Solution](#)

192. State which one is homogeneous or heterogeneous?




Watch Video Solution

193. If a mixture of 3 mol of  $H_2$  and 1 mole of  $N_2$  is completely converted into  $NH_3$ , what would be the ratio of the initial and final volume at same temperature and pressure?





Watch Video Solution

**194.** Calculate the equilibrium constant for the reaction,



equilibrium constants at 1395K for the following are:



 [Watch Video Solution](#)

**195.** For the reaction,  $A + B \rightleftharpoons 2C$ , 2 mol of  $A$  and 3 mol of  $B$  are allowed to react. If the equilibrium constant is 4 at  $400^\circ C$ , what will be the moles of  $C$  at equilibrium?

 [Watch Video Solution](#)

**196.** In which case does the reaction go farthest to completion:  $K = 1$ ,  $K = 10^{-10}$ , and why?


 [Watch Video Solution](#)

**197.** One mole of  $H_2$  two moles of  $I_2$  and three moles of  $HI$  are injected in one litre flask. What will be the concentration of  $H_2, I_2$  and  $HI$  at equilibrium at  $500^\circ C$ .  $K_c$  for reaction  $H_2 + I_2 \rightleftharpoons 2HI$  is 45.9.



[Watch Video Solution](#)

**198.**  $0.5\text{mol}$  of  $H_2$  and  $0.5\text{ mol}$  of  $I_2$  react in  $10L$  flask at  $448^\circ C$ . The equilibrium constant ( $K_c$ ) is 50 for



- What is the value of  $K_p$ ?
- Calculate the moles of  $I_2$  at equilibrium.



[Watch Video Solution](#)

**199.** The activation energy of

$H_2 + I_2 \rightleftharpoons 2HI(g)$  in equilibrium for the forward reaction is  $167\text{kJmol}^{-1}$  whereas for the reverse reaction is  $180\text{kJmol}^{-1}$ . The presence of catalyst lowers the activation energy by  $80\text{kJmol}^{-1}$ . Assuming that the reactions

are made at  $27^\circ\text{C}$  and the frequency factor for forward and backward reactions are  $4 \times 10^{-4}$  and  $2 \times 10^{-3}$  respectively, calculate  $K_c$ .

 [Watch Video Solution](#)

**200.**  $K_c$  for  $\text{CO}(g) + \text{H}_2\text{O}(g) \rightleftharpoons \text{CO}_2(g) + \text{H}_2(g)$  at  $986^\circ\text{C}$  is 0.63. A mixture of 1 mol  $\text{H}_2\text{O}(g)$  and 3 mol  $\text{CO}(g)$  is allowed to react to come to an equilibrium. The equilibrium pressure is 2.0 atm.

- How many moles of  $\text{H}_2$  are present at equilibrium ?
- Calculate partial pressure of each gas at equilibrium.

 [Watch Video Solution](#)

**201.** At 700K,  $\text{CO}_2$  and  $\text{H}_2$  react to form  $\text{CO}$  and  $\text{H}_2\text{O}$ . For this purpose,  $K_c$  is 0.11. If a mixture of 0.45 mol of  $\text{CO}_2$  and 0.45 mol of  $\text{H}_2$  is heated to 700K.

- Find out amount of each gas at equilibrium.
- When equilibrium has been reached, another 0.34 mol of  $\text{CO}_2$  and


0.34 mol of  $H_2$  are added to the reaction mixture. Find the composition of mixture at new equilibrium.

 [Watch Video Solution](#)

**202.** The degree of dissociation of  $N_2O_4$  into  $NO_2$  at 1 atm  $40^\circ C$  is 0.25. Calculate its  $K_p$  at  $40^\circ C$ .

 [Watch Video Solution](#)

**203.**  $N_2O_4$  dissociates as



At  $40^\circ C$  and one atmosphere % decomposition of  $N_2O_4$  is 50.3%. At what pressure and same temperature, the equilibrium mixture has the ratio of  $N_2O_4: NO_2$  as 1:8?

 [Watch Video Solution](#)

**204.** An equilibrium mixture at  $300\text{K}$  contains  $\text{N}_2\text{O}_4$  and  $\text{NO}_2$  at  $0.28$  and  $1.1\text{atm}$ , respectively. If the volume of container is doubles, calculate the new equilibrium pressure of two gases.

 [Watch Video Solution](#)

**205.** At  $25^\circ\text{C}$  and  $1\text{atm}$  pressure, the partial pressure in equilibrium mixture of  $\text{N}_2\text{O}_4$  and  $\text{NO}_2$ , are  $0.7$  and  $0.3\text{atm}$ , respectively. Calculate the partial pressures of these gases when they are in equilibrium at  $25^\circ\text{C}$  and a total pressure of  $10\text{atm}$ .

 [Watch Video Solution](#)

**206.** Prove  $\alpha = \sqrt{\left(\frac{K_p}{P + K_p}\right)}$  for



where  $\alpha$  is the degree of dissociation at temperature when equilibrium constant is  $K_p$ .






[Watch Video Solution](#)

**207.** At some temperature and under a pressure of 4 atm,  $PCl_5$  is 10 % dissociated. Calculate the pressure at which  $PCl_5$  will be 20 % dissociated temperature remaining same.



[Watch Video Solution](#)


**208.** 1 mole of  $N_2$  and 3 moles of  $PCl_5$  are placed in a 100 litre vessels heated at  $227^\circ C$  the equilibrium pressure is 2.05 atm. Assuming ideal behaviour, calculate degree of dissociation of  $PCl_5$  and  $K_p$  for the reaction



[Watch Video Solution](#)

**209.** One "mole" of  $N_2$  is mixed with three moles of  $H_2$  in a 4L vessel. If 0.25 %  $N_2$  is converted into  $NH_3$  by the reaction

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ , calculate  $K_c$ . Also report  $K_c$  for




 [Watch Video Solution](#)

**210.**  $NH_3$  is heated at 15 atm, from  $25^\circ C$  to  $347^\circ C$  assuming volume constant. The new pressure becomes 50 atm at equilibrium of the reaction  $2NH_3 \rightleftharpoons N_2 + 3H_2$ . Calculate % moles of  $NH_3$  actually decomposed.

 [Watch Video Solution](#)


**211.** What is % dissociation of  $H_2S$  if 1 "mole" of  $H_2S$  is introduced into a 1.10L vessel at 1000K?  $K_c$  for the reaction



 [Watch Video Solution](#)

 Watch Video Solution

**212.** Some solid  $NH_4HS$  is placed in flask containing 0.5 atm of  $NH_3$ . What would be the pressure of  $NH_3$  and  $H_2S$  when equilibrium is reached.



Watch Video Solution

**213.** In an experiment starting with 1 mol  $C_2H_5OH$ , 1 mol  $CH_3COOH$ , and 1 mol of water, the equilibrium mixture mixture of analysis shows that 54.3 % of the acid is esterified. Calculate  $K_c$ .



Watch Video Solution

**214.** When  $C_2H_5OH$  and  $CH_3COOH$  are mixed in equivalent proportion, equilibrium is reached when  $2/3$  of acid and alcohol are used. How much ester will be present when 2g "mole"cule of acid were to react with 2g "mole"cule of alcohol.

 [Watch Video Solution](#)

**215.** When  $\alpha$  - *D* glucose is dissolved in water, it undergoes a partial conversion to  $\beta$  - *D* glucose to exhibit mutarotation. This conversion stops when 63.6 % of glucose is in  $\beta$  form. Assuming that equilibrium has been attained, calculate  $K_c$  for mutarotation.

 [Watch Video Solution](#)

**216.** Calculate  $K_c$  for the reaction  $KI + I_2 \rightleftharpoons KI_3$ . Given that initial weight of *KI* is 1.326g weight of  $KI_3$  is 0.105g and number of moles of free  $I_2$  is 0.0025 at equilibrium the volume of solution is 1L.



 [Watch Video Solution](#)

**217.** Sulphide ions in alkaline solution react with solid sulphur to form polyvalent sulphide ions. The equilibrium constant for the formation of

$S_2^{2-}$  and  $S_3^{2-}$  from  $S$  and  $S^{2-}$  ions is 1.7 and 5.3 respectively. Calculate equilibrium constant for the formation of  $S_3^{2-}$  from  $S_2^{2-}$  and  $S$ .

 [Watch Video Solution](#)

**218.** When  $NO$  and  $NO_2$  are mixed, the following equilibria are readily obtained,



In an experiment when  $NO$  and  $NO_2$  are mixed in the ratio of 1:2, the final total pressure was 5.05 atm and the partial pressure of  $N_2O_4$  was 1.7 atm. Calculate

a. the equilibrium partial pressure of  $NO$ .

b.  $K_p$  for  $NO + NO_2 \rightleftharpoons N_2O_3$ .

 [Watch Video Solution](#)

**219.**  $N_2$  and  $O_2$  combine at a given temperature to produce  $NO$ . At equilibrium the yield of  $NO$  is 'x' percent by volume. If  $x = \sqrt{Ka \cdot b} - \frac{K(a+b)}{4}$ , where  $K$  is the equilibrium constant of the given reaction at the given temperature and  $a$  and  $b$  are the volume percentage of  $N_2$  and  $O_2$ , respectively, in the initial state. Report the value of  $K$  at which  $X$  is maximum

 [Watch Video Solution](#)

**220.** A vessel at  $1000K$  contains carbon dioxide with a pressure of  $0.5atm$ . Some of the carbon dioxide is converted to carbon monoxide on addition of graphite. Calculate the value of  $K_p$  if total pressure at equilibrium is  $0.8atm$ .


 [Watch Video Solution](#)

**221.** The equilibrium constant  $K_p$  of the reaction:  $2SO_2 + O_2 \rightleftharpoons 2SO_3$  is  $900atm^{-1}$  at  $800K$ . A mixture containing  $SO_3$  and  $O_2$  having initial

pressure of 1 atm and 2 atm respectively, is heated at constant volume to equilibrate. Calculate the partial pressure of each gas at 800K at equilibrium.

 [Watch Video Solution](#)

**222.** When 0.15 mol of CO taken in a 2.5L flask is maintained at 750K along with a catalyst, the following reaction takes place




Hydrogen is introduced until the total pressure of the system is 8.5 atm at equilibrium and 0.08 mol of methanol is formed.

Calculate

- $K_p$  and  $K_c$
- The final pressure, if the same amount of CO and  $\text{H}_2$  as before are used, but with no catalyst so that the reaction does not take place.

 [Watch Video Solution](#)

**223.** For the reaction



in solution which was originally 0.1M in KCN and 0.03M in AgNO<sub>3</sub>.

 [Watch Video Solution](#)

**224.**  $\Delta G^\ominus = 77.77\text{kJmol}^{-1}$  at 1000K for the reaction  $1/2\text{N}_2(\text{g}) + 1/2\text{O}_2(\text{g}) \rightleftharpoons \text{NO}(\text{g})$ . What is the partial pressure of NO under equilibrium at 1000K for air at 1atm pressure containing 80% N<sub>2</sub> and 20% O<sub>2</sub> volume.

 [Watch Video Solution](#)

**225.** A saturated solution of Mg(OH)<sub>2</sub> has a vapour pressure of 759.5mm at 373K. Calculate the solubility and K<sub>sp</sub> of Mg(OH)<sub>2</sub>. "(Assume molarity equals molality)"

 [Watch Video Solution](#)



**226.** For the reaction

$\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$  was put in to 10L container and heated to  $800^\circ\text{C}$ , what percentage of the  $\text{CaCO}_3$  would remain unreacted at equilibrium. if ' $k_p$ ' is 1.16 atm and ' $\text{CaCO}_3(\text{s})$ ' initially is 20g

 [Watch Video Solution](#)

**227.** Consider the reaction:




When the system is at equilibrium at  $200^\circ\text{C}$ , the concentrations are found to be:

$$[A] = 0.20M, [B] = 0.30M, [C] = 0.30M$$



- If the volume of the container is suddenly doubled at  $200^\circ\text{C}$ , find the equilibrium concentrations.
- If the volume of the container is suddenly halved (instead of being doubled in part (i) at  $200^\circ\text{C}$ , find the equilibrium concentrations.

 [Watch Video Solution](#)

**228.** Calculate the equilibrium constant for the reaction




If the equilibrium constants at 1395K for the following are:



 [Watch Video Solution](#)

**229.** Calculate the total pressure developed in a vessel containing a mixture of three parts  $H_2$  and one part of  $N_2$  to give a mixture containing 10 % ammonia (by moles) at equilibrium at  $450^\circ C$ .



 [Watch Video Solution](#)

**230.** Average value of poisson's ratio for a mixture of 2 mole of each gas A and B is 1.66, then

 [Watch Video Solution](#)

**231.** Consider the following equilibrium:



8.0g of  $SO_3$  are put in a container at  $600^\circ C$ . The equilibrium pressure and density are 1.8 atm and  $1.6gL^{-1}$ , respectively. Find the value of  $K_p$ .

 [Watch Video Solution](#)

**232.** When  $N_2O_5(g)$  is heated it dissociates to give  $N_2O_3$  and  $O_2$ .  $K_c$  for  $N_2O_5 \rightarrow N_2O_3 + O_2$  is 7.75 and  $K_c$  for  $N_2O_3 \rightarrow N_2O + O_2$  is  $4.0molL^{-1}$ . (both  $K_c$  are at same temperature) 4 mol  $N_2O_5$  in 1.0L vessel is kept at a certain temperature. the concentration of  $O_2$  was  $4.5molL^{-1}$ . Find the concentration of  $N_2O_5$ ,  $N_2O_3$ , and  $N_2O$  at equilibrium.

 [Watch Video Solution](#)

**233.** For a reversible reaction:  $X + 2Y \rightarrow 2Z$ , the equilibrium concentrations of  $X, Y$  and  $Z$  are  $0.32, 0.40$  and  $0.35$  moles  $L^{-1}$  respectively at  $25^\circ C$ .

a. If initially the system contained only  $X$  and  $Y$  and then reached the state of equilibrium, what were the initial concentrations of  $X$  and  $Y$ .

b. If at the start only  $X$  and  $Z$  were present, what were the initial concentrations?



[Watch Video Solution](#)

**234.** Under what pressure conditions  $CuSO_4 \cdot 5H_2O$  be efforescent at  $25^\circ C$ . How good a drying agent is  $CuSO_4 \cdot 3H_2O$  at the same temperature? Given



$K_p = 1.086 \times 10^{-4} atm^2$  at  $35^\circ C$ . Vapoure pressure of water at  $25^\circ C$  is  $23.8mm$  of Hg.



[Watch Video Solution](#)



**235.** Under what pressure conditions  $CuSO_4 \cdot 5H_2O$  be efforescent at  $25^\circ C$ . How good a drying agent is  $CuSO_4 \cdot 3H_2O$  at the same temperature? Given




$K_p = 1.086 \times 10^{-4} atm^2$  at  $25^\circ C$ . Vapour pressure of water at  $25^\circ C$  is 23.8 mm of Hg.

 [Watch Video Solution](#)

**236.** From the data given below which of the following reactant is most effective drying agent at  $0^\circ C$ . Given  $P_{H_2O}^\circ = 4.58 mm$  at  $0^\circ C$ .





iii.



 [Watch Video Solution](#)



**237.** Following two equilibria are established on mixing two gases  $A_2$  and  $C$ .



If  $A_2$  and  $C$  mixed in 2:1 molar, ratio calculate the equilibrium partial pressure of  $A_2$ ,  $C$ ,  $A_2C$  and  $K_p$  for the reaction (ii). Given that the total pressure to be 1.4 atm and partial pressure of  $A_6$  to be 0.2 atm at equilibrium

 [Watch Video Solution](#)

**238.** 1 mol of A in 1 litre vessel maintained at constant T shows the equilibrium



If the equilibrium pressure is  $\frac{13}{6}$  times of initial pressure and

$[C]_{eq} = \frac{4}{9}[A]_{eq}$ , Calculate  $K_{C_1}$  and  $K_{C_2}$ .

 [Watch Video Solution](#)

239. One "mole" of  $N_2O_4(g)$  at  $100K$  is kept in a closed container at  $1.0$  atm pressure. It is heated to  $400K$ , where  $30\%$  by mass of  $N_2O_4(g)$  decomposes to  $NO_2(g)$ . The resultant pressure will be

A.  $4.2$

B.  $5.2$

C.  $3.2$

D.  $6.2$



Watch Video Solution

### Concept Application exercise 7.1

1. In a reaction between hydrogen and iodine  $6.84\text{mol}$  of hydrogen and  $4.02$  mol of iodine are found to be in equilibrium with  $42.85$  mol of

hydrogen iodide at  $350^{\circ}\text{C}$ . Calculate the equilibrium constant.

 [Watch Video Solution](#)

2. Calculate the equilibrium constant  $K_p$  and  $K_c$  for the reaction:


$\text{CO}(g) + 1/2\text{O}_2(g) \rightleftharpoons \text{CO}_2(g)$ . Given that the partial pressure at

equilibrium in a vessel at  $3000\text{K}$  are  $p_{\text{CO}} = 0.4\text{atm}$ ,  $p_{\text{CO}_2} = 0.6\text{atm}$ ,  $p_{\text{O}_2} = 0.2$

atm

 [Watch Video Solution](#)

3. The equilibrium composition for the reaction is



0.20          0.10           $0.40\text{molL}^{-1}$


What will be the equilibrium concentration of  $\text{PCl}_5$  on adding  $0.10\text{mol}$  of

$\text{Cl}_2$  at the same temperature?

 [Watch Video Solution](#)



#### 4. For the reaction




Fill in the blanks in the following table for the three solution at equilibrium.

Solution	$[\text{Cu}^{2+}(aq)]$ $\text{molL}^{-1}$	$[\text{Ag}^{\oplus}(aq)]$ $\text{molL}^{-1}$	$KL^{-1}$ $\text{molL}^{-1}$
1.	(a)	$1.0 \times 10^{-9}$	$2.0 \times 10^{15}$
2.	$2.0 \times 10^{-7}$	$1.0 \times 10^{-11}$	(b)
3.	$2.0 \times 10^{-2}$	(c)	$2.0 \times 10^{15}$



[Watch Video Solution](#)

#### 5. The value of $K_c$ for the reaction:



at  $100^\circ\text{C}$  is 49. If 1.0L flask containing one mole of  $\text{A}_2$  is connected with a 2.0L flask containing one mole of  $\text{B}_2$ , how many moles of AB will be formed at  $100^\circ\text{C}$ ?



[Watch Video Solution](#)

6. At  $440^\circ\text{C}$ , the equilibrium constant ( $K$ ) for the following reaction is 49.5,  $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ . If 0.2 mol of  $\text{H}_2$  and 0.2 mol of  $\text{I}_2$  are placed in a 10 - L vessel and permitted to react at this temperature, what will be the concentration of each substance at equilibrium?

 [Watch Video Solution](#)

7. 0.15 mol of  $\text{CO}$  taken in a 2.5L flask is maintained at 750K along with a catalyst so that the following reaction can take place  $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}(\text{g})$ . Hydrogen is introduced until the total pressure of the system is 8.5 atm at equilibrium and 0.08 mol of methanol is formed. Calculate

a.  $K_p$  and  $K_c$

b. The final pressure if the same amount of  $\text{CO}$  and  $\text{H}_2$  as before is used but no catalyst so that the reaction does not take place.

 [Watch Video Solution](#)

8. A vessel at  $1000K$  contains carbon dioxide with a pressure of  $0.5\text{atm}$ . Some of the carbon dioxide is converted to carbon monoxide on addition of graphite. Calculate the value of  $K_p$  if total pressure at equilibrium is  $0.8\text{atm}$ .

 [Watch Video Solution](#)

9. For the reaction,  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ , the concentration of an equilibrium mixture at  $298K$  is  $N_2O_4 = 4.50 \times 10^{-2}\text{molL}^{-1}$  and  $NO_2 = 1.61 \times 10^{-2}\text{molL}^{-1}$ . What is the value of equilibrium constant?

 [Watch Video Solution](#)

10. For an equilibrium reaction, the rate constants for the forward and the backward reaction are  $2.38 \times 10^{-4}$  and  $8.15 \times 10^{-5}$ , respectively. Calculate the equilibrium constant for the reaction.

 [Watch Video Solution](#)

11. In a reaction between  $H_2$  and  $I_2$  at a certain temperature, the amounts of  $H_2$ ,  $I_2$  and HI at equilibrium were found to be 0.45 mol, 0.39 mol, and 3.0 mol respectively. Calculate the equilibrium constant for the reaction at the given temperature.

 [Watch Video Solution](#)

12. At 700K, the equilibrium constant  $K_p$  for the reaction



is  $1.80 \times 10^{-3} kPa$ . What is the numerical value of  $K_c$  in moles per litre for this reaction at the same temperature?


 [Watch Video Solution](#)

13. Two moles of  $PCl_5$  were heated to  $327^\circ C$  in a closed two-litre vessel, and when equilibrium was achieved,  $PCl_5$  was found to be 40% dissociated into  $PCl_3$  and  $Cl_2$ . Calculate the equilibrium constant  $K_p$  and  $K_c$  for this reaction.



Watch Video Solution

14. For the reaction,



the partial pressure of  $N_2$  and  $H_2$  are 0.80 and 0.40 atmosphere, respectively, at equilibrium. The total pressure of the system is 2.80 atm.

What is  $K_p$  for the above reaction?



Watch Video Solution

15. The equilibrium constant at 278 K for



is  $2.0 \times 10^{15}$ . At a particular moment, the concentration of

$Cu^{2+}$  and  $Ag^+$  ions are found to be

$1.8 \times 10^{-2} \text{ mol L}^{-1}$  and  $3.0 \times 10^{-9} \text{ mol L}^{-1}$

respectively. Is the system in equilibrium at that moment?



Watch Video Solution

16.  $AB_2$  dissociates as

$AB_2(g) \rightleftharpoons AB(g) + B(g)$ . If the initial pressure is 500 mm of Hg and the total pressure at equilibrium is 700 mm of Hg. Calculate  $K_p$  for the reaction.

 [Watch Video Solution](#)

17. Under what pressure must an equimolar mixture of  $PCl_5$  and  $Cl_2$  be placed at  $250^\circ C$  in order to obtain  $PCl_5$  at 1 atm? ( $K_p$  for dissociation of  $PCl_5 = 1.78$ ).

 [Watch Video Solution](#)

18.  $XY_2$  dissociates  $XY_2(g) \rightleftharpoons XY(g) + Y(g)$ . When the initial pressure of  $XY_2$  is 600 mm Hg, the total equilibrium pressure is 800 mm Hg. Calculate  $K$  for the reaction Assuming that the volume of the system remains unchanged.

A. 50

B. 100

C. 200

D. 400

**Answer: A**

 [Watch Video Solution](#)

19.  $K_c$  for the reaction  $SO_2(g) + \frac{1}{2}O_2(g) \rightarrow SO_3(g)$  is  $61 \cdot 7$  at  $60^\circ C$ .

What is its unit? Calculate  $K_p$  for the reaction and write its unit. \* \*

 [Watch Video Solution](#)

20. 1 mol of  $H_2$ , 2 mol of  $I_2$  and 3 mol of HI were taken in a 1 - L flask. If the value of  $K_c$  for the equation  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$  is 50 at  $440^\circ C$ , what will be the concentration of each specie at equilibrium?

 [Watch Video Solution](#)

21. For  $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ ,  $K_c$  is equal to .....

A.  $K_c = \frac{1}{[\text{CO}_2]}$

B.  $K_c = [\text{CO}_2]$

C.  $K_c = \frac{[\text{CaO}][\text{CO}_2]}{[\text{CaCO}_3]}$

D.  $K_c = \frac{[\text{CaCO}_3]}{[\text{CaO}][\text{CO}_2]}$

Answer: B

 Watch Video Solution

22. For the reaction  $\text{C}(\text{s}) + \text{CO}_2(\text{g}) \rightleftharpoons 2\text{CO}(\text{g})$ , the partial pressure of  $\text{CO}_2$  and  $\text{CO}$  is 2.0 and 4.0 atm, respectively, at equilibrium. The  $K_p$  of the reaction is



A. 0.5

B. 5.0

C. 30.0

D. 8.0

**Answer: D**

 [Watch Video Solution](#)

**23.** In a chemical equilibrium,  $K_c = K_p$  when

A. The number of molecules entering into a reaction is more than the number of molecules produced.

B. The number of molecules entering into the reaction is equal to the number of molecules produced.

C. the number of molecules entering into the reaction is less to the number of moleculad produced.

D. None of the above

**Answer: B**



[Watch Video Solution](#)

**24.** In a general reaction  $A + B \rightleftharpoons AB$ , which value of equilibrium constant most favours the production of AB?

A. (a)  $9.0 \times 10^{-3}$

B. (b)  $3.0 \times 10^{-3}$

C. (c)  $9.0 \times 10^{-7}$

D. (d)  $9.0 \times 10^{-12}$

**Answer: A::B::C**



[Watch Video Solution](#)

25. During thermal dissociation of a gas, the vapour density.

- A. Remains the same
- B. Increases
- C. Decreases
- D. Increases in some cases and decreases in others

**Answer: C**



[Watch Video Solution](#)

26. The vapour density of fully dissociated  $NH_4Cl$  would be

- A. Less than half of the vapour density of pure  $NH_4Cl$
- B. Double of the vapour density of pure  $NH_4Cl$
- C. Half of the vapour density of pure  $NH_4Cl$
- D. One-third of the vapour density of pure  $NH_4Cl$

Answer: C



Watch Video Solution

27. In the reversible reaction,  $2HI(g) \rightleftharpoons H_2(g) + I_2(g)$ ,  $K_p$  is

A. Greater than  $K_c$

B. Less than  $K_c$

C. Equal to  $K_c$

D. Zero

Answer: C



Watch Video Solution

28. At 500K, the equilibrium constant for reaction  $cis - C_2H_2Cl_2 \rightleftharpoons trans - C_2H_2Cl_2$  is 0.6. At the same temperature, the

equilibrium constant for the reaction  $\text{trans} - \text{C}_2\text{H}_2\text{Cl}_2 \rightleftharpoons \text{cis} - \text{C}_2\text{H}_2\text{Cl}_2$

will be

A. 1.67

B. 1.65

C. 1.06


D. 1.60

**Answer: A::B::C**



**Watch Video Solution**

**29.** 2 mol of  $\text{N}_2$  is mixed with 6 mol of  $\text{H}_2$  in a closed vessel of one litre capacity. If 50 %  $\text{N}_2$  is converted into  $\text{NH}_3$  at equilibrium, the value of  $K_c$  for the reaction



A. 4/27

B. 27/4

C. 2/27

D. 20

**Answer: A::B::C**

 **Watch Video Solution**

**30.** For the reaction

$H_2(g) + CO_2(g) \rightleftharpoons CO(g) + H_2O(g)$ , if the initial concentration of  $[H_2] = [CO_2]$  and  $x$  moles /litres of hydrogen is consumed at equilibrium, the correct expression of  $K_p$  is :

- A.  $\frac{x^2}{(1-x)^2}$
- B.  $\frac{(1+x)^2}{(1-x)^2}$
- C.  $\frac{1+x^2}{(2+x)^2}$
- D.  $\frac{x^2}{1+x^2}$

**Answer: A::B::C**



Watch Video Solution

31. Partial pressure of  $O_2$  in the reaction



A.  $K_p$

B.  $\sqrt{K_p}$

C.  $\sqrt[3]{K_p}$

D.  $(K_p)^2$

Answer: A::B::C



Watch Video Solution

32. Two moles of  $PCl_5$  were heated to  $327^\circ C$  in a closed two-litre vessel, and when equilibrium was achieved,  $PCl_5$  was found to be 40% dissociated into  $PCl_3$  and  $Cl_2$ . Calculate the equilibrium constant  $K_p$  and  $K_c$  for this reaction.

A. 0.530

B. 0.266

C. 0.130

D. 0.170

**Answer: B**

 [Watch Video Solution](#)

**33.** For the reaction,



$$\left( K_c = 1.8 \times 10^{-6} \text{ at } 184^\circ \text{C} \right)$$

$$(R = 0.0083 \text{ kJ} / (\text{mol K}))$$

When  $K_p$  and  $K_c$  are compared at  $184^\circ \text{C}$  it is found that

A.  $K_p$  is greater than  $K_c$

B.  $K_p$  is less than  $K_c$

C.  $K_p = K_c$



D. None of the above

**Answer: A**



[Watch Video Solution](#)

**34.** For a reaction  $NH_4COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$ , the equilibrium pressure is 3 atm.  $K_p$  for the reaction will be

A. 4

B. 20

C. 25

D. 15

**Answer: A::B::C**



[Watch Video Solution](#)

35. For the reaction  $A + B \rightleftharpoons C + D$ , the initial concentrations of A and B are equal. The equilibrium concentration of C is two times the equilibrium concentration of A. The value of equilibrium constant is .....

A. 1

B. 2

C. 3

D. 4

**Answer: D**



[Watch Video Solution](#)

36. 15 mol of  $H_2$  and 5.2 moles of  $I_2$  are mixed and allowed to attain equilibrium at  $500^\circ C$ . At equilibrium, the concentration of HI is found to be 10 mol. The equilibrium constant for the formation of HI is.

A. 50

B. 25

C. 200

D. 15

**Answer: A**



[Watch Video Solution](#)

37. For the reaction:  $2\text{NOCl}(g) \rightleftharpoons 2\text{NO}(g) + \text{Cl}_2(g)$ ,  $K_c$  at  $427^\circ\text{C}$  is  $3 \times 10^{-6} \text{Lmol}^{-1}$ . The value of  $K_p$  is

A. 7.5

B.  $2.5 \times 10^{-5}$

C.  $2.0 \times 10^{-4}$

D.  $1.72 \times 10^{-4}$

**Answer: D**



[Watch Video Solution](#)

38. For the reaction



Which one is the correct representation?

A.  $K_p = [p_{\text{H}_2\text{O}}]^2$

B.  $K_c = [\text{H}_2\text{O}]^2$

C.  $K_p = K_c(RT)^2$


D. All are correct

Answer: D



Watch Video Solution

39. Which one is the correct representation for the reaction



$$\text{A. } K_p = \frac{[p_{SO_3}]^2}{[p_{SO_2}]^2 [p_{O_2}]}$$

$$\text{B. } K_c = \frac{[SO_3]^2}{[SO_2]^2 [O_2]}$$

$$\text{C. } K_p = \frac{[n_{SO_3}]^2}{[n_{SO_2}]^2 [n_{O_2}]} \times \left[ \frac{P}{\text{Total mole}} \right]^{-1}$$

D. All the above

**Answer: D**



**Watch Video Solution**

**40.** For the reactions,

$CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ , the  $\frac{K_p}{K_c}$  is equal to

A.  $1/RT$

B.  $Rt$

C.  $\sqrt{RT}$

D.  $(RT)^2$

**Answer: A**



**Watch Video Solution**

41. The equilibrium constant for the reaction  $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$  at temperature (T)  $4 \times 10^{-4}$ . The value of  $K_c$  for the reaction  $NO(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{1}{2}O_2(g)$  at the same temperature is?

A.  $25 \times 10^2$

B. 50

C.  $4 \times 10^{-4}$


D. 10.00

**Answer: B**



**Watch Video Solution**

42.  $K_p/K_c$  for the reaction



A.  $RT$

B.  $(RT)^{1/2}$

C.  $\frac{1}{(RT)^3}$

D.  $\frac{1}{\sqrt{RT}}$

**Answer: D**



[Watch Video Solution](#)

43. The unit of equilibrium constant  $K_c$  for the reaction  $A + B \rightleftharpoons C$  would be

A.  $\text{mol}^{-1}\text{L}$

B.  $\text{molL}^{-1}$





C. molL

D. No unit

**Answer: A**

 [Watch Video Solution](#)

**44.** For which of the following reaction does the equilibrium constant depend on the units of concentration?



**Answer: D**

 [Watch Video Solution](#)



45. To the system,



already at equilibrium, more water vapour is added without altering temperature or volume of the system. When equilibrium is re-established, the pressure of water vapour is doubled. The pressure of  $\text{HCl}$  present in the system increases by a factor of

A. 2

B.  $2^{1/2}$

C.  $2^{1/3}$

D.  $2^2$

**Answer: B**



[Watch Video Solution](#)

46. For the reaction,  $\text{A}(\text{g}) + 2\text{B}(\text{g}) \rightleftharpoons 2\text{C}(\text{g})$ , the rate constant for forward and the reverse reactions are  $1 \times 10^{-4}$  and  $2.5 \times 10^{-2}$  respectively. The

value of equilibrium constant, K for the reaction would be

A.  $2 \times 10^{-4}$

B.  $3 \times 10^{-2}$

C.  $4 \times 10^{-3}$


D.  $3 \times 10^2$

**Answer: C**




[Watch Video Solution](#)

**47.** The equilibrium constant for the reaction



is 20 at 500K. The equilibrium constant for the reaction



A. 20

B. 0.5


C. 0.05

Answer: C



Watch Video Solution

48. For the reaction




in solution which was originally 0.1M in KCN and 0.03M in  $AgNO_3$ .



Watch Video Solution

49. At a certain temperature, the equilibrium constant ( $K_c$ ) is 16 for the reaction:



If we take one mole of each of the four gases in one litre container then what will be the equilibrium concentration of NO and  $NO_2$ ?

A.  $1.6 \text{ molL}^{-1}$

B.  $0.8\text{molL}^{-1}$

C.  $0.4\text{molL}^{-1}$

D.  $0.6\text{molL}^{-1}$

**Answer: C**



**Watch Video Solution**

50.  $\text{HI}$  was heated in a sealed tube at  $400^\circ\text{C}$  till the equilibrium was reached.  $\text{HI}$  was found to be 22 % decomposed. The equilibrium constant for dissociation is

A. 1.99

B. 0.0199

C. 0.0796

D. 0.282

**Answer: B**



Watch Video Solution

51. For the equilibrium  $AB(g) \rightleftharpoons A(g) + B(g)$  at a given temperature, the pressure at which one-third of AB is dissociated is numerically equal to

- A. 8 times  $K_p$
- B. 16 times  $K_p$
- C. 4 times  $K_p$
- D. 9 times  $K_p$

Answer: A



Watch Video Solution

52. For a reversible reaction, if the concentration of the reactants are doubled, then the equilibrium constant will

- A. change to  $1/4K$

B. change to  $1/2K$

C. change to  $2K$


D. remain the same

**Answer: D**



**Watch Video Solution**

**53.** For the equilibrium



$K_p$  is equal to four times the total pressure. Calculate the number of moles of  $A$  formed.

A. 0.45

B. 0.30

C. 0.60

D. 0.90

**Answer: D**

 [Watch Video Solution](#)

## Ex 7.2


1.  $1.5\text{ mol}$  of  $\text{PCl}_5$  are heated at constant temperature in a closed vessel of  $4\text{ L}$  capacity. At the equilibrium point,  $\text{PCl}_5$  is  $35\%$  dissociated into  $\text{PCl}_3$  and  $\text{Cl}_2$ . Calculate the equilibrium constant.

 [Watch Video Solution](#)

2. Calculate the degree of dissociation of  $\text{HI}$  at  $450^\circ\text{ C}$  if the equilibrium constant for the dissociation reaction is  $0.263$ .

 [Watch Video Solution](#)

3. Calculate the percent dissociation of  $H_2S(g)$  if  $0.1\text{mol}$  of  $H_2S$  is kept in  $0.4\text{L}$  vessel at  $1000\text{K}$ . For the reaction:



The value of  $K_c$  is  $1.0 \times 10^{-6}$

 [Watch Video Solution](#)

4. One mole of  $H_2$  two moles of  $I_2$  and three moles of  $HI$  are injected in one litre flask. What will be the concentration of  $H_2, I_2$  and  $HI$  at equilibrium at  $500^\circ\text{C}$ .  $K_c$  for reaction  $H_2 + I_2 \rightleftharpoons 2HI$  is  $45.9$ .

 [Watch Video Solution](#)

5. At  $700\text{K}$ , hydrogen and bromine react to form hydrogen bromine. The value of equilibrium constant for this reaction is  $5 \times 10^8$ . Calculate the amount of the  $H_2, Br_2$  and  $HBr$  at equilibrium if a mixture of  $0.6\text{mol}$  of  $H_2$  and  $0.2\text{mol}$  of  $Br_2$  is heated to  $700\text{K}$ .

 [Watch Video Solution](#)



[Watch Video Solution](#)

6. At some temperature and under a pressure of 4 atm,  $PCl_5$  is 10 % dissociated. Calculate the pressure at which  $PCl_5$  will be 20 % dissociated temperature remaining same.

[Watch Video Solution](#)


7. 20 %  $N_2O_4$  molecules are dissociated in a sample of gas at  $27^\circ C$  and 760 torr. Calculate the density of the equilibrium mixture.

[Watch Video Solution](#)

8. 0.1 mol of  $PCl_5$  is vaporised in a litre vessel at  $260^\circ C$ . Calculate the concentration of  $Cl_2$  at equilibrium, if the equilibrium constant for the dissociation of  $PCl_5$  is 0.0414.

[Watch Video Solution](#)

9. The equilibrium constant for the reaction



is 4.0 at 25 ° C. Calculate the weight of ethyl acetate that will be obtained when 120g of acetic acid are reacted with 92g of alcohol.

 [Watch Video Solution](#)

10. The vapour density of  $\text{PCl}_5$  at 43K is is found to be 70.2. Find the degree of dissociation of  $\text{PCl}_5$  at this temperature.

 [Watch Video Solution](#)

11. For the equilibrium  $\text{AB}(g) \rightleftharpoons \text{A}(g) + \text{B}(g)$ .  $K_p$  is equal to four times the total pressure. Calculate the number moles of A formed if one mol of AB is taken initially.

 [Watch Video Solution](#)

12. The vapour density of a mixture containing  $NO_2$  and  $N_2O_4$  is 38.3 at 300 K. the number of moles of  $NO_2$  in 100 g of the mixture is approximately

 [Watch Video Solution](#)

13.  $NH_3$  is heated at 15 atm, from  $25^\circ C$  to  $347^\circ C$  assuming volume constant. The new pressure becomes 50 atm at equilibrium of the reaction  $2NH_3 \rightleftharpoons N_2 + 3H_2$ . Calculate % moles of  $NH_3$  actually decomposed.

 [Watch Video Solution](#)

14. The pressure of iodine gas at 1273K is found to be 0.112 atm whereas the expected pressure is 0.074 atm. The increased pressure is due to dissociation  $I_2 \rightleftharpoons 2I$ . Calculate  $K_p$ .

 [Watch Video Solution](#)


15.  $K_c$  for  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$  is 0.00466 at 298K. If a 1L container initially contained 0.8 mol of  $N_2O_4$ , what would be the concentrations of  $N_2O_4$  and  $NO_2$  at equilibrium? Also calculate the equilibrium concentration of  $N_2O_4$  and  $NO_2$  if the volume is halved at the same temperature.

 [Watch Video Solution](#)

16. At a certain temperature,  $K_p$  for dissociation of solid  $CaCO_3$  is  $4 \times 10^{-2}$  atm and for the reaction,  $C(s) + CO_2 \rightleftharpoons 2CO$  is 2.0 atm, respectively. Calculate the pressure of CO at this temperature when solid C, CaO,  $CaCO_3$  are mixed and allowed to attain equilibrium.

 [Watch Video Solution](#)

17. Given below are the values of  $\Delta H^\ominus$  and  $\Delta S^\ominus$  for the reaction given below at 27 °C.




$$\Delta H^\ominus = -98.32 \text{ kJ mol}^{-1}, \Delta S^\ominus = -95 \text{ J mol}^{-1}$$

Find  $K_p$  for the reaction

 [Watch Video Solution](#)

18. The yield of product in the reaction,




would be higher at:

- A. Low temperature and high pressure
- B. High temperature and high pressure
- C. Low temperature and low pressure
- D. High temperature and low pressure

**Answer: A**

 [Watch Video Solution](#)

19. Manufacture of ammonia from the elements is represented by



The maximum yield of ammonia will be obtained when the process is made to take place

- A. At low pressure and high temperature
- B. At low pressure and low temperature
- C. At high pressure and high temperature
- D. At high pressure and low temperature

**Answer: D**

 [Watch Video Solution](#)

20. The reaction  $2SO_2 + O_2 \rightleftharpoons 2SO_3 + \text{Heat}$ , will be favoured by

- A. Low temperature and low pressure
- B. High temperature and low pressure




C. High temperature and high pressure

D. Low temperature and high pressure

**Answer: D**

 [Watch Video Solution](#)

21. In which of the following equilibrium, change in volume of the system does not alter the number of moles:



**Answer: A**

 [Watch Video Solution](#)

22. In the dissociation of  $2HI \rightleftharpoons H_2 + I_2$ , the degree of dissociation will be affected by

- A. Increase of temperature
- B. Addition of an inert gas
- C. Addition of  $H_2$  and  $I_2$
- D. Increase of pressure

**Answer: A**



[Watch Video Solution](#)

23. In line kilns, the following reaction,



proceeds to completion because of

- A. High temperature
- B.  $CO_2$  escapes






C. Low temperature and low pressure

D. molecular mass of  $\text{CaO}$  is less than that of  $\text{CaCO}_3$

**Answer: B**

 [Watch Video Solution](#)

**24.** Which among the following reactions will be favoured at low pressure?



**Answer: C**

 [Watch Video Solution](#)

25. If  $E_f$  and  $E_r$  are the activation energies of forward and backward reactions and the reaction is known to be exothermic, then

A.  $E_f > E_r$

B.  $E_f < E_r$

C.  $E_f = E_r$

D. No relation can be given between  $E_f$  and  $E_r$

**Answer: B**



[Watch Video Solution](#)

26.  $K_p$  for a reaction at  $25^\circ\text{C}$  is 10 atm. The activation energy for forward and reverse reactions are 12 and  $20\text{kJmol}^{-1}$  respectively. The  $K_c$  for the reaction at  $40^\circ\text{C}$  will be:

A.  $4.33 \times 10^{-1}M$

B.  $3.4 \times 10^{-2}M$

C.  $3.4 \times 10^{-1}M$

D.  $4.33 \times 10^{-2}M$

**Answer: C**

 [Watch Video Solution](#)

**27.** The concentration of a pure solid or liquid phase is not include in the expression of equilibrium constant because :

- A. Solid and liquid concentrations are independent of their quantities
- B. Solid and liquid react slowly
- C. Solid and liquids at equilibrium do not interact with gaseous phase
- D. The molecules of solids and liquid cannot migrate to the gaseous phase

**Answer: A**

 [Watch Video Solution](#)

28. For an equilibrium reaction involving gases, the forward reaction is first order while the reverse reaction is second order. The unit of  $K_p$  for forward equilibrium is

A. atm

B.  $\text{atm}^2$

C.  $\text{atm}^{-1}$

D.  $\text{atm}^{-2}$

**Answer: A**



[Watch Video Solution](#)

29. For the reaction,  $\text{PCl}_3(g) + \text{Cl}_2(g) \rightleftharpoons \text{PCl}_5(g)$ , the position of equilibrium can be shifted to the right by:

A. Doubling the volume

B. Increasing the temperature

C. Addition of equimolar quantities of  $PCl_3$  and  $PCl_5$

D. Addition of  $Cl_2$  at constant volume

**Answer: D**



[Watch Video Solution](#)

**30.** What are the favourable conditions for the synthesis of ammonia.

A. High temperature and high pressure

B. Low temperature and low pressure

C. High temperature and low pressure


D. Low temperature and high pressure

**Answer: D**



[Watch Video Solution](#)

31. Which of the following change will shift the reaction in forward direction?



- A. Increase in total pressure
- B. Increase in temperature
- C. Increase in concentration of  $I$
- D. Decrease in concentration of  $I_2$

**Answer: B**



[Watch Video Solution](#)


32. In a vessel containing  $SO_3$ ,  $SO_2$  and  $O_2$  at equilibrium, some helium gas is introduced so that total pressure increases while temperature and volume remain the same. According to Le Chatelier's principle, the dissociation of  $SO_3$ :

- A. Increases
- B. Decreases
- C. Remains unaltered
- D. Changes unpredictably

**Answer: C**

 [Watch Video Solution](#)

**33.** Vapour density of the equilibrium mixture of  $NO_2$  and  $N_2O_4$  is found to be 40 for the equilibrium



Calculate

- A. abnormal molecular weight
- B. degree of dissociation
- C. percentage of  $NO_2$  in the mixture
- D. N/A

**Answer: B**

 [Watch Video Solution](#)

**34.** Calculate the pressure of  $CO_2$  gas at  $700K$  in the heterogenous equilibrium reaction  $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ , if  $\Delta G^\ominus$  for this reaction is  $130.2kJmol^{-1}$ .

 [Watch Video Solution](#)

**35.** The equilibrium constant  $K_{p_2}$  and  $K_{p_2}$  for the reactions  $A \rightleftharpoons 2B$  and  $P \rightleftharpoons Q + R$ , respectively, are in the ratio of 2:3. If the degree of dissociation of A and P are equal, the ratio of the total pressure at equilibrium is,

A. 1:36

B. 1:1

C. 1:3



D. 1:9

Answer: A::C

 [Watch Video Solution](#)

36. For  $I_2(g) \rightleftharpoons 2I(g)$ ,  $K_p = 1.79 \times 10^{-10}$ . The partial pressure of  $I_2 = 1.0$  atm and  $I = 0.5 \times 10^{-6}$  atm after 50 min . Comment on the status of equilibrium process.

 [Watch Video Solution](#)

37. Calculate the volume percent of chlorine gas at equilibrium in the dissociation of  $PCl_5(g)$  under a total pressure of 1.5 atm. The  $K_p$  for its dissociation = 0.3.


 [Watch Video Solution](#)

38.  $N_2O_4 \rightleftharpoons 2NO_2$ ,  $K_c = 4$ . This reversible reaction is studied graphically as shown in the figure. Select the correct statement out of *I*, *II* and *III*.

*I* : Reaction quotient has maximum value at point *A*

*II* : Reaction proceeds left to right at a point when  $[N_2O_4] = [NO_2] = 0.1M$

*III* :  $K = Q$  when point *D* or *F* is reached:



A. *I*, *II*

B. *II*, *III*


C. *II*

D. I, II, III

**Answer: B**

 [Watch Video Solution](#)

**39.** The equilibrium:



is attained by mixing equal moles of  $P_4$  and  $Cl_2$  in an evacuated vessel.

Then at equilibrium:

A.  $[Cl_2] > [PCl_3]$

B.  $[Cl_2] > [P_4]$

C.  $[P_4] > [Cl_2]$

D.  $[PCl_3] > [P_4]$

**Answer: C**

 [Watch Video Solution](#)

40.  $N_2O_4(g)$  is dissociated to an extent of 20 % at equilibrium pressure of 1.0 atm and  $57^\circ C$ . Find the percentage of  $N_2O_4$  at 0.2 atm and  $57^\circ C$ .

 [Watch Video Solution](#)


## Exercises (Subjective)

1. The equilibrium pressure of

$NH_4CN(s) \rightleftharpoons NH_3(g) + HCN(g)$  is 2.98 atm. Calculate  $K_p$

 [Watch Video Solution](#)

2. To 500mL of 0.150M  $AgNO_3$  solution were added 500mL of 1.09M  $Fe^{2+}$  solution and the reaction is allowed to reach an equilibrium at  $25^\circ C$



For 25 mL of the solution, 30mL of 0.0832M  $KMnO_4$  was required for oxidation. Calculate the equilibrium constant for the the reaction  $25^\circ C$ .

 [Watch Video Solution](#)

[Watch Video Solution](#)

3. For the equilibrium




$K_p = 9 \text{ atm}^2$  at  $37^\circ \text{C}$ . A 5 litre vessel contains 0.1 mole of  $\text{LiCl} \cdot \text{NH}_3$ . How many moles of  $\text{NH}_3$  should be added to the flask at this temperature to derive the backward reaction for completion?


Use:  $R = 0.082 \text{ atm} \cdot \text{L} / \text{molK}$

[Watch Video Solution](#)

4. The equilibrium constant of the reaction,



is 0.15 at 900K. Calculate the equilibrium constant for

[Watch Video Solution](#)

5.  $K_c$  for the reaction  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  is  $0.5 \text{ mol}^{-2} \text{ L}^2$  at 400K. Find  $K_p$ .

Given  $R = 0.082 \text{ L} \cdot \text{atm} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$

 [Watch Video Solution](#)

6. The equilibrium constant  $K_c$  for  $A(g) \rightleftharpoons B(g)$  is 1.1. Which gas has a molar concentration greater than 1.

 [Watch Video Solution](#)

7. In an equilibrium  $A + B \rightleftharpoons C + D$ , A and B are mixed in vessel at temperature T. The initial concentration of A was twice the initial concentration of B. After the equilibrium has reached, concentration of C was thrice the equilibrium concentration of B. Calculate  $K_c$ .

A. (a) 1

B. (b) 2

C.

D.

**Answer: A:C**

 [Watch Video Solution](#)

8. For a gaseous phase reaction  $A + 2B \rightleftharpoons AB_2$ ,  $K_c = 0.3475L^2\text{mole}^{-2}$  at  $200^\circ\text{C}$ . When 2 moles of B are mixed with one "mole" of A, what total pressure is required to convert 60 % of A in  $AB_2$ ?

 [Watch Video Solution](#)


9. For a reaction  $2HI \rightleftharpoons H_2 + I_2$ , at equilibrium 7.8g, 203.2g, and 1638.4g of  $H_2$ ,  $I_2$ , and HI, respectively were found. Calculate  $K_c$ .

 [Watch Video Solution](#)

10. 25 moles of  $H_2$  and 18 moles of  $I_2$  vapour were heated in a sealed tube at  $445^\circ\text{C}$  when at equilibrium 30.8 moles of HI were formed. Calculate the degree of dissociation of pure HI at the given temperature,

 [Watch Video Solution](#)

11. In the dissociation of HI, 20 % of HI is dissociated at equilibrium. Calculate  $K_p$  for



 [Watch Video Solution](#)

12. The value of  $K_p$  for dissociation of  $2HI \rightleftharpoons H_2 + I_2$  is  $1.84 \times 10^{-2}$ . If the equilibrium concentration of  $H_2$  is  $0.4789 \text{ mol L}^{-1}$ , calculate the concentration of HI at equilibrium.


 [Watch Video Solution](#)



13. 0.96g of HI were, heated to attain equilibrium  $2HI \rightleftharpoons H_2 + I_2$ . The reaction mixture on titration requires 15.7mL of N/10 hypo solution. Calculate the degree of dissociation of HI.

 [Watch Video Solution](#)

14. An equilibrium mixture



present in a vessel of one litre capacity at 815 °C was found by analysis to contain 0.4 mol of CO, 0.3 mol of  $H_2O$ , 0.2 mol of  $CO_2$  and 0.6 mol of  $H_2$ .

a. Calculate  $K_c$


b. If it is derived to increase the concentration of CO to 0.6 mol by adding  $CO_2$  to the vessel, how many moles must be added into equilibrium mixture at constant temperature in order to get this change?

 [Watch Video Solution](#)

15. A mixture of one mole of  $CO_2$  and "mole" of  $H_2$  attains equilibrium at a temperature of  $250^\circ C$  and a total pressure of 0.1 atm for the change  $CO_2(g) + H_2(g) \rightleftharpoons CO(g) + H_2O(g)$ . Calculate  $K_p$  if the analysis of final reaction mixture shows 0.16 volume percent of CO.

 [Watch Video Solution](#)


16. At a certain temperature, the equilibrium constant ( $K_c$ ) is 16 for the reaction:



If we take one mole of each of the four gases in one litre container then what will be the equilibrium concentration of  $NO$  and  $NO_2$ ?

 [Watch Video Solution](#)

17. The equilibrium mixture for



present in 1L vessel at  $600^\circ C$  contains 0.50, 0.12, and 5.0 moles of


$SO_2$ ,  $O_2$ , and  $SO_3$  respectively.

a. Calculate  $K_c$  for the given change at  $600^\circ C$ .

b. Also calculate  $K_p$ .

 [Watch Video Solution](#)

18. At 273 K and 1atm , 10 litre of  $N_2O_4$  decompose to  $NO_2$  decomposes to  $NO_2$  according to equation



What is degree of dissociation ( $\alpha$ ) when the original volume is 25% less than that of existing volume?

 [Watch Video Solution](#)

19. At 340K and 1 atm pressure,  $N_2O_4$  is 66 % into  $NO_2$ . What volume of 10g  $N_2O_4$  occupy under these conditions?

 [Watch Video Solution](#)

20. How much  $PCl_5$  must be added to a one litre vessel at  $250^\circ C$  in order to obtain a concentration of  $0.1 \text{ mol of } Cl_2$ ?  $K_c$  for  $PCl_5 \rightleftharpoons PCl_3 + Cl_2$  is  $0.0414 \text{ molL}^{-1}$

 [Watch Video Solution](#)

21. The degree of dissociation of  $PCl_5$  at 1 atm pressure is 0.2. Calculate the pressure at which  $PCl_5$  is dissociated to 50 % ?

 [Watch Video Solution](#)

22. At  $473K$ , partially dissociated vapours of  $PCl_5$  are 62 times as heavy as  $H_2$ . Calculate the degree of dissociation of  $PCl_5$ .

 [Watch Video Solution](#)

23. In a mixture of  $N_2$  and  $H_2$  in the ratio 1:3 at 30 atm and  $300^\circ C$ , the % of  $NH_3$  at equilibrium is 17.8. Calculate  $K_p$  for  $N_2 + 3H_2 \rightleftharpoons 2NH_3$ .



[Watch Video Solution](#)

24. A reaction carried out by 1 mol of  $N_2$  and 3 mol of  $H_2$  shows at equilibrium the mole fraction of  $NH_3$  as 0.012 at  $500^\circ C$  and 10 atm pressure. Calculate  $K_p$ . Also report the pressure at which "mole" % of  $NH_3$  in equilibrium mixture is increased to 10.4.




[Watch Video Solution](#)

25. The equilibrium constant  $K_p$ , for the reaction  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$  is  $1.6 \times 10^{-4}$  at  $400^\circ C$ . What will be the equilibrium constant at  $500^\circ C$  if the heat of reaction in this temperature range is  $-25.14$  kcal?




[Watch Video Solution](#)

26. What concentration of  $CO_2$  be in equilibrium with  $2.5 \times 10^{-2} \text{ mol L}^{-1}$  of CO at  $100^\circ \text{C}$  for the reaction:



 [Watch Video Solution](#)

27. Calculate  $K_c$  for the reaction:




if 1.58 mol  $H_2S$ , 1.27 mol  $H_2$  and  $2.78 \times 10^{-6}$  mol of  $S_2$  are in equilibrium in a flask of capacity 180L at  $750^\circ \text{C}$ .

 [Watch Video Solution](#)

28. For  $NH_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$ , the observed, pressure for reaction mixture in equilibrium is 1.12 atm at  $106^\circ \text{C}$ . What is the value of  $K_p$  for the reaction?

 [Watch Video Solution](#)

29. If 50% of  $CO_2$  converts to CO at the following equilibrium:



and the equilibrium pressure is 12 atm. Calculate  $K_p$ .



 [Watch Video Solution](#)

30. For  $A + B \rightleftharpoons C$ , the equilibrium concentration of A and B at a temperature are  $15\text{molL}^{-1}$ . When volume is doubled the reaction has equilibrium concentration of A as  $10\text{molL}^{-1}$ , calculate

a.  $K_c$

 [Watch Video Solution](#)

31. Two solid compounds A and B dissociate into gaseous products at  $20^\circ C$  as




At  $20^\circ C$  pressure over excess solid A is 50 mm and that over excess solid

B is 68 mm. Find:

- The dissociation constant of A and B
- Relative number of moles of A' and B' in the vapour phase over a mixture of the solids A and B.
- Show that the total pressure of gas over the solid mixture would be 84.4 mm.

 [Watch Video Solution](#)

**32.** Consider the heterogeneous equilibrium




Calculate the partial pressure of  $\text{CO}(g)$  when  $\text{CaCO}_3$  and  $\text{C}$  are mixed and allowed to attain equilibrium at the temperature for which the above two equilibria have been studied.

 [Watch Video Solution](#)



**33.** Would 1 %  $CO_2$  in air be sufficient to prevent any loss in weight when  $M_2CO_3$  is heated at  $120^\circ C$ ?



$K_p = 0.0095 \text{ atm}$  at  $120^\circ C$ . How long would the partial pressure of  $CO_2$  have to be to promote this reaction at  $120^\circ C$ ?

 [Watch Video Solution](#)

**34.** Under what pressure conditions  $CuSO_4 \cdot 5H_2O$  be efforescent at  $25^\circ C$ . How good a drying agent is  $CuSO_4 \cdot 3H_2O$  at the same temperature? Given



$K_p = 1.086 \times 10^{-4} \text{ atm}^2$  at  $35^\circ C$ . Vapoure pressure of water at  $25^\circ C$  is  $23.8 \text{ mm}$  of Hg.


 [Watch Video Solution](#)

**35.** For the reaction,  $SnO_2(s) + 2H_2(g) \rightleftharpoons Sn(l) + 2H_2O(g)$  the equilibrium mixture of steam and hydrogen contained 45 % and 24 %  $H_2$  at  $900K$  and

1100K respectively. Calculate  $K_p$  at both the temperature.

 [Watch Video Solution](#)

**36.** For the reaction:



$K_c = 9.14 \times 10^{-6}$  at  $25^\circ C$ . If the initial concentration of the ions are

$Fe^{3+} = 0.5M$ ,  $(Hg_2)^{2+} = 0.5M$ ,  $Fe^{2+} = 0.03M$  and  $Hg^{2+} = 0.03M$ , what

will be the concentration of ions at equilibrium.

 [Watch Video Solution](#)

**37.** 0.1 mol each of ethyl alcohol and acetic acid are allowed to react and at equilibrium, the acid was exactly neutralised by 100mL of 0.85N NaOH.

If no hydrolysis of ester is supposed to have undergone, find  $K_c$ .

 [Watch Video Solution](#)


38. At  $450^{\circ}\text{C}$  the equilibrium constant  $K_p$  for the reaction  $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$  was found to be  $1.6 \times 10^{-5}$  at a pressure of 200 atm. If  $\text{N}_2$  and  $\text{H}_2$  are taken in 1:3 ratio. What is % of  $\text{NH}_3$  formed at this temperature?

 [Watch Video Solution](#)

39.  $K_p$  for the reaction  $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$  at  $400^{\circ}\text{C}$  is  $1.64 \times 10^{-4} \text{ atm}^{-2}$ . Find  $K_c$ .

 [Watch Video Solution](#)

40. Equilibrium constant  $K_p$  for



is 0.0118 atm at  $1065^{\circ}\text{C}$  and heat of dissociation is 42.4 Kcal. Find equilibrium constant at  $1132^{\circ}\text{C}$ .

 [Watch Video Solution](#)

41.  $K_p$  for  $3/2H_2 + 1/2N_2 \rightleftharpoons NH_3$  are 0.0266 and  $0.0129 \text{ atm}^{-1}$ , respectively, at  $350^\circ \text{C}$  and  $400^\circ \text{C}$ . Calculate the heat of formation of  $NH_3$ .

 [Watch Video Solution](#)

42. In a reaction at equilibrium, X moles of the reactant A decomposes to give 1 mole each of C and D. It has been found that the fraction of A decomposed at equilibrium is independent of initial concentration of A. Calculate X.

 [Watch Video Solution](#)

43. For the reaction  $A + B \rightleftharpoons 3. C$  at  $25^\circ \text{C}$ , a 3L vessel contains 1, 2, and 4 moles of A, B and C respectively. Predict the direction of reaction if:


- $K_c$  for the reaction is 10.
- $K_c$  for the reaction is 15.
- $K_c$  for the reaction is 10.66

 [Watch Video Solution](#)

44. An equilibrium mixture at 300K contains  $N_2O_4$  and  $NO_2$  at 0.28 and 1.1atm, respectively. If the volume of container is doubles, calculate the new equilibrium pressure of two gases.


 [Watch Video Solution](#)

45. NO and  $Br_2$  at initial pressures of 98.4 and 41.3 torr respectively were allowed react at 300K. At equilibrium the total pressure was 110.5 torr. Calculate the value of equilibrium constant,  $K_p$  and the standard free energy change at 300K for the reaction:



 [Watch Video Solution](#)

46. In the reaction equilibrium



When 5 mol of each is taken and the temperature is kept at 298K, the total pressure was found to be 20 bar.

Given :  $\Delta_f G_{n_2O_4}^\ominus = 100\text{kJ}$ ,  $\Delta_f G_{NO_2}^\ominus = 50\text{KJ}$

a. Find  $\Delta G^\ominus$  of the reaction at 298K.

b. Find the direction of the reaction.



[Watch Video Solution](#)

## Exercises (Linked Comprehensive)

1. Consider the following equilibrium:



If  $O_2$  is added and volume of the reaction vessel is reduced, the equilibrium

- A. Shift in the product side
- B. Shifts in the reactant side
- C. Cannot be predicted

D. Remains unchanged

**Answer: A**

 [Watch Video Solution](#)

2. If we add  $SO_4^{2-}$  ion to a saturated solution of  $Ag_2SO_4$ , it will result in a//an

A. Increase in  $Ag^{\oplus}$  concentration.

B. Decrease in  $Ag^{\oplus}$  concentration

C. It will shift  $Ag^{\oplus}$  ions from solid  $Ag_2SO_4$  into solution.

D. It will decrease the  $SO_4^{2-}$  ion concentration in the solution.

**Answer: B**

 [Watch Video Solution](#)

3. Physical and chemical equilibrium can respond to a change in their pressure, temperature, and concentration of reactants and products. To describe the change in the equilibrium we have a principle named Le Chatelier principle. According to this principle, even if we make some changes in equilibrium, then also the system even re-establishes the equilibrium by undoing the effect.

In the reaction  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ . If we increase the pressure of the system, the equilibrium is

- A. Shifts in the product side
- B. Shift un reactant side
- C. Remains unchanged
- D. Cannot be predicted


**Answer: A**



**Watch Video Solution**




4. Consider the chemical reaction:




When  $\text{H}^{\oplus}(\text{aq})$  is added, the colour green is favoured. Use one or more of the following interpretations to answer the questions:

- i. Some unreacted  $\text{Ni}^{2+}(\text{aq})$  is present in the solution at equilibrium
- ii. Some unreacted  $\text{NH}_3(\text{aq})$  is present in the solution at equilibrium
- iii. The colour change indicates new equilibrium conditions with reduced



- iv. The colour change indicates new equilibrium conditions with increased



The deepening of blue colour on dissolving more  $\text{Ni}(\text{NO}_3)_2$  supports interpretation (s).


- A. i only
- B. i and iv only
- C. ii and iv only
- D. i and ii only

Answer: B




Watch Video Solution

5. Consider the chemical reaction:




When  $H^{\oplus}(aq)$  is added, the colour green is favoured. Use one or more of the following interpretations to answer the questions:

- Some unreacted  $Ni^{2+}(aq)$  is present in the solution at equilibrium
- Some unreacted  $NH_3(aq)$  is present in the solution at equilibrium
- The colour change indicates new equilibrium conditions with reduced



- The colour change indicates new equilibrium conditions with increased



The deepening of blue colour on addition of more  $NH_3(aq)$  supports interpretation(s).

A. i only

B. i and iv only



C. i and ii only

D. ii and iv only

**Answer: D**

 [Watch Video Solution](#)

6. One "mole" of  $NH_4HS(s)$  was allowed to decompose in a 1 - L container at  $200^\circ C$ . It decomposes reversibly to  $NH_3(g)$  and  $H_2S(g)$ .  $NH_3(g)$  further undergoes decomposition to form  $N_2(g)$  and  $H_2(g)$ . Finally, when equilibrium was set up, the ratio between the number of moles of  $NH_3(g)$  and  $H_2(g)$  was found to be 3.



Answer the following:

What is the "mole" fraction of hydrogen gas in the equilibrium mixture in the gas phase?

A.  $1/4$

B.  $3/4$



C.  $1/8$

D. 4

**Answer: B**

 [Watch Video Solution](#)

7. One "mole" of  $NH_4HS(s)$  was allowed to decompose in a 1 - L container at  $200^\circ C$ . It decomposes reversibly to  $NH_3(g)$  and  $H_2S(g)$ .  $NH_3(g)$  further undergoes decomposition to form  $N_2(g)$  and  $H_2(g)$ . Finally, when equilibrium was set up, the ratio between the number of moles of  $NH_3(g)$  and  $H_2(g)$  was found to be 3.



Answer the following:

To attain equilibrium, how much % by weight of solid  $NH_4HS$  got dissociated?

A. 19 %

B. 30 %

C. 33 %


D. 15 %

**Answer: C**



[Watch Video Solution](#)

8. One "mole" of  $NH_4HS(s)$  was allowed to decompose in a 1 - L container at  $200^\circ C$ . It decomposes reversibly to  $NH_3(g)$  and  $H_2S(g)$ .  $NH_3(g)$  further undergoes decomposition to form  $N_2(g)$  and  $H_2(g)$ . Finally, when equilibrium was set up, the ratio between the number of moles of  $NH_3(g)$  and  $H_2(g)$  was found to be 3.



Answer the following:

What is the "mole" fraction of hydrogen gas in the equilibrium mixture in the gas phase?

A.  $16.83\text{gL}^{-1}$

B.  $16.83\text{gmL}^{-1}$

C.  $18.415\text{gL}^{-1}$


D.  $14.83\text{gL}^{-1}$

**Answer: A**



[Watch Video Solution](#)

9. The percentage of ammonia produced from nitrogen and hydrogen under conditions of temperature and pressure is given in the graph



Use the graph answering the following questions:

What happens to the percentage of ammonia produced when the temperature is increased


- A. The % is decreased
- B. The % is increased
- C. No effect
- D. Cannot be predicted

**Answer: A**



**Watch Video Solution**

10. The percentage of ammonia produced from nitrogen and hydrogen under conditions of temperature and pressure is given in the graph



Use the graph answering the following questions:

What happens to the percentage of ammonia produced when the pressure is increased?


- A. The % is decreased
- B. The % is increased
- C. No effect
- D. Cannot be predicted



Answer: B

 Watch Video Solution

11. The percentage of ammonia produced from nitrogen and hydrogen under conditions of temperature and pressure is given in the graph



Use the graph answering the following questions:

What conditions of pressure produce the highest percentage of ammonia?

A. At least 50 atm

B. At least 150 atm


C. At least 300 atm

D. At least 100 atm

**Answer: C**

 [Watch Video Solution](#)

**12.** The synthesis of ammonia is given as:



temperature ( $T$ ) =  $375^\circ\text{C}$

The expression of equilibrium constant is

$$\text{A. } K_c = \frac{[N_2][H_2]^3}{[NH_3]^2}$$

$$\text{B. } K_c = \frac{[N_2][H_2]}{[NH_3]}$$


$$\text{C. } K_c = \frac{[NH_3]}{[N_2][H_2]^3}$$

$$D. K_c = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

**Answer: D**

 [Watch Video Solution](#)

**13.** The synthesis of ammonia is given as:



temperature ( $T$ ) =  $375^\circ C$


On increasing the temperature, the value of equilibrium constant  $K_c$

- A. Increases
- B. Decreases
- C. Remain unchanged
- D. Cannot be predicted

**Answer: B**

 [Watch Video Solution](#)

14. The synthesis of ammonia is given as:



The relationship between  $K_p$  and  $K_c$  for this reaction is

A.  $K_c = K_p(RT)^{-2}$

B.  $K_p = K_c(RT)^{-1}$

C.  $K_p = K_c(RT)^2$


D.  $K_p = K_c(RT)^4$

**Answer: A**



**Watch Video Solution**

15. The synthesis of ammonia is given as:




Which of the following factors does not alter the yield of  $NH_3$  at equilibrium?

- A. Catalyst
- B. Increase in pressure
- C. Increase in temperature
- D. Decrease in pressure

**Answer: A**

 [Watch Video Solution](#)

**16.** The synthesis of ammonia is given as:



temperature ( $T$ ) =  $375^\circ C$

Starting with 2 mol of each ( $N_2, H_2$  and  $NH_3$ ) in 5.0L reaction vessel at  $375^\circ C$ , predict what is true for the reaction?


- A. The reaction is at equilibrium

- B. The reaction proceed in forward direction.
- C. The reaction proceed in backward direction
- D.  $Q_c$  for the reaction is less then  $K_c$

**Answer: C**

 [Watch Video Solution](#)

17. Phosphorous pentachloride when heated in a sealed tube at 700K it undergoes decomposition as



Vapour density of the mixture is 74.25.

The reaction is


- A. Endothermic
- B. Exothermic
- C. May be endothermic or exothermic
- D. Unpredictable

**Answer: A**



[Watch Video Solution](#)

**18.** Phosphorous pentachloride when heated in a sealed tube at 700K it undergoes decomposition as



Vapour density of the mixture is 74.25.

Percentage dissociation of  $PCl_5$  may be given as


- A. 4.04
- B. 40.4
- C. 44.0
- D. 0.404

**Answer: B**



[Watch Video Solution](#)

19. Phosphorous pentachloride when heated in a sealed tube at 700K it undergoes decomposition as



Vapour density of the mixture is 74.25.

Equilibrium constant  $K_c$  for the reaction will be

A. 0.66M

B. 0.56M

C. 0.46M


D. 0.36M

**Answer: A**



[Watch Video Solution](#)

20. Phosphorous pentachloride when heated in a sealed tube at 700K it undergoes decomposition as





Vapour density of the mixture is 74.25.


If pressure is increased then the equilibrium will

- A. Be unaffected
- B. Shift in backward direction
- C. Shift in forward direction
- D. Cannot be predicted

**Answer: B**

 [Watch Video Solution](#)

21. Phosphorous pentachloride when heated in a sealed tube at 700K it undergoes decomposition as



Vapour density of the mixture is 74.25.

When an inert gas is added to the given reversible process, then the equilibrium will.

- A. Be unaffected
- B. Shift in backward direction
- C. Shift in forward direction
- D. Cannot be predicted

**Answer: C**

 [Watch Video Solution](#)

**22.** Decomposition of ammonium chloride is an endothermic reaction.

The equilibrium may be represented as:



A 6.250g sample of  $\text{NH}_4\text{Cl}$  is placed in an evacuated 4.0L container at  $27^\circ\text{C}$ . After equilibrium the total pressure inside the container is 0.820 bar and some solid remains in the container. Answer the followings

The value of  $K_p$  for the reaction at 300K is

- A. 16.2

B. 0.168

C. 1.68

D. 32.4

**Answer: B**

 [Watch Video Solution](#)

**23.** Decomposition of ammonium chloride is an endothermic reaction.

The equilibrium may be represented as:



A 6.250g sample of  $\text{NH}_4\text{Cl}$  is placed in an evacuated 4.0L container at

27 °C. After equilibrium the total pressure inside the container is 0.820

bar and some solid remains in the container. Answer the followings

The amount of solid  $\text{NH}_4\text{Cl}$  left behind in the container at equilibrium is

A. 2.856

B. 28.56

C. 0.2856

D. 1.320

**Answer: A**



**Watch Video Solution**

**24.** Decomposition of ammonium chloride is an endothermic reaction.

The equilibrium may be represented as:



A 6.250g sample of  $\text{NH}_4\text{Cl}$  is placed in an evacuated 4.0L container at  $27^\circ\text{C}$ . After equilibrium the total pressure inside the container is 0.820 bar and some solid remains in the container. Answer the followings

If the volume of container were doubled at constant temperature, then what would happen to the amount of solid in the container.

A. Decrease

B. Increases

C. Remain unchanged

D. None

**Answer: A**



**Watch Video Solution**

**25.** Decomposition of ammonium chloride is an endothermic reaction.

The equilibrium may be represented as:



A 6.250g sample of  $NH_4Cl$  is placed in an evacuated 4.0L container at  $27^\circ C$ . After equilibrium the total pressure inside the container is 0.820 bar and some solid remains in the container. Answer the followings

The value of  $K_p$  for the reaction at 300K is

A. Increasing the temperature

B. Decreasing the temperature

C. Adding more  $NH_4Cl$

D. Removing  $\text{HCl}(\text{g})$

**Answer: A**

 [Watch Video Solution](#)

**26.** Decomposition of ammonium chloride is an endothermic reaction.

The equilibrium may be represented as:



A  $6.250\text{g}$  sample of  $\text{NH}_4\text{Cl}$  is placed in an evacuated  $4.0\text{L}$  container at

$27^\circ\text{C}$ . After equilibrium the total pressure inside the container is  $0.820$

bar and some solid remains in the container. Answer the followings

The value of  $K_p$  for the reaction at  $300\text{K}$  is

- A. Increase in volume
- B. Decrease in temperature
- C. Decrease in pressure
- D. Increase in temperature

**Answer: B**





 [Watch Video Solution](#)

27.  $K_p$  and  $K_c$  are inter related as

$$K_p = K_c(RT)^{\Delta n}$$

Answer the following questions:

Which of the following have  $K_p = K_c$ ?



**Answer: A::B**




 [Watch Video Solution](#)

28.  $K_p$  and  $K_c$  are inter related as

$$K_p = K_c(RT)^{\Delta n}$$

Answer the following questions:

Which of the following have same units of  $K_p$  ?



**Answer: A::B**



**Watch Video Solution**





29.  $K_p$  and  $K_c$  are inter related as

$$K_p = K_c(RT)^{\Delta n}$$

Answer the following questions:

In which of the following equilibria  $K_p$  is less than  $K_c$ ?





**Answer: B**

 [Watch Video Solution](#)

**30.**  $K_p$  and  $K_c$  are inter related as

$$K_p = K_c(RT)^{\Delta n}$$

Answer the following questions:

For  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ ,  $K_p/K_c$  is equal to:

A.  $RT^3$

B.  $1/RT$

C.  $(RT)^4$

D.  $1/(RT)^2$

**Answer: D**




**Watch Video Solution**

31.  $K_p$  and  $K_c$  are inter related as

$$K_p = K_c(RT)^{\Delta n}$$

Answer the following questions:

The unit of equilibrium constant for



A.  $\text{mol L}^{-2}$

B.  $\text{mol}^2\text{L}^{-2}$

C.  $\text{Lmol}^{-2}$




D. None of these

**Answer: D**



**Watch Video Solution**

**32.** Consider the following reactions:



Which of the following is/are incorrect?

A.  $K_3 = K_1/K_2$

B.  $K_3 = K_1^2/K_2^3$



C.  $K_3 = K_1 \times K_2$

D.  $K_3 = K_1\sqrt{K_2}$

**Answer: C**

 [Watch Video Solution](#)

**33.** The equilibrium constant of the following reactions at 400K are given:



Then, the equilibrium constant  $K$  for the reaction



is

A. 2.04

B. 20.5

C. 0.85

D. 1.4




**Answer: D**



[Watch Video Solution](#)

**34.** The relation between  $K_p$  and  $K_c$  is  $K_p = K_c(RT)^{\Delta n}$  unit of

$$K_p = (\text{atm})^{\Delta n}, \text{ unit of } K_c = (\text{molL}^{-1})^{\Delta n}$$



Which of the following relations is correct?

$$\text{A. } K_3 = K_1/K_2$$

$$\text{B. } K_3 = K_1 \times K_2$$

$$\text{C. } K_3 = K_1 + K_2$$

$$\text{D. } K_3 = K_1/\sqrt{K_2}$$




**Answer: B**

 [Watch Video Solution](#)


**35.** The relation between  $K_p$  and  $K_c$  is  $K_p = K_c(RT)^{\Delta n}$  unit of

$$K_p = (\text{atm})^{\Delta n}, \text{ unit of } K_c = (\text{molL}^{-1})^{\Delta n}$$

$H_3ClO_4$  is a tribasic acid, it undergoes ionisation as



Then, equilibrium constant for the following reaction will be:



A.  $K_1K_2K_3$

B.  $\frac{(K_1K_3)^2}{K_2}$

C.  $\frac{K_1}{K_2}$



D.  $\frac{K_1K_2}{K_3^2}$

**Answer: A**




**Watch Video Solution**

**36.** If  $K_1$  and  $K_2$  are the respective equilibrium constants for the two reactions



The equilibrium constant of the reaction,



A.  $K_1/K_2^2$

B.  $(K_1/K_2)^{1/2}$

C.  $K_1^2/K_2^3$

D.  $K_2/K_1$

**Answer: D**



**Watch Video Solution**



The pressure inside the chamber is 100 atm and temperature at 300K

If  $K_p$  for the given reaction is  $1.44 \times 10^{-5}$ , then the value of  $K_c$  will be:

A.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 500)^{-2}} \text{molL}^{-1}$

B.  $\frac{1.44 \times 10^{-5}}{(8.314 \times 200)^{-2}} \text{molL}^{-1}$

C.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 700)^2} \text{molL}^{-1}$


D.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 300)^{-2}} \text{molL}^{-1}$

**Answer: A::B::D**



The pressure inside the chamber is 100 atm and temperature at 300K

The preparation of ammonia by Haber's process is an exothermic reaction. If the preparation follows the following temperature-pressure relationship for its % yield. Then for temperature  $T_1, T_2$  and  $T_3$  the correct option is:





A.  $T_3 > T_2 > T_1$

B.  $T_1 > T_2 > T_3$

C.  $T_3 < T_2 < T_3$

D.  $T_1 = T_2 = T_3$

**Answer: B**

 [Watch Video Solution](#)



The pressure inside the chamber is 100 atm and temperature at 300K

On adding catalyst the equilibrium of reaction:

- A. Shift in backward direction
- B. Shift in forward direction
- C. Does not affect the equilibrium
- D. Cannot predict.


**Answer: C**

 [Watch Video Solution](#)



The pressure inside the chamber is 100 atm and temperature at 300K

If  $K_p$  for the reaction is  $1.44 \times 10^{-5}$ , then the value of  $K_p$  for the decomposition of  $NH_3$



will be:

A. (a)  $\sqrt{1.44 \times 10^{-5}}$

B. (b)  $(1.44 \times 10^{-5})^4$

C. (c)  $\frac{1}{1.44 \times 10^{-5}}$

D. (d)  $1.00 \times 10^{-3}$

**Answer: C**

 [Watch Video Solution](#)

**41.** Mass action ratio or reaction quotient  $Q$  for a reaction can be calculate using the law of mass action



$$Q = \frac{[C][D]}{[A][B]}$$

The value of  $Q$  decides whether the reaction is at equilibrium or not.


At equilibrium,  $Q = K$

For an equilibrium process,  $Q \neq K$

when  $Q > K$ , reaction will favour backward direction and when  $Q < K$ , it will favour direction.

Answer the following questions:

The reaction quotient  $Q$  for:



is given by  $Q = \frac{[H_2O]^2}{[O_2][H_2]^2}$ . The reaction will proceed in backward

direction, when

A.  $Q = K_c$

B.  $Q < K_c$

$$C. Q > K_c$$

$$D. Q = 0$$

**Answer: C**



**Watch Video Solution**

**42.** Mass action ratio or reaction quotient  $Q$  for a reaction can be calculate using the law of masss action



$$Q = \frac{[C][D]}{[A][B]}$$

The value of  $Q$  decides whether the reaction is at equilibrium or not.


At equilibrium,  $Q = K$

For an equilibrium process,  $Q \neq K$

when  $Q > K$ , reaction will favour backward direction and when  $Q < K$ , it will favour direction.

Answer the following questions:

For the reaction:



A 4L vessel contains 2, 1, and 4 mol of A, B and c, respectively. The reaction at the same temperature

- A. Must proceed in forward direction
- B. Must proceed in backward direction
- C. Must be in equilibrium
- D. Cannot be predicted

**Answer: A**

 [Watch Video Solution](#)

**43.** Mass action ratio or reaction quotient  $Q$  for a reaction can be calculate using the law of masss action



$$Q = \frac{[C][D]}{[A][B]}$$

The value of  $Q$  decides whether the reaction is at equilibrium or not.


At equilibrium,  $Q = K$

For an equilibrium process,  $Q \neq K$

when  $Q > K$ , reaction will favour backward direction and when  $Q < K$ , it will favour forward direction.

Answer the following questions:

In a reaction mixture containing  $H_2$ ,  $N_2$  and  $NH_3$  at partial pressure of 2 atm, 1 atm and 3 atm respectively, the value of  $K_p$  at 700K is  $4.00 \times 10^{-5} atm^{-2}$ . In which direction the net reaction will go?




- A. Forward
- B. Backward
- C. No reaction
- D. Cannot be predicted

**Answer: B**



**Watch Video Solution**

**44.** Mass action ratio or reaction quotient  $Q$  for a reaction can be calculate using the law of masss action



$$Q = \frac{[C][D]}{[A][B]}$$

The value of  $Q$  decides whether the reaction is at equilibrium or not.


At equilibrium,  $Q = K$

For an equilibrium process,  $Q \neq K$

when  $Q > K$ , reaction will favour backward direction and when  $Q < K$ , it will favour direction.

Answer the following questions:

In the following reaction:



the equilibrium is not attained. The rate of forward reaction is greater than that of backward reaction. Thus, which of the following is the correct relation between  $K_p$  and  $Q_p$ ?

A.  $K_p = Q_p$

B.  $Q_p > K_p$

$$C. Q_p < K_p$$

$$D. K_p = Q_p = 1$$

**Answer: C**



**Watch Video Solution**

**45.** Mass action ratio or reaction quotient  $Q$  for a reaction can be calculate using the law of masss action



$$Q = \frac{[C][D]}{[A][B]}$$

The value of  $Q$  decides whether the reaction is at equilibrium or not.

At equilibrium,  $Q = K$

For an equilibrium process,  $Q \neq K$

when  $Q > K$ , reaction will favour backward direction and when  $Q < K$ , it will favour direction.

Answer the following questions:


In the reaction:





a graph is plotted to show that the variation or the rate of forward and backward reaction against time.

Which of following is correct?




- A.  $Q > K = 3, Q = K = 2, Q < K = 1$
- B.  $Q > K = 2, Q = K = 3, Q < K = 1$
- C.  $Q > K = 1, Q = K = 2, Q < K = 3$
- D.  $Q > K = 2, Q = K = 1, Q < K = 3$

**Answer: A**



46. Dehydration of salts is an important class of heterogeneous reactions. The salt hydrates during dehydration often dissociate in steps to form a number of intermediate hydrates according to the prevailing pressure of moisture in contact with the solid hydrates. Thus, copper sulphate pentahydrate on dissociation yield trihydrates, monohydrates and then the anhydrous salt in the above order as follows:



The equilibrium constant  $K_p$  for the equilibrium between pentahydrate and trihydrate is:

A. 7.8

B. 60.84

C. 31.36


D. 5.6

Answer: B

 [Watch Video Solution](#)

47. Dehydration of salts is an important class of heterogeneous reactions.

The salt hydrates during dehydration often dissociate in steps to form a number of intermediate hydrates according to the prevailing pressure of moisture in contact with the solid hydrates. Thus, copper sulphate pentahydrate on dissociation yield trihydrates, monohydrates and then the anhydrous salt in the above order as follows:



The ratio of equilibrium constant between pentahydrate and trihydrate and equilibrium between trihydrate and monohydrate is

A. 1.9

B. 2.9

C. 8.6


D. 5.6

**Answer: A**



[Watch Video Solution](#)

**48.** Dehydration of salts is an important class of heterogeneous reactions. The salt hydrates during dehydration often dissociate in steps to form a number of intermediate hydrates according to the prevailing pressure of moisture in contact with the solid hydrates. Thus, copper sulphate pentahydrate on dissociation yield trihydrates, monohydrates and then the anhydrous salt in the above order as follows:



Which of the following conditions is favourable for dehydration of  $\text{CuCO}_4 \cdot 5\text{H}_2\text{O}$ ?

- i. Low humidity in air
- ii. High temperature
- iii.  $p_{\text{H}_2\text{O}}$  increases

The correct option is:


- A. i
- B. i, ii
- C. ii, iii
- D. i, ii, iii

**Answer: B**




 [Watch Video Solution](#)


49. X, Y and Z react in the 1 : 1 : 1 stoichiometric ratio.

The concentration of X, Y and Z we are found to vary with time as shown in the figure below:



Which of the following equilibrium reaction represents the correct variation of concentration with time?






Answer: C

 [Watch Video Solution](#)

50. X, Y and Z react in the 1 : 1 : 1 stoichiometric ratio.

The concentration of X, Y and Z we are found to vary with time as shown in the figure below:



The value of the equilibrium constant ( $K_c$ ) for the equilibrium represented the in above sketch will be

A.  $\frac{9}{2}$

B.  $\frac{11}{4}$

C.  $\frac{2}{3}$

D.  $\frac{10}{7}$


**Answer: C**

 [Watch Video Solution](#)

51. X, Y and Z react in the 1 : 1 : 1 stoichiometric ratio.

The concentration of X, Y and Z we are found to vary with time as shown in the figure below:





If the above equilibrium is established in a 2.0L container by taking reactants in sufficient amount then how many moles of components Y must have reacted to establish the equilibrium?


- A. 0
- B. 6
- C. 12
- D. 8

**Answer: D**



Watch Video Solution

52. Two solids X and Y dissociate into gaseous products at a certain temperature as follows:



At a given temperature, pressure over excess solid 'X' is 40 mm of Hg and total pressure over solid 'Y(s)' is 60 mm of Hg.

Now, answer the following questions:

Ratio of  $K_p$  for reaction (i) to that of reaction (ii), is:

A. 4:9

B. 2:3

C. 4:9


D. 2:1

**Answer: A**



Watch Video Solution

53. Two solid  $X$  and  $Y$  dissociate into gaseous products at a certain temperature as follows:



At a given temperature, the pressure over excess solid  $X$  is  $40\text{mm}$  and total pressure over solid  $Y$  is  $80\text{mm}$ . Calculate

- The value of  $K_p$  for two reactions.
- The ratio of moles of  $A$  and  $B$  in the vapour state over a mixture of  $X$  and  $Y$ .
- The total pressure of gases over a mixture of  $X$  and  $Y$ .

A. 2:3

B. 2:5

C. 4:9


D. 1:1

**Answer: C**



**Watch Video Solution**

54. Two solid  $X$  and  $Y$  dissociate into gaseous products at a certain temperature as follows:



At a given temperature, the pressure over excess solid  $X$  is  $40\text{mm}$  and total pressure over solid  $Y$  is  $80\text{mm}$ . Calculate

- The value of  $K_p$  for two reactions.
- The ratio of moles of  $A$  and  $B$  in the vapour state over a mixture of  $X$  and  $Y$ .
- The total pressure of gases over a mixture of  $X$  and  $Y$ .

A.  $100\text{ mm}$

B.  $74.84\text{ mm}$

C.  $50\text{ mm}$

D.  $120.74\text{ mm}$

**Answer: B**



**Watch Video Solution**

## Exercises (Multiple Correct)

1. For the reaction  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ , which of the following factors will have no effect on the value of equilibrium constant?

- A. Temperature
- B. Initial concentration of  $N_2O_4$
- C. Pressure of catalyst
- D. Pressure

**Answer: B::C::D**



[Watch Video Solution](#)

2. For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ , the equilibrium can be shifted in favour of product by

- A. Increasing the  $[H_2]$

B. Increasing the pressure

C. Increasing the  $[I_2]$

D. By using the catalyst

**Answer: A::B**

 [Watch Video Solution](#)

3. A reaction  $S_8(g) \rightleftharpoons 4S_2(g)$  is carried out by taking 2 mol of  $S_8(g)$  and 0.2 mol of  $S_2(g)$  in a reaction vessel of 1L. Which one is not correct if  $K_c = 6.30 \times 10^{-6}$

A. Reaction quotient is  $8 \times 10^{-4}$

B. Reaction proceed in backward direction.

C. Reaction proceed is forward direction

D.  $K_p = 2.55 \text{ atm}^3$

**Answer: A::B::D**

 [Watch Video Solution](#)

4. For the equilibrium at 298K,  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ ,  $G_{N_2O_4}^\ominus = 100kJmol^{-1}$  and  $G_{NO_2}^\ominus = 50kJmol^{-1}$ . If 5 mol of  $N_2O_4$  and 2 moles of  $NO_2$  are taken initially in one litre container than which statement are correct.

A. reaction proceeds in forward direction

B.  $K_c = 1$

C.  $\Delta G = -0.55KJ$ ,  $\Delta G^\ominus = 0$

D. At equilibrium  $[N_2O_4] = 4.84M$  and  $[NO_2] = 0.212M$

**Answer: A::B::C::D**

 [Watch Video Solution](#)

5. Which are true for the reaction:  $A_2 \rightleftharpoons 2C + D$ ?

A. if  $\Delta H = 0$ ,  $K_p$  and increases with temperature and dissociation.

B. if  $\Delta H = +ve$ ,  $K_p$  increases with temperature and dissociation of  $A_2$  increases.

C. if  $\Delta H = -ve$ ,  $K_p$  decreases with temperature and dissociation of  $A_2$  decreases..

$$D. K_p = 4\alpha^3 \left[ \frac{P}{1 + 2\alpha} \right]^2$$

**Answer: A::B::C::D**

 [Watch Video Solution](#)

**6. van't Hoff equation is**

A.  $(d/dT)\ln K = -\Delta H/RT^2$

B.  $d/dT(\ln K) = +\Delta H/RT^2$

C.  $(d/dT)\ln K = -\Delta H/RT$

D.  $K = Ae^{-\Delta H/RT}$



**Answer: B::D**





Watch Video Solution

7. For given two equilibria attained in a container which are correct if degree of dissociation of A and A' are  $\alpha$  and  $\alpha'$  .



$$\text{A. } \frac{K_{p_1}}{K_{p_2}} = \left[ \frac{(3\alpha' + 2\alpha)}{(3\alpha + 2\alpha')} \right]^3 \times \frac{\alpha}{\alpha'}$$

$$\text{B. } P'_C / P'_D = 4$$

$$\text{C. } P'_B = 2P'_C + 2P'_D$$

$$\text{D. } \alpha > \alpha'$$

Answer: A::B::C::D



Watch Video Solution

8. In a reaction  $A_2(g) + 4B_2(g) \rightleftharpoons 2AB_4(g)$ ,  $\Delta H < 0$ . The formation of  $AB_4$  is not favoured by

- A. Low temperature and higher pressure
- B. High temperature and low pressure
- C. Low temperature and low pressure
- D. High temperature and high pressure

Answer: B::C::D



Watch Video Solution

9. The reaction which proceeds in the backward direction is





- A. (a)  $Fe_3O_4 + 6HCl \rightleftharpoons 2FeCl_3 + 3H_2O$
- B. (b)  $NH_3 + H_2O + NaCl \rightleftharpoons NH_4Cl + NaOH$
- C. (c)  $SnCl_4 + Hg_2Cl_2 \rightleftharpoons SnCl_2 + 2HgCl_2$



Answer: B::C::D

 [Watch Video Solution](#)

10. For which of the following reaction,  $K_p \neq K_c$ ?



Answer: A::B

 [Watch Video Solution](#)

11. Select the incorrect statements:

A.  $K_p$  or  $K_c$  are dimensionless if pressure or concentration are expressed in standard state.

B. The numerical value of  $K_p$  changes with experimental conditions, i.e. P, T, and C at which equilibrium is attained.

C. Active mass of reactant = concentration of reactant


D. Dissolution of  $NH_3$  in water increases with increasing pressure.

**Answer: A::B::C::D**



**Watch Video Solution**

**12.** For the chemical reaction



the amount of  $X_3Y$  at equilibrium is affected by

A. Temperature and pressure

B. Temperature only

C. Pressure only

D. Temperature, pressure, and catalyst

**Answer: B::C::D**

 [Watch Video Solution](#)

**13.** When two reactants A and B are mixed to give products C and D, the reaction quotient ( $Q$ ) at the initial stages of the reaction

A. Is zero

B. Decreases with time

C. Is independent of time

D. Increases with time

**Answer: A**

 [Watch Video Solution](#)

14. At constant temperature , the equilibrium constant ( $K_p$ ) for the decomposition reaction

$$N_2O_4 \rightleftharpoons 2NO_2, \text{ is expressed by } K_p = \frac{(4x^2p)}{(1 - x^2)}$$

where  $p$  = pressure ,  $x$  = extent of decomposition .which one of the following statements is true?


- A.  $K_p$  increases with increase of  $P$ .
- B.  $K_p$  increases with increases of  $x$ .
- C.  $K_p$  increase with decrease of  $x$ .
- D.  $K_p$  remains constant with change in  $p$  and  $x$

**Answer: D**



[Watch Video Solution](#)

15. Consider the following equilibrium in a closed container,



At a fixed temperature, the volume of the reaction container is halved. For this change which of the following statements holds true regarding the equilibrium constant ( $K_p$ ) and degree of dissociation ( $\alpha$ )?

- A. neither  $K_p$  nor  $\alpha$  changes
- B. Both  $K_p$  and  $\alpha$  change
- C.  $K_p$  changes but  $\alpha$  does not change
- D.  $K_p$  does not change but  $\alpha$  changes

**Answer: B::C::D**



[Watch Video Solution](#)

**16.** Which of the following do not change the value of  $K$  for a reaction?




- A. Addition of catalyst
- B. Increase in temperature
- C. Increase in pressure

D. Removal of one of the products

Answer: A::C::D

 [Watch Video Solution](#)

17. For which of the following reactions at equilibrium at constant temperature, doubling the volume will cause a shift to the right?



Answer: A::B

 [Watch Video Solution](#)



18. Unit of equilibrium constant is:

A.  $(\text{molL}^{-1})^{1-n}$

B.  $(\text{molL}^{-1})^{\Delta n}$

C.  $(\text{atm})^{\Delta n}$

D. All

Answer: B::C



Watch Video Solution

19. Which is/are correct?

A. (a)  $2.303\log K = -\Delta H^\ominus / RT + \Delta S^\ominus / R$

B. (b)  $\Delta G^\ominus = -2.303RT\log K$

C. (c)  $-2.303\log K = -\Delta H^\ominus / RT^2 + \Delta S^\ominus / R$

D. (d)  $2.303\log K = (1/RT)(\Delta H^\ominus + \Delta S^\ominus)$

**Answer: A::B**

 [Watch Video Solution](#)

20. For the reaction,  $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ , which is the correct representation?

A.  $K_p = (p_{\text{CO}_2})$




B.  $K_p = K_c(RT)$

C.  $K_p = (\text{CO}_2)/1$

D. None

**Answer: A::B::C**

 [Watch Video Solution](#)



Correct relation(s) between  $K_1, K_2, K_3$  and  $K_4$  is/are

A.  $K_1 \times K_3 = 1$

B.  $\sqrt{K_1} \times K_4 = 1$

C.  $\sqrt{K_3} \times K_2 = 1$

D. None

Answer: A::B::C



Watch Video Solution

22. The rate of disappearance of A at two temperature is given by  $A \rightleftharpoons B$

i.  $\frac{-d[A]}{dt} = 2 \times 10^{-2}[A] - 4 \times 10^{-3}[B]$  at 300K

ii.  $\frac{-d[A]}{dt} = 4 \times 10^{-2}[A] - 16 \times 10^{-4}[B]$  at 400K

From the given values of heat of reaction

A. 3.86kcal

B. 6.93kcal

C. 1.68kcal

D.  $1.68 \times 10^{-2}$ kcal

**Answer: B::C::D**



**Watch Video Solution**

23. Which of the following factors would favour the formation of ammonia?

A. Increase in temperature

B. Increase in pressure

C. Addition of catalyst

D. Addition of promoter

**Answer: B::C::D**



**Watch Video Solution**

24. Which of the following will not affect the value of equilibrium constant of a reaction?

A. Change in the concentration of the reactants

B. Change in temperature

C. Change in pressure

D. Addition of catalyst

**Answer: A::C::D**



**Watch Video Solution**

25. Which of the following statement is/are wrong?

- A. At equilibrium, concentrations of reactants and products become constant because the reaction stops.
- B. Addition of catalyst speeds up the forward reaction more than the backward reaction.
- C. Equilibrium constant of an exothermic reaction decreases with increase of temperature.
- D.  $K_p$  is always greater than  $K$ .

**Answer: A::B::D**



[Watch Video Solution](#)

26. When  $\text{NaNO}_3$  is heated in a closed vessel, oxygen is liberated and  $\text{NaNO}_2$  is left behind. At equilibrium, which are correct


- A. Addition of  $\text{NaNO}_2$  favours reverse reactions.
- B. Addition of  $\text{NaNO}_2$  favours forward reactions.
- C. Increasing temperature favours forward reaction
- D. Increasing pressure reverse reaction.

Answer: C::D

 [Watch Video Solution](#)

## Exercises (Single Correct)

1. In the dissociation of  $\text{PCl}_5$  as



If the degree of dissociation is  $\alpha$  at equilibrium pressure  $P$ , then the equilibrium constant for the reaction is

A.  $K_p = \frac{\alpha^2}{1 + \alpha^2 P}$

B.  $K_p = \frac{\alpha^2 P^2}{1 - \alpha^2}$


$$C. K_p = \frac{P^2}{1 - \alpha^2}$$

$$D. K_p = \frac{\alpha^2 P}{1 - \alpha^2}$$

**Answer: D**

 [Watch Video Solution](#)

2. For a hypothetical reaction of kind.



More  $AB_3$  could be produced at equilibrium by

- A. Using a catalyst
- B. Removing some of  $B_2$
- C. Increasing the temperature
- D. Increasing the pressure

**Answer: D**

 [Watch Video Solution](#)



3. The equilibrium constant for a reaction  $A + B \rightleftharpoons C + D$  is  $1 \times 10^{-2}$  at 298K and is 2 at 273K. The chemical process resulting in the formation of  $C$  and  $D$  is

- A. Exothermic
- B. Endothermic
- C. Unpredictable
- D. None

**Answer: B**

 [Watch Video Solution](#)

4. The solubility of  $CO_2$  in water increases with

- A. Increasing in temperature
- B. Reduction of gas pressure

C. Increasing in gas pressure

D. Increasing in volume

**Answer: C**

 [Watch Video Solution](#)

5. The equilibrium constant for a reaction

$A + 2B \rightleftharpoons 2C$  is 40. The equilibrium constant for reaction  $C \rightleftharpoons B + 1/2A$  is

A.  $1/40$

B.  $(1/40)^{1/2}$


C.  $(1/40)^2$

D. 40

**Answer: B**

 [Watch Video Solution](#)

6. Inert gas has been added to the following equilibrium system at constant volume



To which direction will the equilibrium shift?

- A. Forward
- B. Backward
- C. No effect
- D. Unpredictable

**Answer: C**

 [Watch Video Solution](#)

7. The equilibrium constant  $K$  for the reaction  $2HI(g) \rightleftharpoons H_2(g) + I_2(g)$  at room temperature is 2.85 and that at 698K is  $1.4 \times 10^{-2}$ . This implies

- A. Exothermic

B. Endothermic

C. Exergonic


D. Unpredictable

**Answer: A**



**Watch Video Solution**

8. The decomposition of  $N_2O_4$  to  $NO_2$  is carried out at  $280^\circ C$  in chloroform. When equilibrium is reached, 0.2 mol of  $N_2O_4$  and  $2 \times 10^{-3}$  mol of  $NO_2$  are present in a 2L solution. The equilibrium constant for the reaction



A.  $1 \times 10^{-2}$

B.  $2 \times 10^{-3}$

C.  $1 \times 10^{-5}$

D.  $2 \times 10^{-5}$

**Answer: C**

 [Watch Video Solution](#)

9. For the reaction  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ , the degree of dissociation at equilibrium is 0.2 at 1 atm pressure. The equilibrium constant  $K_p$  will be

A. 1/2

B. 1/4

C. 1/6

D. 1/8

**Answer: C**

 [Watch Video Solution](#)

10. 4 mol of carbon dioxide was heated in  $1dm^3$  vessel under conditions which produced at equilibrium 25 % dissociation into carbon monoxide

and oxygen. The number of moles of carbon monoxide produced

A. 0.5

B. 1.0

C. 2.0

D. 4.0

**Answer: B**



[Watch Video Solution](#)

**11.** 1 mol of  $N_2$  is mixed with 3 mol of  $H_2$  in a litre container. If 50 % of  $N_2$  is converted into ammonia by the reaction  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ , then the total number of moles of gas at the equilibrium are

A. 1.5

B. 4.5

C. 3.0

D. 6.0

**Answer: C**



**Watch Video Solution**

12. The equilibrium constant of a reaction is 300, if the volume of the reaction flask is tripled, the equilibrium constant will be

A. 100

B. 300

C. 250

D. 150

**Answer: B**



**Watch Video Solution**

13. For reaction :  $H_2(h) + I_2(g) \rightleftharpoons 2HI(g)$  at certain temperature, the value of equilibrium constant is 50. If the volume of the vessel is reduced to half of its original volume, the value of new equilibrium constant will be

- A. 25
- B. 50
- C. 100
- D. Unpredictable

**Answer: B**

 [Watch Video Solution](#)

14. The system  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$  attains equilibrium. If the equilibrium concentration of  $PCl_3(g)$  is doubled, the concentration of  $Cl_2(g)$  would become

- A. 1/4 its original value



B.  $1/2$  its original value

C. Twice its original value

D. Unpredictable

**Answer: D**

 [Watch Video Solution](#)

15.  $XY_2$  dissociates  $XY_2(g) \rightleftharpoons XY(g) + Y(g)$ . When the initial pressure of  $XY_2$  is 600 mm Hg, the total equilibrium pressure is 800 mm Hg. Calculate  $K$  for the reaction Assuming that the volume of the system remains unchanged.

A. 50.0

B. 100.0


C. 166.6

D. 400.0

**Answer: B**

 [Watch Video Solution](#)

**16.** Consider the reaction



Which occurs in one step. The specific rate constant are 0.25 and 5000 for the forward and reverse reaction, respectively. The equilibrium constant is

A.  $2.0 \times 10^{-4}$

B.  $4.0 \times 10^2$


C.  $5.0 \times 10^{-5}$

D.  $2.5 \times 10^{-6}$

**Answer: C**

 [Watch Video Solution](#)

17. For the equilibrium system



the equilibrium constant is  $1.0 \times 10^{-5}$ . What is the concentration of HX if the equilibrium concentration of  $H_2$  and  $X_2$  are  $1.2 \times 10^{-3}$  M, and  $1.2 \times 10^{-4}$  M respectively?

A.  $12 \times 10^{-4}M$

B.  $12 \times 10^{-3}M$

C.  $12 \times 10^{-2}M$



D.  $12 \times 10^{-1}M$

**Answer: C**



**Watch Video Solution**

18. In alkaline solution, the following equilibria exist



$K_1$  and  $K_2$  have values 12 and 11, respectively.

$S_3^{2-} \rightarrow S^{2-} + 2S$ . What is equilibrium constant for the reaction

A. 132

B.  $7.58 \times 10^{-3}$

C. 1.09



D. 0.918

**Answer: B**



[Watch Video Solution](#)

**19.** Given the equilibrium constants



The equilibrium constant for the disproportionation equilibrium



A. (a)  $-3.3 \times 10^5$

B. (b)  $3 \times 10^{-5}$

C. (c)  $3.3 \times 10^5$

D. (d)  $3 \times 10^{-6}$

**Answer: D**



**Watch Video Solution**

**20.** When the reaction,  $2NO_2(g) \rightleftharpoons N_2O_4(g)$  reaches equilibrium at 298K. The partial pressure of  $NO_2$  and  $N_2O_4$  are 0.2Kpa and 0.4Kpa, respectively. What is the equilibrium constant  $K_p$  of the above reaction at 298K?

A. 0.1

B. 0.5

C. 1.0

D. 10

**Answer: D**

 [Watch Video Solution](#)

21. The vapour density of mixture consisting of  $NO_2$  and  $N_2O_4$  is 38.3 at  $26.7^\circ C$ . Calculate the number of moles of  $NO_2$  in 100g of the mixture.

A. 0.2

B. 0.4

C. 0.8

D. 1.6

**Answer: B**

 [Watch Video Solution](#)

22. In the problem number 21, the number of mole of  $N_2O_4$  in 100g of the mixture is:

A. 0.43

B. 0.86

C. 0.57

D. 0.2

**Answer: B**



[Watch Video Solution](#)

**23.** One mole of  $SO_3$  was placed in a litre reaction flask at a given temperature when the reaction equilibrium was established in the reaction.

$2SO_3 \rightleftharpoons 2SO_2 + O_2$  the vessel was found to contain 0.6 mol of  $SO_2$ . The value of the equilibrium constant is

A. 0.36

B. 0.675

C. 0.45

D. 0.54

**Answer: B**



[Watch Video Solution](#)

24. The equilibrium constant for the reaction  $w + x \rightleftharpoons y + z$  is 9. If one mole of each of  $w$  and  $x$  are mixed and there is no change in volume, the number of moles of  $y$  formed is

A. (a) 0.10

B. (b) 0.50

C. (c) 0.75

D. (d) 0.54

**Answer: C**



[Watch Video Solution](#)



25. In the gaseous equilibrium

$A + 2B \rightleftharpoons C + \text{Heat}$ , the forward reaction is favoured:

- A. Low P, High T
- B. Low P, Low T
- C. High P, Low T
- D. High P, High T

**Answer: C**



[Watch Video Solution](#)

26. The active mass of 64g of HI in a 2 - L flask would be

- A. 2
- B. 1
- C. 5
- D. 0.25

**Answer: D**

 [Watch Video Solution](#)

27. For  $N_2 + 3H_2 \rightleftharpoons 2NH_3 + \text{Heat}$

A.  $K_p = K_c$

B.  $K_p = K_c RT$

C.  $K_p = K_c (RT)^{-2}$

D.  $K_p = K_c (RT)^{-1}$

**Answer: C**

 [Watch Video Solution](#)

28. For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ , the equilibrium constant changes with:

- A. Total pressure
- B. Catalyst
- C. The amounts of  $H_2$  and  $I_2$  present
- D. Temperature

**Answer: D**

 [Watch Video Solution](#)

29. The equilibrium constant  $K$  for the reaction  $2HI(g) \rightleftharpoons H_2(g) + I_2(g)$  at room temperature is 2.85 and that at 698K is  $1.4 \times 10^{-2}$ . This implies



- A.  $HI$  is exothermic compound
- B.  $HI$  is very stable at room temperature
- C.  $HI$  is relatively less stable than  $H_2$  and  $I_2$  at room temperature
- D.  $HI$  is resonance stabilised

**Answer: C**



Watch Video Solution

30.  $K_1$  and  $K_2$  are equilibrium constant for reactions (i) and (ii)



Then,

A.  $K_1 = (1/K_2)^2$

B.  $K_1 = K_2^2$

C.  $K_1 = 1/K_2$

D.  $K_1 = (K_2)^\circ$

Answer: A



Watch Video Solution

31. The equilibrium constant  $K_p$  for a homogeneous gaseous reaction is  $10^{-8}$ . The standard Gibbs free energy change  $\Delta G^\ominus$  for the reaction

(using  $R = 2\text{calK}^{-1}\text{mol}^{-1}$ ) is


- A.  $10.98\text{kcal}$
- B.  $-1.9\text{kcal}$
- C.  $-4.1454\text{kcal}$
- D.  $+4.1454\text{kcal}$

**Answer: A**



[Watch Video Solution](#)

**32.** Which of the following will not change the concentration of ammonia in the equilibrium



- A. increase of temperature
- B. increase of volume
- C. decrease of volume

D. addition of catalyst

**Answer: D**

 [Watch Video Solution](#)

**33.** In a chemical reaction, equilibrium is said to have been established when the

A. Concentrations of reactants and products are equal

B. Opposing reactions ceases


C. Velocities of opposing reaction become equal

D. Temperature of opposing reactions are equal

**Answer: C**

 [Watch Video Solution](#)

34. In a chemical reaction



- A. Equal volumes of  $N_2$  and  $H_2$  are reacting
- B. Equal masses of  $N_2$  and  $H_2$  are reacting
- C. The reaction has stopped
- D. The same amount of ammonia is formed as is decomposed into  $N_2$  and  $H_2$

**Answer: D**



[Watch Video Solution](#)

35. The equilibrium constant of a reversible reaction at a given temperature

- A. Depends on initial concentration, of the reactants.
- B. Depends on the concentration of the products at equilibrium.

C. Does not depend on the initial concentration.

D. It is not characteristic of the reaction

**Answer: C**

 [Watch Video Solution](#)

**36.** According to Le- Chatelier's principle. Adding heat to a solid  $\rightleftharpoons$  liquid equilibrium will cause the.

A. Amount of solid to decrease

B. Amount of liquid to decrease

C. Temperature to rise

D. Temperature to fall

**Answer: A**

 [Watch Video Solution](#)



37. In the formation of nitric acid,  $N_2$  and  $O_2$  are made to combine. Thus,  $N_2 + O_2 \rightleftharpoons 2NO + \text{Heat}$  which of the following condition will favour the formation of NO?

- A. low temperature
- B. high temperature
- C. freezing point
- D. all are favourable

**Answer: A**

 [Watch Video Solution](#)

38. Which of the following factors will favour the reverse reaction in a chemical equilibrium?

- A. increase in concentration of one of the reactants
- B. increase in concentration of one of the products

C. removal of one of the products regularly

D. None of these

**Answer: B**

 [Watch Video Solution](#)

**39.** For the system  $A(g) + 2B(g) \rightleftharpoons C(g)$  the equilibrium concentration is

$$A = 0.06 \text{ molL}^{-1}, B = 0.12 \text{ molL}^{-1}$$

$C = 0.216 \text{ molL}^{-1}$  The  $K_{eq}$  for the reaction is

A. 250

B. 416

C.  $4 \times 10^{-3}$

D. 125

**Answer: A**

 [Watch Video Solution](#)

40. When 4mol of A is mixed with 4mol of B, 2mol of C and D are formed at equilibrium, according to the reaction



the equilibrium constant is

A. 4

B. 1

C. 1/2


D. 1/4

**Answer: B**



[Watch Video Solution](#)

41. Consider the reaction



in closed container at equilibrium. What would be the effect of addition of  $\text{CaCO}_3$  on the equilibrium concentration of  $\text{CO}_2$ ?

- A. Increases
- B. Decreases
- C. Data is not sufficient
- D. Remains unaffected

**Answer: D**

 [Watch Video Solution](#)

**42.** The equilibrium constant for the reaction  $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$  is  $4 \times 10^{-4}$  at 200 K. In presence of a catalyst at 200K is:





- A.  $40 \times 10^{-4}$
- B.  $4 \times 10^{-4}$
- C.  $4 \times 10^{-2}$

D. incomplete data

**Answer: B**

 [Watch Video Solution](#)


**43.** In which of the following reaction, the yield of the products does not increase by increase in the pressure?



**Answer: A**

 [Watch Video Solution](#)

44. At a certain temperature, only 50% HI is dissociated at equilibrium in the following reaction:



the equilibrium constant for this reaction is:

- A. 1.0
- B. 3.0
- C. 0.5
- D. 0.25

**Answer: D**



[Watch Video Solution](#)

45. For a reaction  $A(g) \rightleftharpoons B(g) + C(g)$ ,  $K_p$  at  $400^\circ C$  is  $1.5 \times 10^{-4}$  and  $K_p$  at  $600^\circ C$  is  $6 \times 10^{-3}$ . Which statement is incorrect?

- A. The reaction is exothermic

B. Increase in temperature increases the formation of B


C. Increase in pressure increases the formation of A

D. Decrease in temperature and increase in pressure shift the equilibrium towards left

**Answer: A**

 [Watch Video Solution](#)

**46.** 8 mol of gas  $AB_3$  are introduced into a  $1.0\text{dm}^3$  vessel. It dissociates as



At equilibrium, 2 mol of  $A_2$  is found to be present. The equilibrium constant for the reaction is

A. (a)  $2\text{mol}^2\text{L}^{-2}$

B. (b)  $3\text{mol}^2\text{L}^{-2}$

C. (c)  $27\text{mol}^2\text{L}^{-2}$


D. (d)  $36\text{mol}^2\text{L}^{-2}$

**Answer: C**



**Watch Video Solution**

47. 1 mol of  $XY(g)$  and 0.2 mol of  $Y(g)$  are mixed in 1 L vessel. At equilibrium, 0.6 mol of  $Y(g)$  is present. The value of  $K$  for the reaction



A.  $0.04 \text{ molL}^{-1}$

B.  $0.06 \text{ molL}^{-1}$

C.  $0.36 \text{ molL}^{-1}$

D.  $0.40 \text{ molL}^{-1}$


**Answer: D**



**Watch Video Solution**



48. How will the lowering of temperature affect the chemical equilibrium in the system



- A. Relative concentration of products and reactants does not change.
- B. Relative concentration of products and reactants change.
- C. Equilibrium is shift to the left.
- D. Equilibrium is shift to the right.

Answer: B::D



Watch Video Solution

49. For the reaction  $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ , the value of  $K_p$  is  $1.7 \times 10^3$  at 500K and  $1.7 \times 10^4$  at 600K. Which of the following is/are correct ?

- A. The proportions of  $NO_2$  in the equilibrium mixture is increased by decrease in pressure.

B. The standard enthalpy change for the forward reaction is negative

C. Units of  $K_p$  are  $\text{atm}^{-1}$

D. At 500K the degree of dissociation of  $N_2O_4$  decreases by 50 % by increasing the pressure by 100 %

**Answer: A**

 [Watch Video Solution](#)

50. At equilibrium  $X + Y \rightleftharpoons 3Z$ , 1 mol of X, 2 mol of Y and 4 mol of Z are contained in a 3 - L vessel. What will be the value of the reaction coefficient Q,

A. 10

B. 15

C. 10.67


D. None of these

**Answer: C**



**Watch Video Solution**

51. What concentration of  $CO_2$  be in equilibrium with  $0.025M$   $CO$  at  $120^\circ C$  for the reaction



if the value of  $K_c = 5.0$  ?

A.  $0.125M$

B.  $0.0125M$

C.  $1.25M$




D.  $12.5M$

**Answer: A**



**Watch Video Solution**

52. Which of the following reactions will not be affected by increasing the pressure?




Answer: B::D



Watch Video Solution

53. The value of  $K_c = 4.24$  at  $800K$  for the reaction.



Calculate equilibrium concentration of  $CO_2$ ,  $H_2$ ,  $CO$  and  $H_2O$  at  $800K$ . If

only  $CO$  and  $H_2O$  are present initially at concentrations of  $0.10M$  each.

A. Adding a suitable catalyst

B. Adding an inert gas

C. Decreasing the volume of the container


D. Increasing the amount of  $CO(g)$

**Answer: D**



**Watch Video Solution**

**54.** For the chemical reaction



The amount of  $X_3Y$  at equilibrium is not affected by

A. Temperature and pressure

B. Temperature only

C. Pressure only

D. Temperature, pressure, and catalyst

**Answer: A**



Watch Video Solution

55. When two reactants A and B are mixed to give products C and D, the reaction quotient (Q) at the initial stages of the reaction

- A. Is zero
- B. Decreases with time
- C. Is independent of time
- D. Increases with time

**Answer: D**



Watch Video Solution

56. At constant temperature, the equilibrium constant ( $K_p$ ) for the decomposition reaction

$$N_2O_4 \rightleftharpoons 2NO_2, \text{ is expressed by } K_p = \frac{(4x^2p)}{(1-x^2)}$$

where  $p$  = pressure,  $x$  = extent of decomposition. Which one of the following statements is true?

- A.  $K_p$  increase with increase in  $p$
- B.  $K_p$  increases with increase in  $x$
- C.  $K_p$  increases with decrease in  $x$ .
- D.  $K_p$  remains constant with change in  $p$  and  $x$

 [Watch Video Solution](#)

57. The equilibrium constant  $K_{p1}$  and  $K_{p2}$  for the reactions  $X \rightleftharpoons 2Y$  and  $Z \rightleftharpoons P + Q$ , respectively, are in the ratio of 1:9. If the degrees of dissociation of  $X$  and  $Z$  are equal, then the ratio of total pressure at equilibria is

- A. 1:36
- B. 1:9

C. 1:6

D. 1:4

 [Watch Video Solution](#)

58. For the reaction  $X \rightleftharpoons 2Y$  and  $Z \rightleftharpoons P + Q$  occurring at two different pressures  $P_1$  and  $P_2$ , respectively. The ratio of the two pressures is 3:1. What will be the ratio of equilibrium constants  $K_{p2}:K_{p1}$ , if the degrees of dissociation of X and Z are equal.

A. (a) 1:36

B. (b) 1:12

C. (c) 1:9

D. (d) 2:3

 [Watch Video Solution](#)



## Exercises (Assertion-Reasoning)

1. Assertion :  $K_p$  can be less than, greater than or equal to  $K_c$

Reason : Relation between  $K_p$  and  $K_c$  depends on the change in number of moles of gaseous reactants and products ( $\Delta n$ ).

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: A**



[Watch Video Solution](#)

2. Assertion (A) : For  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ , the equilibrium constant is  $K$ . The for  $\frac{1}{2}N_2(g) + \frac{3}{2}H_2(g) \rightleftharpoons NH_3(g)$ , the equilibrium constant will be  $\sqrt{K}$ .

Reason (R) : If concentrations are changed to half, the equilibrium constants will be halved.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: C**



**Watch Video Solution**

3. Assertion (A) : The equilibrium constant is fixed and characteristic for any given chemical reaction at a specified temperature.

Reason (R) : The composition of the final equilibrium mixture at a particular temperature depends upon the starting amount of reactants.

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: A**



**Watch Video Solution**

4. Assertion (A) :  $K_p$  is always greater than  $K_c$ .

Reason (R) : The effect of pressure is greater on the rate of reaction than the effect of concentration.

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If both (A) and (R) are incorrect.

**Answer: D**

 [Watch Video Solution](#)

5. Assertion (A) : A catalyst does not influences the values of equilibrium constant

Reason (R) : Catalyst influences the rate of both forward and backward reactions equally.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: A**

 [Watch Video Solution](#)

6. Assertion (A) : Equilibrium constant of a reaction increases if temperature is increased

Reason (R) : The forward reaction becomes faster with increase of temperature.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: C**



**Watch Video Solution**

7. Assertion (A) : The active mass of pure solid and pure liquid is taken unity.

Reason (R) : The active mass of pure solids and liquids depends on the density and molecular mass. The density and molecular of a mass of pure liquids and solids are constant.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: A**



[Watch Video Solution](#)

8. Assertion (A) : For  $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ , if more  $Cl_2$  is added the equilibrium will shift in backward direction. Hence, equilibrium constant will decrease.

Reason (R) : Addition of inert gas to the equilibrium mixture at constant volume does not alter the equilibrium.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: C**



**Watch Video Solution**

9. Assertion (A) : Adding inert gas to dissociation equilibrium of  $N_2O_4$  at constant pressure and temperature increases the dissociation.

Reason (R) : molar concentration of the reactants and products decreases.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)



- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: A**

 [Watch Video Solution](#)

**10.** Assertion (A) : The value of  $K$  for a reaction may increase or decrease with increase in temperature depending upon whether the reaction is exothermic or endothermic.

Reason (R) : With increase in temperature, the extent of reaction increases.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)

- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)
- C. If (A) is correct, but (R) is incorrect
- D. If (A) is incorrect, but (R) is correct.

**Answer: B**

 [Watch Video Solution](#)

**11.** Assertion (A) : When a catalyst is added to a reaction mixture in equilibrium the amount of the products increases.

Reason (R) : The forward reaction becomes faster on adding the catalyst.

- A. If both (A) and (R) are correct, and (R) is the correct explanation for (A)
- B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A)


C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: C**

 [Watch Video Solution](#)

**12. Assertion (A) :** For the reaction



**Reason (R) :** In this reaction, the sum of stoichiometric coefficient of reactants is equal to the sum of stoichiometric coefficients of products.

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect


D. If (A) is incorrect, but (R) is correct.

**Answer: A**




**Watch Video Solution**

**13.** Assertion (A) : A change of pressure has no effect in case of the equilibrium,



Reason (R) : The reaction,



A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: C**



[Watch Video Solution](#)

**14.** Assertion (A) : The value of  $K$  increases with increase in temperature in case of endothermic reaction

Reason (R) : The increase in temperature shifts the equilibrium in the backward direction in case of exothermic reaction.

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: D**



[Watch Video Solution](#)

15. Assertion (A) : Greater the value of  $K$ , more is the fraction of initial concentration of reactants converted to products at equilibrium.

Reason (R) : The value of  $K$  depends on the initial concentration of reactants.

A. If both (A) and (R) are correct, and (R) is the correct explanation for

(A)

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A)

C. If (A) is correct, but (R) is incorrect

D. If (A) is incorrect, but (R) is correct.

**Answer: C**



**Watch Video Solution**

**Exercises (Integer)**

1. A reaction attains equilibrium, when the free energy change is

A. (a) 1

B. (b) 2

C. (c) 3

D. (d) 0

**Answer: D**



[Watch Video Solution](#)

2. For a homogeneous chemical reaction,  $K_p = K_c$  when

A.  $\Delta n = 0$

B.  $\Delta n = 1$

C.  $\Delta n = 2$

D.  $\Delta n = \infty$

**Answer: A**



[Watch Video Solution](#)

3. For the reaction  $A + B \rightleftharpoons C$ , the rate constants for the forward and the reverse reactions are  $4 \times 10^2$  and  $2 \times 10^2$  respectively. The value of equilibrium constant  $K$  for the reaction would be

A. 1

B. 2

C. 3

D. 4

**Answer: B**



[Watch Video Solution](#)



4. The equilibrium constant for the reactions

$A + B \rightleftharpoons AB$  is 0.5 at 200K. The equilibrium constant for the reaction

$AB \rightleftharpoons A + B$  would be

A. 1

B. 2

C. 3

D. 4

**Answer: B**



[Watch Video Solution](#)

5. One mole of ethanol is treated with one mole of ethanoic acid at 25 ° C.

Half of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be

A. 1

B. 2

C. 3

D. 4

**Answer: A**



**Watch Video Solution**

6. In the reaction  $A + B \rightleftharpoons AB$ , if the concentration of A and B is increased by a factor of 2, it will cause the equilibrium concentration of AB to change to

A. Two times to original value

B. Three times to original value

C. Same

D. Zero

**Answer: A**

 [Watch Video Solution](#)

7. At equilibrium, the value of equilibrium constant  $K$  is

- A. 1
- B. 2
- C. 3
- D. 0

**Answer: A**

 [Watch Video Solution](#)

## Exercises (Fill In The Blanks)

1. At equilibrium stage, the rate of forward reaction is ..... to the rate of backward reaction.

 [Watch Video Solution](#)

2. The equilibrium constant does not depend on the initial concentrations of the reactants but depends on ..... of various reactants and products at .....

 [Watch Video Solution](#)

3. In the reaction  $2NO(g) \rightleftharpoons N_2(g) + O_2(g)$ , the values of  $K_c$  and  $K_p$  are ..... at a given temperature.

 [Watch Video Solution](#)

4. Number of moles when divided by the total volume in litre gives ..... of the respective species.

 [Watch Video Solution](#)

5. The equilibrium state is attained when the reversible reaction is carried out in a..... space.

 [Watch Video Solution](#)

6. The chemical equilibrium is ..... in nature.

 [Watch Video Solution](#)

7. A catalyst ..... the equilibrium state but helps to attain in lesser time.

 [Watch Video Solution](#)

8. The equilibrium concentration of x, y and z are 4, 2 and 2 mol  $L^{-1}$ , respectively, at equilibrium of the reaction  $2x + y \rightleftharpoons z$ . The value of  $K_c$  is .....



 [Watch Video Solution](#)

9. At equilibrium , the amount of each constituent of reaction mixture becomes .....

 [Watch Video Solution](#)

10. The equilibrium constant has no unit if  $\Delta n = \dots\dots\dots$

 [Watch Video Solution](#)

11. The relation between  $K_p$  and  $K_c$  of a reversible reaction at constant temperature is  $K_p = \dots\dots\dots$

 [Watch Video Solution](#)

12. For the reaction,  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ , the units of  $K_p$  are .....



 [Watch Video Solution](#)

13. In the reaction  $A + B \rightleftharpoons C + D$ , the value of equilibrium constant is 10. If the rate constant of forward reaction is 80, the rate constant of backward reaction is .....

 [Watch Video Solution](#)

14. A tenfold increase in pressure on the reaction  $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$  at equilibrium result in ..... in  $K_p$ .

 [Watch Video Solution](#)

15. The equilibrium constant for the reaction  $2A + 2B \rightleftharpoons 2C + 2D$  is 200. The equilibrium constant for the reaction  $A + B \rightleftharpoons C + D$ , at the same temperature is .....

 [Watch Video Solution](#)

16. If the activation energies of the forward and backward reactions of a reversible reaction are  $E_a(f)$  and  $E_a(b)$ , respectively. The  $\Delta E$  of the reaction is .....

 [Watch Video Solution](#)

17. If the value of equilibrium constant is large, ..... Are more stable.

 [Watch Video Solution](#)

18. The magnitude of equilibrium constant is a measure of ..... to which the reversible reaction proceeds in a particular direction at a given .....

 [Watch Video Solution](#)

19. Le Chatelier's principle is applicable to both ..... and ..... equilibria.





[Watch Video Solution](#)

20. Low temperature is favourable for ..... reactions.



[Watch Video Solution](#)

21. Low pressure is favourable for those reversible reactions in which there is ..... in the number of molecules.



[Watch Video Solution](#)

22. If the temperature of the system at equilibrium is increased, the equilibrium will shift in the direction which ..... heat.



[Watch Video Solution](#)

23. An endothermic reaction which proceeds with decrease in volume will give maximum yield of the products at ..... and .....

 [Watch Video Solution](#)

24. The formation of ammonia by Haber's process is favoured by ..... pressure.

 [Watch Video Solution](#)

25. Low pressure favours those reactions which occur with ..... in the number of molecule.

 [Watch Video Solution](#)

26. For a system of gases A, B, C, and D at equilibrium  $A + 2B \rightleftharpoons C + 3D$ , the partial pressures are found to be  $A = 2.0$ ,  $B = 2.0$ ,  $C = 3.0$ , and

$D = 5.0 \text{ atm}$ . The value of equilibrium constant is .....

 [Watch Video Solution](#)

27.  $A + B \rightleftharpoons C + D$

If initially the concentration of A and B are both equal but at equilibrium concentration of D will be twice that of A, then what will be the equilibrium constant of the reaction?

 [Watch Video Solution](#)

28. For reaction  $PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g)$ ,  $K_c$  is 30 at 300K. The value of  $K_p$  at 300K is .....

 [Watch Video Solution](#)

29. The equilibrium constant for the reaction

$CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$  is:

 [Watch Video Solution](#)

30. The vapour density of  $PCl_5$  is 104.16 but when heated to  $230^\circ C$ , its vapour density is reduced to 62. The degree of dissociation of  $PCl_5$  at  $230^\circ C$  is .....

 [Watch Video Solution](#)

31. In line kilns, the following reaction,



proceeds to completion because of

 [Watch Video Solution](#)

32. The degree of dissociation of  $PCl_5$  will be more at ..... pressure.

 [Watch Video Solution](#)

33. When the system  $2HI(g) \leftrightarrow H_2(g) + I_2(g)$  is at equilibrium, inert gas is introduced. Dissociation of  $HI$  is .....

 [Watch Video Solution](#)

34. When a product is removed from the system which is at equilibrium ....., reaction is favoured.

 [Watch Video Solution](#)

35. The melting of ice is favoured by ..... pressure and ..... temperature.

 [Watch Video Solution](#)

## Exercises (True/False)

1. The dissociation of  $CaCO_3$  is suppressed at high pressure

 [Watch Video Solution](#)

2. More of  $SO_3$  decompose at low temperature.

 [Watch Video Solution](#)

3. Addition of inert gas to system at equilibrium changes only  $K_p$  not  $K_c$ .

 [Watch Video Solution](#)

4. The melting of ice in water decreases with increase in pressure.

 [Watch Video Solution](#)

5. The evaporation of liquid with increase in pressure.

 [Watch Video Solution](#)

6. If equilibrium constant for the reaction



the equilibrium constant  $k'$  is  $1/K$ .

 [Watch Video Solution](#)

7.  $K_p$  is equal to  $K_c$  if  $\Delta n$  is positive.

 [Watch Video Solution](#)

8. The value  $K_c$  of a reaction has a higher value at higher temperature.

The reaction is exothermic in nature.

 [Watch Video Solution](#)

9. The reaction having higher value of equilibrium constant is faster than the reaction having lower value of equilibrium constant.(T/F)



[Watch Video Solution](#)

10. Ammonium chloride dissociates as,



The vapour density becomes half the initial value when degree of dissociation is 0.5.



[Watch Video Solution](#)

11. The law of mass action applicable to heterogeneous equilibria.



[Watch Video Solution](#)

12. Equilibrium can be achieved only in an open vessel. (T/F)



[Watch Video Solution](#)



13. The solubility of sodium hydroxide increases with increase of temperature.

 [Watch Video Solution](#)

14. The degree of dissociation of  $PCl_5$  decreases with increase in pressure.

 [Watch Video Solution](#)

15. High pressure and low temperature are favourable conditions for the synthesis of ammonia.

 [Watch Video Solution](#)

16. An endothermic reaction proceeds faster in the forward reaction with decrease in temperature.





[Watch Video Solution](#)

17. A catalyst increases the rate of forward reaction and decrease the rate of backwark reaction.



[Watch Video Solution](#)

18. The value of  $K$  does not depends upon pressure.



[Watch Video Solution](#)

19. For any reaction, greater the value of equilibrium constant greater is the extent of reaction.



[Watch Video Solution](#)

20. Solid  $\leftrightarrow$  liquid equilibrium can be achieved only at melting point of the substance.

 [Watch Video Solution](#)

21. Assertion : The additions of an inert gas at constant volume to a system at equilibrium does not affect the state of equilibrium

Reason: The inert gas does not react with any of the reactants or products.

 [Watch Video Solution](#)

22. For a reversible system at a constant temperature, the value of  $K_c$  increases if the concentrations are changed at equilibrium.

 [Watch Video Solution](#)

23. The equilibrium constant is 10 at 100K. Hence,  $\Delta G$  will be negative.

 [Watch Video Solution](#)

24. Unit of  $K_p$  is  $(\text{atm})^{\Delta n}$

 [Watch Video Solution](#)

25. The value of equilibrium constant is independent of the speed with which the equilibrium is attained.

 [Watch Video Solution](#)

26. In Haber's process, once the equilibrium is established, addition of nitrogen decreases the yield of ammonia.

 [Watch Video Solution](#)

27. At chemical equilibrium , the concentration of all reactants and products are equal

 [Watch Video Solution](#)

28. The equilibrium state can be attained from both sides of the chemical reaction.

 [Watch Video Solution](#)

29. A reaction continues even after the attainment of equilibrium.


 [Watch Video Solution](#)

30. The equilibrium can be attained at a faster rate if one of the products is allowed to escape from the reaction mixture.

 [Watch Video Solution](#)

## Archives (Multiple Correct)

1. For the gas phase reaction



carried out in a vessel, the equilibrium concentration of  $C_2H_4$  can be increased by

- A. Increasing the temperature
- B. Decreasing the temperature
- C. Removing some  $H_2$
- D. Adding some  $C_2H_6$

**Answer: A::C::D**



[Watch Video Solution](#)

2. When  $\text{NaNO}_3$  is heated in a closed vessel, oxygen is liberated and  $\text{NaNO}_2$  is left behind. At equilibrium, which are correct

- A. Addition of  $\text{NaNO}_2$  favours reverse reaction.
- B. Addition of  $\text{NaNO}_3$  favours forward reaction
- C. Increasing the temperature favours forward reaction.
- D. Increasing the pressure favours reverse reaction.

**Answer: C::D**

 [Watch Video Solution](#)

3. The equilibrium  $\text{SO}_2\text{Cl}_2(g) \rightleftharpoons \text{SO}_2(g) + \text{Cl}_2(g)$  is attained at  $25^\circ\text{C}$  in a closed container and an inert gas, helium, is introduced. Which of the following statement is/are correct?

- A. The concentrations of  $\text{SO}_2$ ,  $\text{Cl}_2$  and  $\text{SO}_2\text{Cl}_2$  change.
- B. More chlorine is formed.

C. The concentration of  $\text{SO}_2$  is reduced.

D. All are incorrect.

**Answer: D**

 [Watch Video Solution](#)

4. For the reaction  $\text{PCl}_{5(g)} \rightleftharpoons \text{PCl}_{3(g)} + \text{Cl}_{2(g)}$ , the forward reaction at constant temperature is favoured by

A. Introducing an inert gas at constant volume.

B. Introducing chlorine gas at constant volume,

C. Introducing an inert gas at constant pressure


D. Increasing the volume of the container.

**Answer: C::D**

 [Watch Video Solution](#)



5. For the reaction



at a given temperature, the equilibrium amount of  $\text{CO}_2(g)$  can be increased by

- A. Adding a suitable catalyst
- B. Adding an inert gas
- C. Decreasing the volume of the container
- D. Increasing the amount of  $\text{CO}(g)$

**Answer: D**

 [Watch Video Solution](#)

6. The equilibrium  $2\text{Cu}^I \rightleftharpoons \text{Cu} + \text{Cu}^{II}$

In aqueous medium at  $25^\circ\text{C}$  shifts towards the left in the presence of

- A.  $\text{NO}_3^\ominus$

B.  $Cl^{\ominus}$

C.  $SCN^{\ominus}$

D.  $CN^{\ominus}$

**Answer: B::C::D**



**Watch Video Solution**

## Archives (Single Correct)

1. For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ , the equilibrium constant changes with:

A. Total pressure

B. Catalyst

C. The amounts of  $H_2$  and  $I_2$  present

D. Temperature

**Answer: D**

 [Watch Video Solution](#)





2. Pure ammonia is placed in a vessel at a temperature where its dissociation constant ( $\alpha$ ) is appreciable. At equilibrium,

- A.  $K_p$  does not change significantly with pressure
- B.  $\alpha$  does not change with pressure
- C. The concentration of  $NH_3$  does not change with pressure.
- D. The concentration of hydrogen is less than that of nitrogen.

**Answer: A**

 [Watch Video Solution](#)

3. An example of a reversible reaction is



**Answer: D**

 [Watch Video Solution](#)

4. One "mole" of  $N_2O_4(g)$  at 100K is kept in a closed container at 1.0 atm pressure. It is heated to 400K, where 30 % by mass of  $N_2O_4(g)$  decomposes to  $NO_2(g)$ . The resultant pressure will be

A. 1.2 atm

B. 2.4 atm


C. 2.0 atm

D. 1.0 atm

**Answer: B**

 [Watch Video Solution](#)

5. For the chemical reaction




the amount of  $X_3Y$  at equilibrium is affected by

- A. Temperature and pressure
- B. Temperature only
- C. Pressure only
- D. Temperature, pressure, and catalyst

**Answer: A**

 [Watch Video Solution](#)

6. For the reversible reaction



at  $500^\circ C$ , the value of  $K_p$  is  $1.44 \times 10^{-5}$  when the partial pressure is measured in atmosphere. The corresponding value of  $K_c$  with concentration in  $\text{mol L}^{-1}$  is

A.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 500)^{-2}}$

B.  $\frac{1.44 \times 10^{-5}}{(8.314 \times 773)^{-2}}$

C.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 773)^2}$

D.  $\frac{1.44 \times 10^{-5}}{(0.082 \times 773)^{-2}}$

**Answer: D**



**Watch Video Solution**


7. When two reactants A and B are mixed to give products C and D, the reaction quotient Q at the initial stages of the reaction

- A. Is zero
- B. Decreases with time
- C. Is independent of time
- D. Increases with time

**Answer: D**

 [Watch Video Solution](#)

8. At constant temperature , the equilibrium constant ( $K_p$ ) for the decomposition reaction



where  $p$ = pressure , $x$  = extent of decomposition .which one of the following statements is true?

- A.  $K_p$  increases with increase in  $p$ .
- B.  $K_p$  increases with increase in  $x$ .


C.  $K_p$  increases with decrease in x.

D.  $K_p$  remains constant with change in p and x

**Answer: D**

 [Watch Video Solution](#)

9. Consider the following equilibrium in a closed container,



At a fixed temperature, the volume of the reaction container is halved.

For this change which of the following statements holds true regarding the equilibrium constant ( $K_p$ ) and degree of dissociation ( $\alpha$ )?

- A. Neither  $K_p$  nor  $\alpha$  changes
- B. Both  $K_p$  and  $\alpha$  change
- C.  $K_p$  changes but  $\alpha$  does not change
- D.  $K_p$  does not change but  $\alpha$  changes



**Answer: D**



**Watch Video Solution**


## Archives (Fill In The Blanks)

1. For a given reversible reaction at a fixed temperature, equilibrium constants  $K_p$  and  $K_c$  are related by .....



**Watch Video Solution**

2. A 10-fold increase in pressure on the reaction



at equilibrium results in ..... in  $K_p$ .



**Watch Video Solution**

3. For a gaseous reaction  $2B \rightarrow A$ , the equilibrium constant  $K_p$  is ..... to/ than  $K_c$ .


 [Watch Video Solution](#)

## Archives (True/False)


1. When a liquid and its vapour are at equilibrium and the pressure is suddenly decreased, cooling occurs.

 [Watch Video Solution](#)

2. If equilibrium constant for the reactions.



is  $K$ , then the backward reactions.



its value is  $1/K$ . Is it true or false? If false then write the correct constant.

 [Watch Video Solution](#)

3. Catalyst makes a reaction more exothermic.


 [Watch Video Solution](#)

4. The rate of an exothermic reactions increases with increase in temperature.

 [Watch Video Solution](#)


## Archives (Subjective)

1. One mole of nitrogen is mixed with three moles of hydrogen in a 4 litre container. If 0.25 per cent of nitrogen is converted into ammonia by the following reaction




calculate the equilibrium constant of the reaction in concentration units.

What will be the value of K for the following reaction?




 [Watch Video Solution](#)

2. 1 mole of  $N_2$  and 3 moles of  $PCl_5$  are placed in a 100 litre vessels heated at  $227^\circ C$  the equilibrium pressure is 2.05 atm Assuming ideal behaviour, Calculate degree of dissociation of  $PCl_5$  and  $K_p$  for the reaction



 [Watch Video Solution](#)

3. The equilibrium constant of the reaction




at  $100^\circ C$  is 50. If a 1L flask containing 1 mol of  $A_2$  is connected to a 2L flask containing 2 mol of  $B_2$ , how many moles of AB will be formed at 373K?



 [Watch Video Solution](#)

4. At a certain temperature, equilibrium constant ( $K_c$ ) is 16 for the reaction:




If we take 1 mol of each of the four gases in a 1L container, what would be the equilibrium concentrations of NO and  $NO_2$ ?

 [Watch Video Solution](#)

5.  $N_2O_4$  is 25 % dissociated at  $37^\circ C$  and  $1atm$ . Calculate  $K_p$

 [Watch Video Solution](#)

6. For the reaction



Hydrogen gas is introduced into a five-litre flask at  $327^\circ C$ , containing 0.2 mol of  $CO(g)$  and a catalyst, until the pressure is  $4.92atm$ . At this point,

0.1 mol of  $CH_3OH(g)$  is formed. Calculate the equilibrium constants  $K_p$  and  $K_c$ .

 [Watch Video Solution](#)


7. 0.15 mol of  $CO$  taken in a 2.5 L flask is maintained at 750 K along with a catalyst so that the following reaction can take place  $CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$ . Hydrogen is introduced until the total pressure of the system is 8.5 atm at equilibrium and 0.08 mol of methanol is formed. Calculate

a.  $K_p$  and  $K_c$

b. The final pressure if the same amount of  $CO$  and  $H_2$  as before is used but no catalyst so that the reaction does not take place.

 [Watch Video Solution](#)

8. The progress of the reaction  $A \rightleftharpoons nB$  with time is presented in the figure given below:



Determine

- The value of  $n$ .
- The equilibrium constant  $K$ .
- The initial rate of concentration of A.



[Watch Video Solution](#)

9. The degree of dissociation is 0.4 at 400K and 1.0 atm for the gaseous reaction



assuming ideal behaviour of all gases, calculate the density of equilibrium


mixture at 400K and 1.0 atm (relative atomic mass of P is 31.0 and of Cl is 35.5).

 [Watch Video Solution](#)

**10.** When 3.06g of solid  $NH_4HS$  is introduced into a two-litre evacuated flask at  $27^\circ C$ , 30 % of the solid decomposes into gaseous ammonia and hydrogen sulphide. (i) Calculate  $K_c$  and  $K_p$  for the reaction at  $27^\circ C$ . (ii) What would happen to the equilibrium when more solid  $NH_4HS$  is introduced into the flask?

 [Watch Video Solution](#)

**11.** In the reaction equilibrium



When 5 mol of each is taken and the temperature is kept at 298K, the total pressure was found to be 20 bar.

Given :  $\Delta_f G_{N_2O_4}^\ominus = 100kJ$ ,  $\Delta_f G_{NO_2}^\ominus = 50KJ$



a. Find  $\Delta G^\ominus$  of the reaction at 298K.

b. Find the direction of the reaction.

 [Watch Video Solution](#)

## Subjective type

1. The equilibrium constant  $K_p$  of the reaction:  $2SO_2 + O_2 \rightleftharpoons 2SO_3$  is  $900 \text{ atm}^{-1}$  at 800K. A mixture containing  $SO_3$  and  $O_2$  having initial pressure of 1 atm and 2 atm respectively, is heated at constant volume to equilibrate. Calculate the partial pressure of each gas at 800K at equilibrium.

 [Watch Video Solution](#)