



CHEMISTRY

BOOKS - CENGAGE CHEMISTRY (ENGLISH)

ISOMERISM

Illustration

1. Give various functional isomers of butanol and give their isomeric relations.



[Watch Video Solution](#)

2. Give various functional isomers of pentanal and give their isomeric relations.



[Watch Video Solution](#)

3. Give various functional isomers of butanoic acid and give their isomeric relations.



[Watch Video Solution](#)

4. (I) Give various functional isomers of butan-1-amine and give their isomeric relations.

(II) Give various isomers of butan -1-amine and their isomeric relations.



Watch Video Solution

5. Give the various isomers of pent-1-yne and their isomeric relations.



Watch Video Solution

6. Write various functional isomers of 2-phenylethanol and give their isomeric relations.



[Watch Video Solution](#)

7. I. Give various functional isomers of butanenitrile and give their isomeric relations.



[Watch Video Solution](#)

8. Write more stable tautomeric structures of the following:

(a) Phenol

(b) Methyl-3-oxobutane

(c) Cyclohexan-1,3,5-trione



Watch Video Solution

9. Keto-enol tautomerism is observed in :

(i) a. $PhCHO$, b. $PhCOCH_3$

c. $PhCOPh$, d. $PhCOCH_2COCH_3$





[Watch Video Solution](#)

10. . Which of the following is more easily enolised?

- a. Cyclobutenone
- b. Triketo cyclobutane



[Watch Video Solution](#)

11. Write all the acyclic and cyclic isomers of a compound having molecular formula C_3H_6O .



[Watch Video Solution](#)

12. Write all the ring-chain isomers of (a) pent-1-ene and (b) Pent-1-yne.



Watch Video Solution




13. Arrange the following ligands in the decreasing order of priority.

a. i. $-OH$, ii. $-COOH$

iii $-CH_2OH$, iv. $CH(OH)CH_3$





b. i. $-NH_2$ ii. $-C \equiv N$

iii. $-NO_2$ iv. $-CH_2NH_2$







Watch Video Solution

14. Which of the following alkenes have diastereomers?



Watch Video Solution

15. Assign $E - Z$ configuration to each of the following:








a.



Watch Video Solution

16. Give the decreasing order of net dipole moments of the following:






 [Watch Video Solution](#)

17. Give the structural and diastereomers of pent-1-ene.

 [Watch Video Solution](#)

18. Specify the configuration of following compounds in *D* of *L*.



 [Watch Video Solution](#)

19. The optical rotation of an optically active compound is $+20^\circ$. The length of tube is 10cm and the density of solution is 0.4gml^{-1} . The specific rotation of the

compound is:

i. $+50^\circ$ ii. $+500^\circ$ iii. $+5^\circ$ iv. $+0.5^\circ$



[Watch Video Solution](#)

20. Explain the following: What is stereocentre and how it is related to a chiral centre?



[Watch Video Solution](#)

21. Write the name and structure of the following optically active compounds with

lowest molecular weight.

i. alkane ii. Alkene

iii. Alkyne iv. unsaturated hydrocarbon

v. alkyl halide vi. Alcohol

vii. Acid viii. Amine.



[Watch Video Solution](#)

22. How many stereoisomers are possible for the compound $CH_3CH = CHCHClCH_3$?



[Watch Video Solution](#)

23. How many structural isomers and stereoisomers of bromopiperidine are possible?



Watch Video Solution

24. The specific rotation of a pure enantiomer is $+10^\circ$. What will be its observed rotation if its is isolated from a reaction with

i. 30 % racemisation and 70 % retention.

ii. 70 % racemisation and 30 % inversion.





Watch Video Solution

25. An aqueous solution containing 10gm of optically pure fructose was diluted to 500ml with water and placed in a polarimeter tube 20cm long. The measured rotation was -5.0° . To this solution, 500ml of a solution containing 10gm of racemic fructose is added. What will be the change in specific rotation?







Watch Video Solution

26. Which of the following will show dipole moment (μ)?





[Watch Video Solution](#)

27. Draw Fischer projection formulae for the following molecules:




Watch Video Solution

28. a. Convert Wedge formula to Fischer formula.





b. Convert Fischer formula to Wedge formula.



 [Watch Video Solution](#)

29. Convert Sawhorse formula to Fischer formula.





 [Watch Video Solution](#)

30. Assign the *R* and *S* configurations to the enantiomers of 2-chlorobutane.

 [Watch Video Solution](#)

31. Assign the *R* and *S* designation to the following compounds:





Watch Video Solution

32. Give the effect of the configuration of (*S*) – 2 – bromo butane on performing the following operations:


- a. Exchanging ligands across the horizontal bond.
- b. Exchanging ligands across the vertical bond.
- c. Making both switches (a) and (b)
- d. Exchanging a horizontal and vertical ligand.
- e. A 180° vertical or horizontal rotation

outside the plane of paper.

f. A 90° rotation on the plane of paper.

 [Watch Video Solution](#)

33. Assign *R* and *S* configuration of the following compound.



 [Watch Video Solution](#)

Example



1. State whether the following statements are true or false.

a. *D* and *L* stereoisomers are enantiomers.



[Watch Video Solution](#)




2. Write the stable tautomer form the following:



 [Watch Video Solution](#)

3. Some possible structures of a compound A

($C_{10}H_{16}$) are:





Write the possible stereoisomers of (a), (b),
and (c)



Watch Video Solution

4. Some possible structures of a compound A

($C_{10}H_{16}$) are:




Write the possible stereoisomers of (a) and (b).



Watch Video Solution

5. Consider the compound (A) given below:



Give the total number of stereoisomers possible for (A).



Watch Video Solution

6. How many geometrical isomers are possible for the following?

a. Decalin b. 1 – Methyldecalin

c. 9 – Methyldecalin



[Watch Video Solution](#)

7. How many geometrical isomers are possible for the following ?

a. 1, 2 – Dimethylcyclobutane

b. 1, 3, 5 – Trimethylcyclohexane

c. 1, 1, 2 – Trimethylcyclopropane



[Watch Video Solution](#)

8. How many geometrical isomers are possible for 1, 2, 4-Trimethyl cyclohexane.



[Watch Video Solution](#)

9. How many isomeric dienes with a six-membered ring are possible of the compound

with the molecular formula C_7H_{10} ?



[Watch Video Solution](#)

10. Give the structures of six C atom cyclic diene with lowest molecular mass that shows only geometrical isomers (diastereomers) and not optical isomers.



[Watch Video Solution](#)

11. Give the structural formula of an unsaturated hydrocarbon with the lowest number of C atoms (or with lowest molecular mass) which shows:

a. Optical isomers

b. Geometrical isomers

c. Both optical and geometrical isomers



Watch Video Solution

12. Give the structural formula of a cyclic alkyne with the lowest number of C atoms and showing:

a. Both geometrical and optical isomerisms.

b. Geometrical isomerism with meso stereoisomers.



Watch Video Solution

13. Give the type of isomerism exhibited by each of the following pairs:

- a. Acetic acid and methyl formate
- b. *o* – Nitrophenol and *m* – nitrophenol
- c. *n* – Butane and isobutane
- d. Maleic and fumaric acid
- e. Benzyl alcohol and *o* – methyl phenol
- f. Propionic acid and methyl acetate
- g. *n* – Butyl alcohol and diethyl ether
- h. 1, 1 – Dibromoethane and 1, 2 –
dibromoethane



[Watch Video Solution](#)

14. 2, 2' - difluoro-6, 6' - dimethylbiphenyl is non-resolvable, whereas 2, 2' - dinitro-6, 6' - disulfonic acid biphenyl is resolvable. Account for this fact.



[Watch Video Solution](#)

15. What is the observed rotation when $0.1M$ solutions of (*R*) - 2 butanol is mixed with an equal volume of $0.05M$ solution of racemic 2 - butanol and the resulting solution is taken

in a cell of 5cm long tube in a polarimeter?

The specific rotation of ($R - 2$) butonal is

$+13.9^\circ$



Watch Video Solution

Exercise

1. 0.90gm of an organic compound $C_4H_{10}O_2(A)$ when treated with sodium gives 224ml of hydrogen at NTP . Compound (A) can be separated into fraction (B) and

(C), by crystallisation of which the fraction (B) is resolved into isomers (D) and (E). Write down the structural formula of (A) to (E) with proper reasoning.



[Watch Video Solution](#)

2. Write down the structures of close homologues of heptane having one quaternary C atom in their molecule and the other having two tertiary carbon atoms.



[Watch Video Solution](#)

3. An alkane (A) C_5H_{12} on chlorination at $300^\circ C$ gives a mixture of four different monochlorinated derivatives (B), (C), (D), and (E). Two of these derivatives give the same stable alkene (F) on dehydrohalogenation. On oxidation with hot alkaline $KMnO_4$ followed by acidification, (F) gives two products (G) and (H). Give the structures of (A) to (H) with proper reasoning.



Watch Video Solution

4. Which alkane, having a molecular weight of 86, will form only two monobromo alkanes?



Watch Video Solution

5. A hydrocarbon (A) was found to have vapour density 36. It forms only a single monochloro substitution product. Suggest (A).

A. a) Iso-pentane

B. b) Neo-pentane

C. c) Cyclo-hexane

D. d) Methyl cyclo-hexane

Answer:



Watch Video Solution

6. Write the appropriate structural formulae for:

a. A cyclic molecule that is a constitutional isomer of cyclohexane

b. Molecules with the formula C_6H_{12} that contain one ring and are enantiomers of each other.

c. Molecules with formula C_6H_{12} that contain one ring and are diastereomers of each other.

d. Molecules with formula C_6H_{12} that contain no ring and are enantiomers of each other.

e. Molecules with formula C_6H_{12} that contain no ring and are diastereomers of each other.



[Watch Video Solution](#)

7. There are four dimethylcyclopropane isomers.

a. Write the three-dimensional formulas for these isomers.

b. Which of these isomers is chiral?

c. If a mixture consisting of 1 mol of each of these isomers is subjected to simple gas chromatography, how many fractions would be obtained and how many compounds each fraction will have?

d. How many of these fractions would be optically active?



Watch Video Solution

8. Which of the following objects listed below possess a plane of symmetry and are, therefore, achiral?

- a. A hammer
- b. A car
- c. A woodscrew
- d. An ear
- e. A screwdriver
- f. A tennis shoe


g. A golf club

h. A baseball bat





Watch Video Solution

9. Assign (*R*) or (*S*) designations to each of the following compounds




a.






Watch Video Solution

10. Consider the following pair of structures and tell whether they represent enantiomers or two molecules of the same compound in different orientation.



[Watch Video Solution](#)

11. Tell whether the two structures in each pair represent enantiomers or two molecules of the same compound in different orientations.



 Watch Video Solution

12. A sample of optically active alcohol has a specific rotation, $[\alpha]_D^{25}$, equal to $+1.151^\circ$. Specific rotation for pure form $[\alpha]_D^{25}$ is $+5.756^\circ$.


What is the percentage enantiomeric excess of the sample?



Watch Video Solution

13. The following are the formulae for three compounds 2, 3 – dichlorobutane written in

non-eclipsed conformations. In each instance, tell which compounds are enantiomers and which are meso.



[Watch Video Solution](#)




14. Write the three-dimensional formulas for all of the stereoisomers of each of the following compounds. Label pairs of enantiomers and meso compounds.










Watch Video Solution






15. Consider the following pairs of structure. Identify the relationship between them by describing them as presenting enantiomers, diastereomers, constitutional isomers, or two



molecule of the same compound.














 [View Text Solution](#)

16. Discuss the anticipated stereochemistry of each of the following compounds.





Watch Video Solution

17. Write formulae for all the isomers of each of the following. Designate pairs of enantiomers and achiral compounds where they exist.

a. 1-Bromo-4- chlorocyclohexane

b. 1-Bromo-3-chlorocyclohexane

c. 1-Bromo-2- chlorocyclohexane



18. Which of the following are chiral and capable of existing as enantiomers?

a. 1 – 3 – Dibromobutane

b. 1, 2 – Dichloropropane

c. 1, 5 – Dibromopentane

c. 3 – Ethylpentane

e. 2 – Chlorobicyclo [1.1.0] butane

f. 2 – Iodobicyclo [2.2.2] octane

g. 2 – Bromobicyclo [2.1.1] hexane

h. 5 – Bromobicyclo [2.1.1] hexane



Watch Video Solution

19. a. Write the structure of 2,2 – dibromobicyclo [2.2.1] heptane.
- b. How many stereocenters does it contain?
- c. How many stereoisomers are predicted by the 2^n rule?
- d. Only one pair of enantiomers is possible for 2 – 2, dibromobicyclo [2.2.1] heptane. Explain.







Watch Video Solution

Linked Comprehension Type Exercise

1. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2-methylbutanal. Based on this information, answer the following question.

1. Compound (A) is



Answer: A



Watch Video Solution

2. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2 – methyl butanal. Based on this information, answer the following question.

The total number of stereoisomers possible for (A) is :

A. 8

B. 7

C. 6

D. 4

Answer: C



Watch Video Solution

3. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2 – methyl butanal. Based on this information, answer the following question.

If the stereochemistry about the double bond in (A) is cis, the total number of stereoisomers for (A) is :

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

4. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2 – methyl butanal. Based on this information, answer the following question.

If the stereochemistry about the double bond in (A) is cis, the total number of enantiomers for (A) is :

A. None

B. 2

C. 4

D. 6

Answer: B



Watch Video Solution

5. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2 – methyl butanal. Based on this information, answer the following question.

If the stereochemistry about the double bond in (A) is trans, the total number of stereoisomers for (A) is:

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

6. An organic compound (A) ($C_{10}H_{20}$) on reductive ozonolysis gives 2 – methyl butanal. Based on this information, answer the

following question.

If the stereochemistry about the double bond in (A) is trans, the total number of enantiomers for (A) is :

A. None

B. 2


C. 4

D. 6

Answer: B



Watch Video Solution



7.

If the both compounds (A) and (B) are optically inactive, the possible structures of (B) are:

A. *cis* – I

B. *trans* – I


C. *cis* – II

D. *trans* – II

Answer: C::D



Watch Video Solution




↓ Three equivalent
of H₂+Pd at 120°C

(C)


8.

If the both compounds (A) and (B) are optically inactive, the possible structures of (B) are:





B.




C.

D. All




Answer: A



Watch Video Solution



If the both compounds (A) and (B) are optically inactive, the possible structures of (B) are:



D. All

Answer: B::C



Watch Video Solution

10. How many isomeric dienes with a six-membered ring are possible of the compound with the molecular formula C_7H_{10} ?

A. 5

B. 6

C. 7

D. 8




Answer: D



Watch Video Solution

11. In this paragraph, some statements are given based on isomerism. Read the following statements given for every question and provide the answer.

Which of the following cyclic dienes does not show geometrical isomerism?





Answer: D



Watch Video Solution

12. In this paragraph, some statements are given based on isomerism. Read the following statements given for every question and provide the answer.

Which of the following structures of a cyclic alkene with the lowest number of C atoms shows both geometrical and optical isomerism?



C. Both

D. None

Answer: A



Watch Video Solution

13. In this paragraph, some statements are given based on isomerism. Read the following statements given for every question and provide the answer.

Which of the following cyclic dienes does not show geometrical isomerism?

A. (I)

B. (II)

C. Both

D. None





Answer: B



Watch Video Solution

Multiple Choice Questions Exercise

1. Which of the following show stable or major form of tautomerism?



Answer: B::D



Watch Video Solution

2. Keto enol tautomerism is not observed in:

A. Phenol

B. Glycerol

C. HCN

D. Benzophenone

Answer: D



Watch Video Solution

3. Arrange the following in decreasing order of enol content:

i. Diethyl malonate

ii. Acetoacetic ester (AAE or EAA)

iii. Acetyl acetone

iv. $PhCOH_2COH_3$

A. $(iv) > (iii) > (ii) > (i)$

B. $(iv) > (iii) > (i) > (ii)$

C. $(iii) > (iv) > (ii) > (i)$


D. $(iii) > (iv) > (i) > (ii)$

Answer: A



Watch Video Solution

4. Which of the following will not show geometrical isomerism?



B. 2, 4 – Dinitro phenyl hydrazone of acetone

C. Oxime of formaldehyde


D. Cyclohexan 1, 2 – diol

Answer: B::C



Watch Video Solution

5. Which of the following will show geometrical isomerism ?



D.





Answer: B::C::D




Watch Video Solution

6. Which of the following are not resolvable?

A. 2, 3 – Pentadiene









Answer: C::D



Watch Video Solution

7. Which of the following are resolvable ?









Answer: B::D

 **Watch Video Solution**

8. Which of the following is/are optically active?






Answer: B



Watch Video Solution

9. The compounds



are optically inactive because

A. Both compounds have the plane of symmetry.

B. Both compounds have the centre of symmetry.

C. Compound (I) has the plane of symmetry, while compound (II) has the centre of

symmetry.

D. Compound (I) has the centre of symmetry, while compound (II) has the plane of symmetry.

Answer: C



Watch Video Solution

10. The stable conformer (*s*) of trans-1,4-dimethyl] cyclohexane is/are:

A. 1 – axial – 4 – equatorial form

B. 1 – axial – 4 – axial form

C. 1 – equatorial – 4 – axial form

D. 1 – equatorial – 4 – equatorial form

Answer: D



Watch Video Solution

11. The stable conformer (*s*) of cis-cyclohexane

1 – 3 – diol is/are:

A. 1 – axial – 3 – axial form

B. 1 – axial – 3 – equatorial form

C. 1-equatorial- 3- axial form

D. 1- equatorial -3- equatorial form

Answer: A



Watch Video Solution

12. Which form (*s*) of cyclohexane is/are free from angle strain?

A. Chair form

B. Boat form

C. Twist boat form

D. All

Answer: D



Watch Video Solution

13. Which of the following statements is/are wrong, about the more stability of chair form than boat form?

A. In chair conformation, all the ($C - H$) bonds in adjacent carbons are in the skew position.

B. In boat conformation, there are four skew interactions and two eclipsed interaction.

C. In a boat conformation, there are three skew interactions and three eclipsed interactions.

D. Boat form has an angle strain

Answer: C::D



Watch Video Solution

14. Which of the following methods are used for resolution ?

A. Biological methods by using special enzymes

B. By making their diastereomers

C. Chromatographic method using special
absorbents

D. Azetotropic distillation

Answer: A::B::C



Watch Video Solution

15. According to Baeyer's strain theory, which of the following is/are most stable cyclic compounds?

A. Cyclopropane

B. Cyclobutane

C. Cyclopentane

D. Cyclohexane

Answer: C



Watch Video Solution

16. The angle strain in cyclohexane is :

A. $9^{\circ}, 44'$

B. $0^\circ, 44'$

C. $-5^\circ, 36'$

D. $5^\circ, 36'$

Answer: C



Watch Video Solution

17. Which of the following statements are correct?

A. Methyl cyclopropane and methyl cyclobutane do not show stereoisomerism.

B. Both show stereoisomerism.

C. 1,2-Dimethyl cyclopropane shows stereoisomerism, but methyl cyclobutane does not.

D. 1,2-Dimethyl cyclopropane does not show stereoisomerism, but methyl cyclobutane shows.

Answer: A::C



Watch Video Solution

18. Which of the following statement regarding 1, 2- dimethylcyclopropane (I) and 1, 2- dimethyl cyclobutane (II) are wrong

A. Both of them show three stereoisomers.

B. The cis form of both is optically inactive (meso) and the trans form the both has a pair of enantiomers.

C. The cis form of both has a pair of enantiomers and the trans form of both is optically inactive (meso).

D. The meso form of both is optically inactive due to the presence of the centre of symmetry.

Answer: C::D



Watch Video Solution

19. Which of the following statements regarding 1, 3- dimethyl cyclobutane is/are correct?

A. Both cis and trans forms are optically active.

B. Both cis and trans forms are optically inactive.

C. The cis form is optically active, while the trans form is optically inactive.

D. The trans form is optically active, while the cis form is optically inactive.

Answer: B



Watch Video Solution

20. Which of the following statements regarding 1,3-dimethylcyclobutane is/are correct?

- A. The cis has plane of symmetry, while the trans form has both plane and centre of symmetry.
- B. The cis form has both plane and centre of symmetry, while the trans form has only plane of symmetry.
- C. Both have only plane of symmetry.
- D. Neither of them has any element of symmetry.

Answer: A



Watch Video Solution

21. Which of the following statements regarding 1, 2-dimethyl cyclo-pentane and 1, 3-dimethyl cyclopentane is/are correct?

A. In both, cis form is meso, while trans form is resolvable

B. In both, trans form is meso, while cis form is resolvable

C. In both cis and trans forms are meso.

D. In both, cis and trans forms are resolvable.

Answer: A



Watch Video Solution

22. The configuration of sugars is related to glyceraldehyde and that of amino acids is related to:

A. Serine

B. Leucine

C. Alanine

D. Glycine

Answer: A



Watch Video Solution

23. Which of the following statement are true?

A. 2-Butanone on reaction with

2, 4 – *DNP* forms two compounds

which are geometrical isomers and can be separated.

B. Acetophenone on reaction with HCN forms two compounds which are geometrical isomers.

C. Acetone on reaction with NH_3 forms two compounds which are resolvable.

D. Acetaldehyde on reaction with NH_2OH forms two compounds which have different melting points.

Answer: A::D



Watch Video Solution

24. Which of the following statements are correct?

A. Butan-2 — one shows tautomerism.

B. Compounds containing asymmetric C atoms are always optically active.

C. Members belonging to the same class of compounds are called isomers.

D. Isomers have the same molecular formula.

Answer: A::D



Watch Video Solution

25. Which of the following statements are wrong?

- A. *m*-Chlorobromo benzene is an isomer of *m*-bromochloro benzene.
- B. All alkenes show geometrical isomerism.
- C. Dimethyl ether and ethanol are functional isomers.
- D. Geometrical isomers have different physical properties.

Answer: A::B



Watch Video Solution

26. Which of the following statements are correct?

A. 2, 3, 4- Tribromo pentane has three chiral C atoms..

B. Tartaric acid has two asymmetric C atoms.

C. d and l forms of an optically active compound have different specific rotations with opposite signs.

D. Staggered and eclipsed forms of ethane
have different stabilities.

Answer: B::D



Watch Video Solution

27. Which of the following statements are wrong?

A. Isobutane and *n*-butane are chain isomers.

B. Ethyl cyanide and ethyl isocyanide are functional isomers.

C. $H - C \equiv N$ and $H - \overset{\oplus}{N} \equiv \overset{\ominus}{C}$ are tautomers.

D. Maleic and fumaric acids are enantiomers.

Answer: D



Watch Video Solution

28. Which of the following statements are correct?

A. A reaction in which different stereoisomers produced different products or act a different rates is called stereospecific reaction.

B. A reaction in which a given substrate produces diastereoisomeric products in different amounts and where one

diastereomer predominates very much over the other is called stereoselective reaction.

C. If the replacement of one group at an achiral centre by a new substituent generates a chiral centre by a new substituent generates a chiral centre, the original molecule is said to be enantiotopic.

D. The *E* and *Z* system of naming geometrical isomers is based on the *CIP* sequence rule.

Answer: A::B::C::D



Watch Video Solution

29. Which of the following statements are correct?

A. (a) Diastereomers have different physical properties but not identical chemical properties.

B. (b) Enantiomers have the same physical and chemical properties but different physiological properties.

C. (c) Polarimeter is used in measuring the optical rotation of compound.

D. Only organic molecules show optical isomerism.

Answer: A::B::C



Watch Video Solution

30. Which of the following statements are wrong?

A. Ethyl benzene is the chain isomer of xylene.

B. Alkenes can exhibit the position, chain, functional, and geometrical isomerism.

C. Esters and carboxylic acids are examples of metamerism.

D. Metamers belong to different classes of compounds.

Answer: C::D



Watch Video Solution

31. Only two isomeric monochloro derivatives are possible for

A. *n* Butane

B. 2, 4-Dimethyl pentane

C. Benzene


D. 2-Methyl propane



Answer: A::D



Watch Video Solution

32. Keto-enol tautomerism is observed in









Answer: B::D



Watch Video Solution

33. Which of the following has/have asymmetric carbon atom?



Answer: C::D



Watch Video Solution

34. The molecule (s) that will have dipole moment is/are:

A. 2, 2 Dimethyl propane

B. trans-2-Pentene

C. cis -3- Hexene

D. 2, 2, 3, 3-Tetramethyl butane

Answer: B::C



Watch Video Solution

35. Which one of the following compounds will show geometrical isomerism ?

A. 2-Butene

B. Propene

C. 1-phenyl propane





D. 2-Methyl-2-butene

Answer: A::C



Watch Video Solution

36. Tautomerism is exhibited by :




Answer: A::C::D



Watch Video Solution

37. The correct statements about the compound given below is :



A. The compound is optically active

B. The compound possesses plane of symmetry.

C. The compound possesses plane of symmetry.


D. The compound possesses axis of symmetry.

Answer: A::D



Watch Video Solution

38. In the Newman projection for 2,2-dimethylbutane



X and *Y* can, respectively, be

- A. *H* and *H*
- B. *H* and *C*₂*H*₅
- C. *C*₂*H*₅ and *H*
- D. *CH*₃ and *CH*₃

Answer: B::D



Watch Video Solution

Single Correct Answer Type

1. Which of the following will not be able to show optical isomerism (enantiomerism)?

A. 1, 2-Propadiene

B. 2, 3-Pentadiene

C. sec-Butyl alcohol

D. All exhibit enantiomerism

Answer: A



Watch Video Solution

2. The type of isomerism exhibited by the compound with formula $C_4H_{10}O$ is :

- A. Chain and position
- B. Functional and position
- C. Metamerism

D. Chain, position, functional, and metamerism

Answer: D



Watch Video Solution

3. Acids and ester having the same number of carbon atoms are:

A. Functional isomers

B. Tautomers

C. Metamers

D. Not isomers

Answer: A



Watch Video Solution

4. Which of the following kinds of isomerism can nitroethane exhibit?

A. Metamerism

B. Optical activity

C. Tautomerism


D. Position isomerism



Answer: C



Watch Video Solution

5. Which among the following is likely to show geometrical isomerism?





Answer: B



Watch Video Solution

6. How many gem dihalides with different formulas are possible for $C_3H_6Cl_2$?

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

7. Which of the following statement is correct?

A. The presence of chiral carbon is essential condition for enantiomerism.

B. Functional isomerism is a kind of stereoisomerism.

C. The compounds containing one chiral carbon only are always chiral.

D. All statements are wrong.

Answer: C



Watch Video Solution

8. The total number of halogenated products likely to be formed by ethane is:

A. 4

B. 6

C. 9


D. 8

Answer: C



Watch Video Solution

9. If the optical rotation produced by the compound (A) is $+52^\circ$, the one produced by compound (B) is :



A. -52°

B. $+52^\circ$

C. 0°





D. Unpredictable

Answer: A



Watch Video Solution

10. The *E*-isomer among the following is:







Answer: B



Watch Video Solution

11. Which of the following will not show geometrical isomerism?



Answer: B



Watch Video Solution

12. The dihedral angle between the hydrogen atoms of two methyl groups in staggered conformation of ethane is

A. 120°

B. 180°

C. 90°

D. 60°

Answer: D



[Watch Video Solution](#)

13. Which of the following object is chiral?

A. Nail

B. Blade

C. Tennis racket

D. Laced football

Answer: D



[Watch Video Solution](#)

Single Correct Answer Type Exercise

1. Predict the number of stereoisomers in $CH_2OH \cdot (CHOH)_4 \cdot CHO$.

A. 16

B. 8

C. 4

D. 2

Answer: A

 [Watch Video Solution](#)

2. Predict the number of stereoisomers for 2, 5 heptadiene.

A. 4

B. 3

C. 2

D. 5

Answer: B

 [Watch Video Solution](#)

3. Symbol *D* stands for:

A. (a) Dextrorotatory, which rotates

P. P. L. towards right.

B. (b) Dextrorotatory, which rotates

P. P. L. towards left.

C. (c) Relative configuration with respect to

lactic acid taken as standard.

D. (d) Relative configuration with respect to glyceraldehyde taken as standard and (OH) group is on the right side.

Answer: D



Watch Video Solution

4. Which of the following compounds will not show geometrical isomerism?

A. 3- Phenyl-2 propenoic acid

B. 2 Butene

C. 3- Methyl-2 butenoic acid

D. 3- Methyl-2 pentenoic acid.

Answer: C



Watch Video Solution

5. The enolic form of acetone contains:

A. 9σ -bond, 1π -bond, and 2 lone pairs.

B. 8σ -bonds, 2π -bonds, and 2 lone pairs.

C. 10σ - bonds 2π -bonds and 1 lone pair.

D. 9σ - bonds, 2π -bonds, and 2 lone pairs.

Answer: A



Watch Video Solution

6. Arrange the following in the increasing order of expected enol content.



iii. CH_3CHO

iv. $CH_3COCH_2COCH_3$

A. $iii < ii < iv$

B. $iii < ii < i < iv$

C. $i < iv < ii < iii$

D. $iv < i < ii < iii$

Answer: B



Watch Video Solution

7. The decreasing order of reactivity of the following alkenes is:

i. 2, 3 -Dimethyl-2 butene

ii. 2- Methyl-2 butene

2- Butene

iv. Ethene

A. $(iv) > (iii) > (ii) > (i)$

B. $(i) > (ii) > (iii) > (iv)$

C. $(iv) > (ii) > (iii) > (i)$

D. $(i) > (iii) > (ii) > (iv)$

Answer: A



Watch Video Solution

8. Which of the following is the least stable form of cyclohexane?

A. Boat-form

B. Chair form

C. Skew- boat form

D. Crown form

Answer: A



Watch Video Solution

9. The minimum number of carbon atoms an alkane should contain in order to exhibit optical activity is:

A. 5

B. 6

C. 7

D. 8

Answer: C



Watch Video Solution

10. A compound whose molecule is superimposable on its mirror image despite containing chiral carbon atoms is called:

- A. Threo isomer
- B. Meso compound
- C. Enantiomer
- D. No special name

Answer: B



Watch Video Solution

11. The *IUPAC* name of the compound with formula C_nH_{2n+2} , having the lowest possible molecular mass and capable of showing enantiomerism, is:

- A. 3- Methyl hexane
- B. 2, 3-Dimethyl pentane
- C. Methane

D. Both (a) and (b)

Answer: D



Watch Video Solution

12. A racemic mixture is optically inactive due to :



- A. The presence of plane of symmetry
- B. External compensation
- C. Internal compensation

D. None of these

Answer: B

 [Watch Video Solution](#)

13. The pair of structures given below represents:



A. (a) Enantiomers

B. (b) Position isomers

C. (c) Conformers

D. (d) None of these

Answer: B



Watch Video Solution

14. The degree of unsaturation or index of hydrogen deficiency in the following is :

i. C_6H_{14}



A. 0,2,4

B. 1,0,2

C. 4,1,0

D. 2,0,4

Answer: A



Watch Video Solution

15. The degree of unsaturation in

i. $C_3H_3Cl_3$, ii. C_3H_4O , iii. C_4H_5N is :

A. 2,1,3

B. 1,2,3

C. 3,2,1



D. 2,3,1

Answer: B



Watch Video Solution

16. Which of the following alkenes is most stable?



Answer: C



Watch Video Solution

17. Which of the following will have zero dipole moment?

A. (a) cis, - 1, 2-Dichloroethene

B. (b) trans - 1, 2-Dichloroethene

C. (c) Dichloromethane

D. (d) Dibromomethane

Answer: B



Watch Video Solution

18. The number of conformations exhibited by ethane is

A. Infinite

B. Two

C. Three

D. Four

Answer: A



Watch Video Solution

19. Which of the following is not an isomer of butanal?

A. (a) 2-Butanone

B. (b) 2-Methyl propanal

C. (c) 2- Butanol

D. (d) But-2-en-1-ol

Answer: C



Watch Video Solution

20. The least energetic conformation of cyclohexane is:

- A. Boat form
- B. Half chair form
- C. Chair form
- D. Twisted boat form

Answer: C



Watch Video Solution

21. How many chiral carbons are present in glucose molecule $CHO(CHOH)_4CH_2OH$?

A. 4

B. 3

C. 2

D. 1

Answer: A



Watch Video Solution

22. The process of separation of a racemic mixture into d- and l- components is called

A. Resolution

B. Dehydration

C. Revolution

D. Dehydrohalogenation

Answer: A



Watch Video Solution

23. If a compound has n asymmetric carbon atoms with different terminal groups, the number of stereoisomers is given by the formula:

A. $(1/2)^n$

B. 2^n

C. $2\sqrt{n}$

D. $\sqrt{2n}$

Answer: B



Watch Video Solution

24. How many chiral compounds are possible on monochlorination of 2-methyl butane?

A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

25. A hydrocarbon with formula C_8H_{18} gives one monochloro derivative. The hydrocarbon can be:

A. *n*-octane

B. 2 Methyl heptane

C. 2, 2, 4-Trimethyl pentane

D. 2, 2, 3, 3-Tetramethyl butane

Answer: D



Watch Video Solution

26. The most strained cycloalkane is :

A. Cyclopropane

B. Cyclobutane

C. Cyclopentane

D. Cyclohexane

Answer: A



Watch Video Solution

27. The total number of acyclic isomers, including the stereoisomers, with formula C_4H_7Cl is

A. 12

B. 11

C. 10

D. 9

Answer: A



Watch Video Solution

28. Mesotartaric acid is optically inactive due to the presence of:




- A. Geometrical isomerism
- B. Two chiral carbon atoms
- C. Molecular symmetry
- D. External compensation

Answer: C



Watch Video Solution

29. Butene when treated with chlorine at about $500^{\circ}C$ forms:







Answer: B



Watch Video Solution

30. Which of the following are diastereomers ?



A. (I) and (III)

B. (II) and (IV)

C. (I) and (II)

D. None

Answer: C



Watch Video Solution

31. An organic compound contains 66 % *C* and 13.3 % *H*. Its vapour density is 37. The possible number of isomers of resonance is not correct?

A. 6

B. 7

C. 5

D. 8

Answer: B



Watch Video Solution

32. Which of the following statements regarding the concept of resonance is not correct?

A. The different resonating structures of a molecule have fixed arrangement of

atomic nuclei

B. The different resonating structures differ in the arrangement of electrons.

C. None of the individual resonating structure explains the various characteristics of the molecule.

D. The hybrid structure have equal contribution from all the resonating structures.

Answer: D



Watch Video Solution

33. The keto form of phenol contains:





- A. 3π , 13σ , 4 non-bonding electrons.
- B. 3π , 9σ , 4 non-bonding electrons.
- C. 3π , 9σ , 4 non-bonding electrons.
- D. 3π , 8σ , 4 non-bonding electrons.

Answer: A



Watch Video Solution

34. Tautomerism is not exhibited by :



Answer: B



Watch Video Solution

35. The number of isomers of the compound $C_2FClBrI$ is :

A. 3

B. 4

C. 5

D. 6

Answer: D



Watch Video Solution

36. Pure enantiomeric acid + optically active alcohol having chiral C atom \rightarrow ?

The product will be:

A. An optically active mixture

B. A meso compound

C. A racemic mixture

D. A pure enantiomer

Answer: A



Watch Video Solution

37. Racemic acid + optically active alcohol
having chiral C atom \rightarrow ?


The product will be:

- A. Optically active mixture
- B. Meso compound
- C. Diastereomeric mixture
- D. Racemic mixture

Answer: C



Watch Video Solution



How many product are possible?

A. 1

B. 2


C. 3

D. 4

Answer: B



Watch Video Solution



39. formula $\text{C}_4\text{H}_8\text{Br}_2$.

Find compound with formula $\text{C}_4\text{H}_8\text{Br}_2$

How many structure of (X) are possible?

A. 2

B. 3


C. 4

D. 5

Answer: B



Watch Video Solution




Hydrogenation of the above compound in the presence of poisoned *Pd* catalyst gives:

- A. An optically active compound
- B. An optically inactive compound
- C. A racemic mixture

D. A diastereomeric mixture

Answer: B

 Watch Video Solution



Hydrogenation of the above compound in the presence of poisoned *Pd* catalyst gives:

- A. An optically active compound
- B. An optically inactive compound
- C. A racemic mixture
- D. A diastereomeric mixture

Answer: C



Watch Video Solution

42. An S_N2 reaction at an asymmetric carbon of a compound always gives

- A. A enantiomer of the substrate
- B. A product with opposite optical rotation
- C. A mixture of diastereomers
- D. A single stereoisomers

Answer: D



Watch Video Solution

43. The nodal plane in the π -bond of ethene is located in :

- A. The molecular plane
- B. A plane parallel to the molecular plane
- C. A plane perpendicular to the molecular plane which bisects the carbon-carbon σ -bond at right angle
- D. A plane perpendicular to the molecular plane which contains the carbon-carbon σ bond

Answer: A



Watch Video Solution

44. Which of the following compounds exhibits stereoisomerism?

A. 2-Methylbutane

B. 3- Methylbutane

C. 3-Methylbutanoic acid


D. 2-Methylbutanoic acid

Answer: D







Watch Video Solution

45. Consider the following reaction



Identify the structure of the (X)





Answer: B



Watch Video Solution

46. Which of the following hydrocarbons has the lowest dipole moment?







Answer: B



Watch Video Solution

47. The geometrical isomerism is shown by:




Answer: D



Watch Video Solution

48. The number of geometrical isomers in



A. 2

B. 4

C. 5

D. 6

Answer: D



Watch Video Solution


49. The smallest aldehyde and its next homologue are treated with NH_2OH to form oxime. Find out the correct answer out of the following.

- A. Two different oximes are formed
- B. Three different oximes are formed.
- C. Two oximes are optically active.
- D. All oximes are optically active.

Answer: B



Watch Video Solution



↓ Fractional distillation

M (Isomeric products)

50.

What are the numerical values of N and M ?

A. 6, 6

B. 5, 4

C. 4, 4

D. 3, 3

Answer: B



Watch Video Solution

51. Molecule in which the distance between two adjacent carbon atom is largest is

A. Ethane

B. Ethene

C. Ethyne

D. Benzene

Answer: A



Watch Video Solution

52. The compound which is not isomeric with diethyl ether is:

A. *n*- Propyl methyl ether

B. Butan-1 – *ol*

C. 2- Methyl propan-2 – *ol*

D. Butanone

Answer: D



Watch Video Solution

53. Among the following, the compound that can be most readily sulphonated is:

A. Benzene

B. Nitrobenzene

C. Toluene

D. Chlorobenzene

Answer: C



Watch Video Solution

54. Which of the following compounds will exhibit cis-trans (geometrical) isomerism ?

A. 2-Butene

B. 2-Butyne

C. 2-Butanol

D. Butanal

Answer: A



Watch Video Solution

55. An isomer of ethanol is:

- A. Methanol
- B. Diethyl ether
- C. Acetone
- D. Dimethyl ether

Answer: D



Watch Video Solution

56. Which of the following will have the least hindered rotation about carbon-carbon bonds?

A. Ethane

B. Ethylene

C. Acetylene

D. Hexachlorethane

Answer: A



Watch Video Solution

57. The number of isomers of C_6H_{14} is:

A. 4

B. 5

C. 6

D. 7

Answer: B



Watch Video Solution

58. The enolic form of acetone contains:

- A. 9σ -bonds, 1π -bond, and 2 lone pairs.
- B. 8σ -bonds, 2π -bonds, and 2 lone pairs.
- C. 10σ -bonds, 1π -bond, and 1 lone pair.
- D. 9σ -bonds, 2π -bonds, and 2 lone pairs.

Answer: A



Watch Video Solution

59. Three dimensional arrangements which can be interconverted into one another due to rotation along a single bond are known as

- A. Conformers
- B. Diastereomers
- C. Enantiomers
- D. Positional isomerisms

Answer: A



Watch Video Solution

60. How many optically active stereoisomers are possible for butane-2, 3-diol

1

2

3

4

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

61. The optically active tartaric acid is named as $D - (+) -$ tartaric acid because it has a positive

A. Optical rotation and is derived from D -glucose.

B. pH in organic solvent.

C. Optical rotation and is derived from

$D - (+) -$ glyceral-dehyde.

D. Optical rotation when substituted by

deuterium.

Answer: C



Watch Video Solution

62. Which of the following compounds will exhibit geometrical isomerism ?

A. 1- Phenyl-2 butene

B. 3-Phenyl1-butene

C. 2-Phenyl-1-butene

D. 1, 1-Diphenyl 1 – 1-propene

Answer: A



Watch Video Solution

63. The number of isomers of the compound

C₂FClBrI is :

A. 3

B. 4

C. 5

D. 6

Answer: D



Watch Video Solution

64. Which of the following compounds exhibits stereoisomerism?

A. 2-Methyl butene

B. 3-Methyl butyne


C. 3-Methyl butanoic acid

D. 2-Methyl butanoic acid

Answer: D



Watch Video Solution



65.

In the above compound, C_2 is rotated clockwise 120° about $C_2 - C_3$ bond. The resulting conformer is:

A. Partially eclipsed

B. Eclipsed

C. Gauche

D. Staggered

Answer: C



Watch Video Solution

Assertion Reasoning Type Exercise

1. Assertion (A): Metamers can also be position or chain isomers.

Reason (R): Tautomerism was introduced by *C. P. Laar* to explain the chemical reactivity of a substance according to two possible structures.

A. If both (A) and (R) are true and (R) is the correct explanation for (A).

B. If both (A) and (R) are true and (R) is not the true explanation for (A)

C. If (A) is true, but (R) is false.

D. If both (A) and (R) are false

Answer: B



Watch Video Solution

2. Assertion (A): Pentane and 3 methyl pentane are chain isomers.

Reason (R): Pentane is a straight-chain alkane while 3-methyl pentane is branched-chain alkane.

A. both (A) and (R) are true and (R) is the correct explanation for (A).

B. both (A) and (R) are true and (R) is not the true explanation for (A)

C. (A) is true, but (R) is false.

D. (A) is false and (R) is true.

Answer: D



Watch Video Solution

3. Assertion (A) . Pent-1-ene and pent-2-ene are position isomers.

Reason (R). Position isomers differ in the position of functional group or a substituent.

A. If both (A) and (R) are true and (R) is the correct explanation for (A).

B. If both (A) and (R) are true and (R) is not the true explanation for (A)

C. If (A) is true, but (R) is false.

D. If (A) is false and (R) is true.

Answer: D



Watch Video Solution

4. Assertion (A): Both cis 1, 3-dimethyl cyclobutane and trans 1 – 3, dimethyl cyclobutane are optically inactivity

Reason (R): cis-1,3- Dimethyl cyclobutane has the plane of symmetry, wheras trans form has the centre of symmetry.

A. If both (A) and (R) are true and (R) is the correct explanation for (A).

B. If both (A) and (R) are true and (R) is not the true explanation for (A)

C. If (A) is true, but (R) is false.


D. If both (A) and (R) are false

Answer: A



Watch Video Solution

5. Assertion (A): The cis form of



exist in three-diastereomers.

Reason (R): One form is optically inactive due to the presence of centre of symmetry.

A. If both (A) and (R) are true and (R) is the correct explanation for (A).

B. If both (A) and (R) are true and (R) is not the true explanation for (A)

C. If (A) is true, but (R) is false.

D. If both (A) and (R) are false

Answer: C



[Watch Video Solution](#)

Fill In The Blanks Exercise

1. Isomers which are mirror images are known as (superimposable, non-superimposable, enantiomers, diastereomers, epimers)



[Watch Video Solution](#)

2. The structure of the enol form of



intermolecular hydrogen bonding is.....



[Watch Video Solution](#)

3. The total number of cyclic isomers possible for a hydrocarbon with the molecular formula

C_4H_6 is



[Watch Video Solution](#)

True False Exercise

1. *m*-Chloro bromobenzene is an isomer of *m* bromo chloro benzene.



[Watch Video Solution](#)

2. 2, 3, 4- Trichloropentane has three asymmetric carbon atoms.



[Watch Video Solution](#)

Analytical And Descriptive Type Exercise

1. Write the structure of the isomers of alcohols with molecular formula $C_4H_{10}O$.

Which of these exhibits optical activity ?



[Watch Video Solution](#)

2. Write the structural of all the possible isomers of dichloroethene. Which of them will have zero dipole moment?



[Watch Video Solution](#)

3. Write more stable tautomeric structures of the following:

(a) Phenol

(b) Methyl-3-oxobutane

(c) Cyclohexan-1,3,5-trione



[Watch Video Solution](#)


4. Discuss the hybridisation of carbon atoms in allene (C_3H_4) and show the π -orbital

overlap.



Watch Video Solution

5. Identify the pairs of enantiomers and diastereomers from the following:



Watch Video Solution

6. Draw Newman projection of the less stable staggered form of butane.

b. Give reason for the relatively less stability of the staggered form.



[Watch Video Solution](#)

Assertion Reasoning Type

1. Statement I: Molecules that are non-superimposable on their mirror images are

chiral.

Statement II: All chiral molecules have chiral centres.

A. Statement I and II are true and

Statement II is the correct explanation for Statement I.

B. Statement I and II are true, but

Statement II is not the correct explanation of Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

Answer: C



Watch Video Solution