



CHEMISTRY

BOOKS - CENGAGE CHEMISTRY (ENGLISH)

PERIODIC CLASSIFICATION OF ELEMENTS AND GENERAL INORGANIC CHEMISTRY

Illustration

1. What would be the IUPAC name and symbol for the element with atomic number 120 ?



Watch Video Solution

2. There are 2, 8 and 8 elements in the first, second and third periods of periodic table respectively.

Explain.



[Watch Video Solution](#)

3. Which of the following is correct:

(a) The element Mendelebium (d) has been named in the honour of Mendeleev. What is the atomic number of that element?

(i) 100 , (ii) 101 , (iii) 102 , (iv) 103

(b) The element Seaborgium (Sg) has been named in

the honour of Glenn *T.* Seaborg. What is the atomic number of that elements?

(i) 104 , (ii) 105 , (iii) 106 , (iv) 107

(c) Glenn *T.* Seaborg was awarded Nobel Prize in 1951 for the discovery of which element / elements?

(i) Uranium (*U*)

(ii) Elements from 90 to 93

(iii) Elements from 94 to 102

(iv) elements from 103 to 106



Watch Video Solution

4. (a) Which of the atomic numbers and the *IUPAC* name and symbol for the elements Mendeleevium

(*Md*) and Seaborgium (*Sg*)?

(b) What is the atomic number of the element for which both the American and Soviet scientists claimed credit for the discovery?

(c) Refer to the problem (b) above, what name is given to the element by the American and Soviet scientists?



[Watch Video Solution](#)

5. How would you justify the presence of 18 elements in the 5th period of the Periodic Table?



[Watch Video Solution](#)

6. Considering the atomic number and position in the periodic table, arrange the following elements in the increasing order of metallic character : Si, Be, Mg, Na, P.



[Watch Video Solution](#)

7. Predict the period, group number and block of the following elements. A (at.no = 8), B (at.no. = 11), C (at.no. = 28), D (at.no. = 54).



[Watch Video Solution](#)

8. What is the atomic number of the element having maximum number of unpaired $2p$ electrons? To which group it belongs?



[Watch Video Solution](#)

9. The elements $Z = 117$ and 120 have not yet been discovered. In which family / group would you place these elements and also give the electronic configuration in each case.



[Watch Video Solution](#)

10. Write the electronic configuration of the following elements and predict the block, period and group to which each element belongs.

E ($Z = 47$)



Watch Video Solution

11. Write the names and the atomic numbers (Z) of the following elements:

- a. The fourth alkaline earth metal
- b. The fifth alkali metal
- c. The sixth element of the first transition series

d. The second inner transition elements and

e. The third noble gas



[Watch Video Solution](#)

12. How do the electronic configuration of the elements with $Z = 106 - 108$ differ from one another?



[Watch Video Solution](#)

13. Predict the name and position of the element in the periodic table with the electronic configuration

$(n - 1)d^8ns^2$ for $n = 5$.



[Watch Video Solution](#)

14. Elements A to E have the following electronic configuration:

$A: [He]2s^22p^2$, $B: [Ne]3s^23p^2$, $C: [Ne]3s^23p^3$,

$D: [Ne]3s^23p^4$, $E: [Ar]4s^2$.

Which of the above will belong to the same group in the periodic table?



[Watch Video Solution](#)

15. Which of the following species will have the largest and the smallest size Mg , Mg^{2+} , Al , Al^{3+} ?

 [Watch Video Solution](#)

16. a. Compare the size of Cl , and Cl^{\ominus} ion.

b. the radii of Ar is greater than the radii of chlorine.

Explain.

 [Watch Video Solution](#)

17. a. Give and explain the decreasing order of atomic radius of fluorine (F), nitrogen (N) and oxygen (O).

b. Give the decreasing order of van der Waals radii:

N, O, H, Cl, Br.



Watch Video Solution

18. a. In the sixth period, after filling of $6p$ orbitals, the next electron (i.e. 57^{th}) enters the $5d$ -orbital against aufbau principal and there after the filling of seven $4f$ -orbitals starts with cerium ($Z = 71$). Explain this anomalous behaviour.



Watch Video Solution

19. which of the following species are isoelectronic ?

(i) Ne , (ii) O^{2-} , (iii) Mg^{2+} , (iv) F

(v) Al^{3+} , (vi) Cl^{\ominus} , (vii) K , (viii) Na

Arrange them in decreasing order of their size.



Watch Video Solution

20. Which of the following species has the smallest size?

(a) K^{\oplus} , Sr^{2+} , Ar , (b) Si , P , Cl , (c) O , O^{\ominus} , O^{2-}



Watch Video Solution

21. Arrange the following in order of decreasing radii?

a. F^{\ominus} , O^{2-} , N^{3-} , S^{2-} , b. P , Si , N , C , c. I^{\ominus} , I^{\oplus} , I



Watch Video Solution

22. Calculate the effective nuclear charge experienced by the $4s$ -electron in potassium atom ($Z = 19$).



Watch Video Solution

23. Calculate the effective nuclear charge of the last electron in an atom. The electronic configuration is $1s^2, 2s^2 2p^6, 3s^2 3p^5$.



[Watch Video Solution](#)

24. Calculate the screening constant in Zn .

a. For a $4s$ -electron b. For a $3d$ -electron



[Watch Video Solution](#)

25. The first ionization enthalpy ($\Delta_i H$) values of the third period elements, Na, Mg and Si are respectively

496, 737 and 786 kJ/mol. Predict whether the first $\Delta_i H$ value for Al will be more close to 575 or 760 kJ/mol. Justify your answer.

 [Watch Video Solution](#)

26. Calculate the energy required to convert all the atoms of M (atomic number : 12) to M^{2+} ions present in 12 mg of metal vapours. First and second ionization enthalpies of M are 737.77 and 1450.73 kJ mol^{-1} respectively.

 [Watch Video Solution](#)

27. The first (IE_1) and second (IE_2) ionization energies (KJ/mol) of a few elements designated by roman numerals are given below. Which of these would be an alkali metal?



[Watch Video Solution](#)

28. The electronic configuration for some neutral atoms are given below.

A: $1s^2 2s^2$, B: $1s^2 2s^2 2p^3$

C: $1s^2 2s^2 2p^4$, D: $1s^2 2s^2 2p^6 3s^1$

In which of this electronic configuration would you expect to have highest (a) IE_1 and (b) IE_2 .



[Watch Video Solution](#)

29. The ionisation potential of hydrogen is 13.60eV .

Calculate the energy required to produce one mole

of H^{\oplus} ion ($1\text{eV} = 96.3\text{kJmol}^{-1}$).



Watch Video Solution

30. Which of the following electronic configurations

has the lowest value of ionisation energy? Explain.

a. $1s^2 2s^2 2p^6 3s^1$

b. $1s^2 2s^2 2p^6$



Watch Video Solution

31. For each of the following pairs which has greater IE and why?

a. Li, Li^{\oplus} , b. K, Br , c. Br, I , d. Na^{\oplus}, Ne





Watch Video Solution

32. Calculate the percentage of $Mg_{(g)}^{\oplus}$ and $Mg_{(g)}^{2+}$ if 2.4g of Mg absorbs 120kJ of energy. The IE_1 and IE_2 of $Mg_{(g)}$ are 740 and 1450kJmol⁻¹.



Watch Video Solution

33. The electronic configuration for the following atoms are given below :



i. From the above configuration, arrange them in decreasing IE .

ii. Which of the electronic configuration given above will have the lowest IE ?

iii. Which of the electronic configuration given above will be for noble gases ?



Watch Video Solution

34. The IE_1 and IE_2 ($kJmol^{-1}$) of three elements A,

B and C are given below :

	<i>A</i>	<i>B</i>	<i>C</i>
IE_1	400	550	1150
IE_2	2650	1070	2090

Identify the elements which represent (a) an alkali metal, (b) an alkaline earth metal or (c) non-metal.



Watch Video Solution

35. The sum of IE_1 and IE_2 and those of IE_2 and

those of IE_3 and IE_4 in ($kJmol^{-1}$) of *Ni* and *Pt*

are :

	$(IE_1 + IE_2)$	$(IE_2 + IE_4)$	Total
<i>Ni</i>	2.5×10^3	8.8×10^3	11.3×10^3
<i>Pt</i>	2.7×10^3	6.7×10^3	9.4×10^3

a. What is the most common oxidation state (O.S.) of *Ni* and *Pt*.



Watch Video Solution

36. The IE_1 of *C* atom is greater than that of boron (*B*) atom, whereas the reverse is true for IE_2 . Explain?



Watch Video Solution

37. The second IE for alkali metals shows a jump while the third IE for alkaline metals shows a jump.

Explain.




Watch Video Solution

38. Which of the following will have the most negative and the least negative electron gain enthalpy ? P,S,Cl,F. Explain your answer.



Watch Video Solution

39. The amount of energy released when 10^{12} atoms of Cl vapours are converted to Cl^{\ominus} ions, according to the equation :



Calculate the $\Delta_{eg}H^{\ominus}$ of Cl atom in $kJmol^{-1}$ and $eVatom^{-1}$.

 [Watch Video Solution](#)

40. The electron gain enthalpy of chlorine $3.7eV$.

How much energy in kJ and $kcal$ is released when $1g$ of chlorine is converted completely to Cl^{\ominus} ion in the gaseous state. ($1eV = 96.3kJmol^{-1}$)

 [Watch Video Solution](#)

41. Write the electronic configuration of the element with atomic number of 9, 11, 21 and 36. Predict the following from these configurations:

(a) Which of them has the lowest ionisation potential ?

(b) Which of them has the highest electron gain enthalpy ?



(c) Which of them are non-metals?

(d) Which of them has zero electron gain enthalpy ?





Watch Video Solution

42. Which of the following has the highest electron gain enthalpy. Give reasons.



Watch Video Solution

43. Arrange the elements with the following electronic configuration of valence electron in decreasing order of $\Delta_{eg}H^\ominus$.



Watch Video Solution

44. Which one of the following pairs has higher $\Delta_{eg}H^\ominus$?

(a) S^\ominus, O , (b) N^\ominus, P

(c) S, O^\ominus , (d) O^\ominus, S^\ominus



Watch Video Solution

45. Give the decreasing order of $\Delta_{eg}H^\ominus$ of the following elements: F, C, N, O .



Watch Video Solution

46. How many Cl atoms can you ionise in the process $Cl \rightarrow Cl^+ + e$, by the energy liberated for the process $Cl + e \rightarrow Cl^-$ for one Avogadro's number of atoms ? (Given : $IP = 13.0eV$ and $EA = 3.60eV$).

 [Watch Video Solution](#)

47. Calculate the percentage of ionic character in $Cs - Cl$ bond in $CsCl$ molecule. The electronegativity values of Cs and Cl are 0.7 and 3.0 respectively.

 [Watch Video Solution](#)

48. The ionic resonance energy of $C - H$ bond is $\approx 6.0 \text{ kcal mol}^{-1}$. The EN of H is 2.1. Calculate EN of carbon ? Given : $\Delta_{C-H} \approx 6.0 \text{ kcal mol}^{-1}$

 [Watch Video Solution](#)

49. The ionic resonance energy of $C - H$ bond is $\approx 6.0 \text{ kcal mol}^{-1}$. The EN of H is 2.1. Calculate EN of carbon ? Given : $\Delta_{C-H} \approx 6.0 \text{ kcal mol}^{-1}$

 [Watch Video Solution](#)

50. Arrange the following compounds in order of their decreasing stabilities:



(Given EN values of element as below)

$$H = 2.1, F = 4, Cl = 3.0, Br = 2.8, I = 2.3, N = 3.0$$



[Watch Video Solution](#)

51. Predict which of the following hydroxy compounds is acidic and which is basic in aqueous solution :

(i). $CsOH$, ii. IOH .

EN

values

are:

$$C_s = 0.7, O = 3.5, H = 2.1, I = 2.5.$$



[Watch Video Solution](#)

52. $NaOH$ behaves as a base while $Zn(OH)_2$ is amphoteric why ?



[Watch Video Solution](#)

53. Calculate the $d_{(N-C)}$ in $(CH_3)_3N$ molecule by using Eqs. (1.18) and (1.19) above and show from which equation $d_{(N-C)}$ closely resembles with the

experimental value of $d_{(M|N-C)} = 1.47\text{\AA}$.

(Given:

$$r_N = 0.75\text{\AA}, r_C = 0.77\text{\AA}, \chi_A = 3.0, \chi_C = 2.5$$



[View Text Solution](#)

54. In which of the following triatomic molecules, the observed bond angle is $116^\circ, 49'$?

a. H_2O b. OF_2 c. CO_2 d. O_3



[Watch Video Solution](#)

55. Using the periodic table, predict the formulas of compounds which might be formed by the following pairs of elements :

(a) Silicon and bromine

(b) Aluminium and sulphur

 [Watch Video Solution](#)

56. Are the oxidation state and covalency of Al in

$[AlCl(H_2O)_5]^{2+}$ same ?

 [Watch Video Solution](#)

57. Show by a chemical reaction with water that

Na_2O is a basic oxide and Cl_2O_7 is an acidic oxide.



[Watch Video Solution](#)

58. Predict the formulae of the stable binary compounds that would be formed by the following pairs of compounds :

(a) Al and Cl , (b) Mg and I

(c) Element 113 and F , (d) Si and S (e) Element 119 and oxygen.



[Watch Video Solution](#)

59. Give the decreasing order of magnetic moment of the following:

(a) Ca , (b) Al , (c) N , (d) O



Watch Video Solution

60. Which bivalent ion of first transition series show the highest paramagnetism?



Watch Video Solution

61. The following species are in the increasing order of their acidic property: ZnO , Na_2O_2 , P_2O_5 , MgO .



Watch Video Solution

62. Give the decreasing order of the basic properties of oxides.

a. Tl_2O , b. Al_2O_3 , c. Tl_2O_3 , d. Ga_2O_3



Watch Video Solution

63. Give the increasing order of melting points of the following compounds:

a. i. NaF , ii. $NaCl$, iii. $NaBr$, iv. NaI

b. i. CaI_2 , ii. $CaBr_2$, iii. $CaCl_2$, iv. CaF_2 , v. $BeCl_2$

d. I. $NaCl$, II. $MgCl_2$, III. $AlCl_3$

e. I. CCl_4 , II. BCl_3 , III. $BeCl_2$, IV. $LiCl$



Watch Video Solution

64. Give the decreasing order of covalent character of the following compounds.

a. I. $GeCl_2$ II. $GeCl_4$ III. $SnCl_2$ IV. $SnCl_4$

V. $PbCl_2$ VI. $PbCl_4$

b. I. CH_4 II. NH_3 III. H_2O IV. HF

c. I. HF II. HCl III. HBr IV. HI

d. I. AgI II. NaI III. CuI IV. NaCl



Watch Video Solution

65. Arrange the following in increasing order of covalent character - $NaCl$, $MgCl_2$, $AlCl_3$.

 [Watch Video Solution](#)

66. Calculate the ratio of lattice energies of $CaCl_{2(s)}$ and $NaCl_{(s)}$, if the inter-ionic distance in $CaCl_2$ is twice that of $NaCl_{(s)}$.

 [Watch Video Solution](#)

67. Give the decreasing order of melting points of the following compounds:

i. NaF ii. BeO iii. MgO iv. SrO

Given: The inter-ionic distance in Å as

$$NaF = 2.31, BeO = 1.65, MgO = 2.106, SrO = 2.58$$

b. Given the decreasing order of hardness of the following compounds:

i. CaO ii. BeO iii. TiC



Given : The inter-ionic distances in Å as

$$CaO = 2.405, BaO = 2.762, TiC = 2.159$$



[Watch Video Solution](#)

68. Identify the complexes which are expected to be coloured.



[Watch Video Solution](#)

69. ($p\pi - p\pi$) back bonding occurs in the halides of boron but not in those of aluminium. Explain.







[Watch Video Solution](#)

70. The Lewis acid character of boron trihalides decreases as: Explain ?



[Watch Video Solution](#)

71. The correct order of relative basic character of NaOH , $\text{Mg}(\text{OH})_2$ and $\text{Al}(\text{OH})_3$ is



Answer: C



Watch Video Solution

72. Which of the oxides behave both as neutral oxide and suboxide ?

(a) N_2O , b. NO , c. C_3O_2 , d. CO



Watch Video Solution

73. Which of the following is not amphoteric oxide ?

A. ZnO

B. BeO

C. Al_2O_3




D. CrO_3

Answer: D



Watch Video Solution

74. Which of the following is superoxide ?





Answer: C



Watch Video Solution

75. Which of the oxides is coloured and contains $3e^-$ bond ?



Answer: C



Watch Video Solution

76. The least stable hydride is

A. Staanane

B. Silane

C. Plumbane




D. Germane

Answer: C



Watch Video Solution

77. The most stable hydride is




Answer: A



Watch Video Solution

78. The strongest reducing hydride is







Answer: A



Watch Video Solution

79. Give the decreasing order of acidic strength of the following :



(e) LiH

(f) BeH_2 .



Watch Video Solution

80. Arrange the following acids in the decreasing order of their acid strength:

HF, HCl, HBr, HI



Watch Video Solution

81. Arrange the order of decreasing/increasing properties given below:

1. Decreasing order of atomic and ionic radii

a. Mg^{2+} , O^{2-} , Na^{\oplus} , F^{\ominus}



[Watch Video Solution](#)

Solved Examples

1. Predict the group number and block to which the given elements belong.

$A(Z = 9)$, $B(Z = 20)$ and $C(Z = 29)$



[Watch Video Solution](#)

2. The first (IE_1) and second (IE_2) Ionization energies (KJ/mol) of a few elements designated by roman numerals are given below. Which of these would be an alkali metal?

 [Watch Video Solution](#)

3. Identify the three elements A , B and C from the data given below

a. The elements have successive atomic numbers

b. A forms a stable anion, A^\ominus

c. C forms a stable cation, C^\oplus

 [Watch Video Solution](#)

4. Among the elements, *Ar*, *Si*, *Na* and *Cl*. Select an elements with

a. Highest *IE* b. Highest *EA*

c. Smallest size d. Highest electrical conductivity



[Watch Video Solution](#)

5. Arrange in decreasing order as directed.

(a) Decreasing order of *EN*: *C*, *O*, *N*, *F*



[Watch Video Solution](#)

6. Predict from each set, the atom/ion which has the greatest IE_1 with explanation:

a. Cl or F , b. S or Cl



Watch Video Solution

7. The $\Delta_{eg}H^\ominus$ of Br is $3.4eV$. How much energy in kcal is released when $0.8g$ of $Br(g)$ is completely converted to $Br^\ominus(g)$ ions.

$$(1eV = 23.06kcalmol^{-1})$$

b. The energy released when 10^7 atoms of $I(g)$ is converted to $I^\ominus(g)$ ions, is $5 \times 10^{-13}J$. Calculate $\Delta_{eg}H^\ominus$ of $I(g)$ in (i) $eVatom^{-1}$ and (ii) $kJmol^{-1}$.

 [Watch Video Solution](#)

8. Predict from each set, the element which has the more negative electron gain enthalpy ($\Delta_{eg}H^\ominus$).

Give reasons:

a. *C* or *Si* , b. *F* or *Cl* , c. *O* or *S*

 [Watch Video Solution](#)

9. The electronegativity of caesium is 0.7 and that of fluorine is 4.0 . The bond formed between the two is

 [Watch Video Solution](#)

10. Among the elements with $Z = 9, 12$ and 36 ,

identify by atomic number of an element which is

(a) Highly electropositive

(b) Highly EN

(c) An inert gas



[Watch Video Solution](#)

11. Explain the question (based on ionisation energy):


(a) Why IE_1 of N is higher than that of O atom ?



[Watch Video Solution](#)

12. Answer the following question (Based on EA , $\Delta_{eg}H^\ominus$ and IE).

IE_1 of Li is $5.4eV\text{atom}^{-1}$ and the EA of Cl is $3.6eV\text{atom}^{-1}$. Calculate Δ_rH^\ominus in kcal mol^{-1} and kJmol^{-1} for the reaction



formed at such a low pressure that resulting ions do not combine with each other.



[Watch Video Solution](#)


13. a. Arrange the following species in decreasing order of their sizes//ionic radii.

i. Ar , K^{\oplus} , Cl^{\ominus} , S^{2-} and Ca^{2+}



Watch Video Solution

14. For the gaseous reaction




$\Delta H = 19 \text{ kcal mol}^{-1}$ under the condition when cations and anions are prevented by electrostatic separation from combining with each other. The IE_1 of K is 4.3 eV . Calculate $\Delta_{\text{eg}} H^{\ominus}$ of F .



Watch Video Solution

15. For the gaseous reaction



$\Delta H = 19 \text{ kcal mol}^{-1}$ under the condition when cations and anions are prevented by electrostatic separation from combining with each other. The IE_1 of K is 4.3 eV . Calculate $\Delta_{\text{eg}} H^{\ominus}$ of F .

 [Watch Video Solution](#)

16. From N atoms of an element A , when half the atoms transfer one electron to the another atom.

405 kJ mol^{-1} of energy was found to be consumed.

An additional energy of 745 kJ mol^{-1} was further

required to convert all the A^\ominus ions to A^\oplus . Calculate the ionisation energy and the electron gain enthalpy of atom A in eV ($1eV = 96.48kJ$).

 [Watch Video Solution](#)

17. The conservation of gaseous atoms K and F to K^\oplus and F^\ominus absorbs $0.85eV$ of energy. If the IE and $\Delta_{eg}H^\ominus$ of K and F have magnitudes in the ratio of 7:6, what is the electron gain enthalpy ($\Delta_{eg}H^\ominus$) of fluorine ?

 [Watch Video Solution](#)

18. Which of the following Na , Mg , Si and P would have the greatest difference between the IE_1 and IE_2 ?



Watch Video Solution

19. Classify the following oxides as

a. Strongly acidic b. Weakly acidic

c. Neutral d. Amphoteric

e. Weakly basic and f. Strongly basic

i. SnO_2 ii. SnO iii. CO iv. PbO

v. MnO_2 vi. RaO vii. N_2O viii. FeO

ix. Ag_2O x. OsO_4 xi. Al_2O_3 xii. Fe_2O_3

xiii. CeO_2 xiv. CO_2 xv. MgO xvi. K_2O



Watch Video Solution

20. Select the strongest and weakest acid in each of the following sets:

a. HBr , HF , H_2Te , H_2Se , PH_3 , H_2O

b. $HClO$, HIO , H_3PO_3 , H_2SO_3 , H_2AsO_3



Watch Video Solution

21. A $0.10M$ aqueous solution of which salt in the following pair would have the higher pH ?

$NaNO_2$ or $NaAsO_2$



Watch Video Solution

22. Identify:

a. The good oxidising agent

b. The good reducing agent

c. The good dehydrating agent (s) among the following substances:

H_2SO_3 , HNO_3 , P_4O_{10} , H_2S , H_2SO_4 .



Watch Video Solution

23. Which of the following has the greatest affinity

for water: P_4O_{10} , Cl_2O_7 , I_2O_5 .



Watch Video Solution

24. Why Al^{3+} is the only stable oxidation state of Al in its compounds while Tl has +1 and +3 oxidation states?

 [Watch Video Solution](#)

25. Give the name and atomic number of the inert gas atom in which the total number of d -electrons is equal to the difference in number to the p - and s -electrons.

 [Watch Video Solution](#)

26. Classify the elements having atomic numbers (9, 12, 16, 34, 53, 56) into three separate pairs on the basis of similar chemical properties.



Watch Video Solution

27. Calculate the electronegativity of fluorine from the following data:

$$E_{H-H} = 104.4 \text{ kcal mol}^{-1}, E_{F-F} = 36.6 \text{ kcal mol}^{-1}$$

$$E_{H-F} = 134.3 \text{ kcal mol}^{-1}, \chi_H = 2.1$$



Watch Video Solution

28. Calculate electronegativity of carbon at Pauling scale Given that :

$$E_{H-H} = 104.2 \text{ kcal mol}^{-1} \quad E_{C-C} = 83.1 \text{ kcal mol}^{-1} ,$$

$$E_{C-H} = 98.8 \text{ kcal mol}^{-1} .$$

Electronegativity of hydrogen = 2.1.

 [Watch Video Solution](#)

29. Ionisation potential and electron affinity of fluorine are 17.42 and 3.45 eV respectively .Calculate the electronegativity of fluorine on Mulliken scale and Pauling scale .

 [Watch Video Solution](#)

30. Calculate the electronegativity of silicon using Allred -Rochow method .(Covalent radius of silicoan = 1.175\AA).

 [Watch Video Solution](#)

31. Calculate the electronegativity value of chlorine on Mulliken's scale, given that $IP = 13.0eV$ and $EA = 4.0eV$.

 [Watch Video Solution](#)

32. Find the electronegativity of lead with the help of the given values. Screening constant (σ) of $Pb = 76.70$. Atomic number of lead = 82 and covalent radius of $Pb = 5.3\text{\AA}$

 [Watch Video Solution](#)

33. The ionisation potentials of atoms A and B are 400 and 300kcalmol^{-1} respectively. The electron gain enthalpy of these atoms are 80.0 and 85.0kcalmol^{-1} respectively. Which of the atoms has higher electronegativity.

 [Watch Video Solution](#)

34. Give the names of seven f -orbitals and how they are represented.



Watch Video Solution

Ex 1.1

1. $3d -$, $4d -$ and $5d -$ series consists of 10 elements each? Explain.



Watch Video Solution

2. Why the f -block elements are called inner transition elements?

 [Watch Video Solution](#)





3. Transition elements show horizontal as well as vertical relationship.

 [Watch Video Solution](#)

4. Be and Al are placed in different periods and groups but they show the similar properties. Explain.

 [Watch Video Solution](#)

5. The outer electronic configuration of some elements are given below:



State to which of the periodic table each of these elements belongs.



Watch Video Solution

6. Arrange the following elements in decreasing order of metallic character:

K, Mg, B, Al.



Watch Video Solution

7. Name one species that will be isoelectronic with the following atoms or ions.

(a) Ca^{2+} , (b) Ne , (c) Cl^{\ominus} , (d) Rb^{\oplus}



Watch Video Solution

8. Which of the following pairs would have a smaller size. Explain.

(a) Na^{\oplus} or Mg^{2+} , (b) O^{2-} or F^{\ominus}

(c) P or As



Watch Video Solution

9. Arrange the following ions in the decreasing order of their sizes and explain.

Li^+ , K^+ , Mg^{2+} , Al^{3+}



Watch Video Solution

10. Which important property did Mendeleev use to classify the elements in his periodic table and did he stick to that?



Watch Video Solution

11. Elements with $Z = 107, 108$ and 109 have been made recently. Indicate the groups to which they belong.



Watch Video Solution

12. Assertion : Zn, Cd and Hg are not regarded as transition elements.

Reason : The electronic configurations of Zn, Cd and Hg are represented by the general formula $(n - 1)d^{10}ns^2$.



[Watch Video Solution](#)

13. Why *Cu*, *Ag* and *Au* are transition elements although they have completely filled *d*-orbitals ?



[Watch Video Solution](#)

14. Stability Of Completely Filled And Half Filled Orbitals

 [Watch Video Solution](#)

Ex 1.1 (Very Short)

1. What are super heavy elements?

 [Watch Video Solution](#)

2. How many anomalous pairs were present in original Mendeleev periodic table ?

 [Watch Video Solution](#)

3. How many *s*-block elements are known ?

 [Watch Video Solution](#)

4. Which block of elements consists of metals, non metals and metalloids ?

 [Watch Video Solution](#)

5. Which is the hardest elements ?



[Watch Video Solution](#)

6. Which elements is lowest melting point liquid metal ?



[Watch Video Solution](#)

7. Which element is heaviest melting and boiling point metal ?



[Watch Video Solution](#)

8. Write the name the liquid non-metal element.

 [Watch Video Solution](#)

9. Name the best and the poorest conductor of current among metals.

 [Watch Video Solution](#)

10. Name the heaviest solid metal.

 [Watch Video Solution](#)

11. Name the most poisonous element.



Watch Video Solution

12. Name of the heaviest naturally occurring element.



Watch Video Solution

13. Among the radioactive elements, which is a liquid element?



Watch Video Solution

14. Name the element having the lowest electronegative, lightest and liquid metal.

 [Watch Video Solution](#)

15. Name the lightest solid non-metal having the highest tensile strength.

 [Watch Video Solution](#)

16. Which element has highest catenation property?

 [Watch Video Solution](#)

17. Which non-metal sublime on heating and have metallic lusture.

 [Watch Video Solution](#)

18. Which is the most stable element?

 [Watch Video Solution](#)

19. Name the element which is the poorest conductor of current among non-metals.

 [Watch Video Solution](#)

20. Name the elements with highest EN and with highest EA or $\Delta_{eg}H^\ominus$.

 [Watch Video Solution](#)

21. Which is used in high temperature thermometry ?

 [Watch Video Solution](#)

22. Which element is used in the making of infrared (IR) windows, prism and lenses ?

 [Watch Video Solution](#)

Ex 1.1 (Objective)

1. Which of the following statement is wrong?

A. Among the following elements :

K, Mn, Ca, Cs, Fe, Cu, Pb, Os, Y

The number of transition element is 4.

B. All the lanthanides and actinides belong to

IIIB or the *3rd* group

in the periodic table.

C. The inner transition elements belong to the *f*-

block of the periodic table and are shown

separately at the bottom of the periodic table.

D. The d -block elements have variable valency.

Answer: A



Watch Video Solution

2. Which of the following statement is wrong?

A. In the lanthanide series the electrons occupy

$4f$ -orbitals in

preference to $5d$ the and $6p$ -orbitals.

B. Zero group was not present in the periodic table when Mendeleev presented it.

C. Law of octave was presented by Newland.

D. Cuprous compounds are coloured while cupric compounds are colourless.

Answer: D



Watch Video Solution

3. Which of the following triads have approximately equal size ?

A. Na^{\oplus} , Mg^{2+} , Al^{3-} (isoelectronic)

B. F^{\ominus} , Ne , O^{2-} (isoelectronic)

C. Fe , Co , Ni

D. Mn^{1+} , Fe^{2+} , Cr (isoelectronic)

Answer: C



Watch Video Solution

4. Which is the correct order of size ?

$(O^{\ominus}, O^{2-}, F^{\ominus}$ and $F)$

A. $O^{2-} > O^{\ominus} > F^{\ominus} > F$

B. $O^\ominus > O^{2-} > F^\ominus > F$

C. $O^{2-} > F^\ominus > F > O^\ominus$

D. $O^{2-} > F^\ominus > O^\ominus > F$

Answer: A


 [Watch Video Solution](#)

5. Select the correct statement :


Which of the following graph represents Mosely's experiment.

(ν = frequency of X-rays)


A.




B.



C.



D.



Answer: D



Watch Video Solution

6. Calculate the screening constant of alkali metals for valency electrons.

 [Watch Video Solution](#)

7. Calculate the screening constant of Li for valency electrons.

 [Watch Video Solution](#)

8. Answer the following :

(a) Name of the element not known at time of Mendeleev.

(b) Name any two typical elements.

(c) Name any two bridge elements.

(d) Name two pairs showing diagonal relationship.

(e) Name two transition elements.

(f) Name two rare earth elements.

(g) Name two transuranic elements.

(h) In which case compounds of Cr show maximum radius.

(i) K_2CrO_7 , (ii) CrO_2Cl_2 , (iii) $Cr_2(SO_4)_3$

(iv) $CrCl_2$



Watch Video Solution

9. In s - and p -block elements the OS . changes in 2 units, but in transition elements it changes it chnages in units of one.(T/F)



Watch Video Solution

10. Mn_2O_7 is an acidic oxide, why ?



Watch Video Solution

11. Why Ar (argon) (at, wt. = 39.94) has been placed before K (at. Wt. = 39.10) in the periodic table?



Watch Video Solution

 Watch Video Solution

12. Why Ag is a noble metal and K is a highly reactive metal ?



Watch Video Solution

Ex 1.2

1. Answer the following :

(a) Why inert gases are monoatomic ?

(b) Comment on 'Iodine possesses some metallic lusture'.

(c) Of all noble metals, gold (Au) has a relatively high EA . Explain.

(d) Cl and be converted to Cl^\ominus ion easily than F to F^\ominus ion. Explain.

(e) Why $Al(OH)_3$ is amphoteric in nature ?





Watch Video Solution

Ex 1.2(Objective)

1. Which has maximum polarising power in cation ?





Answer: B



Watch Video Solution

2. Which has the maximum IE ?



Answer: A



Watch Video Solution

3. The correct order of hydration enthalpies of alkali metal ions is:



Answer: B





Watch Video Solution

4. Inert pair effect is shown by

A. *s*-block

B. *p*-block

C. *d*-block

D. *f*-block

Answer: B



Watch Video Solution

5. Which is / are amphoteric oxide ?

A. ZnO

B. BeO

C. SnO

D. All of these

Answer: D



Watch Video Solution

6. EA is positive when

A. O^{\ominus} is formed from O

B. O^{\oplus} is formed from O

C. O^{2-} is formed from O

D. EA is always a negative value

Answer: C





Watch Video Solution

7. Which has the maximum covalent character?

A. $MgCl_2$

B. $NaCl$






Answer: C



Watch Video Solution

8. In which solvent KBr has maximum solubility?



Answer: D



Watch Video Solution

9. Lattice energy of $BeCO_3(I)$, $MgCO_3(II)$ and $CaCO_3(III)$ is in order.

A. $I > II > III$

B. $III > II > I$

C. $II > III > I$

D. $III > I > II$

Answer: A



10. NO_2 and N_2O_4 are two forms of nitrogen dioxide. One exists in gaseous state while other in liquid state. The nature of NO_2 and N_2O_4 forms are

A. Both are diamagnetic

B. Both are paramagnetic

C. NO_2 is diamagnetic while N_2O_4 is paramagnetic

D. NO_2 is paramagnetic while N_2O_4 is diamagnetic

Answer: D



Watch Video Solution

11. Magnetic moment of $V(Z = 23)$, $Cr(Z = 24)$, and $Mn(Z = 25)$ are x , y , z respectively hence

A. $x = y = z$

B. $x < y < z$

C. $x < z < y$

D. $z < y < x$

Answer: C



Watch Video Solution

12. Solubility of groups 1 and 2 fluorides increases down the group'. Which of the following is correct explanation for the above given statement ?

A. Both the hydration and lattice energies decrease down the group (\downarrow) but decrease in lattice energy is rapid.

B. Both the energies increase down the group but increase in hydration energy is rapid.

C. Both the energies decrease down the group
but decrease in hydration energy is rapid.

D. Hydration energy increases and lattice energy
decreases down the group.




Answer: A



Watch Video Solution

13. Which of the following molecule is theoretically
not possible ?

A. OF_4





Answer: A



Watch Video Solution

14. Which of the following triads have approximately equal size ?



C. F^{\ominus} , Ne , O^{2-} (isoelectronic)

D. Fe , Co , Ni

Answer: D



Watch Video Solution

15. Which pair is different from the others ?

A. $Na - K$

B. $Ca - Mg$

C. $Li - Mg$

D. $B - Al$

Answer: C



Watch Video Solution

16. Compound XY is predominantly ionic as $X^{\oplus}Y^{\ominus}$ if

A. $(IE)_X < (IE)_Y$

B. $(EA)_X < (EA)_Y$

C. $(EN)_X < (EN)_Y$

D. $(IE)_Y < (IE)_X$.

Answer: C



17. (X) , (Y) , (Z) are elements in third short period. Oxide of (X) is ionic, (Y) is amphoteric and (Z) is a giant molecule. (X) , (Y) and (Z) will have atomic number in the order :

A. $(X) < (Y) < (Z)$

B. $(Z) < (Y) < (X)$





C. $(X) < (Z) < (Y)$

D. $(Y) < (X) < (Z)$

Answer: A

18. Which of the correct order of size ?

$(O^\ominus, O^{2-}, F^\ominus \text{ and } F)$



Answer: A



Watch Video Solution

19. F has the highest electronegativity among the group 17 elements (i.e. $ns^2np^2np^5$ type), on the Pauling scale, but the EA of F is less than that of Cl because

A. F being the first member of the family behaves in an unusual manner.

B. The atomic number of F is less than that of Cl .

C. Cl can accommodate an electron better than F by utilising its vacant $3d$ orbital.





D. Small size, high EN and an increased electron-electron repulsion makes addition of an electron to F less favourable.

Answer: D



Watch Video Solution

20. The correct order of decreasing ionic character is







Answer: A



Watch Video Solution

21. The correct order of decreasing polarisability of ion is



Answer: C



Watch Video Solution

22. Which of the following has the smallest bond length ?



Answer: B



Watch Video Solution

1. Arrange the following in order of decreasing ionic character.

a. ClF_3 , SO_2 , N_2 , K_2O and LiF

b. $Cl - H$, $F - H$, $Br - H$, $Na - I$, $K - F$ and $Li - Cl$

c. AlF_3 , $AlCl_3$, $AlBr_3$



Watch Video Solution

2. Arrange the following in order of decreasing bond angle.

CO_2 , H_2O , CH_4



Watch Video Solution

3. $CaCO_3$ dissolves in HCl but not in water. Why ?

 [Watch Video Solution](#)

4. Why MgO exist as $Mg^{2+}O^{2-}$ not as $Mg^{\oplus}O^{\ominus}$ whereas the formation of Mg^{2+} from Mg requires more energy than formation of Mg^{\oplus} and formation of O^{\ominus} from O is exothermic whereas the formation of O^{2-} is endothermic .

 [Watch Video Solution](#)

5. Anhydrous $AlCl_3$ is covalent. From the data given below, predict whether it would remain covalent or become ionic in aqueous solution. (Ionisation energy for Al is $1537kJmol^{-1}$)

$$\Delta_{\text{hydration}} f \text{ or } Al^{3+} = -4665kJmol^{-1}$$

$$\Delta_{\text{hydration}} f \text{ or } Cl^{\ominus} = -381kJmol^{-1}.$$



[Watch Video Solution](#)

6. Which compound for each of the following pairs is more ionic and why?

a. $BeBr_2$ or $MgBr_2$, b. $PbCl_2$ or $PbCl_4$

c. $AgBr$ or AgI , d. CuO or CuS

 [Watch Video Solution](#)


7. $NaBr$ gives pale yellow precipitate with $AgNO_3$ solution but CBr_4 does not. Why ?

 [Watch Video Solution](#)

8. Copper is conducting as such while $CuSO_4$ is conducting only in molten state or in aqueous solution. Why ?

 [Watch Video Solution](#)

9. Explain the observed bond angle order.



 [Watch Video Solution](#)

10. NH_4^\oplus has bond angle identical to CH_4 but NH_3 has different bond angle. Why?

 [Watch Video Solution](#)

11. Electronegativities of F, O, N, Cl, H are 4.0, 3.5, 3, 3 and 2.1 respectively. In which atoms there is strongest bond.



Watch Video Solution

12. The IE_1 of Li is $5.4eV$ and IE_1 of H is $13.6eV$.

Calculate the charge acting on the outermost electron of Li atom.



Watch Video Solution

13. $SnCl_2$ is solid but $SnCl_4$ is liquid. Why ?



Watch Video Solution

1. The correct order of IE_2 of C , N , O and F is

A. $O > F > N > C$

B. $F > O > N > C$

C. $C > N > O > F$

D. $O > N > F > C$

Answer: A



Watch Video Solution

2. The least stable ion among the following is



Answer: D



Watch Video Solution

3. Which has the most stable + 2 oxidation state ?



C. *Pb*



D. *Ag*



Answer: C



Watch Video Solution

4. Amongst the element with following electronic configurations, which one of them may have the highest ionization energy ?





Answer: A



Watch Video Solution

5. The magnitude of lattice energy of a solid increases if

A. The ions are large

B. The ions are small

C. The ions are of equal sizes

D. Charges on the ions are small

Answer: B



Watch Video Solution

6. Bond angle in PH_3 is





- A. Much less than NH_3
- B. Much less than PF_3
- C. Slightly more than NH_3
- D. Much more than PF_3

Answer: A



Watch Video Solution

7. The correct order of decreasing bond angles in H_2S , NH_3 , BF_3 and SiH_4 is





Answer: B



Watch Video Solution

8. The bond angle around central atom is maximum for




Answer: A



Watch Video Solution

Ex 1.3 (Multiple Correct)

1. Which of the following can conduct electricity in?



Answer: A::B::C



Watch Video Solution

2. Which of the following are expected to be covalent?




Answer: A::B



Watch Video Solution

3. Which of the following does not exist?





Answer: A::B::C



Watch Video Solution

4. Which of the following relation is/are correct?

A. Covalent character \propto Pseudo inert configuration

B. Ionic character \propto Inert configuration

C. Covalent character $\propto \frac{1}{\text{Dipole moment}}$

$$\text{D. Ionic character} \propto \frac{1}{\text{Dipole moment}}$$

Answer: A::B::C

 **Watch Video Solution**

5. Which of the following are correct?

A. As^{5+} salts are better oxidising agents

B. Tl^{3+} salts are better oxidising agents

C. Ga^{\oplus} salts are better oxidising agents

D. Pb^{4+} salts are better oxidising agents

Answer: B::C::D



[Watch Video Solution](#)

Ex 1.29 (Integer)

1. On the basis of quantum number, which period of the periodic table should have 32 elements.



[Watch Video Solution](#)

2. What is the atomic number of the element present in the second period and group 15.



[Watch Video Solution](#)

3. $\Delta_f H^\ominus$ of hypothetical MX is -150 kJ mol^{-1} and for MX_2 is -600 kJ mol^{-1} . The enthalpy of disproportionation of MX is $= -100x \text{ kJ mol}^{-1}$.

Find the value of x.



[Watch Video Solution](#)

Ex 1.32 (Assertion-Reasoning)

1. Assertion (A) : IE_1 of Mg is greater than that of Al.

Reason (R) : It is easier to remove an electron from 3s

orbital than

from $3p$ orbital.

A. If both A and R are correct and R is the correct explanation for A.

B. If both A and R are correct and R is not the correct explanation for A.

C. If A is correct and R is incorrect

D. If A is incorrect and R is correct

Answer: C



Watch Video Solution

2. Assertion (A): s- block elements can form ionic hydrides. Which on electrolysis in fused state give H_2 gas at cathode.

Reason (R) : s-block elements have low IE and low EN as compared to p-block elements.

- A. If both A and R are correct and R is the correct explanation for A.
- B. If both A and R are correct and R is not the correct explanation for A.
- C. If A is correct and R is incorrect
- D. If A is incorrect and R is correct

Answer: D



Watch Video Solution

Exercises (Linked Comprehension)

1. Effective nuclear charge (Z_{eff}) is the net attractive force on electrons under consideration and is equal to:

$$Z_{eff} = Z - \sigma \text{ (nuclear charge - screening constant).}$$

Z_{eff} or σ is calculated by Slater's formula, as given.

If one electron is present in the outermost orbit, there will be no screening in that orbital. Each

electrons contribute, 0.35 (total electrons minus one electron) present in the outermost shell.

A contribution of 0.85 for each electrons is taken in the $(n - 1)th$ shell.

The screening constant (σ) for $4s$ electron of $Mn(Z = 25)$ will be

A. 18.00

B. 4.25

C. 18.35

D. 22.6

Answer: D



Watch Video Solution

2. Effective nuclear charge (Z_{eff}) is the net attractive force

on electrons under consideration and is equal to:

$$Z_{eff} = Z - \sigma \text{ (nuclear charge - screening constant).}$$

Z_{eff} or σ is calculated by Slater's formula, as given.

If one electron is present in the outermost orbit, there will be no screening in that orbital. Each electron contributes 0.35 (total electrons minus one electron) present in the outermost shell.

A contribution of 0.85 for each electron is taken in the $(n - 1)th$ shell.

Which of the following statements is wrong?

- A. IE_1 of $Ga > Al$, due to imperfect shielding of $3d$ -orbitals in Ga .
- B. IE_1 of $Ga < Al$, due to perfect shielding of $3d$ -orbitals in Ga .
- C. The atomic size of Ga and Al are almost same because of poor shielding effect of electrons in d -orbitals as the effective nuclear charge increases in Ga .
- D. IE_1 of group 16 elements is less than that of group 15 elements.

Answer: B



3. Effective nuclear charge (Z_{eff}) is the net attractive force on electrons under consideration and is equal to:

$$Z_{eff} = Z - \sigma \text{ (nuclear charge - screening constant).}$$

Z_{eff} or σ is calculated by Slater's formula, as given.

If one electron is present in the outermost orbit, there will be no screening in that orbital. Each electron contributes 0.35 (total electrons minus one electron) present in the outermost shell.

A contribution of 0.85 for each electron is taken in

the $(n - 1)th$ shell.

Which of the following statement is wrong?

- A. The number of lobes in f -orbitals are 8.
- B. IE_1 of elements increases along the period.
- C. IE_1 of the group 3 elements is more than that of the group 2 elements.
- D. IE_1, IE_2 and IE_3 of an element are 9.5, 18.5 and $154.4eV$. Predict that the element has either two s -electrons or two p -electrons in the valence shell.

Answer: C



4. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, s - p - d -and f -. Each period begins with the filling of new energy shell. Two series of f -block elements are placed at the bottom of the periodic table.

The element with $Z = 39$ belongs to

- A. s -block
- B. p -block
- C. d -block

D. *f*-block

Answer: C



Watch Video Solution

5. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, *s* – *p* – *d*-and *f*-. Each period begins with the filling of new energy shell. Two series of *f*-block elements are placed at the bottom of the periodic table.

The element with $Z = 113$ has been discovered. Its

block, group number, period and outer shell electronic configuration are

A. *s*-block, group 2, period 7, $7s^2$

B. *p*-block, group 13, period 7, $7s^2 7p^1$

C. *p*-block, group, 13, period 6, $6s^2 6p^1$

D. *d*-block, group 12, period 6, $5d^{10}, 6s^2$

Answer: B



Watch Video Solution

6. In the long or modern form of the periodic table, the elements in the periodic table have been divided

into four blocks, $s - p - d$ -and f -. Each period begins with the filling of new energy shell. Two series of f -block elements are placed at the bottom of the periodic table.

Which of the element whose atomic numbers are given below cannot be accommodated in the present set-up of the modern periodic table?

A. 109

B. 118

C. 120

D. 125

Answer: D



Watch Video Solution

7. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, s – p – d - and f -. Each period begins with the filling of new energy shell. Two series of f -block elements are placed at the bottom of the periodic table.

What is the maximum number of electrons that can be accommodated with $n = 5$ ($n =$ principal quantum number)?

A. 10

B. 18

C. 36

D. 50

Answer: C



Watch Video Solution

8. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, s - p - d -and f -. Each period begins with the filling of new energy shell. Two series

of

f-block elements are placed at the bottom of the periodic table.

Which of the following ions is most stable?





Answer: A



Watch Video Solution

9. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, s – p – d -and f -. Each period begins with the filling of new energy shell. Two series of f -block elements are placed at the bottom of the periodic table.

The last element of the p -block in the present periodic table is represented by the configuration, where $[X]$ represents inert gas is



C. $[X]4f^{14}, 5d^{10}, 6s^26p^6$

D. $[X]$ None of the above

Answer: C



Watch Video Solution

10. In the long or modern form of the periodic table, the elements in the periodic table have been divided into four blocks, $s - p - d$ -and f -. Each period begins with the filling of new energy shell. Two series of f -block elements are placed at the bottom of the periodic table.

If the elements were discovered in future in which

the electrons would be present in g -orbitals. Then g -orbital starts, and ends up with what atomic number?

A. 121, 139

B. 121, 138

C. 122, 140

D. 122, 139

Answer: B



Watch Video Solution

11. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, diapositive ion and tripositive ion are known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2eV\text{atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9eV$.

The correct order of arrangement of the first ionisation energies of C , N , O and F (in decreasing values) is

A. $C > N > O > F$

B. $O > N > F > C$

C. $O > F > N > C$

D. $F > N > O > C$

Answer: D



Watch Video Solution

12. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, diapositive ion and tripositive ion are known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2eV\text{atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9eV$.

Four elements have the following first ionization energies in KJmol^{-1} : 762, 709, 59 and 558. The elements are *Ca*, *Ge*, *In* and *Sn* (not in order). Which of these elements has the ionisation energy of 762KJmol^{-1} ?

A. *In*

B. *Ga*

C. *Sn*

D. *Ge*


Answer: D



Watch Video Solution

13. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, diapositive ion and tripositive ion are known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2eV\text{atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9eV$.

Which of the following are isoelectronic species?



A. *I*, *II* and *III*

B. *II*, *III* and *IV*

C. *I*, *II* and *IV*

D. *II* and *I*

Answer: B







Watch Video Solution

14. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is

known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, dipositive ion and tripositive ion are known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2 eV \text{ atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9 eV$.

Among the following ionisation reactions, which one will have the maximum value of ionisation energy?








Answer: B


 [Watch Video Solution](#)

15. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, diapositive ion and tripositive ion are

known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2eV\text{atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9eV$.

Consider the IE_1 of the element whose electronic configurations correspond to the following





Answer: C



Watch Video Solution

16. The energy required to remove an electron from the outermost shell of an isolate gaseous atom is known as IE_1 of that atom. Similarly, the energy required for the removal of the electron from the unipositive ion, diapositive ion and tripositive ion are known as IE_2 , IE_3 and IE_4 respectively, and are called successive ionisation energies. The magnitude of the charge depends on the size of the orbital of

electron. Electrons in smaller orbitals are on average close with each other and have more repulsion. Thus for $Be(2s^2)$, the IE_1 and IE_2 are 9.3 and $18.2eV\text{atom}^{-1}$, whereas for $Ca(4s^2)$, the values are 6.1 and $11.9eV$.

The relationship between IE_4 and IE_2 of an element is

A. $IE_4 > IE_2$

B.

C. $IE_4 = IE_2$

D. None of the above

Answer: A



17. Energy is released when an electron is added to neutral isolated gaseous atom in its ground state to give monoanion and this is known as EA , or $\Delta_{eg}H_1^\ominus$. The greater the amount of energy released the greater is the EA . EA is expressed in $eV\text{atom}^{-1}$ or $kcal$ or $KJmol^{-1}$.

The EA values of element depends on the following:

- i. Nuclear charge
- ii. Electroniv configuration
- iii. Atomic size
- iv. chemical environment

A. I,iii, iv

B. I,ii,iii

C. ii,iii,iv

D. All

Answer: B



Watch Video Solution

18. Energy is released when an electron is added to neutral isolated gaseous atom in its ground state to give monoanion and this is known as EA , or $\Delta_{eg}H_1^\ominus$. The greater the amount of energy released the

greater is the EA . EA is expressed in $eV\text{atom}^{-1}$ or $kcal$ or $KkJmol^{-1}$.

EA values of N and P are exceptionally low, because

A. Both N and P have half-filled p -orbitals in the valence shell.

B. The atom is more stable than the corresponding anion.

C. The electronic configuration of the anion N^{\ominus} and P^{\ominus} is relatively more stable than the corresponding atom.

D. Both (b) and C.

Answer: A



Watch Video Solution

19. Energy is released when an electron is added to neutral isolated gaseous atom in its ground state to give monoanion and this is known as EA , or $\Delta_{eg}H_1^\ominus$. The greater the amount of energy released the greater is the EA . EA is expressed in $eV\text{atom}^{-1}$ or $kcal$ or $KJmol^{-1}$.

Select the correct statements (more than one correct)

A. EA and $\Delta_{eg}H_1^\ominus$ of an atom of element have same magnitude

B. $\Delta_{eg}H_1^\ominus$ of $Al > B$

C. $\Delta_{eg}H_1^\ominus$ of $P > N$

D. $\Delta_{eg}H_1^\ominus$ of $S > O$

Answer: A::B::C::D



Watch Video Solution

20. Energy is released when an electron is added to neutral isolated gaseous atom in its ground state to give monoanion and this is

known as EA , or $\Delta_{eg}H_1^\ominus$. The greater the amount of energy

released the greater is the EA . EA is expressed in $eV\text{atom}^{-1}$ or $kcal$ or $kJ\text{mol}^{-1}$.

Select the correct statement (more than one correct)

A. $\Delta_{eg}H^\ominus$ of noble gases have large positive values.

B. $\Delta_{eg}H^\ominus$ of noble gases have large negative values.

C. $\Delta_{eg}H^\ominus$ of helium (He) is the lowest of all the noble gases.

D. $\Delta_{eg}H^\ominus$ of Ar is lower than that of Ne .


Answer: A::C::D



Watch Video Solution

21. Along the period (\rightarrow) atomic / ionic radii and metallic character decreases while IE , EN , non-metallic character and oxidising power increases. Down the group (\downarrow), atomic / ionic radii, metallic character and reducing character increase while IE and EN decrease. However, $\Delta_{eg}H^\ominus$ becomes less negative down a group but more negative along a period.

Which of the following isoelectronic species has lowest IE_1 ?



Answer: C



Watch Video Solution

22. Along the period (\rightarrow) atomic / ionic radii and metallic character decreases while IE , EN , non-

metallic character and oxidising power increases. Down the group (\downarrow), atomic/ionic radii, metallic character and reducing character increase while IE and EN decrease. However, $\Delta_{eg}H^\ominus$ becomes less negative down a group but more negative along a period.

If the ionic radii of M^\oplus and X^\ominus are about $135pm$, then expected values of metallic radii of M and X should be respectively.





- A. 65 and $230pm$
- B. 230 and $60pm$
- C. 230 and $135pm$
- D. 135 and $135pm$

Answer: B

 [Watch Video Solution](#)

23. Along the period (\rightarrow) atomic / ionic radii and metallic character decreases while IE , EN , non-metallic character and oxidising power increases. Down the group (\downarrow), atomic / ionic radii, metallic character and reducing character increase while IE and EN decrease. However, $\Delta_{eg}H^\ominus$ becomes less negative down a group but more negative along a period.

In which of the following pairs, both species have nearly the same size ? (more than one correct)



Answer: B::C::D



Watch Video Solution

24. Along the period (\rightarrow) atomic/ionic radii and metallic character decreases while IE , EN , non-metallic character and oxidising power increases.

Down the group (\downarrow), atomic/ionic radii, metallic

character and reducing character increase while IE and EN decrease. However, $\Delta_{eg}H^\ominus$ becomes less negative down a group but more negative along a period.

Correct order of IE_2 of the following is

A. $F > O > N > C$

B. $O > N > F > C$

C. $O > F > N > C$

D. $C > N > O > F$.

Answer: C



Watch Video Solution

25. Along the period (\rightarrow) atomic/ionic radii and metallic character decreases while IE , EN , non-metallic character and oxidising power increases. Down the group (\downarrow), atomic/ionic radii, metallic character and reducing character increase while IE and EN decrease. However, $\Delta_{eg}H^\ominus$ becomes less negative down a group but more negative along a period.

Correct order of IE_2 of the following is

A. IE and EA are defined at absolute zero temperature. At any other temperature, heat

capacities for the reactants and product have to be taken into account.

B. The ionisation enthalpy ($\Delta_r H^\ominus$) and IE related to each other by the equation.

$$\Delta_i H^\ominus \quad (\text{ionisation enthalpy}) \\ = \left[IE \left(\text{ionisation energy} + \frac{5}{2} RT \right) \right]$$

C. The electron gain enthalpy ($\Delta_{eg} H^\ominus$) and EA are related to each other by the equation:

$$\Delta_{eg} H^\ominus \quad (\text{ionisation enthalpy}) \\ = \left[- IE \left(\text{electron affinity} - \frac{5}{2} RT \right) \right]$$

D. The value of C_p (heat capacity at constant pressure and C_v (heat capacity at constant volume) are $\frac{3}{2}R$ and $\frac{5}{2}R$ respectively)

Answer: A::B::C

 [Watch Video Solution](#)

Exercises (Multiple Correct) General Electronic Configuration And Periodicity

1. Which of the following statement (*s*) is/are correct ?

- A. Mendeleev's periodic law was based on atomic number of the element.
- B. Zero group was not present in the periodic table when Mendeleev presented it.
- C. The effective nuclear charge (Z_{eff}) is the atomic number minus shielding effect.
- D. There are four transition series in the periodic table each one consists of 10 elements.

Answer: B::C::D



Watch Video Solution

2. Which of the following statement (*s*) is/are correct ?

A. Similar electrons configuration is repeated

after intervals of 2, 8, 8,

18 and 32 in the extended form of periodic

table.

B. In a period from right ot left (\leftarrow), reducing

nature increases.

C. Along the period (\rightarrow) the number of valency electrons increases from 1 to 8.

D. The first member of lanthanide series is lanthanum.

Answer: A::B::C::D



Watch Video Solution

3. Which of the following statement (*s*) is/are correct ?

A. All the members of the actinide series are man made.

B. Zero group elements are paramagnetic.

C. Third group of the periodic table accommodates maximum number of elements.

D. All elements of zero groups are non-metals.

Answer: C::D



Watch Video Solution

4. Which of the following statement (s) is/are correct ?

- A. In Mendeleev periodic table, all groups are divided into two subgroups.
- B. There is no relationship between electronic configuration of the elements and their position in the extended form of periodic table.
- C. *s*-block elements have one or two electrons in their outermost shell.
- D. Osmium has the maximum density among metals.

Answer: C::D



Watch Video Solution

5. Which of the following statement (*s*) is/are correct ?

A. The last member of the *7th* period of the periodic table will have atomic number 118 if observed.

B. All the transition elements are metals and paramagnetic .

C. The maximum number of elements are present in the *5th* period of the periodic table.

D. Every period of the periodic table starts with a member of alkali group.



Answer: A::B



Watch Video Solution

Exercises (Multiple Correct) Atomic And Ionic Radii

1. Which of the following is correct in order of increasing size ?





D. All of these

Answer: A::C



Watch Video Solution

2. Reason for diagonal relationship is

A. Same size

B. Same electronegativity

C. Same electron affinity

D. Same polarisability

Answer: A::B::D



Watch Video Solution

3. Which of the following pairs of elements have almost similar atomic radii ?

A. *Zr, Hf*

B. *Mo, W*

C. *Co, Ni*

D. *Nb, Ta*

Answer: A::B::C::D





4. Which of the following statements is / *are* correct

?

A. An anion is larger than a cation if they are

isoelectronic.

B. Out of Na^{\oplus} and Al^{3+} , Na^{\oplus} has the largest

size.

C. The ionic radii of trivalent lanthanides

$[La^{3+}, Ce^{3+}, Pr^{3+}, \dots]$ decreases with

increasing atomic number.

D. Out of P^{3-} , S^{2-} , and Cl^{\ominus} , Cl^{\ominus} ion has the largest size.

Answer: A::B::C



Watch Video Solution

Multiple Correct Answer type

1. Which of the following statements is / *are* correct ?

A. Z_{eff} of elements increases along the period
(\rightarrow).

B. Z_{eff} of elements increases down the group

(↓).

C. Isoelectronic species have the same nuclear charge.

D. Screening constant (σ) increases down the group (↓).

Answer: A::D



Watch Video Solution

Exercises (Multiple Correct) Ionisation Energy (Ie)

1. Which of the following statements is /are correct ?

A. Successive IE 's are lower.

B. Less energy is required to remove an electron from a

half-filled shell or completely filled shell.

C. IE_1 of $Al < IE_1$ of Mg

D. IE_1 of $Be > IE_1$ of B

Answer: C::D



Watch Video Solution

2. Explain what do you understand by screening effect. How does this effect influence the values of ionisation energies ?

A. Size of the atom

B. Charge on the nucleus

C. The inner electrons which effectively screen the nuclear charge



D. Atomic number of the element

Answer: A::B::C



Watch Video Solution

3. Which is the correct increasing order of ionisation energy ?



Answer: A::D



Watch Video Solution

4. Which is correct about ionisation potential ?

A. IE_1 of $N > IE_1$ of O

B. IE_2 of $N > IE_2$ of O

C. IE_2 of $Li > IE_2$ of Ne

D. IE_1 of $Al > IE_1$ of Ca

Answer: A::C



Watch Video Solution

5. The first eight ionisation energies for a particular neutral atom is as given below. All values are expressed in $kJmol^{-1}$. Which oxidation states (s) is

/are not possible of the atom ?

1st	2nd	3rd	4th	5th	6th	7th	8th
1.31	3.39	5.30	7.47	10.99	13.33	71.33	84.01

A. - 2

B. - 3

C. - 6

D. 6

Answer: B::C::D



Watch Video Solution

6. IE_2 for an element is invariably higher than IE_1

because


- A. The size of cations is smaller than its atom.
- B. It is difficult to remove electron from cations.
- C. Ionisation energy is endothermic.
- D. All of the above

Answer: A::B



Watch Video Solution

7. Which sequence is correct regarding the first ionisation potential of coinage metals ?



B. $Cu < Ag < Au$

C. $Cu > Ag < Au$

D. $Ag > Cu > Au$

Answer: C



Watch Video Solution

8. Mark the correct statements out of the following:

A. He has highest IE_1 in the periodic table.

B. Cl has the highest EA out of all elements in the periodic table.

C. Hg and Br are liquid at room temperature.

D. In any period, the atomic radius of the noble gas is lowest.

Answer: A::B::C

 [Watch Video Solution](#)

9. Ionization energy is influenced by :

A. Size of the atom

B. Charge on the nucleus

C. Electrons present in inner shells

D. None of the above

Answer: A::B::C



Watch Video Solution

10. Which of the following statement (*s*) is/*are* correct ?

A. The element which has higher EA acts as a strong oxidising agent.

B. The element with higher EA has higher IE .





C. Along a period halogen has maximum EA .

D. The decreasing order of EA is $F > Cl > Br$.

Answer: A::B::C

 [Watch Video Solution](#)

11. Which of the following process do not involve absorption of energy?






Answer: A::B::C



Watch Video Solution

Exercises (Multiple Correct) Electronegativity (En)

1. Born-Haber cycle cannot be used to estimate



Answer: B::C



Watch Video Solution

2. Select the correct statement (*s*).

A. On Mulliken scale, the average of *IP* and *EA*

(in eV_{atom}^{-1}) is known as *EN*.

B. The maximum *EN* is shown by *Cl*.

C. *H*, *P* and *Te* have similar value of *EN*.

D. *H*, *S* and *Te* have similar value of *EN*.

Answer: A::C

3. Select the correct statement (s).

A. Mulliken's values of EN are about 2.8 times more than the Pauling scale.

B. Mulliken's value of EN are about 2.8 times less than the Pauling scale.

C. On Mulliken's scale if IP and EA are in

$$kJmol^{-1}, \text{ then } EN = \frac{IP + EN}{540}$$

D. On Mulliken's scale if IP and EA are in kcal

$$mol^{-1}, \text{ then } EN = \frac{IP + EA}{2 \times 62.5}$$

Answer: A::C::D

 Watch Video Solution

4. Select the correct statement (*s*)

A. On Pauling scale, the difference in EN of two atoms A and B in SI units is.

$$(EN_A - EN_B) = 0.1017\sqrt{\Delta_{A-B}}$$

B. On Pauling scale, the difference in EN of two atoms A and B in $kcal\ mol^{-1}$ is.

$$(EN_A - EN_B) = 0.208\sqrt{\Delta_{A-B}}$$

C. The Mulliken's EN values are scaled down to match the Pauling

value by dividing $\left(\frac{IP + EA}{2}\right)$ in eV by 2.8.

D. The Maulliken's EN values are scaled down to match the Pauling

value by multiplying $\left(\frac{IP + EA}{2}\right)$ in eV by 3.17

Answer: A::B::C



Watch Video Solution

5. Select the correct statement (s).

A. EN of Ga and $Ge > EN$ of Al and Si , due to d -block contraction.

B. EN of Ga and $Ge < EN$ of Al and Si , due to d -block contraction.

C. EN of $Pb > EN$ of Tl and Bi , due to d -block contraction

D. EN of $Pb < EN$ of Tl and Bi , due to d -block contraction

Answer: A::C



Watch Video Solution

Exercises (Multiple Correct) Miscellaneous

1. Transition metals are characterised by which of the following properties ?

- A. Variable valency
- B. Coloured compounds
- C. High melting and boiling points
- D. Tendency to form complexes

Answer: A::B::C::D



Watch Video Solution

2. Which is correct statement regarding BOH (where χ is electronegativity) ?

- A. If $\chi_O - \chi_B > \chi_O - \chi_H$ will be basic.
- B. If $\chi_O - \chi_B < \chi_O - \chi_H$ will be acidic
- C. If $\chi_O - \chi_B > \chi_O - \chi_H$ will be acidic.
- D. If $\chi_O - \chi_B < \chi_O - \chi_H$ will be basic

Answer: A::B



Watch Video Solution

3. The elements which are radioactive and have been named after the name of planet are

A. *Hg* (Mercury) (Hergentium)

B. *Np* (Neptunium)

C. *Pu* (Plutonium)

D. *Ra*(Radium)

Answer: B::C



Watch Video Solution

4. The properties which are common to the elements belonging to group 1 and 17 of periodic tables are:

A. Electropositive character increases down the group.

B. Reactivity decreases from a top to bottom

C. Atomic radii increases as atomic number increases

D. Electronegativity decreases on moving down a group.

Answer: A::C::D



 Watch Video Solution

5. The number of which subatomic particle is same in case of chlorine atom and chloride ion?




- A. Electron
- B. Proton
- C. Neutrons
- D. All of the above

Answer: B::C



Watch Video Solution

6. Which of the following show amphoteric behaviour?





Answer: A::B::C::D



Watch Video Solution

7. Which is correct in increasing order of ionic character ?



D. None of the above

Answer: A::B



Watch Video Solution

8. Highly pure dilute solution of sodium in ammonia

A. Shows blue colouration due to solvated electrons

B. Shows electrical conductivity due to both solvated electrons and solvated sodium ions

C. Shows red colouration due to solvated electrons but a bad conductor of electricity


D. Produces hydrogen gas or carbonate

Answer: A::B

 [Watch Video Solution](#)

9. Which of the following are ionic carbides ?

A. CaC_2



Answer: A::B::D



Watch Video Solution

10. Which of the following substance (s) is/are used in laboratory for drying purposes ?



B. Graphite

C. Anhydrous $CaCl_2$

D. Na_3PO_4

Answer: A::C



Watch Video Solution

11. Which of the following properties can be determined by using Born-Haber cycle ?

A. Electronegativity

B. Hydration energy

C. Lattice energy of ionic crystals



D. Binding energy of electrons

Answer: A::B::D



Watch Video Solution

12. The compound (*s*) which have $-O-O-$ bonds
(*s*) is / are





Answer: A::B::C



Watch Video Solution

13. Which of the following compounds are paramagnetic in nature ?



Answer: A::D





Watch Video Solution

14. Select the correct statement (*s*).

- A. Cr^{2+} compounds are ionic.
- B. They are oxidised to Cr^{3+} by air.
- C. They are reducing agent in aqueous solution.
- D. None is correct.

Answer: A::B::C



Watch Video Solution

15. On moving down the group from F to I , which of the following properties decreases ?

A. Ionic radius

B. IE

C. Oxidising power

D. EN

Answer: B::C::D



Watch Video Solution

16. Select the correct statement (s)

A. Alkali metals have lowest IE in respective period.

B. Noble gas have highest IE is respective period.

C. EA_1 of $N < EA_1$ of O .

D. F^\ominus is the strongest reducing agent among halide ions.

Answer: A::B::C



Watch Video Solution

17. The electronic configuration of given species (X)

is $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5, 4s^1$. This can be its

A. Cationic form X^{\oplus}

B. Anionic form X^{\ominus}

C. Excited state





D. Ground state

Answer: A::D



Watch Video Solution

18. Which of the following sets contain only isoelectronic ions?





Answer: A::C::D



Watch Video Solution

19. In which of the following arrangements, the order is according to the property indicated against it?



Answer: B::C::D



Watch Video Solution

20. In which of the following arrangements, the order is according to the property indicated against it?

A. Basic strength: $SbH_3 > AsH_3 > PH_3 > NH_3$

B. $IE_1 : N > O > C > B$.

C. Oxidising power :

$PbO_2 > SnO_2 > SiO_2 > CO_2$

D. Acid strength: $HI > HCl > HF$

Answer: B::C::D



Watch Video Solution

21. The bond dissociation energy of B-F in BF_3 is 646 kJ mol^{-1} whereas that of C-F in CF_4 is 515 kJ mol^{-1} .

The correct reason for higher B-F bond dissociation energy as compared to that of C-F bond is

A. Lower degree of $p\pi - p\pi$ interaction between

B and F in BF_3 than that between C and F in

CF_4

B. Significant $p\pi - p\pi$ interaction between B and

F in BF_3 whereas there is no possibility of

such interaction between C and F in CF_4

C. Smaller size of B -atoms as compared to that of C -atom and the stronger bond between B and F in BF_3 as compared to that between C and CF_4

D. Lewis acid character of boron trihalides is as:



Answer: B:

 [Watch Video Solution](#)

1. According to the periodic law of elements, the variation in properties of elements is related to their :

atomic masses

nuclear masses

atomic numbers

nuclear neutron-proton number ratios.

A. Atomic mass

B. Atomic number

C. Density

D. Mass number

Answer: B



Watch Video Solution

2. Newland's law of octave applies to which of the following set elements ?

A. *Be, Mg, Ca*

B. *As, K, Ca*

C. *B, N, C*

D. None of these

Answer: A





Watch Video Solution

3. The element whose electronic configuration is $1s^2, 2s^2 2p^6, 3s^2$ is a / an

- A. Metal
- B. Metalloid
- C. Inert gas
- D. Non-metal

Answer: A



Watch Video Solution

4. The number of periods and groups in the long form of periodic table are

A. 7 and 9

B. 8 and 18

C. 7 and 18

D. 6 and 10

Answer: C



Watch Video Solution

5. The elements of group 1, 2, 13, 14, 15, 16, 17, 18 are collectively called

- A. Noble elements
- B. Typical elements
- C. Transition elements
- D. Representative elements

Answer: D



Watch Video Solution

6. The statement that is false for the long form of the periodic table is :

1. it reflects the sequence of filling of electrons in the order of sub energy levels s, p, d, f

2. it helps to predict the stable valency states of the elements

3. it reflects trends in physical and chemical properties of the elements

4. it helps to predict the relative ionocity of the bond between any two elements.

A. It reflects the sequence of filling the electrons

in the order of sub-energy levels s, p, d and f .

B. It helps to predict the stable valency states of the elements.

C. It reflects trends in physical and chemical properties of the elements.

D. It helps to predict the relative ionic character of the bond between any two elements

Answer: D



Watch Video Solution

7. In the periodic table, going down in fluorine group

- A. Reactivity will increase
- B. Electronegativity will increase
- C. Ionic radius will increase
- D. Ionisation potential will increase

Answer: C



Watch Video Solution

8. In the long form of periodic table, the non-metals are placed in

- A. s-block

B. p -block

C. d -block

D. f -block

Answer: B



Watch Video Solution

9. Alkali metals in each period have

A. Smallest size

B. Highest EN

C. Lowest IE

D. Highest IE

Answer: C

 Watch Video Solution

10. Which one pair of atoms or ions will have same configuration ?

A. F^{\oplus} and Ne

B. Li^{\oplus} and He^{\ominus}

C. Na and K

D. Cl^{\ominus} and Ar

Answer: D



Watch Video Solution

11. In the modern periodic table, the period indicates the value of

- A. Atomic number
- B. Atomic mass
- C. Principal quantum number
- D. Azimuthal quantum number

Answer: C





Watch Video Solution

12. Which of the following does not reflect periodicity of elements ?

A. Bonding behaviour

B. EN

C. IE

D. Neutron / proton ratio

Answer: D



Watch Video Solution

13. What is the number of elements in the 3rd period of the modern periodic table ?

A. 8 elements

B. 32 elements

C. 3 elements

D. 18 elements

Answer: A



Watch Video Solution

14. Which of the following set have almost similar atomic radii ?

A. *Hf, Zr*

B. *K, Rb*

C. *Be, Al*

D. *B, Al*

Answer: C



Watch Video Solution

15. Which of the following belongs to the category of transition metal?

A. *K*

B. *Ra*

C. *Fe*

D. All of the above

Answer: C



Watch Video Solution

16. Without looking at the periodic table, select the elements belonging to same from the following list.

A. $Z = 12, 38, 4, 88$

B. $Z = 9, 16, 3, 35$

C. $Z = 5, 11, 27, 19$

D. $Z = 24, 47, 42, 55$

Answer: A



Watch Video Solution

17. The elements of same group of the periodic table have

- A. Same number of protons
- B. Same valence shell
- C. Same valence electrons
- D. Same electrons affinity

Answer: C



Watch Video Solution

18. The elements which are characterised by the outer shell configuration ns^1 to np^6 are collectively called

- A. Transition elements
- B. Representative elements
- C. Lanthanides
- D. Inner transition elements

Answer: B



Watch Video Solution

19. General electronic configuration of transition metals is

A. ns^2nd^{1-10}

B. $(n - 1)d^{1-10}ns^{1-2}$

C. $(n - 1)d^2ns^{0-2}$

D. $(n - 1)d^{1-10}ns^2$

Answer: B



Watch Video Solution

20. An element with atomic number 20 will be placed in which period of the periodic table

A. 4

B. 3

C. 2

D. 1

Answer: A



Watch Video Solution

21. The statement that is not correct for periodic classification of elements is

A. The properties of elements are the periodic functions of their atomic numbers.

B. Non-metallic elements are less in number than metallic elements.

C. The first ionisation energies of elements along a period do not vary in a regular manner with an increase in atomic number.

D. For transition elements the ionisation energies increase gradually with increase in atomic

number.

Answer: D



Watch Video Solution

22. In the modern periodic table, elements are arranged in

- A. Increasing mass
- B. Increasing volume
- C. Increasing atomic number
- D. Alphabetically

Answer: C



Watch Video Solution

23. Name of the heaviest naturally occurring element.

A. *U*

B. *Ra*

C. *Pb*

D. *Hg*

Answer: A





Watch Video Solution

24. The screening effect of d -electron is



- A. Equal to p -electron
- B. Much more than p -electron
- C. Same as f -electrons
- D. Less than p -electrons

Answer: D



Watch Video Solution

25. Which of the following represents the electronic configuration of the electropositive element ?




Answer: B



Watch Video Solution

1. Which of the following has the largest ionic radius ?



Answer: D



Watch Video Solution

2. The size of species I , I^+ and I^\ominus decrease in the order.

A. $I^\oplus > I^\ominus > I$

B. $I^\ominus > I > I^\oplus$

C. $I^\ominus > I^\oplus > I$



D. $I > I^\oplus > I^\ominus$

Answer: B



Watch Video Solution

3. Which one of the following is the smallest in size?




Answer: A



Watch Video Solution

4. Which of the following represent increasing order of size of 4th period element ?



B. *Kr, Br, Ca, K*

C. *K, Ca, Br, Kr*

D. *Br, Kr, Ca, K*

Answer: D



Watch Video Solution

5. Which of the following van der Waals radii is the largest ?

A. *Ne*

B. *Cl*

C. O

D. F

Answer: B



Watch Video Solution

6. The correct order of the size of C , N , P and S is

A. $N < C < P < S$

B. $C < N < P < S$

C. $N < C < S < P$

D. $C < N < S < P$

Answer: C



Watch Video Solution

7. The correct order of the size of Be , C , F and Ne is

A. $Be > C > F > Ne$

B. $Be < C < F < Ne$

C. $F < C < S < P$




D. $Be > C > F < Ne$

Answer: D



Watch Video Solution

8. The correct order of increasing radii are



Answer: A



Watch Video Solution

9. The correct arrangement of decreasing order of atomic radius among Na , K , Mg and Rb is

A. $Rb > Na > N > Mg$

B. $K > Rb > Na > Mg$

C. $Rb > K > Na > Mg$

D. $Mg > Rb > K > Na$

Answer: C



Watch Video Solution

10. Which of the following pairs of elements have almost similar atomic radii ?

A. *Zr, Hf*

B. *Cu, Ag*

C. *Sc, Ti*

D. *Pd, Pt*

Answer: A



Watch Video Solution

11. The radius of isoelectronic species

- A. Increases with increase in nuclear charge
- B. Decreases with increases in nuclear charge
- C. Same for all
- D. First increases and then decreases

Answer: B



Watch Video Solution

12. Atomic radii of fluorine and neon in Angstrom units are respectively given by

- A. 0.72, 1.60

B. 1.60, 1.60

C. 0.72, 0.72

D. 1.60, 0.72

Answer: A

 [Watch Video Solution](#)

13. Anything that influences the valence electrons will affect the chemistry of the element. Which one of the following factors does not affect the valence shell ?

A. Valence principal quantum number (n)

B. Nuclear charge (Z)

C. Nuclear mass

D. Number of core electrons

Answer: C

 [Watch Video Solution](#)

14. The size of isoelectronic species F^\ominus , Ne , and Na^\oplus is affected by

A. Nuclear charge (Z)

B. Valence principal quantum number (n)

C. Electron-electron interaction in the outer orbitals


D. None of the factors because their size the same



Answer: C



Watch Video Solution

15. Ionic radii of





Answer: D



Watch Video Solution

Exercises (Single Correct) Effective Nuclear Charge (Z_{Eff})

1. Which of the following statement is most correct ?

Effective nuclear charge of atom depends on

A. The charge on the ion

B. The atomic number of an atom

C. The screening effect

D. Both (a) and (c)

Answer: D



Watch Video Solution

2. The screening effect of d-electrons is

A. More than s -electrons

B. More than p -electrons

C. Less than s -electrons

D. Same as f -electrons

Answer: C

 [Watch Video Solution](#)

3. The chemistry of lithium is very similar to that to that of magnesium even though they are placed in different groups. Its reason is

- A. Both are found together in nature
- B. Both have nearly the same size
- C. Both have similar electronic configuration

D. The ratio of their charge and size (i.e. charge density) is nearly the same

Answer: D



Watch Video Solution

4. In a given energy level, the order of penetration effect of different orbitals is

A. $f < d < p < s$

B. $s = p = d = f$



C. $s < p < d < f$

$$D. p > s > d > f$$

Answer: A

 **Watch Video Solution**

5. Which one of the following group of atoms or ions is not isoelectronic ?



Answer: D



Watch Video Solution

6. The correct order of relative stability of half filled and completely filled sub-shell is

A. $p^3 > d^5 < d^{10} < p^6$

B. $d^5 > p^3 < d^{10} < p^6$

C. $d^5 > p^3 > d^{10} > p^6$

D. $p^3 < d^{10} < d^5 < p^6$

Answer: C









Watch Video Solution

Exercises (Single Correct) Ionisation Energy (Ie)

1. From the ground state electronic configuration of the elements given below, pick up the one with the highest value of second ionisation energies







Answer: B



Watch Video Solution

2. Which of the following process refers to IE_2 ?



Answer: B



Watch Video Solution

3. Which of the following statement concerning ionisation energy is not correct ?

A. The IE_2 is always more than the first.

B. Within a group, there is a gradual increase in ionisation energy because nuclear charge increases.

C. Ionisation energies of Be is more than B .


D. Ionisation energies of noble gases are high.

Answer: B



Watch Video Solution

4. The graph of IE_1 or $\Delta_1 H_1^\ominus$ versus atomic number (Z) is given below:



Which of the following statement is correct ?

A. Alkali metals are at the maxima and noble gases at the minima.

B. Noble gases are at the maxima and alkali metals at the minima.

C. Transition elements are at the maxima.



D. Minima and maxima do not show any regular behaviour.

Answer: B



Watch Video Solution

5. Which of the following isoelectronic ions have the lowest ionisation enthalpy ?



Answer: D



Watch Video Solution

6. The second ionisation potential is

A. less than First IE

B. higher than 1st IE

C. same




D. depends on the elements

Answer: B



Watch Video Solution

7. Which of the following process requires the largest amount of energy ?







D. All require same amount of energy

Answer: B

 **Watch Video Solution**

8. Which of the following is an energy consuming process ?



Answer: C



Watch Video Solution

9. Arrange S , P and As in order of increasing ionisation energy.

A. $S < P < As$

B. $P < S < As$

C. $As < S < P$

D. $As < P < S$

Answer: C





Watch Video Solution

10. The five successive ionization enthalpies of an element are 800, 2427, 3658, 25024 and 32824 KJ mol^{-1} . The number of valence electrons in the element is :

A. 3

B. 5

C. 1





D. 2

Answer: A



Watch Video Solution

11. Which of the following transitions involves maximum amount of energy?



Answer: D



12. Which of the elements show least values of ionisation within their periods ?

A. Alkaline earth metals

B. Alkali metals

C. Noble gases

D. Chalcogens

Answer: B



Watch Video Solution

13. Which of the following has the largest ionisation energy.

A. ${}_{11}\text{Na}$

B. ${}_{19}\text{K}$

C. ${}_{12}\text{Mg}$

D. ${}_{37}\text{Rb}$




Answer: C



Watch Video Solution

14. Which one of the following elements has the highest ionisation energy?

A. $[\text{Ne}]3s^23p^1$




Answer: B



Watch Video Solution

15. The correct order of second ionisation potentials of carbon, nitrogen, oxygen and fluoroine is



C. $O > F > N > C$

D. $F > O > N > C$

Answer: C



Watch Video Solution

16. Which has the largest first ionisation energy ?

A. Na

B. K

C. Rb

D. Li

Answer: D



Watch Video Solution

17. Which of the following element has the highest ionisation energy ?

A. Carbon

B. Boron

C. Oxygen

D. Nitrogen

Answer: D





Watch Video Solution

18. Ionisation enthalpy of nitrogen is more than oxygen because of

- A. Greater attraction of electrons by the nucleus
- B. Extra stability of the half-filled p -orbitals
- C. Smaller size of nitrogen
- D. More penetrating effect

Answer: B



Watch Video Solution

19. The set representing the correct order of the first ionisation potential is

A. $K > Na > Li$

B. $Be > Mg > Ca$

C. $B > C > N$

D. $Ge > Si > C$

Answer: B



Watch Video Solution

20. The first ionisation potential of which of the element is highest

A. *Na*

B. *Mg*

C. *Al*

D. *Si*

Answer: D



Watch Video Solution

21. Highest ionisation potential in a period is shown by

- A. Alkali metals
- B. Transition elements
- C. Halogens
- D. Alkaline earth metals

Answer: C



Watch Video Solution

22. The first ionisation energy is maximum for

A. Na

B. Mg

C. K

D. Kr

Answer: D



Watch Video Solution

23. Which sequence is correct regarding the first ionisation potential of coinage metals ?

A. $Cu > Ag < Au$

B. $Cu > Ag > Au$

C. $Cu < Ag < Au$

D. $Au > Ag < Cu$

Answer: A

 [Watch Video Solution](#)

24. The second ionisation potentials in electron volts of oxygen and fluorine atoms are respectively given by

A. 35.1, 38.3

B. 38.3, 38.3

C. 38.3, 35.1

D. 35.1, 35.1

Answer: C



Watch Video Solution

25. The value of IE_1 , IE_2 , IE_3 and IE_4 of an atom are respectively $7.5eV$, $25.6eV$, $48.6eV$ and $170.6eV$.

The electronic configuration of the atom will be

A. $1s^2, 2s^2 2p^6, 3s^1$

B. $1s^2 2s^2 2p^6 3s^2 3p^1$

C. $1s^2, 2s^2 2p^6 3s^2 3p^3$

D. $1s^2, 2s^2 2p^6 3s^2$

Answer: B



Watch Video Solution

26. IE_1, IE_2 and IE_3 values are 100, 150 and $1500eV$ respectively. The element can be

A. Na

B. B

C. Be

D. F

Answer: C



Watch Video Solution

27. $N_0/2$ atoms of $X_{(g)}$ are converted into $X_{(g)}^{\oplus}$ by energy E_1 , $N_0/2$ atoms of $X_{(g)}$ are converted into $X_{(g)}^{\ominus}$ by energy E_2 . Hence ionisation potential and electron affinity of $X_{(g)}$ per atom are

A. $\frac{2E_1}{N_0}, \frac{2(E_2 - E_1)}{N_0}$

B. $\frac{2E_1}{N_0}, \frac{2E_2}{N_0}$

C. $\frac{(E_1 - E_2)}{N_0}, \frac{2E_2}{N_0}$

D. None is correct.

Answer: A



Watch Video Solution

28. Which of the following ionisation energy values for calcium show a sudden increase?

A. Third

B. Second

C. First

D. Fourth

Answer: A

 [Watch Video Solution](#)

29. Which one of the following statements is incorrect in relation to ionization enthalpy?

A. Ionisation enthalpy increases for each successive electron.

B. The greatest increase in ionisation enthalpy is experienced on removal of electron from core

noble gas configuration.

C. End of valence electrons is marked by a big jump in ionisation enthalpy.

D. Removal of electron from orbitals bearing lower n value is easier than from orbitals having higher than n value.

Answer: D

 [Watch Video Solution](#)

30. IE_2 for an element is invariably higher than IE_1 because

- A. It is difficult to remove electron from cation
- B. The size of the cation is smaller than its atoms
- C. Z_{eff} is more for cation
- D. All the above

Answer: D



Watch Video Solution

31. Which of the following metal requires radiation of the lowest wavelength to cause emission of electrons?

A. Na

B. K

C. Mg

D. Ca


Answer: C




Watch Video Solution

Exercises (Single Correct) Isoelectronic Species

1. Which of the following are isoelectronic?








D. None of these

Answer: B



Watch Video Solution

2. Consider the isoelectronic series, K^\oplus , S^{2-} , Cl^\ominus , Ca^{2+} , the radii of the ions decrease as



Answer: C



Watch Video Solution

3. Which of the following are isoelectronic species ?



A. II, III and IV

B. I, II and III

C. I, II and IV

D. II and I

Answer: A



Watch Video Solution

4. Two p -block elements x (outer configuration ns^2np^3) and z (outer configuration ns^2np^4) occupy neighbouring positions in a period. Using this information which of the following is correct with respect to their ionisation potential I_x and I_z .

A. $I_x > I_z$

B. $I_z > I_x$

C. $I_z = I_x$

D. Relation between I_x and I_x is uncertain

Answer: A



Watch Video Solution

Exercises (Single Correct) Matallic-Non Metallic Character

1. Considering the elements B , Al , Mg and K , the correct order of their metallic character is

A. $B > Al > Mg > K$

B. $Al > Mg > B > K$

C. $Mg > Al > K > B$

D. $K > Mg > Al > B$


Answer: D



Watch Video Solution

2. IE_1 and IE_2 of Mg are 178 and 348kcalmol^{-1} .

The energy required for the reaction



A. $+170\text{kcal}$

B. $+526\text{kcal}$

C. -170kcal

D. -526kcal

Answer: B



Watch Video Solution

3. Considering the elements B , C , N and Si , the correct order of their non-metallic character is ?

A. $B > C > Si > N > F$

B. $Si > C > B > N > F$

C. $F > N > C > B > Si$

D. $F > N > C > Si > B$

Answer: C



Watch Video Solution

Exercises (Single Correct) Electron Affinity (Ea) And Electron Gain Enthalpy ($\Delta_{Eg}H^{\ominus}$)

1. Which of the following will have lowest electron affinity?

- A. Oxygen
- B. Fluorine
- C. Nitrogen
- D. Carbon

Answer: C



Watch Video Solution

2. Second and successive electron affinity of an element

A. is always successive (energy is released)

B. is always positive (energy is absorbed)

C. can be positive or negative

D. is always zero

Answer: B



Watch Video Solution

3. Which one of the following statements is incorrect?

A. Greater is the nuclear, greater is the electron affinity.

B. Neon has zero electron affinity.

C. Electron affinity decreases from fluorine to iodine in the group

D. Electron affinity decreases in going down a group and increases across from the left to the right (\rightarrow).

Answer: C



Watch Video Solution

4. The lower electron affinity of fluorine than that of chlorine is due to

- A. Smaller size
- B. Smaller nuclear charge
- C. Difference in their electronic arrangement
- D. Its highest reactivity

Answer: A





Watch Video Solution

5. The EA order for halogen is

A. $F > Cl > Br > I$

B. $F < Cl < Br < I$

C. $F < Cl > Br > I$

D. $F > Cl < Br > I$

Answer: C



Watch Video Solution

6. The EA for inert gases is likely to be

- A. High
- B. Small
- C. Zero
- D. Positive

Answer: C



Watch Video Solution

7. Ionisation of energy F^\ominus is $320kJmol^{-1}$. The electronic gain enthalpy of fluorine would be

A. -320kJmol^{-1}

B. -160kJmol^{-1}

C. $+320\text{kJmol}^{-1}$

D. $+160\text{kJmol}^{-1}$

Answer: A



Watch Video Solution

8. Which of the following represents the correct order of electron affinities?

A. $F > Cl > Br > I$

B. $C < N < O < F$

C. $N < C < O < F$

D. $C < Si < P < N$

Answer: C



Watch Video Solution

9. Fluorine has the highest electronegativity among the group on the Pauling scale, but the electron affinity of fluorine is less than that of chlorine because

A. The atomic number of fluorine is less than that of chlorine

B. Fluorine being the first member of the family behaves in an unusual manner

C. Chlorine can accommodate an electron better than fluorine by utilising its vacant $3d$ orbital

D. Small size, high electron density and an increase in electron repulsion make addition of an electron to fluorine less favourable than that in the case of chlorine





Answer: D



Watch Video Solution

Exercises (Single Correct) Electronegativity (En)

1. Which is true about the electronegative order of the following elements ?



Answer: A

 [Watch Video Solution](#)

2. The electronegativity of the following elements increases in the order

A. *C, N, Si, P*

B. *N, Si, C, P*

C. *Si, P, C, N*

D. *P, Si, N, C*

Answer: C

 [Watch Video Solution](#)

3. An atom with high EA generally has

A. Tendency to form $+ve$ ions

B. High ionisation energy

C. Large atomic size

D. Low electron affinity

Answer: B



Watch Video Solution

4. The electronegativity of the following elements increases in the order

A. $S < P < N < O$

B. $P < S < N < O$

C. $N < O < P < S$

D. $N < P < S < O$

Answer: B



Watch Video Solution

5. Downwards in a group, the electropositive character of elements

A. Increases

B. decreases

C. Remain same




D. None

Answer: A



Watch Video Solution

6. What is the correct order of electronegativity?



$$D. M^{4-} < M^{2-} < M^{3-} < M^{1-}$$

Answer: B



Watch Video Solution

7. Due to screening effect of electrons in an atom

A. IE decreases

B. IE increases

C. No change in IE

D. Attraction of nucleus on the valence electron
increases

Answer: A



Watch Video Solution

8. Select the group where EN increases down the group

A. F, Cl, Br

B. Li, Na, K

C. Ca, Sr, Ba

D. Zn, Cd, Hg

Answer: D





Watch Video Solution

9. Which of the following element has the highest EN ?

A. As

B. Sb

C. P

D. S

Answer: D



Watch Video Solution

10. In C , N , O and F the electronegativity

A. Increases from carbon to fluorine

B. Decreases from carbon to fluorine

C. Increases up to oxygen and is minimum at
fluorine

D. Is minimum at nitrogen and then increases
continuously



Answer: A



Watch Video Solution

Exercises (Single Correct) Acidic, Basic and Amphoteric Character

1. Among the following oxides, which is least acidic ?





Answer: A



Watch Video Solution

2. Which of the following oxides is most basic ?



Answer: A



Watch Video Solution

3. The order in which the following oxides are arranged according to decreasing basic nature is

A. Na_2O , MgO , Al_2O_3 , CuO

B. CuO , Al_2O_3 , MgO , Na_2O

C. Al_2O_3 , CuO , MgO , Na_2O

D. CuO , MgO , Na_2 , Al_2O_3

Answer: A






Watch Video Solution

4. Identify the correct order of acidic strength of

CO_2 , CuO , CaO and H_2O .

A. $CaO < CuO < H_2O < CO_2$



Answer: B



Watch Video Solution

5. What is the nature of Al_2O_3 and B_2O_3 ?

A. Acidic, acidic

B. Acidic, amphoteric

C. Amphoteric, amphoteric

D. Amphoteric, acidic

Answer: D



Watch Video Solution

6. which one of the following oxides is neutral?

A. SiO_2

B. CO

C. ZnO

D. SnO_2

Answer: B



Watch Video Solution

7. Which of the following oxides is amphoteric in nature ?







Answer: C



Watch Video Solution

8. The correct order of acidic strength of the following is



Answer: A



Watch Video Solution

9. Which of the oxides behave both as neutral oxide and suboxide ?

(a) N_2O , b. NO , c. C_3O_2 , d. CO

A. CO

B. CO_2

C. C_3O_2




D. N_2O

Answer: D



Watch Video Solution

10. Which of the oxides is not a mixed oxide ?



Answer: D



Watch Video Solution

11. Which of the oxides is basic as well as normal oxide ?

A. N_2O

B. Na_2O

C. NO

D. H_2O


Answer: B



Watch Video Solution

Exercises (Single Correct) Bond Angle

1. Decreasing order of bond angle of (NH_3, PH_3, AsH_3) is



Answer: A



Watch Video Solution

2. The correct order of bond angle of NO_2^+ , NO_2 and NO_2^- is









Answer: B

 **Watch Video Solution**

3. Decreasing order of bond angle of

$\left(NH_3, NH_4^{\oplus}, PCl_5, SCl_2 \right)$ is









Answer: C

 **Watch Video Solution**

4. Increasing order of bond angle of $(Cl_2O, ClO_2, Cl_2O_7, I_3^\ominus)$ is








Answer: D

 [Watch Video Solution](#)

Exercises (Single Correct) Lattice And Hydration Energy

1. Decreasing order of hydration energy of the following is









Answer: A

 **Watch Video Solution**

2. Extent of hydrolysis of the following is







Answer: D



Watch Video Solution

3. Give the decreasing order of thermal stability of the following .



Answer: B



Watch Video Solution

4. Lattice energy of an ionic compound depends on

A. Change density of the ions only

B. Packing of ions only

C. Size of the ion only

D. Charge on the ions only

Answer: A



Watch Video Solution

5. Na_2SO_4 is soluble in water while $BaSO_4$ is insoluble. Which of the reason is correct about the

above statement.

A. Lattice energy of $BaSO_4$ exceeds its hydration energy.

B. Hydration energy of $BaSO_4$ exceeds its lattice energy.

C. The solubility in H_2O of a compound depends only on its hydration energy.

D. The solubility in H_2O of a compound depends only on its lattice energy.

Answer: A



Watch Video Solution

6. Calculate the lattice energy from the following data (given $1eV = 23.0kcalmol^{-1}$)

i. $\Delta_f H^\ominus (KI) = -78.0kcalmol^{-1}$

ii. IE_1 of $K = 4.0eV$

iii. $\Delta_{diss} H^\ominus (I_2) = 28.0kcalmol^{-1}$

iv. $\Delta_{sub} H^\ominus (K) = 20.0kcalmol^{-1}$

v. EA of $I = -70.0kcalmol^{-1}$

vi. $\Delta_{sub} H^\ominus$ of $I_2 = 14.0kcalmol^{-1}$

A. $+14.1kcalmol^{-1}$

B. $-14.1kcalmol^{-1}$

C. $-141kcalmol^{-1}$

$$D. +141\text{kcalmol}^{-1}$$

Answer: C



Watch Video Solution

7. Calculate the EA of O atom to O^{2-} ion from the following data:

i. $\Delta_f H^\ominus [MgO(s)] = -600\text{kJmol}^{-1}$

ii. $\Delta_u H^\ominus [MgO(s)] = -3860\text{kJmol}^{-1}$

iii. $IE_1 + IE_2$ of $Mg(g) = 2170\text{kJmol}^{-1}$

iv. $\Delta_{\text{diss}} H^\ominus$ of $Mg(s) = +494\text{kJmol}^{-1}$

v. $\Delta_{\text{sub}} H^\ominus$ of $Mg(s) = +150\text{kJmol}^{-1}$

A. $+693\text{kJmol}^{-1}$

B. -693kJmol^{-1}

C. $+69.3\text{kJmol}^{-1}$

D. -69.3kJmol^{-1}

Answer: A



Watch Video Solution

Exercises (Single Correct)Miscellaneous

1. Which of the following is incorrect ?

- A. With increase in atomic size, ionisation energy increases
- B. With increase in atomic size, electron affinity increases
- C. With increase in atomic size, metallic character increases
- D. With increase in atomic size, electronegativity increases

Answer: C



Watch Video Solution

2. Which of the following is incorrect?

A. An element which has high electronegativity always has high electron gain enthalpy.

B. Electron gain enthalpy is the property of an isolated atom.

C. Electronegativity is the property of bonded atom.

D. Both electronegativity and electron gain enthalpy are usually directly related to nuclear charge and inversely related to atomic size.

Answer: A



Watch Video Solution

3. The ionisation of hydrogen atom would give rise to

- A. Hydride ion
- B. Hydronium ion
- C. Proton
- D. Hydroxyl ion

Answer: C





Watch Video Solution

4. Chloride of an element A gives neutral solution in water. In the periodic table, the elements A belong to

- A. First group
- B. Third group
- C. Fifth group
- D. First transition group

Answer: A



Watch Video Solution

5. In a period, density first ___ then _____ and in a group it_ down the group

A. Decreases, increases, remains constant

B. Increases, decreases, decreases

C. Increases, remains constant , increases

D. Increases, decreases, increases

Answer: D



Watch Video Solution

6. In the transformation of $Na(s) \rightarrow Na^{\oplus}(g)$, the energies involved are

- A. Ionisation energy
- B. Sublimation energy
- C. Ionisation energy and sublimation energy
- D. Bond dissociation energy

Answer: C



Watch Video Solution

7. Beryllium and aluminium exhibit many properties which are similar . But, the two elements differ in





- A. Forming covalent halides
- B. Forming covalent hydrides
- C. Exhibiting maximum covalency in compounds
- D. Exhibiting amphoteric nature in their oxides

Answer: C



Watch Video Solution

8. Among $LiCl$, $BeCl_2$, BCl_3 and CCl_4 the covalent bond character varies as .







Answer: B



Watch Video Solution

9. The correct order of decreasing polarisability of ion is



Answer: C



Watch Video Solution

10. Diagonal relationship is shown by

A. All elements with their diagonally opposite elements.

B. All elements of *3rd* and *4th* periods

C. Some of the elements of *2nd* and *3rd* periods

D. Elements of d-block

Answer: C



Watch Video Solution

Exercises (Assertion Reasoning)

1. Assertion: Helium and beryllium have similar outer electronic configuration of the type ns^2 .

Reason: Both are chemically inert.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: C



Watch Video Solution

2. Assertion (A) : EA of O is less than that of F but greater than that of N .

Reason (R) : IE is as follows: $N > O > F$.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: C



Watch Video Solution

3. Assertion (A): IE_1 of N is lower than O .

Reason (R) : Across a period Z_{eff} decrease.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of

Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If both (A) and (R) are correct

Answer: C



Watch Video Solution

4. Assertion(A) : IE_2 of Li is the highest in the second period.

Reason (R) : Li^{\oplus} has noble gas i.e., Ne gas configuration.

- A. If both Assertion (A) and (R) are correct and Reason (R) is the correct explanation of Assertion (A).
- B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).
- C. If (A) is correct but (R) is not correct.
- D. If (A) and (R) are correct

Answer: C



Watch Video Solution

5. Assertion(A) : IE_1 of C is greater than that of B .

Reason (R) : Due to penetration effect the IE_1 of C

is greater than

that of B

A. If both Assertion (A) and (R) are correct and

Reason (R) is the

correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not

the correct explanation

for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) is incorrect and (R) is correct

Answer: D

 [Watch Video Solution](#)

6. Assertion (A) : In the modern periodic table period indicates the value of azimuthal quantum number.

Reason (R) : Each period begins with the filling of new shell.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the

correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) is incorrect and (R) is correct

Answer: D

 [Watch Video Solution](#)

7. Assertion (A) : Anything that influences of the valence electrons will affect the chemistry of the

element. So, the nuclear molar mass does not affect the valence shell.

Reason (R) : Nucleus contains protons and neutrons, whereas protons i.e. nuclear charge affects the valence shell but neutrons do not.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).


C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: A

 **Watch Video Solution**

8. Assertion (A) : Metallic character order is



Reason (R) : Along the period (\rightarrow) metallic character increases and decreases down the group (\downarrow).

A. If both Assertion (A) and (R) are correct and Reason (R) is the correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: C



Watch Video Solution

9. Assertion (A) : When the transition element ionises, the $4s$ -orbital electrons are removed before the $3d$ -orbital electrons.

Reason (R) : The energy of $3d$ -orbital electrons is lower than that of $4s$ -orbital electrons.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of Assertion (A).

B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: A



Watch Video Solution

10. Assertion (A) : Mercurous ion is paramagnetic.

Reason (R): It contains one unpaired electron.

A. If both Assertion (A) and (R) are correct and

Reason (R) is the correct explanation of

Assertion (A).

B. If both (A) and (R) are correct but (R) is not

the correct explanation for (A).

C. If (A) is correct but (R) is not correct.

D. If both (A) and (R) are incorrect



11. Assertion (A): Lithium chloride is predominantly covalent compound.

Reason (R): electronegativity difference between Li and Cl is small.

- A. If both Assertion (A) and (R) are correct and Reason (R) is the correct explanation of Assertion (A).
- B. If both (A) and (R) are correct but (R) is not the correct explanation for (A).
- C. If (A) is correct but (R) is not correct.

D. If (A) and (R) are correct

Answer: C

 [Watch Video Solution](#)

Exercises (Integer)

1. Among the following oxides how many of them are suboxides ?

(a) C_3O_2 , (b) N_2O , (c) NO_2 , (d) CO

(e) Fe_3O_4 , (f) KO_2

 [Watch Video Solution](#)

2. The number of factors that influence the IE are

a. Size of the atom , b. Charge on the nucleus

c. Shielding effect , d. The atomic mass



Watch Video Solution

3. The number of process (es) requiring the absorption of energy/are

a. $Cl \rightarrow Cl^\ominus$, b. $O^\ominus \rightarrow O^{2-}$

c. $Fe^{+3} \rightarrow Fe^{+2}$, d. $Ar \rightarrow Ar^\ominus$



Watch Video Solution

4. The five successive ionization enthalpies of an element are 800, 2427 , 3658 , 25024 abd 32824 KJ mol⁻¹ . The number of valence electrons in the elements is :

 [Watch Video Solution](#)

5. The diagonal relationship is shown by the elements upto how many groups only ?

 [Watch Video Solution](#)

6. The number of the following pairs contains elements with similar atomic radii

a. *Co, Ni* , b. *Zn, Mo* , c. *Rh, Ir* , d. *Hf, Ti*



[View Text Solution](#)

7. How many of the following energies are involved in the transformation of $Na(s) \rightarrow Na^{\oplus}(aq)$?

(a) $1E$, (b) $\Delta_{\text{sub}}H^{\ominus}$, (c) $\Delta_{\text{diss}}H^{\ominus}$, (d) $\Delta_{\text{hyd}}H^{\ominus}$

(e) $\Delta_u H^{\ominus}$



[View Text Solution](#)

8. Among the following oxides how many of them are mixed oxides ?

(a) H_2O , (b) C_3O_2 , (c) Fe_3O_4 , (d) Fe_2O_3

(e) Pb_3O_4 , (f) PbO_2 , (g) $Co(AlO_2)_2$, (h) Mn_3O_4



[View Text Solution](#)

9. Number of species that are isoelectronic with Ar is.



[Watch Video Solution](#)

10. Among the following oxides, how many of them are amphoteric oxides ?

(a) B_2O_3 , (b) Al_2O_3 , (c) CaO , (d) ZnO

(e) Ga_2O_3 , (f) SnO_2 , (g) PbO_2 , (h) BeO

(i) CuO , (j) Fe_2O_3



Watch Video Solution

11. Among the following elements how many of them are inner transition elements ?

(a) Sg , (b) Bk , (c) Er , (d) Fm

(e) Fe , (f) Pb , (g) Cr , (h) Ca

(i) Ar , (j) Zr , (k) Ce

 [Watch Video Solution](#)

12. How many number of pairs of elements exhibit diagonal relationship.

 [Watch Video Solution](#)

13. Among $BeCO_3$, $CaCO_3$, $SrCO_3$ and $BaCO_3$, how many number than of these compounds are thermally more stable than $MgCO_3$?

 [Watch Video Solution](#)

Exercises (Fill In The Blanks)

1. *IUPAC* name for the element with $Z = 117$ is _____ and its symbol is _____.



[Watch Video Solution](#)

2. In the long form of the periodic table, physical and chemical properties of the elements are a periodic function of their _____.



[Watch Video Solution](#)

3. The first periodic law stated by Mendeleev was

 [Watch Video Solution](#)

4. The elements which constitute $5f$ -block are called _____ with atomic numbers from _____ to _____.

 [Watch Video Solution](#)

5. The elements of group 1, 2, 13, 14, 15, 16, 17, 18 are collectively called

 [Watch Video Solution](#)

 Watch Video Solution

6. Fill in the blanks by picking the correct option.

There are ___ groups and ___ periods in the extended form of periodic table. The group, all members of which are in gaseous state under ordinary condition is ___ group. Most electropositive elements belongs to ___ group.



Watch Video Solution

7. Lather Meyer drew a graph showing the relation between atomic _____ and atomic _____ .



Watch Video Solution

 [Watch Video Solution](#)

8. Ionic radii _____ with increases in atomic number in a period and _____ in a group.



[Watch Video Solution](#)

9. The electron gain enthalpy of oxygen is _____ that of sulphur.



[Watch Video Solution](#)

10. On Mulliken scale the average of IP and EA is known as _____ .

 [Watch Video Solution](#)

11. On the Pauling's EN scale, the element next to F is _____

 [Watch Video Solution](#)

12. The IE of Be is _____ than that of B .

 [Watch Video Solution](#)

13. The bond angle in SO_4^{2-} ion is _____ .

 [Watch Video Solution](#)

14. The angle between two covalent bonds is maximum for (CH_4 , H_2O , CO_2) _____ .

 [Watch Video Solution](#)

15. Second element of group 1 shows diagonal relationship with the second element of group _____ .

 [Watch Video Solution](#)

 Watch Video Solution

16. The EN of the elements C, N, Si and P increases in the order of _____ .

 Watch Video Solution

17. The decreasing order IE for elements Li, Be, CB is _____ .

 Watch Video Solution

18. The type of magnetism exhibited by $[Mn(H_2O)_6]^{2+}$ ion is _____.

 [Watch Video Solution](#)

19. Among the ions Cl^{\ominus} , S^{2-} and Na^{\oplus} , the largest ion is _____.

 [Watch Video Solution](#)

20. The inner electrons are shielded to a _____ extent than the outer electrons.

 [Watch Video Solution](#)

 Watch Video Solution

Exercises (True/False)

1. Name five species that are isoelectronic with F^{\ominus} ion



Watch Video Solution

2. Number of species that are isoelectronics with Sr^{2+} ion are four.



Watch Video Solution

3. Among the elements , Li , K , Ca , Cl and Kr , the element K has the lowest IE and Kr has the highest IE .



[Watch Video Solution](#)

4. Why do beryllium and magnesium not impart colour to the flame in the flame test?



[Watch Video Solution](#)

5. Energy released when electron is added to an isolated gases anion.



[Watch Video Solution](#)

6. van der Waals radius of chlorine is less than that of covalent radius.



[Watch Video Solution](#)

7. The oxides and hydroxides of alkali metals are strong bases.



[Watch Video Solution](#)

8. Lithium is the lightest metal.



[Watch Video Solution](#)

9. As the s character of the hybrid orbital decreases, the EN increases.



[Watch Video Solution](#)

10. Assertion : Ionic bonds are directional in nature whereas covalent bonds are non-directional.

Reason : According to orbital overlap concept , the formation of a covalent bond between two atoms results by pairing of electrons present in the valence shell having same spins.



Watch Video Solution

Exercises (Archives) Multiple Correct

1. Which of the following statements is/are true for the long form of the periodic table?

- A. It refers the sequence of filling the electrons in the order of sub-energy level s , p , d and f
- B. It helps to predict the stable valency states of the elements.

C. It refers trends in physical and chemical properties of the elements.

D. It helps to predict the relative ionicity of the bond between any two elements.

Answer: A::C::D



Watch Video Solution

2. Sodium sulphate is soluble in water, whereas barium sulphate is sparingly soluble because

A. the hydration energy of sodium sulphate is more than its lattice energy

B. the lattice energy of barium sulphate is more than its hydration energy

C. the lattice energy has no role to play in solubility





D. the hydration energy of sodium sulphate is less than its lattice energy.

Answer: A::B



Watch Video Solution

3. Ionic radii of



Answer: D



Watch Video Solution

Exercises (Archives) Single Correct

1. The correct order of second ionisation potentials of carbon, nitrogen, oxygen and fluorine is

A. $C > N > O > F$

B. $O > N > F > C$

C. $O > F > N > C$

D. $F > O > N > C$

Answer: C



Watch Video Solution

2. Which of the following element has the highest ionisation energy ?

A. Boron

B. carbon

C. nitrogen

D. oxygen

Answer: C



Watch Video Solution

3. The first ionisation potential in electron volts of nitrogen and oxygen atoms are respectively given by

A. 14.6, 13.6

B. 13.6, 14.6

C. 13.6, 13.6

D. 14.6, 14.6

Answer: A



Watch Video Solution

4. Atomic radii of fluorine and neon (\AA) respectively are given as

A. 0.72, 1.60

B. 16.0, 1.60

C. 0.72, 0.72

D. None of these

Answer: A



Watch Video Solution

5. The electronegativity of the following elements increases in the order

A. C,N,Si,P

B. N,Si,C,P

C. Si,P,C,N

D. P,Si,N,C

Answer: C



Watch Video Solution

6. The first ionisation enthalpies of Na, Mg, Al and Si are the order

A. $Na < Mg > Al < Si$

B. $Na > Mg > Al > Si$

C. $Na < Mg < Al > Si$



D. $Na > Mg > Al < Si$

Answer: A



Watch Video Solution

7. Which one of the following is the smallest in size?







Answer: D



Watch Video Solution

8. Which of the following is not correct statement for periodic classification of elements?





Answer: B



Watch Video Solution

9. Which of the following is not correct statement for periodic classification of elements?

A. The properties of elements are the periodic functions of their atoms numbers.

B. Non-metallic elements are lesser in number than metallic elements.

C. The first ionisation energies of elements along a period do not vary in a regular manner with the increase in atomic number.

D. For transition elements the d-subshells are filled with electrons monotonically with the increase in atomic number.

Answer: D



Watch Video Solution

10. Which has the most stable + 2 oxidation state ?

A. *Sn*

B. *Pb*

C. *Fe*




D. *Ag*

Answer: B



Watch Video Solution

11. Which of the following has the maximum number of unpaired electrons?



Answer: D



Watch Video Solution

12. Which of the following statements is wrong?

A. The first ionisation potential of Al is less than the first ionisation potential of Mg .


- B. The second ionisation potential of Mg is greater than the second ionisation potential of Na
- C. The first ionisation potential of Na is less than the first ionisation potential of Mg .
- D. The third ionisation potential of Mg is greater than the third ionisation potential of Na

Answer: B



Watch Video Solution

13. The correct order of acidic strength is




Answer: A



Watch Video Solution

14. The correct order of radii is



Answer: B



Watch Video Solution

15. Amongst H_2O , H_2S , H_2Se and H_2Te , the one with the highest boiling point is :

A. H_2O because of hydrogen bonding

B. H_2Te because of higher molecular weight

C. H_2S because of hydrogen bonding

D. H_2Se because of lower molecular weight

Answer: A



Watch Video Solution

16. The set representing the correct order of the first ionisation potential is

A. $K > Na > Li$

B. $Be > Mg > Ca$

$$c. B > C > N$$

$$D. Ge > Si > C$$

Answer: B



Watch Video Solution

17. Which of the following represents correct order of increasing first ionization enthalpy for Ca, Ba, S, Se, and Ar?

$$A. S < Se < Ca < Ba < Ar$$

$$B. Ba < Ca < Se < S < Ar$$

C. $Ca < Ba < S < Se < Ar$

D. $Ca < S < Ba < Se < Ar$

Answer: B



Watch Video Solution

Exercises (Archives) Assertion Reasoning

1. Assertion: F atom has less negative electron gain enthalpy than Cl atom.

Reason: Additional electrons are repelled more

effectively by 3 p-electrons in Cl than by 2 p-electrons in F atom.

A. Statement-I is true, Statement-II is true

,Statement -II is the correct explanation for

Statement-I

B. Statement-I is true ,Statement -II is true,

Statement -II is not the correct explanation for

statement -II

C. Statement-I is true. Statement -II is false.

D. Statement-I is false. Statement -II is true.

Answer: C



2. Statement I : The first ionisation energy of Be is greater than that of B.

Statement II : 2p-orbital is lower in energy than 2s.

A. Statement-I is true, Statement-II is true

,Statement -II is the correct explanation for

Statement-I

B. Statement-I is true ,Statement -II is true,

Statement -II is not the correct explanation for

statement -II

C. Statement-I is true. Statement -II is false.

D. Statement-I is false. Statement -II is true.

Answer: C



Watch Video Solution

Exercises (Archives) Fill In The Blanks

1. The energy released when an electron is added to a neutral gaseous atom is called. of atom.



Watch Video Solution

2. On Mulliken scale the average of IP and EA is known as _____ .



[Watch Video Solution](#)

3. Ca^{2+} has a smaller ionic radius than K^+ because it has



[Watch Video Solution](#)

4. Compounds that formally contain Pb^{4+} are easily reduced to Pb^{+2} . The stability of the lower oxidation state is due to



[Watch Video Solution](#)

Exercises (Archives) True/False

1. The softness of group IA metals increase down the group with increasing atomic number.



[Watch Video Solution](#)

2. In group IA of alkali metals, the ionisation potential decrease down the group,Therefore ,lithium is a poor reducing agent.



[Watch Video Solution](#)

3. F, Cl, Br (Increasing electron affinity)



[Watch Video Solution](#)

4. The basic nature of the hydroxides of group 13 decreases progressively down the group.(T/F)



[Watch Video Solution](#)

Exercises (Archives)Subjective

1. Arrange the following in the given order

(a) Decreasing ionic size, Mg^{2+} , O^{2-} , Na^{\oplus} , F^{\oplus}

(b) Increasing bond length F_2 , N_2 , Cl_2O_2



Watch Video Solution


2. The IE_1 of C atom is greater than that of boron (B) atom, whereas the reverse is true for IE_2 .

Explain?



Watch Video Solution

3. Arrange the following as stated: Increasing order of ionic size



[Watch Video Solution](#)

4. Arrange the following ions in order of their decreasing ionic radii.



[Watch Video Solution](#)