



BIOLOGY

BOOKS - MAXIMUM PUBLICATION

Anatomy of Flowering Plants

Exercise

1. Both apical and intercalary meristems are
which meristems

A. Primary

B. Secondary

C. Lateral

D. None of these

Answer: A



Watch Video Solution

2. Sclerieds present in

A. fruit wall of nuts

B. pulp of guava

C. seed coat of legumes

D. All of these

Answer: D



Watch Video Solution

3. Which of the following condition of xylem is present in both monocot and dicot stems

A. exarch

B. endarch

C. polyarch

D. mesaarch

Answer: B



Watch Video Solution

4. Bark does not include

A. secondary xylem

B. secondary phloem

C. periderm

D. both a and b

Answer: A



Watch Video Solution

5. Which is not a living tissue?

A. collenchyma

B. parenchyma

C. sclerenchyma

D. none of these

Answer: A



Watch Video Solution

6. What is the cells that make the leaves curl in plants during water stress?



Watch Video Solution

7. What part of the plant would show the following:

Radial vascular bundle

polyarch xylem

Well developed pith



[Watch Video Solution](#)

8. Fill in the blank

Apical meristem for elongation of stem, Lateral

meristem for _____





[Watch Video Solution](#)

9. While analysing the anatomy of a given slide, phloem, cambium and xylem are present. Identify the material.



[Watch Video Solution](#)

10. Tissues are specialised to perform mechanical function. They have thickening at corners, living and cellulose. Name it.



[Watch Video Solution](#)

11. What does the fascicular cambium gives rise to?



Watch Video Solution

12. Which type of meristems can be classified on the basis of positions in the plant body?



Watch Video Solution

13. Find the odd one and give:

Phellem, Phellogen, Phloem, Phelloderm



Watch Video Solution

14. Fill in the blank

Stem : Endarch :: Root : _____



Watch Video Solution

15. Fill in the blank

interfascicular cambium: secondary xylem::

cork cambium: _____



Watch Video Solution

16. Activity of cambium is different in two seasons and xylem vessels formed have narrow and wide lumens. Give the type of wood formed during the different seasons.



Watch Video Solution

17. What use are phloem fibres put to?



Watch Video Solution

18. Identify the plant grow from the data given below:

habitat tree

xylem vessels present

absence of companion cell in phloem

no fruits



Watch Video Solution

19. Ramu observed the growth of a plant in his garden .He realised that the plant grow both in length and width. Name the tissue responsible for the growth of the plant



Watch Video Solution

20. A cross-section of of plant materials show the following anatomical features under microscope.

Vascular bundles are radially arranged .

To which organ should it be assigned? Four xylem strands with exarch condition of the protoxylem. To which organ should it be assigned? Edit



[Watch Video Solution](#)

21. A cross-section of of plant materials show the following anatomical features under microscope.

Vascular bundles are radially arranged .

To which organ should it be assigned? Four xylem strands with exarch condition of the protoxylem. To which organ should it be assigned? Edit



[Watch Video Solution](#)

22. the transverse section of a plant material shows the following anatomical feature:

Vascular bundles are conjoint, scattered and surrounded by sclerenchymatous bundle sheaths.

Phloem parenchyma is absent . What will you identify it as?



[Watch Video Solution](#)

23. the transverse section of a plant material shows the following anatomical feature:

Vascular bundles are conjoint, scattered and surrounded by sclerenchymatous bundle sheaths.

Phloem parenchyma is absent . What will you identify it as?



[Watch Video Solution](#)

24. A cross-section of of plant materials show the following anatomical features under microscope.

Vascular bundles are radially arranged .

To which organ should it be assigned? Four xylem strands with exarch condition of the protoxylem. To which organ should it be assigned? Edit



[Watch Video Solution](#)

25. A cross-section of of plant materials show the following anatomical features under microscope.

Vascular bundles are radially arranged .

To which organ should it be assigned? Four xylem strands with exarch condition of the protoxylem. To which organ should it be assigned? Edit



Watch Video Solution

26. During the secondary thickening of a dicot stem, from the outer cortical region, a meristematic tissue is developed. Name this tissue and write the function of it.



[Watch Video Solution](#)

27. Analyse the table and arrange the matter in an appropriate order.

A	B	C
Monocot stem	Bulliform cells	Casparian thickening
Dicot stem	Radial	Protoxylem lacunae
Isobilateral leaf	Closed	Secondary thickening
Dicot root	Open	Epidermits



Watch Video Solution

28. Differentiate the following characters of dicot stem and monocot stem.

Differentiated cortex, Protoxylem lacuna present, open bundle, bundle Sheath, starch sheath, sclerenchymatous hypodermis, scattered bundles, V.Bs. arranged in ring, closed bundle, phloem parenchyma absent, presence of pith, collenchymatous hypodermis.



Watch Video Solution

29. Xylem is the complex tissue that transport water in plants.

Name the main components of xylem. Which of these is mostly likely suitable for conducting water?



Watch Video Solution

30. Xylem is the complex tissue that transport water in plants.

Name the main components of xylem. Which of these is mostly likely suitable for conducting water?



[Watch Video Solution](#)

31. Examine the anatomy of two plants.

Xylem vessels and tracheids are seen.

Xylem vessels absent and tracheids present.



[Watch Video Solution](#)

32. Examine the anatomy of two plans.

Xylem vessels and tracheids are seen.

Xylem vessels absent and tracheids present.



Watch Video Solution

33. Identify the figure A and B and differentiate between A and B.



[Watch Video Solution](#)

34. Phloem is called the complex tissue while parenchyma is a simple tissue. Make the differences.



[Watch Video Solution](#)

35. When the cross section of a stem was examined, a student could notice collenchymatous hypodermis, Limited number of open vascular bundles, cortex, endodermis and pith.

(1) Identify the material.

(2) Draw the diagram of the material showing these parts and label.



[Watch Video Solution](#)

36. When the cross section of a stem was examined, a student could notice collenchymatous hypodermis, Limited number of open vascular bundles, cortex, endodermis and pith.


(1) Identify the material.

(2) Draw the diagram of the material showing these parts and label.



Watch Video Solution

37. Observe the diagrams and identify the tissue types.



Watch Video Solution

38. Analyse the table and arrange the matter in an appropriate order.

A	B	C
Monocot stem	Bulliform cells	Casparian thickening
Dicot stem	Radial	Protoxylem lacunae
Isobilateral leaf	Closed	Secondary thickening
Dicot root	Open	Epidermits



Watch Video Solution

39. While doing anatomy practical, students observe ten following characters:

(i) Undifferentiated ground tissue.

(ii) Numerous vascular bundles and are scattered.

(iii) Sclerenchymatous hypodermis.

(iv) Presence of bundle sheath.

Identify the material. Write any one example of a plant having such characters.



[Watch Video Solution](#)

40. While doing anatomy practical, students observe ten following characters:

(i) Undifferentiated ground tissue.

(ii) Numerous vascular bundles and are scattered.

(iii) Sclerenchymatous hypodermis.

(iv) Presence of bundle sheath.

Identify the material. Write any one example of a plant having such characters.



[Watch Video Solution](#)

41. What is the function of phloem parenchyma?



[Watch Video Solution](#)

42. Activity of cambium is different in two seasons and xylem vessels formed have narrow and wide lumens. Give the type of wood formed during the different seasons.



[Watch Video Solution](#)

43. Mention the characteristic features of sieve tube members.



[Watch Video Solution](#)

44. It is difficult to open and close wooden doors and windows during rainy season.

What is the reason?



Watch Video Solution

45. Heart wood is resistant to micro organisms. Justify the statement.



Watch Video Solution

46. Simple tissues that perform mechanical support in plants and usually dead tissues.

Name the living simple tissue that provides mechanical support to plant organs. In which category of plants(dicots/monocots) do they found?



Watch Video Solution

47. What is present on the surface of the leaves which helps the plant prevent loss of water but is absent in roots?



[Watch Video Solution](#)

48. What is the epidermal cell modification in plants which prevents water loss?



[Watch Video Solution](#)

49. Find the odd one and give:

Phellem, Phellogen, Phloem, Phelloderm



[Watch Video Solution](#)

50. A transverse section of the trunk of a tree shows concentric rings which are known as growth rings. How are these rings formed? What is the significance of these rings?



Watch Video Solution

51. While eating peach or pear it is usually seen that some stone like structures get entangled in the teeth, what are these stone like structures called?



Watch Video Solution

52. We cannot find the woody trunk in monocot plants. Why?



Watch Video Solution

53. Differentiate the following characters of dicot stem and monocot stem.

Differentiated cortex, Protoxylem lacuna present, open bundle, bundle Sheath, starch sheath, sclerenchymatous hypodermis,

scattered bundles, V.Bs. arranged in ring, closed bundle, phloem parenchyma absent, presence of pith, collenchymatous hypodermis.



[Watch Video Solution](#)

54. Differentiate the following characters of dicot stem and monocot stem.

Differentiated cortex, Protoxylem lacuna present, open bundle, bundle sheath, starch sheath, sclerenchymatous hypodermis,

scattered bundles, V.Bs. arranged in ring, closed bundle, phloem parenchyma absent, presence of pith, collenchymatous hypodermis.



[Watch Video Solution](#)

55. The product of photosynthesis is transported from the leaves to various parts of the plants and stored in some cell before being utilised. What are cells tissues that store them?



[Watch Video Solution](#)

56. Name the kind of vascular bundle in Fig. a and b.



[Watch Video Solution](#)

57. Heart wood is resistant to micro organisms. Justify the statement.



[Watch Video Solution](#)

58. Cambial ring is responsible for the formation of secondary vascular tissues both in Dicot stem and Dicot root. Differentiate the origin of cambial ring in Dicot stem and Dicot root.



Watch Video Solution

59. Give reason

(a) Grittiness nature of sapola fruit pulp.

(b) Increase thickness of stem in Teak and not in coconut.



[Watch Video Solution](#)

60. Give reason

(a) Grittiness nature of sapola fruit pulp.

(b) Increase thickness of stem in Teak and not in coconut.



[Watch Video Solution](#)

61. Match the following.

Match the following.

A	B	C
1. guard cell	Suberin	Stem
2. conjoint	Stomata	Casparian stripe
3. endodermis	Vascular bundle	Subsidiary cell



[Watch Video Solution](#)

62. You may have observed droplets of water at the tip of the leaves of grasses during early morning. Your sister told its dew drop.

As a science student do you agree with her?

Explain. Compare this with the process of transpiration.



Watch Video Solution

63. Activity of cambium is different in two seasons and xylem vessels formed have narrow and wide lumens. Give the type of wood formed during the different seasons.



Watch Video Solution

64. Differentiate the following characters of dicot stem and monocot stem.

Differentiated cortex, Protoxylem lacuna present, open bundle, bundle sheath, starch sheath, sclerenchymatous hypodermis, scattered bundles, V.Bs. arranged in ring, closed bundle, phloem parenchyma absent, presence of pith, collenchymatous hypodermis.



Watch Video Solution

65. Fill in the blank

interfascicular cambium: secondary xylem::

cork cambium: _____



Watch Video Solution

66. Identify the following diagram and write its function.



Watch Video Solution

67. Where are companion cells located in the flowering plants? What are their functions?



Watch Video Solution

68. Xylem is the complex tissue that transport water in plants.

Name the main components of xylem. Which of these is mostly likely suitable for conducting water?



Watch Video Solution

69. Fill in the blank

Stem :Endarch :: Root : _____



Watch Video Solution

70. If one debarks a tree, what parts of the plant is being removed?



Watch Video Solution

71. Some anatomical details of a material are given below

vascular bundles are conjoint and open, medullary rays are present, hypodermis collenchymatous, cortex heterogenous.

Identify the material. Name the tissue given above from which interfascicular cambium develops during secondary growth.



[Watch Video Solution](#)

72. Some anatomical details of a material are given below

vascular bundles are conjoint and open, medullary rays are present, hypodermis collenchymatous, cortex heterogenous.

Identify the material. Name the tissue given above from which interfascicular cambium develops during secondary growth.



Watch Video Solution

73. During the secondary thickening of a dicot stem, from the outer cortical region, a meristematic tissue is developed. Name this tissue and write the function of it.



Watch Video Solution

74. Distinguish between heartwood and sapwood.



Watch Video Solution

75. A portion of transverse section of maize stem is shown in the diagram. Label a,b,c,d,e,f,g,h and i



Watch Video Solution

76. Distinguish between heartwood and sapwood.



Watch Video Solution

77. Fill in the blank

Stem :Endarch :: Root : _____



[Watch Video Solution](#)

78. Match the column A with B and C .

A (Collenchyma , Cambium, Sieve tube)

B(Lateral meristem, Companion cell, Thickened

at corners) C (Food Conduction, Hypodermis

of dicot stem, secondary growth)



[Watch Video Solution](#)

79. Match the following :

(Column A) - Nostoc, Gonyaulax, Chlorella,
Cycas, Rhizophora, Funaria

(Column B) - Protonema, respiratory root, red
tide, Space travelers, heterocyst, Coralloid root



Watch Video Solution

80. Grasses are capable of regeneration in spite of continuous grazing by cattle. Suggest the reason for it.



Watch Video Solution

81. In most woody trees, epidermis breaks to form a lens shape opening. Name this opening that permits gaseous exchange.



Watch Video Solution

82. You may have observed droplets of water at the tip of the leaves of grasses during early morning. Your sister told its dew drop.

As a science student do you agree with her?

Explain. Compare this with the process of transpiration.



[Watch Video Solution](#)

83. You may have observed droplets of water at the tip of the leaves of grasses during early morning. Your sister told its dew drop.

As a science student do you agree with her?

Explain. Compare this with the process of transpiration.



[Watch Video Solution](#)

84. What is the cells that make the leaves curl in plants during water stress?



Watch Video Solution

85. Teacher asked Gopi to name the layer from which root hairs and lateral roots arise respectively.



Watch Video Solution

86. In an anatomy lab, Ramu and Salim were taking transverse sections of two specimens A and B respectively. Their observations are given in the table. Complete the table.

Specimen A	Specimen B
1a. Closed vascular bundle വാസ്കുലാർ ബണ്ടിലുകൾ ക്ലോസ്ഡ് ആണ്	1b. Open vascular bundle വാസ്കുലാർ ബണ്ടിലുകൾ ഓപ്പൺ ആണ്
2a.	2b.
3a.	3b.



[Watch Video Solution](#)

87. The following is a list of characteristics features of angiosperms and gymnosperms. Those characters that belong to gymnosperms.

- (a) Tracheids alone from the conducting elements in xylem
- (b) production of fruits.
- (c) naked seeds
- (d) cones are seen
- (e) flowers absence
- (f) xylem mainly contains vessel
- (g) double fertilization present



[Watch Video Solution](#)

88. Cambial ring is responsible for the formation of secondary vascular tissues both in Dicot stem and Dicot root. Differentiate the origin of cambial ring in Dicot stem and Dicot root.



[Watch Video Solution](#)

89. Cambial ring is responsible for the formation of secondary vascular tissues both in Dicot stem and Dicot root. Differentiate the origin of cambial ring in Dicot stem and Dicot root.



Watch Video Solution

90. Some tissues in plants are not able to divide further.

(a) Suggest the name of such tissues.

(b) Give any three examples.

(c) List the difference between epidermal tissue system of roots and leaves.



[Watch Video Solution](#)

91. Some tissues in plants are not able to divide further.

(a) Suggest the name of such tissues.

(b) Give any three examples.

(c) List the difference between epidermal tissue system of roots and leaves.



[Watch Video Solution](#)

92. Some tissues in plants are not able to divide further.

(a) Suggest the name of such tissues.

(b) Give any three examples.

(c) List the difference between epidermal tissue system of roots and leaves.



[Watch Video Solution](#)

93. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

a) Radial and polyarch xylem bundles

b) Parenchymatous (homogenous) cortex

c) Large pith

d) Epidermis with unicellular hairs

e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct

sequence.

iii) Give an account of casparian strips.



Watch Video Solution

94. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

- a) Radial and polyarch xylem bundles
- b) Parenchymatous (homogenous) cortex
- c) Large pith
- d) Epidermis with unicellular hairs
- e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct sequence.

iii) Give an account of casparian strips.



Watch Video Solution

95. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

a) Radial and polyarch xylem bundles

b) Parenchymatous (homogenous) cortex

c) Large pith

d) Epidermis with unicellular hairs

e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct sequence.

iii) Give an account of casparian strips.



Watch Video Solution

96. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

a) Radial and polyarch xylem bundles

b) Parenchymatous (homogenous) cortex

c) Large pith

d) Epidermis with unicellular hairs

e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct sequence.

iii) Give an account of casparian strips.



Watch Video Solution

97. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

- a) Radial and polyarch xylem bundles
- b) Parenchymatous (homogenous) cortex
- c) Large pith
- d) Epidermis with unicellular hairs
- e) Pericycle
- f) Endodermis with casparian strips.
- i) Identify the plant part.
- ii) Re-arrange the given regions from the

periph-ery to the center in their correct sequence.

iii) Give an account of casparian strips.



[Watch Video Solution](#)

98. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

- a) Radial and polyarch xylem bundles
- b) Parenchymatous (homogenous) cortex
- c) Large pith
- d) Epidermis with unicellular hairs

e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct sequence.

iii) Give an account of casparian strips.



[Watch Video Solution](#)

99. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

- a) Radial and polyarch xylem bundles
- b) Parenchymatous (homogenous) cortex
- c) Large pith
- d) Epidermis with unicellular hairs
- e) Pericycle
- f) Endodermis with casparian strips.
- i) Identify the plant part.
- ii) Re-arrange the given regions from the periphery to the center in their correct sequence.
- iii) Give an account of casparian strips.



Watch Video Solution

100. In the anatomy lab. Eugin observed the following features in the T.S of a plant part.

a) Radial and polyarch xylem bundles

b) Parenchymatous (homogenous) cortex

c) Large pith

d) Epidermis with unicellular hairs

e) Pericycle

f) Endodermis with casparian strips.

i) Identify the plant part.

ii) Re-arrange the given regions from the periph-ery to the center in their correct

sequence.

iii) Give an account of casparian strips.



[Watch Video Solution](#)

101. Match the following columns A and B.



[Watch Video Solution](#)

102. Stomata are small openings present in a epidermis of leaves. The stomata are bound by

guard cells. Mention the role off guard cells in stomatalmechanism.



[Watch Video Solution](#)

103. How does periderm develop in dicot stem and replace the outer broken cortical and epidermal layers?



[Watch Video Solution](#)

104. Name the three basic tissue systems in the flowering plants. Give the tissue names under each system.



Watch Video Solution

105. the transverse section of a plant material shows the following anatomical feature:

Vascular bundles are conjoint, scattered and surrounded by sclerenchymatous bundle sheaths.

Phloem parenchyma is absent . What will you identify it as?



[Watch Video Solution](#)

106. the transverse section of a plant material shows the following anatomical feature:

Vascular bundles are conjoint, scattered and surrounded by sclerenchymatous bundle sheaths.

Phloem parenchyma is absent . What will you identify it as?



[Watch Video Solution](#)

107. Phloem is called the complex tissue while parenchyma is a simple tissue. Make the differences.



[Watch Video Solution](#)

108. Interfascicular cambium is a

A. primary meristematic tissue

B. primordial meristem

C. type of protodenn

D. secondary meristematic tissue

Answer: D



Watch Video Solution

109. In an annual ring, the light coloured part is known as

A. early wood

B. latewood

C. heartwood

D. sapwood

Answer: A



View Text Solution

110. The layer of cell outside the phloem meant for giving rise to the root branches is called

A. cambium

B. cork

C. endodermis

D. pericycle

Answer: D



Watch Video Solution

111. Cork cambium of dicot originates from

A. epiblema

B. pericycle

C. cambium of vascular bundles

D. endodermis

Answer: B



View Text Solution

112. Alburnum is otherwise known as

A. periderm

B. heartwood

C. sapwood

D. cork

Answer: B



Watch Video Solution

113. Duramen is present in

- A. inner region of secondary wood
- B. part of sapwood
- C. outer region of secondary wood
- D. region of pericycle

Answer: A



[View Text Solution](#)

114. Which one of the following is a current sequence of tissue present in dicot stem during secondary growth

A. phellogen, cork, primary cortex, secondary cortex

B. Cork, primary cortex, secondary cortex, phellogen

C. primary cortex, secondary cortex,
phellogen, cork

D. secondary cortex, phellogen, cork,
primary cortex

Answer: D



Watch Video Solution

115. The meristem responsible for extra stelar secondary growth in dicot stem is

A. interfascicular cambium

B. intrafascicular cambium

C. intercalary meristem

D. phellogen

Answer: D



Watch Video Solution

116. The following is a list of characteristics features of angiosperms and gymnosperms.

Those characters that belong to

gymnosperms.

(a) Tracheids alone from the conducting elements in xylem

(b) production of fruits.

(c) naked seeds

(d) cones are seen

(e) flowers absence

(f) xylem mainly contains vessel

(g) double fertilization present

A. are the dominant cell type of xylem in

angiosperm

B. are primarily found in mosses and liverworts

C. are responsible for water conduction and support in many land plants

D. first appeared during Palaeozoic era

Answer: C



Watch Video Solution

117. In the following, how sapwood is converted into heart wood?

- A. By degeneration of protoplast of living cell
- B. Tyloses formation
- C. By deposition of resin, oil, gums etc
- D. All of the above

Answer: D



Watch Video Solution

118. As secondary growth proceeds in a dicot stem, the thickness of

A. sapwood increases

B. heartwood increases

C. both sapwood and heartwood increases

D. both heartwood and sapwood remains
the same

Answer: C



Watch Video Solution

119. Which is not a characteristic of plants cell wall?

A. found only in the sporophyte phase of a life cycle

B. among other compounds contains compounds built of simple sugar

C. may contain enzyme that are biologically active

D. often contain strengthening polymer

Answer: A



Watch Video Solution

120. Identify the correct order of the component reference to their arrangement from outer side to inner side in a woody dicot stem.

1.secondary cortex 2. Autumn wood

3.secondary phloem 4. phellem

A. 2,3,1,4

B. 3,4,2,1

C. 4,1,3,2

D. 1,2,4,3

Answer: C



Watch Video Solution

121. Removal of ring wood of tissue outside the vascular cambium from the tree trunk kill it because

A. water cannot move up

B. food does not travel down and root
become starved

C. shoot become starved

D. annual rings are not produced

Answer: B



View Text Solution

122. Which tissue gives rise to secondary growth?

- A. Apical meristem
- B. Adventitious root
- C. Germinating seed
- D. vascular cambium

Answer: D



View Text Solution

123. Growth rings are formed due to activity of

A. extrastelar cambium

B. intrastelar cambium

C. interstelar cambium

D. both(b) and (c)

Answer: D



View Text Solution

124. A nail is driven into the trunk of a 30 years old tree at a point I in above the soil level. The tree grows in height at the rate of 0.5 m a year. After 3 year nail will be

- A. 1m above the soil
- B. 1.5 m above the soil
- C. 2m above the soil
- D. 2.5 m above the soil

Answer: A



Watch Video Solution

125. In which of the following there is no differentiation of bark, sapwood and heartwood?

A. ashok

B. neem

C. mango

D. date palm

Answer: D





126. A tree grows at the rate of 0.5m per year. What will be the height of the board fixed at 1.5m above the base five years ago?

A. 4.0m

B. 3.5m

C. 1.5m

D. 4.5m

Answer: C



Watch Video Solution

127. Secondary phloem remains functional generally

- A. for one year
- B. for less than one year
- C. for many year
- D. as long as plant is alive

Answer: D



Watch Video Solution

