



PHYSICS

BOOKS - TARGET PHYSICS (MARATHI ENGLISH)

FRICTION IS SOLIDS AND LIQUIDS

Mcq

1. Friction is always

- A. parallel to the motion of the body.
- B. perpendicular to the surface of contact.
- C. tangential to the surface of contact.
- D. inclined to the surface of contact.

Answer: C



Watch Video Solution

2. When two surfaces are coated with lubricant, friction between the surfaces _____

A. becomes zero

B. decreases

C. remains constant

D. increases

Answer: B



Watch Video Solution

3. Out of the following which is NOT a friction

?

A. contact friction

B. fluid friction

C. viscous force

D. collision

Answer: D



Watch Video Solution

4. Friction is caused by

- A. interlocking between the irregularities on the contact surface.
- B. apparent area of contact.
- C. repulsive force between air and surface in earth .
- D. gravitational force of attraction towards the earth.

Answer: A



Watch Video Solution

5. Which of the following is a CORRECT statement ?

A. Apparent contact area is equal to actual contact area.

B. Apparent contact area is half of the actual contact area.

C. Apparent contact area is less than actual contact area.

D. Apparent contact area is greater than actual contact area.

Answer: D



Watch Video Solution

6. The elevation of one surface _____ depression in other surface.

- A. gets interlocked with
- B. makes free movement over
- C. transfers free electrons to
- D. gets unlocked with

Answer: A



Watch Video Solution

7. A spherical soap bubble is expanding so that its radius is increasing at the rate of 0.02 cm/sec. At what rate is the surface area is increasing, when its radius is 5cm?

A. vertically upwards called normal reaction.

B. vertically downward called normal reaction.

C. tangential force acting along the motion

.

D. tangential force acting opposite to the motion.

Answer: A



Watch Video Solution

8. In equilibrium, limiting force of static friction (F_s) acting between two surface in contact and applied force (F) is related by

A. $F_s > F$

B. $F_s < F$

C. $F_s = F$

D. $F_s = \frac{1}{2}F$

Answer: C



Watch Video Solution

9. Friction between two bodies in contact when one body just moves or tends to move over the other is called as

- A. static friction
- B. kinetic friction
- C. dynamic friction
- D. rolling friction

Answer: A



Watch Video Solution

10. Choose the CORRECT relation.

A. static friction $<$ kinetic friction $>$
rolling friction.

B. static friction $>$ kinetic friction $<$
rolling friction.

C. static friction $<$ rolling friction $>$
kinetic friction.

D. static friction $>$ kinetic friction $>$
rolling friction.

Answer: D



Watch Video Solution

11. The limiting force of static friction is approximately independent of

- A. apparent area of surfaces in contact.
- B. nature of surfaces in contact.
- C. materials of the surfaces in contact.

D. normal reaction between two surface in contact.

Answer: A



Watch Video Solution

12. Coefficient of static friction does not depend upon

A. nature and materials of the surfaces.

B. normal reaction

C. limiting force of static friction.

D. apparent area of contact.

Answer: B



Watch Video Solution

13. According to 1st law of static friction, the relation between F_s and N is given by

A. $F_s \propto N^2$

B. $F_s \propto \frac{1}{N}$

C. $F_s \propto \frac{1}{N^2}$

D. $F_s \propto N$

Answer: D



Watch Video Solution

14. If the normal reaction is doubled, the frictional force is _____

A. halved

B. unchanged

C. doubled

D. triple

Answer: C



Watch Video Solution

15. A block of 50 kg rests on a table. A horizontal force of 294 N is required to just move the block, the coefficient of static friction between the surfaces in contact (μ_s) is

A. 5.88

B. 1.67

C. 0.6

D. 0.17

Answer: C



Watch Video Solution

16. The friction that exists between the surface of two bodies in contact when one body is sliding over the other is called _____.

A. rolling friction

B. kinetic friction

C. static friction

D. steady friction

Answer: B



Watch Video Solution

17. Choose the CORRECT statement.

A. The magnitude of the force of kinetic friction depends upon shape of surface in contact.

B. The magnitude of force of kinetic friction depends upon apparent area of the surface in contact.

C. Force of kinetic friction is independent of material of the surfaces in contact.

D. The magnitude of the force of kinetic friction is approximately independent of

the relative velocity provided it is not too large or small.

Answer: D



Watch Video Solution

18. A block of mass 4 kg resting on horizontal surface can be kept in uniform motion on the horizontal surface by the application of the force 16 N, the coefficient of kinetic friction between the two surfaces is

A. 0.51

B. 0.41

C. 0.31

D. 0.21

Answer: B



Watch Video Solution

19. why are liquids and gases categorised as fluids ?

A. elastic bodies

B. plastic bodies

C. fluids

D. semisolids

Answer: C



Watch Video Solution

20. Static and homogeneous fluids are _____.

A. isotropic

B. heterogeneous

C. stagnant

D. colloidal

Answer: A



Watch Video Solution

21. The _____ exerted by a liquid at rest per unit area normal to the surface in contact with the liquid is called pressure.

A. velocity

B. displacement

C. thrust

D. work

Answer: C



Watch Video Solution

22. Liquid of density ρ in a container exerts pressure P given by (height of liquid h and area A)

A. $P = Ah\rho g$

B. $p = h\rho g$

C. $P = \frac{h\rho g}{A}$

D. $P = \frac{\rho g}{Ah}$

Answer: B



Watch Video Solution

23. Weight of the liquid column is given by

A. volume of the liquid \times density of liquid

\times g

B. mass of the liquid \times density of liquid

C. volume of the liquid \times height of the
liquid

D. cross sectional area of the liquid \times
density of liquid \times g

Answer: A



Watch Video Solution

24. Choose the CORRECT relation.

A. Absolute pressure =

Gauge pressure + Atmospheric pressure

B. Atmospheric pressure =

Absolute pressure + Gauge pressure

C. Atmospheric pressure =

Gauge pressure - Absolute pressure

D. Absolute pressure =

Atmospheric pressure - Gauge pressure

Answer: A



Watch Video Solution

25. At what depth in fresh water the pressure on a diver is one atmosphere ?

[Density of water = 10^3 kgm^{-3} , Normal pressure = 10^5 Pa , $g = 10 \text{ ms}^{-2}$]

A. 11 m

B. 10.5 m

C. 10 m

D. 9.8 m

Answer: C



Watch Video Solution

26. What is the pressure at the bottom of the ocean at a place where it is 3 km deep ?

[atmospheric pressure = $1.01 \times 10^5 Pa$,

Density of sea water = $1030 kgm^{-3}$]

A. $1.01 \times 10^5 Pa$

B. $3 \times 10^5 Pa$

C. $1.01 \times 10^7 Pa$

D. $3 \times 10^7 Pa$

Answer: D



Watch Video Solution

27. The shape of the vessel containing the liquid does not affect the pressure. This is known as

- A. hydraulic pressure.
- B. hydrostatic pressure.
- C. hydrostatic paradox.
- D. hydrostatic parallax.

Answer: C



Watch Video Solution

28. Pascal's law is NOT applicable to _____.

- A. fluids

B. gases

C. solids

D. liquids

Answer: C



Watch Video Solution

29. The pressure applied to any part of the enclosed fluid at rest is transmitted undiminished to every portion of the fluid and to the walls of the vessel. This is _____.

A. Stock's law

B. Newton's law

C. Pascal's law

D. Torricelli's law

Answer: C



Watch Video Solution

30. To measure pressure difference, a device used is _____.

A. barometer

B. venturimeter

C. open tube manometer

D. aneroid barometer

Answer: C



Watch Video Solution

31. Liquid in the open tube monometer has _____ For measureing small pressure differende.

A. low density

B. high density

C. alcohol

D. water

Answer: A



Watch Video Solution

32. In a hydraulic lift, F_1 and F_2 are the force acting on the small piston and large piston having radii r_1 and r_2 respectively, then

$$\text{A. } F_1 = \frac{r_2^2}{r_1^2} F_2$$

$$\text{B. } F_2 = \frac{r_2^2}{r_1^2} F_1$$

$$\text{C. } F_1 = \frac{r_1}{r_2} F_2$$

$$\text{D. } F_1 = \frac{r_2}{r_1} F_2$$

Answer: B



Watch Video Solution

33. Which of the following is NOT an application of Pascal's law ?

A. Hydraulic brakes

B. Hydraulic lift

C. Hydraulic press

D. Aerodynamic lift

Answer: D



Watch Video Solution

34. The two femurs each of cross-sectional area 10cm^2 support the upper part of a human body of mass 40 kg. the average

pressure sustained by the femurs is (take

$$g = 10ms^{-2})$$

A. $3.92 \times 10^5 Pa$

B. $1.96 \times 10^5 Pa$

C. $3.92 \times 10^2 Pa$

D. $1.96 \times 10^2 Pa$

Answer: B



Watch Video Solution

35. What is the barometric height of a liquid of density 3.4 g cm^{-3} at a place, where that for mercury barometer is 70 cm?

- A. 70 cm
- B. 140 cm
- C. 228 cm
- D. 280 cm

Answer: D



Watch Video Solution

36. Viscosity is the property of the liquids and gases which is more closely related to _____.

A. elasticity

B. inertia

C. tension

D. friction

Answer: D



Watch Video Solution

37. In streamline flow of liquid through a pipe of uniform cross sectional area, all streamlines are

- A. divided into plane layers.
- B. divided into rectangular blocks.
- C. parallel to the axis of the tube.
- D. circular in shape.

Answer: C



Watch Video Solution

38. In streamline flow, the velocity of a liquid at a given point is

A. constant in magnitude only.

B. constant in direction but not constant in magnitude.

C. not constant in direction but constant in magnitude.

D. always constant in magnitude and direction.

Answer: D



Watch Video Solution

39. For streamline flow of an incompressible fluid, If A is area and v is speed then equation of continuity is

A. $Av = \text{constant}$

B. $A + v = \text{constant}$

C. $\frac{A}{v} = \text{constant}$

D. $\frac{v}{A} = \text{constant}$

Answer: A



Watch Video Solution

40. Select the CORRECT statement.

A. Two streamlines are always perpendicular.

B. Two streamlines will intersect at an angle of 30° between them.

C. Two streamlines will never intersect.

D. Two streamlines will not exist.

Answer: C



Watch Video Solution

41. An incompressible fluid flows steadily through a cylindrical pipe which has radius $2R$ at point A and radius R at point B farther along the flow direction. If the velocity at point A is v , its velocity at point B is

A. $\frac{v}{2}$

B. v

C. $2v$

D. $4v$

Answer: D



Watch Video Solution

42. A pipe 2 cm in diameter has a constriction of diameter 1 cm. What is the velocity of flow at the constriction. If velocity of flow in the broader region of the pipe is 5 cm/s?

A. 10 cm/s

B. 20 cm/s

C. 25 cm/s

D. 30 cm/s

Answer: B



Watch Video Solution

43. Flooded river is an example of _____

A. streamline flow

B. turbulent flow

C. laminar flow

D. viscos flow

Answer: B



Watch Video Solution

44. When the velocity of flow of liquid is greater than critical velocity, the liquid is said to have

A. streamline flow

B. laminar flow

C. turbulent flow

D. viscos flow

Answer: C



Watch Video Solution

45. The viscous force acting on adjacent layers of liquid is

A. perpendicular to it.

B. vertically downward.

C. vertically upward.

D. tangential to it.

Answer: D



Watch Video Solution

46. Velocity of liquid layer kept in a vessel is maximum at _____.

A. top

B. bottom

C. middle

D. cannot be predicted

Answer: A



Watch Video Solution

47. The velocity of liquid flowing through a tube at certain distance from the axis of tube

A. increases with distance.

B. remains constant.

C. decreases with distance.

D. depends upon length of the tube.

Answer: C



Watch Video Solution

48. The rate of change of velocity with distance measured from stationary layer is called _____.

A. acceleration

B. force

C. time

D. velocity gradient

Answer: D



Watch Video Solution

49. The S.I. unit and dimensions of velocity gradient in streamline flow is

A. m / s , $[M^0 L^1 T^{-1}]$

B. m , $[M^0 L^1 T^0]$

C. M^{-1} , $[M^0 L^{-1} T]$

D. S^{-1} , $[M^0 L^0 T^{-1}]$

Answer: D



Watch Video Solution

50. An incompressible liquid flows through a uniform cross section tube with velocity 12

cm/s. the thickness of liquid layer is 0.8 cm

then velocity of gradient of flow is

A. $15s^{-1}$

B. $12s^{-1}$

C. $18s^{-1}$

D. $5s^{-1}$

Answer: A



Watch Video Solution

51. The velocity gradient of certain liquid is $5s^{-1}$. If thickness of liquid layer is 2.5 cm, then velocity of flow of liquid will be

A. 10 cm/s

B. 2.5 cm/s

C. 12.5 cm/s

D. 5 cm/s

Answer: C



Watch Video Solution

52. The tangential force or viscous force on any layer of the liquid is directly proportional to the velocity gradient dv/dx . Then the direction of velocity gradient is

- A. perpendicular to the direction of flow of the liquid.
- B. parallel to the direction of the flow of the liquid.
- C. opposite to the direction of the flow of the liquid.

D. independent of the direction of the flow
of liquid.

Answer: A



Watch Video Solution

53. Coefficient of viscosity of a liquid does not
depend upon _____

A. velocity gradient

B. area of layer

C. direction of liquid flow

D. nature of the liquid

Answer: C



Watch Video Solution

54. A metal plate having an area of $0.04m^2$ is placed on a horizontal wooden surface. Oil of coefficient of viscosity $2Ns / s^2$ is introduced between the plate and the surface till the thickness of the oil layer is 0.5 mm. The

horizontal force needed to drag the plate along the surface with a velocity of 5 cm/s is

A. 80 N

B. 60 N

C. 8 N

D. 6 N

Answer: C



Watch Video Solution

55. The force of 2000 dyne is required to move a flat glass plate of surface area 10cm^2 with a velocity of 1 cm/s over a surface of glycerine 1 mm thick. The coefficient of viscosity of glycerine is

A. 0.2 poise

B. 0 poise

C. 20 poise

D. 20.5 poise

Answer: C



Watch Video Solution

56. Viscous force on a small sphere of radius r moving in a fluid varies directly with

A. r^2

B. r

C. $\frac{1}{r}$

D. $\left(\frac{1}{r}\right)^2$

Answer: B



57. Two metal ball of radius R and $2R$ falling through a fluid have same velocity at some point. The viscous drag acting on them at that instant are in the ratio

A. 1 : 4

B. 1 : 2

C. 2 : 1

D. 4 : 1

Answer: B



Watch Video Solution

58. A rain drop of radius 0.5 mm has a terminal velocity in air 2 m/s. If the coefficient of viscosity of air is 1.8×10^{-4} poise, the viscous drag on the rain drop will be

A. 0.014 dyne

B. 0.02 dyne

C. 0.034 dyne

D. 0.04 dyne

Answer: C



Watch Video Solution

59. When a spherical body falls through a viscous fluid, it experiences a viscous force, the motion of the body is

A. initially accelerated then becomes constant.

B. continuously accelerated.

C. continuously moving with different velocity.

D. initially constant then decreases till it becomes zero.

Answer: A



Watch Video Solution

60. Which of the following is correct formula for terminal velocity of a body in a viscous liquid ?

A. $\frac{9}{2} \cdot \frac{r^3(\sigma - \rho)g}{\eta}$

B. $\frac{2}{9} \cdot \frac{r^3(\rho - \sigma)g}{\eta}$

C. $\frac{2}{9} \cdot \frac{r^2g(\rho - \sigma)}{\eta}$

D. $\frac{2}{9} \cdot \frac{r^2(\rho - \sigma)}{\eta g}$

Answer: C



Watch Video Solution

61. find the terminal velocity of a steel ball 2mm in diameter falling through glycerine. Relative density of steel =8, relative density of glycerine =1.3 and viscosity of glycerine=8.3 poise.

A. $15.14 \times 10^{-3} m / s$

B. $17.43 \times 10^{-3} m / s$

C. $14.43 \times 10^{-4} m / s$

D. $17.43 \times 10^{-2} m / s$

Answer: B



Watch Video Solution

62. A drop of radius 2×10^{-5} and density $1.2 \times 10^3 \text{ kg/m}^3$ falls through air. The viscosity of air is $1.8 \times 10^{-5} \text{ N s/m}^2$.

Neglecting buoyancy due to air, the terminal speed of the drop is

A. 2.8 cm/s

B. 3.8 cm/s

C. 4.8 cm/s

D. $5.8\text{cm} / \text{s}$

Answer: D



Watch Video Solution

63. The terminal velocity of a water drop of radius 0.01 mm falling through air is 1.12 cm/s . If the density of air is neglected, the coefficient of viscosity of air is

[density _____ of _____ water

$$= 10^3\text{ kg} / \text{m}^3, g = 9.8\text{ m} / \text{s}^2]$$

A. $2 \times 10^{-5} \text{Ns} / \text{m}^2$

B. $1.9 \times 10^{-5} \text{Ns} / \text{m}^2$

C. $1.8 \times 10^{-5} \text{Ns} / \text{m}^2$

D. $1.7 \times 10^{-5} \text{Ns} / \text{m}^2$

Answer: C



Watch Video Solution

64. Critical velocity is given by

A. $v_C = \frac{N\eta}{\rho D}$

$$\text{B. } v_C = \frac{N\eta\rho}{D}$$

$$\text{C. } v_C = \frac{N\rho D}{\rho}$$

$$\text{D. } v_C = \frac{N}{\eta\rho D}$$

Answer: A



Watch Video Solution

65. If Reynold's number is greater than 3000, the flow of liquid is

A. laminar

B. turbulent

C. regular

D. irregular

Answer: B



Watch Video Solution

66. Water is flowing through a tube of diameter 1 cm at 8 cm/s. Taking $\eta = 10^{-2}$ poise the flow of liquid and Reynold's number are

A. streamline, 80

B. streamline, 800

C. strbulent, 8000

D. turbulent, 9000

Answer: B



Watch Video Solution

67. Bernoulli's principle is based on the law of conservation of

A. conservation of linear momentum.

B. conservation of energy.

C. conservation of kinetic energy.

D. conservation of angular momentum.

Answer: B



Watch Video Solution

68. Pain gun is based on _____.

A. Archimede's principle

B. Boyle's law

C. Bernoulli's principle

D. Newton's laws of motion

Answer: C



Watch Video Solution

69. Bernoulli's principle is true under which of the following (conditions) assumptions ?

A. The fluid is non-viscous and its flow is turbulent.

B. The fluid is non-viscous and its flow is streamline.

C. The fluid is viscous and its flow is turbulent.

D. The fluid is viscous and its flow is streamline.

Answer: B



Watch Video Solution

70. In case of the streamline flow of non-viscous and incompressible fluid, which of the following statement is CORRECT ?

A. The sum of pressure energy, K.E. per unit mass and P.E. per unit mass always remains constant at every point.

B. The sum of pressure energy, K.E. per unit mass and P.E. per unit mass is constant in the beginning.

C. The sum of pressure energy, K.E. per unit mass and P.E. per unit mass is not constant at every point.

D. The pressure energy is equal to sum of K.E. per unit mass and P.E. per unit mass at every point.

Answer: A



Watch Video Solution

71. If P is the pressure on a Liquid of area A and the liquid moves through a certain distance the work done per unit volume is _____.

A. Force

B. Density

C. Pressure

D. Pressure \times Area

Answer: C



Watch Video Solution

72. The velocity of efflux of liquid through orifice is equal to velocity which a body attains while falling freely from the free surface of liquid to the orifice. This is known as _____.

- A. Bernoulli's theorem
- B. Torricelli's law
- C. Stoke's law
- D. Pascal's law

Answer: B



Watch Video Solution

73. A cylinder of height 20 m is completely filled with water. The velocity of efflux of water through a hole on the side wall of the cylinder near its bottom is (Take $g = 10ms^{-2}$)

A. 10

B. 20

C. 25.5

D. 5

Answer: B



Watch Video Solution

74. In a vessel containing water, a hole is made at a depth of 0.10 m from the free surface.

What would be the velocity of efflux ?

A. $14m / s$

B. $4m / s$

C. $20.8m / s$

D. $1.4m / s$

Answer: D



Watch Video Solution

75. A device used for measuring the rate of flow of liquid through pipes is called.....

A. calorimeter q

B. speedometer

C. venturimeter

D. thermometer

Answer: C



Watch Video Solution

76. The venturimeter measures the pressure difference by measuring _____.

A. height difference

B. temperature difference

C. velocity difference

D. volume difference

Answer: A



Watch Video Solution

77. What is the minimum pressure required to force the blood from the heart to the top of the head (vertical distance $0.5m$) ? [density of blood is $1040 \text{ kg } m^{-3}$. Fraction is to be neglected and $g = 9.8ms^{-2}$]

A. $1050Nm^{-2}$

B. $2080Nm^{-2}$

C. $5096Nm^{-2}$

D. $6096Nm^{-2}$

Answer: C



Watch Video Solution

78. Air is streaming past a horizontal airplane wing such that its speed is 120 metre per sec over the upper surface and 90 metre per sec

at the lower surface. If the density of air is $1.3 \text{ kg per metre}^3$ and the wing is 10 metre long and has an average width of 2 metre , then the difference of the pressure on the two sides of the wing is :

A. 4095.0 pascal

B. 409.50 pascal

C. 40.950 pascal

D. 4.0950 pascal

Answer: A



79. In airfoil, the streamline are crowded above the wings of aeroplane more than those below it. This causes (during take off)

A. pressure to drop above wings and results in dynamic lift.

B. pressure to drop below wings and results in dynamic lift.

C. no difference in pressure.

D. to push the aeroplane.

Answer: A



Watch Video Solution

80. An airfoil has _____.

A. convex shape

B. concave shape

C. concavo-convex shape

D. plan e shape

Answer: A



Watch Video Solution

81. Which of the following is NOT an example of application of Bernoulli's principle ?

A. Venturimeter

B. Dynamic lift

C. Air purifier

D. Barometer

Answer: D



Watch Video Solution

82. Maximum acceleration of the train in which a box lying on its floor will remain stationary, is

$$[\mu_s = 0.15, g = 10m / s^2]$$

A. $70m / s^2$

B. $10.5m / s^2$

C. $7.1m / s^2$

D. $1.5m / s^2$

Answer: D



Watch Video Solution

83. If μ , N and S represent coefficient of friction, normal reaction and distance moved, then the general expression for work against friction is

A. μNS

B. $\frac{\mu N}{S}$

C. $\frac{\mu S}{N}$

D. $\frac{S}{\mu N}$

Answer: A



Watch Video Solution

84. Water is filled in a flask up to a height of 20 cm. The bottom of the flask is circular with radius 10 cm. If the atmospheric pressure is 1.01×10^5 Pa, find the force exerted by the

water on the bottom. Take $g = 10\text{ms}^{-2}$ and density of water $= 1000\text{kgm}^{-3}$.

A. $1.03\pi N$

B. $1.03\pi \times 10^3 N$

C. $1.03\pi \times 10^4 N$

D. $1.03\pi \times 10^5 N$

Answer: D



Watch Video Solution

85. In streamline flow velocity of liquid at the bottom layer is _____.

A. zero

B. maximum

C. mean of velocities of all layers

D. infinity

Answer: A



Watch Video Solution

86. Sudden fall of atmospheric pressure by a large amount indicates _____.

A. storm

B. fair weather

C. cold weather

D. calm weather

Answer: A



Watch Video Solution

1. Frictional force is _____.

- A. conservation force
- B. gravitational force
- C. electrostatic force
- D. non-conservative force

Answer: D



Watch Video Solution

2. Assertion: Friction is a conservative force.

Reason: Friction does not depend upon mass of the body.

A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.

Answer: D



Watch Video Solution

3. When one body tends to move or move over the another body, the opposition force depends upon _____.

A. surrounding temperature

B. temperature of the bodies

C. density of material

D. nature of the surfaces in contact

Answer: D



Watch Video Solution

4. When a moving body is suddenly stopped

A. frictional force increases.

B. frictional force gradually reduces to one.

C. frictional force becomes infinite.

D. the frictional force reduces to zero as it

is a self adjusting force.

Answer: D



Watch Video Solution

5. The ball bearings used in a machine are of different materials, this is because

A. adhesive forces are greater than cohesive forces.

B. adhesive forces and cohesive forces are equal in magnitude, opposite in

direction ,

C. adhesive forces are less than cohesive forces.

D. machine looks good, attractive.

Answer: C



Watch Video Solution

6. Assertion : When a bicycle is in motion, the force of friction exerted by the ground on the two wheels is always in forward direction.

Reason : The frictional force acts only when the bodies are in contact

A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.


Answer: D



Watch Video Solution

7. A block of mass M is resting on an inclined plane as shown in the figure. The inclination of the plane to the horizontal is gradually increased. It is found that when the angle of inclination is θ the block just begins to slide down the plane. What is the minimum force F applied parallel to the plane that would just

make the block move up the plane ?



- A. $Mg \sin \theta$
- B. $mg \cos \theta$
- C. $2Mg \cos \theta$
- D. $2Mg \sin \theta$


Answer: D



Watch Video Solution

8. A block of mass $m=2\text{kg}$ is resting on a rough inclined plane of inclination 30° as shown in figure. The coefficient of friction between the block and the plane is $\mu = 0.5$. What minimum force F should be applied perpendicular to the plane on the block, so that blocks does not slip on

the plane? ($g = 10\text{m} / \text{s}^2$)



A. zero

B. 2.67N

C. 4.34N

D. 6.24N

Answer: B



Watch Video Solution

9. If F_i and F_g are the frictional force between surface of contact from iron and glass material, then which of the following relation is correct ? Assume that there are same normal reaction on both the surfaces.

A. $G_g = F_i$

B. $F_g \neq F_i$

C. $F_g = 2F_i$


D. $F_i = 2F_g$

Answer: B



Watch Video Solution

10. A 4 kg block A is placed on the top of 8 kg block B which rests on a smooth table.



A just slips on B when a force of 12 N is applied

on A. Then the maximum horizontal force F applied on B to make both A and B move together, is

A. 12 N

B. 24 N

C. 36 N

D. 48 N

Answer: C



Watch Video Solution

11. Wooden block of weight W_1 is kept on a horizontal surface. A pan is attached to a string which passed over the pulley and other end of the string is attached to block. The weight F_1 is placed in the pan. If the block still remains stationary then

A. $F_1 = F_{2_1}$ and $W_1 = F_{2_1}$

B. $F_1 = N$ and $W_1 = \mu_s N$


C. $F_1 = F_{s_1}$ and $W_1 = N$

D. $F_1 = N$ and $W_1 = F_{s_1}$

Answer: C



Watch Video Solution



12.

A rough vertical board has an acceleration a so that a 2 kg block pressing against it does

not fall. The coefficient of friction between the block and the board should be

A. $> g/a$

B. $< g/a$

C. $= g/a$


D. $> ag$

Answer: A



Watch Video Solution

13. A block of mass m lying on a rough horizontal plane is acted upon by a horizontal force P and another force Q inclined at an angle θ to the vertical. The block will remain in equilibrium, if the coefficient of friction between it and the surface is



- A. $\frac{(P + Q \sin \theta)}{(mg + Q \cos \theta)}$
- B. $\frac{(P \cos \theta + Q)}{(mg - Q \sin \theta)}$

- C. $\frac{(P + Q \cos \theta)}{(mg + Q \sin \theta)}$
- D. $\frac{(P \sin \theta - Q)}{(mg - Q \cos \theta)}$

Answer: A



Watch Video Solution

14. A block of mass 10kg is placed on a rough horizontal surface having coefficient of friction $\mu = 0.5$. If a horizontal force of 100N is acting on it, then acceleration of the will be.

A. zero

B. $10ms^{-2}$

C. $5ms^{-2}$

D. $5.2ms^{-2}$

Answer: C



Watch Video Solution

15. The pressure at a point in a fluid is independent of _____.

A. area

B. force

C. height of liquid

D. direction

Answer: D



Watch Video Solution

16. Assertion : To empty an oil tank, two holes are made.

Reason : Oil will come out two holes so it will emptied faster.

A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.

Answer: C



Watch Video Solution

17. Two vessels have different base area. They are filled with water to the same height. If the amount of water in one be 6 times that in the other, then the ratio of pressure on their bottom will be

A. 16:1

B. 8:1

C. 4:1

D. 1:1

Answer: D



Watch Video Solution

18. The manual of a car instructs the owner to inflate the tyers to a pressure of 200 kPa, then absolute pressure is

A. 301 kPa

B. 200 kPa

C. 101 kPa

D. 99kPa

Answer: A



Watch Video Solution

19. What is the pressure on a swimmer 10m below the surface of lake? $g = 10ms^{-2}$, atmospheric pressure = $1.01 \times 10^5 Pa$

A. 4 atm

B. 1 atm

C. 2 atm

D. 3.8 atm

Answer: C



Watch Video Solution

20. In car lift compressed air exerts a force F_1 on a small piston having a radius of 5 cm. This pressure is transmitted to a second piston of radius 15 cm. If the mass of the car to be lifted is 1350 kg, what is F_1 ? What is the pressure necessary to accomplish this task?

A. $2.03 \times 10^5 Pa$

B. $1.87 \times 10^5 Pa$

C. $1.2 \times 10^5 Pa$

D. $2.42 \times 10^5 Pa$

Answer: A



Watch Video Solution

21. The densities vary very little over a wide range of pressure and temperature in case of _____ hence treated as incompressible.

A. gases

B. liquids

C. fluids

D. LPG


Answer: B



Watch Video Solution

22. Two pistons (areas A_1 and A_2) exerting forces F_1 and F_2 on a liquid exerts pressures P_1 and P_2 as shown in figure. The pistons are

at the same horizontal level and do not move under the influence of force, then



A. $F_2 = \frac{A_1}{A_2} F_1$

B. $F_2 = F_1$

C. $P_2 = \frac{A_1}{A_2} P_1$

D. $P_2 = P_1$

Answer: D



23. hydraulic brakes work on principle of

A. change in pressure results in stopping of wheel motion.

B. change in pressure is transmitted equally to wheels.

C. pressure is tensor quantity.

D. higher applied pressure stop the motion.

Answer: A



Watch Video Solution

24. Choost the CORRECT statement.

A. The liquid pressure is not the same at all points at the same depth.

B. The liquid pressure at all points at the same depth depends on shape of the liquid.

C. The liquid pressure at all points at the same depth depends upon surrounding environment.

D. The liquid pressure is the same at all points at the same depth.

Answer: C



Watch Video Solution

25. A liquid distributed by stirring comes to rest after some time due to its property of

A. surface tension

B. stability

C. viscosity

D. attraction forces between molecules

Answer: D



Watch Video Solution

26. When a fluid is through a tube, then the reason of viscous force acting between its different layers will be

A. transfer of energy from one layer to another.

B. transfer to momentum from one later to another.

C. equal velocity of the molecules.

D. changing density along with tube.

Answer: B



[Watch Video Solution](#)

27. Deep water runs slow. Explain.

A. rain

B. still

C. turbulent

D. river

Answer: C



[Watch Video Solution](#)

28. Water enters through end A with a speed v_1 and leaves through end B with a speed v_2 of cylindrical tube AB. The tube is always completely filled with water. In case I the tube is horizontal, in case II it vertical with the end A upward and in case III it is vertical with the end B upward. We have $v_1 = v_2$ for

A. Case I

B. Case II

C. Case III

D. Each case

Answer: D




Watch Video Solution

29. A pipe GB is fitted with two pipes C and D as shown in the figure. The pipe has area

$A = 24m^2$ at G and velocity of water at G is 10 m/s, and at C is 6 m/s. The velocity of water at

D is



A. $21m/s$

B. $3.3m/s$

C. $30m/s$

D. $2.1m/s$

Answer: A



Watch Video Solution

30. Two drops of equal size are falling vertically through air with a constant terminal velocity of $0.15\text{cm} / \text{s}$. What should be the velocity if these drops coalesce to form one drop ?

A. $0.15\text{cm} / \text{s}$

B. $0.15\sqrt{2}\text{cm} / \text{s}$

C. $0.15 \times 2^{1/3} \text{ cm / s}$

D. $0.15 \times 2^{2/3} \text{ cm / s}$

Answer: D



Watch Video Solution

31. In turbulent flow the velocity of the liquid molecules in contact with the walls of the tube.

A. is zero

B. is maximum

C. is equal to critical velocity

D. may have any value

Answer: D



Watch Video Solution

32. The tangential force of viscous drag on any layer of the liquid is directly proportional to the velocity gradient $\frac{dv}{dx}$. Then the direction of velocity gradient is

A. perpendicular to the direction of flow of the liquid.

B. parallel to the direction of the flow of the liquid.

C. opposite to the direction of the flow of the liquid.

D. independent of the direction of the flow of liquid.

Answer: A



Watch Video Solution

33. The relative velocity of two parallel layers of water is 8 cm/sec. If the perpendicular distance between the layers is 0.1 cm, then velocity gradient will be

- A. 40 per second
- B. 60 per second
- C. 80 per second
- D. 100 per second

Answer: C



[Watch Video Solution](#)

34. A good lubricant must have

- A. highly viscous only
- B. volatile in nature only
- C. low viscous
- D. highly viscous and low volatile

Answer: D



[Watch Video Solution](#)

35. Machine parts are jammed in winter. Why?

A. of low temperature

B. viscosity of lubricant increases

C. viscosity of lubricant decreases

D. of low pressure

Answer: B



Watch Video Solution

36. A sphere is dropped gently into a medium of infinite extent. As the sphere falls, the force acting downwards on it.

A. remains constant.

B. decreases for sometime and then becomes constant.

C. increases for sometime and then becomes constant.

D. decreases for sometime and then becomes zero.

Answer: D



Watch Video Solution

37. Rain drops fall from a great height under gravity. Check the correct statement

A. Their velocity continuously increases till they hit the earth will the same final velocity.

B. They fall with a terminal velocity which is different for drops of different sizes.

C. They fall with a terminal velocity which is the same for every drop.

D. Their velocity goes on increasing continuously till they hit the earth, and the final velocity of each drop is different.

Answer: B



Watch Video Solution

38. A solid sphere falls with a terminal velocity V in CO_2 gas. If its is allowed to fall in vacuum

A. terminal velocity of sphere $= v$.

B. terminal velocity of sphere $< v$.

C. terminal velocity of sphere $> v$.

D. sphere never attains terminal velocity.

Answer: D



Watch Video Solution

39. The maximum average velocity of water in a tube of diameter 2 cm so that the flow becomes laminar is [Viscosity of water is $10^{-3} Nm^{-2} s^{-1}$]

A. $1ms^{-1}$

B. $0.1ms^{-1}$

C. $10ms^{-1}$

D. $100ms^{-1}$

Answer: B



Watch Video Solution

40. Bernoulli's equation is ideally valid for

A. Constant, viscous, incompressible, temperature-dependent flow.

B. Variable, non-viscous, incompressible, temperature-dependent flow.

C. Constant, non-viscous, incompressible, temperature-independent flow.

D. Variable, non-viscous, incompressible,
temperature-independent flow.

Answer: C



Watch Video Solution

41. The pans of a physical balance are in equilibrium. Air is blown under the right hand pan, then the right hand pan will

A. move up

B. move down

C. move erratically

D. remain at the same level

Answer: B



Watch Video Solution

42. Assertion : Aeroplanes are made to run on the runway before take off, so that they acquire the necessary lift.

Reason : According to Bernoulli's theorem, as

velocity increases pressure decreases and viceversa.

- A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion
- B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion
- C. Assertion is True, Reason is False
- D. Assertion is False, Reason is False.

Answer: A



Watch Video Solution

43. Why two boats moving in parallel directions close to each other get attracted?

A. a streamline flow sets between the two.

B. the boats experience attractive pull towards each other.

C. the boats experience force of repulsion.

D. the whirlpool is formed at the mid way between two boats.

Answer: B



Watch Video Solution

44. Water flowing through a horizontal pipe line having a constriction. Then,

A. pressure will be the same throughout the length of the pipe.

B. pressure will be greater at the constriction.

C. pressure will be smaller in the wider portion.

D. pressure will be smaller at the constriction.

Answer: D



Watch Video Solution

45. A vessel is filled with water and kerosene oil. The vessel has a small hole in the bottom. Neglecting viscosity if the thickness of water layer is h_1 and kerosene layer is h_2 then the

velocity v of flow of water will be (density of water is ρ_1 g/cc and that of kerosene is ρ_2 g/cc)

A. $v = \sqrt{2g(h_1 + h_2)}$

B. $v = \sqrt{2g\left(h_1 + h_2 \frac{\rho_2}{\rho_1}\right)}$

C. $v = \sqrt{2g(h_1\rho_1 + h_2\rho_2)}$

D. $v = \sqrt{2g\left(h_1 \frac{\rho_2}{\rho_1} + h_2\right)}$

Answer: B



Watch Video Solution

46. The reading of pressure meter attached with a closed pipe is $3.5 \times 10^5 \text{ Nm}^{-2}$. On opening the valve of the pipe, the reading of the pressure meter is reduced to $3.0 \times 10^5 \text{ Nm}^{-2}$. Calculate the speed of the water flowing in the pipe.

A. $100 \text{ m} / \text{s}$

B. $50 \text{ m} / \text{s}$

C. $10 \text{ m} / \text{s}$

D. $0.1 \text{ m} / \text{s}$

Answer: C



Watch Video Solution

47. In a venturimeter, _____ remains unchanged along the axis.

A. K.E. of flowing liquid

B. pressure energy of flowing liquid

C. P.E. of flowing liquid

D. Both K.E. and P.E.

Answer: C



Watch Video Solution

48. The speed of the efflux from an open tank is identical to that of _____

- A. a freely falling body
- B. a body moving with varying velocity
- C. a body moving tangentially
- D. a body at rest

Answer: A



Watch Video Solution

49. The velocity of efflux of an ideal liquid does not depend on

- A. acceleration due to gravity
- B. height of the liquid level in the vessel
- C. viscosity of the liquid
- D. Both (B) and (C)


Answer: D



Watch Video Solution

50. An L-shaped tube with a small orifice is held in a water stream as shown in fig. the upper end of the tube is 10.6cm above the surface of water. What will be the height of the jet of water coming from the orifice?

(Velocity water steam is 2.45 m/s)



- A. Zero
- B. 10.6 cm
- C. 19.4 cm
- D. 20.0 cm

Answer: C



Watch Video Solution

51. A cylindrical tank has a hole of 1cm^2 in its bottom. If the water is allowed to flow into the tank from a tube above it at the rate of $70\text{cm}^3 / \text{sec}$, then the maximum height up to which water can rise in the tank is

A. 0.25cm

B. 2.5cm

C. 5cm

D. 10cm

Answer: B



Watch Video Solution

52. With increase in temperature, friction _____.

A. increases

B. decreases

C. remains unchanged

D. may increases or decrease

Answer: B



Watch Video Solution

53. The wheels are circular in shape because

A. they required less material.

B. circular wheels are frictionless.

C. they are attractive.

D. rolling friction is least.

Answer: D



Watch Video Solution

54. A body is moving along a straight horizontal road with a velocity of 21 m/s comes to rest after moving a certain distance. If μ between the body and the surface of road is 0.3, the distance covered is $[g = 9.8ms^{-2}]$

A. 100 m

B. 80 m

C. 75 m

D. 60 m

Answer: C



Watch Video Solution

55. A stone weighing 1 kg and sliding on ice with a velocity of 2 m / s is stopped by friction in 10 sec . The force of friction (assuming it to be constant) will be

A. -20 N

B. -0.2 N

C. 0.2 N

D. 20 N

Answer: B



Watch Video Solution

56. The coefficient of viscosity for hot air is

- A. greater than the coefficient of viscosity for cold air.
- B. smaller than the coefficient of viscosity for cold air.
- C. same as the coefficient of viscosity for cold air.
- D. increases or decreases depending on the external pressure.

Answer: A



Watch Video Solution

57. A car is moving along a straight horizontal road with a speed v_0 . If the coefficient of friction between the tyres and the road is μ , the shortest distance in which the car can be stopped is

A. $\frac{v}{2\mu g}$

B. $\frac{v^2}{2\mu g}$

C. $\left(\frac{v^2}{2\mu g}\right)^{1/2}$

D. $\left(\frac{v^2}{2\mu g}\right)^2$


Answer: B



Watch Video Solution

58. The coefficient of friction between m_2 and inclined plane is μ (shown in the figure). If

$$\frac{m_1}{m_2} = \sin \theta \text{ then}$$



- A. no motion takes place.
- B. m_1 moves downward.
- C. m_1 moves upward.
- D. no sufficient information.

Answer: A



Watch Video Solution

59. A vehicle of mass m is moving on a rough horizontal road with momentum P . If the

coefficient of friction between the tyres and the road be μ , then the stopping distance is:

A. $\frac{P}{2\mu mg}$

B. $\frac{P^2}{2\mu mg}$

C. $\frac{P}{2\mu m^2 g}$

D. $\frac{P^2}{2\mu m^2 g}$

Answer: D



Watch Video Solution

60. A 60 kg body is pushed with just enough force to start it moving across a floor and the same force continues to act afterwards. The coefficient of static friction and sliding friction are 0.5 and 0.4 respectively. The acceleration of the body is

A. $6m / s^2$

B. $4.9m / s^2$

C. $3.92m / s^2$

D. $1m / s^2$

Answer: D



Watch Video Solution

61. A ball of radius r and density ρ falls freely under gravity through a distance h before entering water. Velocity of ball does not change even on entering water. If viscosity of water is η the value of h is given by

A. $\frac{2}{9}r^2g\left(\frac{1-\rho}{\eta}\right)$

B. $\frac{2}{81}r^2g\left(\frac{\rho-1}{\eta}\right)$

C. $\frac{2}{81} r^4 g \left(\frac{\rho - 1}{\eta} \right)^2$

D. $\frac{2}{9} r^4 g \left(\frac{\rho - 1}{\eta} \right)^2$

Answer: C



Watch Video Solution

62. In the houses far away from the municipal water tanks often water does not rise to the top floor. This happens because

A. viscosity of water is vary high.

B. water wets the pipes.

C. of changes in the cross-sectional area of
the pipe.

D. of loss of pressure during the flow of
water.

Answer: D



Watch Video Solution

63. A plane is in level flight at constant speed and each of its two wings has an area of $20m^2$. If the speed of the air is 180 km/h over the lower wing and 216 km/h over the upper wing surface, determine the plane's mass. (Take air density to be $1kgm^{-3}$ and $g = 10m / s^2$).

A. 2200 kg

B. 1100 kg

C. 1785 kg

D. 2750 kg

Answer: A



Watch Video Solution

64. For a fluid in a steady flow, the increase in flow speed at a constriction follows _____.

A. conservation of mass and angular momentum

B. conservation of mass and Bernoulli's principle

C. conservation of velocity

D. Torricelli's theorem

Answer: B



Watch Video Solution

65. A cylindrical tank is filled with water to a level of 4 m. A hole is opened at a height of 60 cm from bottom. The ratio of the area of the

hole to that at cross-sectional area at cylinder is 0.2. Then the velocity with which water is coming out is ($g = 10 \text{ m/s}^2$)

A. 7.9 m/s

B. 9.2 m/s

C. 8.4 m/s

D. 8.9 m/s

Answer: C



Watch Video Solution

66. Assertion: The acceleration of a body down a rough inclined plane is greater than the acceleration due to gravity.

Reason: The body is able to slide on an inclined plane only when its acceleration is greater than acceleration due to gravity.

A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.

Answer: D



Watch Video Solution

Mcq Competitive Thinking

1. Which one of the following is NOT used to reduce friction ?

A. Oil

B. Ball bearings

C. Sand

D. Graphite

Answer: C



Watch Video Solution

2. STATEMENT-1: It is easier to pull a heavy object than to push it on a level ground and

STATEMENT-2: The magnitude fo frictional

force depends on the nature of the two surfaces in contact.

A. Assertion is True, Reason is True, Reason

is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason

is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.

Answer: B



Watch Video Solution

3. If a ladder weighting $250N$ is placed against a smooth vertical wall having coefficient of friction between it and floor 0.3 , then what is the maximum force of friction available at the point of contact between the ladder and the floor?

A. 75 N

B. 50 N

C. 35 N

D. 25 N


Answer: A



Watch Video Solution

4. The coefficient of static friction, μ_s between block A of mass 2 kg and the table as shown in the figure is 0.2. What would be the maximum mass value of block B so that the two blocks do not move? The string and the pulley are assumed to be smooth and massless.

$$(g = 10 \text{ m/s}^2)$$



A. 2.0 Kg

B. 4.0 Kg

C. 0.2 Kg

D. 0.4 Kg

Answer: D



5. A lift is moving down with an acceleration equal to the acceleration due to gravity. A body of mass M kept on the floor of the lift is pulled horizontally. If the coefficient of friction is μ then the frictional resistance offered by the body is .

A. mg

B. μmg

C. $2\mu mg$

D. Zero

Answer: D



Watch Video Solution

6. Maximum acceleration of the train in which a 50 kg box lying on its floor will remain stationary is (Given: Co-efficient of static friction between the box and the train's floor is 0.3 and $g = 10m s^{-2}$)

A. $5.0m s^{-2}$

B. $3.0ms^{-2}$

C. $1.5ms^{-2}$

D. $15ms^{-2}$


Answer: B



Watch Video Solution

7. A block B is pushed momentarily along a horizontal surface with an initial velocity v . If μ is the coefficient of sliding friction between B and the surface, block B will come

to rest after a time:



A. $\frac{v}{\mu g}$

B. $\frac{vg}{\mu}$

C. $\frac{v\mu}{g}$

D. $\frac{\mu g}{v}$

Answer: A



Watch Video Solution

8. A block rests on a rough inclined plane making an angle of 30° with the horizontal. The coefficient of static friction between the block and the plane is 0.8. If the frictional force on the block is 10N, the mass of the block (in kg) is

A. 1 kg

B. 2 kg

C. 3 kg

D. 4 kg

Answer: B



Watch Video Solution

9. A wooden box of mass $8kg$ slides down an inclined plane of inclination 30° to the horizontal with a constant acceleration of $0.4ms^{-2}$. What is the force of friction between the box and inclined plane? ($g = 10m/s^2$).

A. $12.2N$

B. $24.4N$

C. $36.8N$

D. $48.8N$

Answer: C



Watch Video Solution

10. A wooden box lying at rest on an inclined surface of a wet wood is held at static equilibrium by a constant force \vec{F} applied perpendicular to the incline. If the mass of the box is 1 kg, the angle of inclination is 30° and

the coefficient of static friction between the box and the inclined plane is 0.2, the minimum magnitude of \vec{F} is (Use $g = 10\text{m/s}^2$)


- A. 0N , as 30° is less than angle of repose
- B. $\geq 1\text{N}$
- C. $\geq 3.3\text{N}$
- D. $\geq 16.3\text{N}$

Answer: D



Watch Video Solution

11. What is the maximum value of the force F such that the block shown in the arrangement, does not move?



A. 20 N

B. 10 N

C. 12 N

D. 15 N

Answer: A



Watch Video Solution


12. A system consists of three masses m_1 , m_2 and m_3 connected by a string passing over a pulley P. The mass m_1 hangs freely and m_2 and m_3 are on a rough

horizontal table (the coefficient of friction= μ).

The pulley is frictionless and of negligible

mass. The downward acceleration of mass m_1

is (assume, $m_1 = m_2 = m_3 = m$)



A. $\frac{g(1 - g\mu)}{9}$

B. $\frac{2g\mu}{3}$

C. $\frac{g(1 - 2\mu)}{3}$

D. $\frac{g(1 - 2\mu)}{2}$

Answer: C



Watch Video Solution

13. Maximum value of static friction is .

- A. limiting friction.
- B. rolling friction.
- C. normal reaction.
- D. coefficient of friction.

Answer: A



Watch Video Solution

14. Which of the following statements is incorrect ?

A. Rolling friction is smaller than sliding friction.

B. Limiting value of static friction is directly proportional to normal reaction.

C. Frictional force opposes the relative motion.

D. Coefficient of sliding friction has dimensions of length.

Answer: D



Watch Video Solution

15. A body of mass M is kept on a rough horizontal surface (friction coefficient $= \mu$).

A person is trying to pull the body by applying

a horizontal force but the body is not moving.

The force by the surface on A is F , where

A. $F = mg$

B. $F = \mu Mgf$

C. $Mg \leq F \leq Mg\sqrt{1 + \mu^2}$

D. $Mg \geq F \geq Mg\sqrt{1 + \mu^2}$

Answer: C



Watch Video Solution

16. Consider a frictionless ramp on which a smooth object is made to slide down from an initial height 'h'. The distance 'd' necessary to stop the object on a flat track (or coefficient of friction ' μ '), kept at the ramp end is

A. h / μ

B. μh

C. $\mu^2 h$


D. $h^2 h$

Answer: A



Watch Video Solution

17. Block B lying on a table weighs w . The coefficient of static friction between the block and the table is μ . Assume that the cord between B and the knot is horizontal. The maximum weight of the block A for which the system will be stationary is



A. $\frac{W \tan \theta}{\mu}$

B. $\mu W \tan \theta$

C. $\mu W \sqrt{1 + \tan^2 \theta}$

D. $\mu W \sin \theta$

Answer: B



Watch Video Solution

18. A heavy uniform chain lies on a horizontal table-top. If the coefficient of friction between the chain and table surface is 0.25, then the

maximum fraction of length of the chain, that
can hang over one edge of the table is

A. 20 %

B. 25 %

C. 30 %


D. 40 %

Answer: A



Watch Video Solution

19. A block A with mass 100kg is resting on another block B of mass 200kg . As shown in figure a horizontal rope tied to a wall hold it. The coefficient of friction between A and B is 0.2 while coefficient of friction between B and the ground is 0.3 . the minimum required force F to start moving B will be.



A. 900 N

B. 100 N

C. 1100 N

D. 1200 N

Answer: C




Watch Video Solution

20. A block of mass 2kg is kept on the floor.

The coefficient of static friction is 0.4 . If a

force F of 2.5N is applied on the block as

shown in the figure, the frictional force between the block and the floor will be.



A. 2.5 N

B. 5 N

C. 7.84 N

D. 10 N


Answer: A



Watch Video Solution

21. Given in figure are two blocks A and B of weight 20N and 100N, respectively. These are being pressed against a wall by a force F as shown. If the coefficient of friction between the blocks is 0.1 and between block B and the wall is 0.15, the frictional force applied by the

wall on block B is:



A. 100 N

B. 80 N

C. 120 N

D. 150 N

Answer: C



Watch Video Solution

22. A block of mass m is lying on horizontal surface of coefficient of friction μ . A force is applied to the block at an angle θ with the horizontal. The block will move with a minimum force F if

A. $\mu = \tan \theta$

B. $\mu = \cos \theta$

C. $\mu = \sin \theta$

D. $\mu = \cot \theta$

Answer: A



Watch Video Solution

23. A marble block of mass 2 kg lying on ice when given a velocity of $6m/s$ is stopped by friction in 10s. Then the coefficient of friction is

A. 0.01

B. 0.02

C. 0.03

D. 0.06

Answer: D



Watch Video Solution

24. A block of mass m_2 is placed on a horizontal table and another block of mass m_1 is placed on top of it. An increasing horizontal

force $F = \alpha t$ is exerted on the upper block but the lower block never moves as a result. If the coefficient of friction between the blocks is μ_1 coefficient of friction between the blocks is μ_1 and that between the lower block and the table is μ_2 , then what is the maximum possible value of μ_1 / μ_2 ?

A. $\frac{m_2}{m_1}$

B. $1 + \frac{m_2}{m_1}$

C. $\frac{m_1}{m_2}$


D. $1 + \frac{m_1}{m_2}$

Answer: B



Watch Video Solution

25. A block of mass $m = 5\text{kg}$ is resting on a rough horizontal surface for which the coefficient of friction is 0.2 . When a force $F = 40\text{N}$ is applied, the acceleration of the block will be ($g = 10\text{m/s}^2$).



A. $5.73m / s^2$

B. $8.0m / s^2$

C. $9.73m / s^2$

D. $10.0m / s^2$

Answer: A




Watch Video Solution

26. A body of mass 20 kg is moving on a rough horizontal plane. A block of mass 3 kg is connected to the 20 kg mass by a string of

negligible mass through a smooth pulley as shown in the figure. The coefficient of kinetic friction between the heavier mass and the surface is

$(g = 10 \text{ m/s}^2)$.



A. 0.025

B. 0.035

C. 0.35

D. 0.25

Answer: B



Watch Video Solution

27. A block A of mass m_1 rests on a horizontal table. A light string connected to it passes over a frictionless pulley at the edge of table and from its other end another block B of mass m_2 is suspended. The coefficient of kinetic friction between the block and table is

μ_k . When the block A is sliding on the table, the tension in the string is.

A. $\frac{(m_2 + \mu_k m_1)g}{(m_1 + m_2)}$

B. $\frac{(m_2 - \mu_k m_1)g}{(m_1 + m_2)}$

C. $\frac{m_1 m_2 (1 + \mu_k)g}{(m_1 + m_2)}$

D. $\frac{m_1 m_2 (1 - \mu_k)g}{(m_1 + m_2)}$


Answer: C



Watch Video Solution

28. Two masses $m_1 = 5\text{kg}$ and $m_2 = 10\text{kg}$, connected by an inextensible string over a frictionless pulley, are moving as shown in the figure. The coefficient of friction of horizontal surface is 0.15. The minimum weight m that should be put on top of m_2 to stop the

motion is :-



A. 43.3 kg

B. 10.3 kg

C. 18.3 kg

D. 27.3 kg

Answer: D



Watch Video Solution

29. Air is blown through a hole on a closed pipe containing liquid. Then the pressure will

- A. increase on sides.
- B. increase downwards.
- C. increase in all directions.
- D. never increase.

Answer: C



Watch Video Solution

30. The pressure at the bottom of a tank of liquid is not proportional to

- A. Acceleration due to gravity
- B. Density of the liquid
- C. Height of the liquid
- D. Area of the liquid surface

Answer: D



Watch Video Solution

31. If the atmospheric pressure is P_a then the pressure P at depth a below the surface of a liquid of density ρ open to the atmosphere is

A. $P_a - \frac{\rho gh}{2}$

B. $P_a - \rho gh$

C. $P_a + \rho gh$

D. $P_1 + \frac{\rho gh}{2}$

Answer: C



Watch Video Solution

32. By sucking a straw a student can reduce the pressure in his lungs to 750mm of Hg (density) $= 13.6\text{kg}/\text{cm}^3$) Using the straw, he can drink water from a glass up to a maximum depth of :

A. 10 cm

B. 75 cm

C. 13.6 cm

D. 1.36 cm

Answer: C



Watch Video Solution

33. If pressure at half the depth of a lake is equal to $\frac{2}{3}$ pressure at the bottom of the lake then what is the depth of the lake ?

A. 10 m

B. 20 m

C. 60 m

D. 30 m

Answer: B



Watch Video Solution

34. The pressure on a swimmer 10 m below the surface of lake is _____ ($g = 10\text{m} / \text{s}^2$, atmospheric pressure = $1.01 \times 10^5 \text{Pa}$)

A. 3

B. 1

C. 2

D. zero


Answer: C



Watch Video Solution

35. A U-tube with both ends open to the atmosphere is partially filled with water. Oil, which is immiscible with water. Is poured into

one side until it stands at a distance of 10mm above the water level on the other side. Meanwhile the water rises by 65mm from its original level (see diagram). The density of the oil is:



A. 650kgm^{-3}

B. 425kgm^{-3}

C. 800kgm^{-3}

D. 928kgm^{-3}


Answer: D



Watch Video Solution

36. A vertical U-tube of uniform inner cross section contains mercury in both sides of its arms. A glycerin (density = 1.3g/cm^3) column of length 10 cm is introduced into one

of its arms. Oil of density 0.8 gm/cm^3 is poured into the other arm until the upper surfaces of the oil and glycerin are in the same horizontal level. Find the length of the same horizontal level. Find the length of the oil column, Density of mercury = 13.6 g/cm^3



A. 10.4 cm

B. 8.2cm

C. 7.2cm

D. 9.6cm

Answer: D



Watch Video Solution

37. A manometer connected to a closed tap reads 4.5×10^5 pascal. When the tap is opened the reading of the manometer falls is

4×10^5 pascal. Then the velocity of flow of water is

A. $7ms^{-1}$

B. $8ms^{-1}$

C. $9ms^{-1}$

D. $10ms^{-1}$

Answer: D



Watch Video Solution

38. Clouds appear to float in air due to

A. viscosity of air.

B. surface tension.

C. gravity.

D. elasticity.

Answer: A



Watch Video Solution

39. Viscosity is a transport phenomenon explained using the concept of transfer of

A. mass alone is transported.

B. energy alone is transported.

C. mass and energy are transported.

D. momentum is transported.

Answer: D



Watch Video Solution

40. More viscous oil is used in summer than in winter in motors due to

A. the rise in temperature in summer, the viscosity of oil decreases.

B. the rise in temperature in summer, the viscosity of oil increases.

C. the decrease in surface tension of oil.





D. the increase in surface tension of oil.

Answer: A



Watch Video Solution

41. A viscous fluid is flowing through a cylindrical tube. The velocity distribution of the fluid is best represented by the diagram



Answer: C



Watch Video Solution

42. In stream line flow of liquid, the total energy of liquid constant at _____.

- A. all points
- B. inner points
- C. outer points
- D. none of these

Answer: A



Watch Video Solution

43. Statement-1 : The stream of water flowing at high speed from a garden hose pipe tends to spread like a fountain when held vertically up, but tends to narrow down when held vertically down.

Statement-2 : In any steady flow of an incompressible fluid, the volume flow rate of the fluid remain constant.


- A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion
- B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion
- C. Assertion is True, Reason is False
- D. Assertion is False, Reason is False.

Answer: A



Watch Video Solution

44. When a body falls in air, the resistance of air depends to a great extent on the shape of the body, 3 different shapes are gives. Identify the combination of air resistances which truly represents the physical situation. (the cross sectional areas are the same).



(1)

Disc



(2)

Ball



(3)

Cigar shaped

A. $1 < 2 < 3$

B. $2 < 3 < 1$

C. $3 < 2 < 1$

D. $3 < 1 < 2$

Answer: C



Watch Video Solution

45. Water is flowing through a tube of non-uniform cross-section ratio of the radius at entry and exit end of the pipe is 3:2. Then the ratio of velocities at entry and exit of liquid is

A. 8 : 27

B. 4 : 9

C. 1 : 1

D. 9 : 4


Answer: B



Watch Video Solution

46. An ideal fluid flows through a pipe of circular cross section with diameters 5 cm and 10 cm as shown. The ratio of velocities of fluid

at A and B is



A. 4 : 1

B. 1 : 4

C. 2 : 1

D. 1 : 2

Answer: A



Watch Video Solution

47. The cylindrical tube of a spray pump has radius R , one end of which has n fine holes, each of radius r . If the speed of the liquid in the tube is V , the speed of the ejection of the liquid through the holes is:

A. $\frac{V^2 R}{nr}$

B. $\frac{VR^2}{n^2 r^2}$

C. $\frac{VR^2}{nr^2}$

D. $\frac{VR^2}{n^3 r^2}$

Answer: C



Watch Video Solution

48. Water is flowing continuously from a tap having an internal diameter $8 \times 10^{-3} \text{m}$. The water velocity as it leaves the tap is 0.4ms^{-1} . The diameter of the water stream at a distance $2 \times 10^{-1} \text{m}$ below the tap is close to ($g = 10 \text{m} / \text{s}^2$)

A. $5.0 \times 10^{-3} \text{m}$

B. $7.5 \times 10^{-3} \text{m}$

C. $9.6 \times 10^{-3} \text{m}$

D. $3.6 \times 10^{-3}m$

Answer: D



Watch Video Solution

49. Velocity of water in a river is

A. same everywhere.

B. more in the middle and less near its
banks.

C. less in the middle and more near its banks.

D. increases from one bank to other bank.

Answer: B



Watch Video Solution

50. The velocity of water in river is 9 km/h of the upper surface . The river is 10 m deep . If the coefficient of viscosity of water is 10^{-2}

poise then the shearing stress between horizontal layers of water is

A. $0.25 \times 10^{-2} N/m^2$

B. $0.25 \times 10^{-3} N/m^2$

C. $0.5 \times 10^{-3} N/m^2$

D. $0.75 \times 10^{-3} N/m^2$

Answer: B



Watch Video Solution

51. A metal plate of area 500cm^2) is kept on a horizontal surface with a layer of oil of thickness 0.5 mm between them. The horizontal force required to drag the plate with a velocity of 2 cm/s is
(coefficient of viscosity = 0.9 kg/ms)

A. 180 N

B. 18 N

C. 0.018 N

D. 1.8 N

Answer: D



Watch Video Solution

52. Spherical balls of radius 'R' are falling in a viscous fluid of viscosity ' η ' with a velocity 'v'. The retarding viscous force acting on the spherical ball is

- A. directly proportional to r but inversely proportional to v.
- B. directly proportional to both r and v.

C. inversely proportional to both r and v .

D. inversely proportional to r but directly proportional to velocity v .

Answer: B



Watch Video Solution

53. A gas bubble of 2 cm diameter rises through a liquid 1.75gcm^{-3} with a fixed speed of 0.35cm s^{-1} . Neglect the density of

the gas. The coefficient of viscosity of the liquid is

A. 870 poise

B. 1120 poise

C. 982 poise

D. 1089 poise

Answer: D



Watch Video Solution

54. What will be the approximate terminal velocity of a rain drop of diameter $1.8 \times 10^{-3} m$, when density of rain water $\approx 10^3 kgm^{-3}$ and the co-efficient of viscosity of air $\approx 1.8 \times 10^{-5} Nsm^{-2}$? (Neglect buoyancy of air.)

A. $49ms^{-1}$ —

B. $98ms^{-1}$

C. $392ms^{-1}$

D. $980ms^{-1}$

Answer: B



Watch Video Solution

55. A solid ball of volume V is dropped in a viscous liquid. It experiences a viscous force F . If the solid ball of volume $2V$ of same material is dropped in the same fluid, then the viscous force acting on it will be

A. $\eta F / 2$

B. $F / 2$

C. $2F$

D. $2nF$


Answer: C



Watch Video Solution

56. A small spherical solid ball is dropped from a great height in a viscous liquid . Its journey in the liquid is best described in the diagram

given below by the



A. Curve A

B. Curve B

C. Curve C

D. Curve D

Answer: D



Watch Video Solution

57. A small metal ball of mass m is dropped in a liquid contained in a vessel, attains a terminal velocity v . If a metal ball of same material but of mass $8m$ is dropped in same liquid then the terminal velocity will be

A. v

B. $2v$

C. $4v$

D. $8v$

Answer: A



Watch Video Solution

58. A small sphere of mass m is dropped from a height H . After it has fallen 100 m it has attained its terminal velocity and continues to fall at that speed. The work done by air friction against the sphere during the first 100 m of fall is-

- A. greater than the work done by air friction in the second 100 m.
- B. less than the work done by air friction in the second 100 m.
- C. equal to 100 mg.
- D. greater than 100 mg.

Answer: B



Watch Video Solution

59. The terminal speed of a sphere of gold (density = 19.5 kg m^{-3}) is 0.2 ms^{-1} in a viscous liquid (density = 1.5 kg m^{-3}). Then, the terminal speed of a sphere of silver (density = 10.5 kg m^{-3}) of the same size in the same liquid is

A. 0.2 ms^{-1}

B. 0.4 ms^{-1}

C. 0.133 ms^{-1}

D. 0.1 ms^{-1}

Answer: D



Watch Video Solution

60. Two solid spheres of same metal but of mass M and $8M$ fall simultaneously on a viscous liquid and their terminal velocities are v and $n v$, then value of n is

A. 16

B. 8

C. 4

D. 2

Answer: C



Watch Video Solution

61. Two spheres of equal masses but radii r_1 and r_2 are allowed to fall in a liquid of infinite are allowed to fall in a liquid of infinite column. The ratio of their terminal velocities are

A. 1

B. $r_1 : r_2$

C. $r_2 : r_1$

D. $\sqrt{r_1} : \sqrt{r_2}$

Answer: C



Watch Video Solution

62. Two drops of the same radius are falling through air with a steady velocity of 5cm s^{-1} .

If the two drops coalesce, the terminal velocity would be

A. 2.5cm s^{-1}

B. 10cm s^{-1}

C. $5\sqrt{2}\text{cm s}^{-1}$

D. $5 \times 4^{1/3}\text{cm s}^{-1}$

Answer: D



Watch Video Solution

63. The onset of turbulence in a liquid is determined by

- A. pascal's law.
- B. Stocke's law.
- C. Reynold's number.
- D. Torriclli's law.

Answer: C



Watch Video Solution

64. The dimensions of Reynold's constant are

A. $[L^0 m^0 T^0]$

B. $[L^{-1}M^1T^{-1}]$

C. $[L^{-1}M^1T^{-2}]$

D. $[L^{-2}M^1T^{-1}]$

Answer: A



Watch Video Solution

65. The speed at which the flow of water in a long cylindrical pipe of diameter 2 cm becaomes turbulent is

(The viscosity of water $= 1 \times 10^{-3}$ Pa. s and

for the onset of turbulent flow in a long cylindrical pipe, Reynolds number =3000)

A. $0.6m / s$

B. $0.45m / s$

C. $0.3m / s$

D. $0.15m / s$

Answer: D



Watch Video Solution

66. A horizontal pipeline carrying gasoline has a cross-sectional diameter of 5 mm. If the viscosity and density of the gasoline are 6×10^{-3} poise and 720 kg/m^3 respectively, the velocity after which the flow becomes turbulent is

- A. $> 1.66 \text{ m/s}$
- B. $> 3.33 \text{ m/s}$
- C. $> 1.6 \times 10^{-3} \text{ m/s}$
- D. $> 0.33 \text{ m/s}$

Answer: D



Watch Video Solution

67. The water flows from a tap of diameter 1.25 cm with a rate of $5 \times 10^{-5} m^3 s^{-1}$. The density and coefficient of viscosity of water are $10^3 kg m^{-3}$ and $10^{-3} Pa \cdot s$ respectively. The flow of water is

A. steady with Reynold's number 5100

B. turbulent with Reynold's number 5100

C. steady with Reynold's number 3900

D. turbulent with Reynold's number 3900

Answer: B



Watch Video Solution

68. Consider the following equation of Bernoulli's theorem.

$$P + \frac{1}{2}\rho v^2 + \rho gh = K \text{ (constant)}$$

The dimensions of K/P are same as that of which of the following

A. Thrust

B. Pressure

C. Angle

D. Velocity

Answer: C



Watch Video Solution

69. Water flows steadily through a horizontal pipe of variable cross-section. If the pressure of water is p at a point where the velocity of

flow is v , at another point (pressure p'), where the velocity of flow is $2v$: The following statements are given below for P' is

A. $P + 2\rho v^2$

B. $P - 2\rho v^2$

C. $P + \frac{3}{2}\rho v^2$

D. $P - \frac{3}{2}\rho v^2$

Answer: D



Watch Video Solution

70. A manometer connected to a closed tap reads 4.5×10^5 pascal. When the tap is opened the reading of the manometer falls to 4×10^5 pascal. Then the velocity of flow of water is

A. $7ms^{-1}$

B. $8ms^{-1}$

C. $9ms^{-1}$

D. $10ms^{-1}$

Answer: D



Watch Video Solution

71. A wind with speed 40 m/s blows parallel to the roof of a house. The area of the roof is 250 m^2 . Assuming that the pressure inside the house is atmospheric pressure, the force exerted by the wind on the roof and the direction of the force will be :

$$(\rho_{\text{air}} = 1.2\text{ kg/m}^3)$$

A. $(\rho_{\text{air}} = 1.2\text{ kg/m}^3)$

B. $4.8 \times 10^5\text{ N}$, upwards.

C. $2.4 \times 10^5\text{ N}$, upwards.

D. $2.4 \times 10^5 N$, downwards.


Answer: C



Watch Video Solution

72. Water is filled in a cylindrical container to a height of $3m$. The ratio of the cross-sectional area of the orifice and the beaker is 0.1 . The square of the speed of the liquid coming out

from the orifice is ($g = 10\text{m} / \text{s}^2$).



- A. $50\text{m}^2 / \text{s}^2$
- B. $50.5\text{m}^2 / \text{s}^2$
- C. $51\text{m}^2 / \text{s}^2$
- D. $52\text{m}^2 / \text{s}^2$

Answer: A



Watch Video Solution

73. A cylindrical tank having cross-sectional area A is filled with water to a height of 2.0 m. A circular hole of cross-section area a is opened at a height of 75 cm from the bottom. If $a/A = \sqrt{0.2}$, then velocity with which water emerges from the hole is

$$(g = 9.8 \text{ m s}^{-2})$$

A. 4.9 m s^{-1}

B. 4.95 m s^{-1}

C. $5.0ms^{-1}$

D. $5.5ms^{-1}$

Answer: D



Watch Video Solution

74. A cylindrical vessel of 100 cm height is kept filled upto the brim. It has four holes 1,2,3,4 which are respectively at heights of 27 cm, 30 cm , 50 cm and 80 cm from the horizontal floor. The water falling at the maximum

horizontal distance from the vessel comes from

A. Hole number 4

B. Hole number 3

C. Hole number 2

D. Hole number 1

Answer: B




Watch Video Solution

75. A cylinder of height h is filled with water and is kept on a block of height $\frac{h}{2}$. The level of water in the cylinder is kept constant. Four holes numbered 1, 2, 3, and 4 are at the side of the cylinder and at heights 0 , $\frac{h}{4}$, $\frac{h}{2}$ and $\frac{3h}{4}$ respectively. When all four holes are opened together, the hole from which water will reach farthest distance on the plane

PQ is the hole no.

- (A) 1
- (B) 2
- (C) 3
- (D) 4



A. 1

B. 2

C. 3

D. 4


Answer: B



Watch Video Solution

76. A roller is made by joining together two cones at their vertices O , it is kept on two rails AB and CD , which are placed asymmetrically with its axis perpendicular to CD and its center O at the centre of line joining AB and Cd it is given a light push so that it starts rolling with its centre O moving parallel to CD in the direction shown As it moves, the roller will

tand to:



A. turn right.

B. go straight.

C. turn left and right alternately.

D. turn left.

Answer: D



Watch Video Solution

77. The coefficient of static and dynamic friction between a body and the surface are 0.75 and 0.5 respectively. A force is applied to the body to make it just slide with a constant acceleration which is

A. $\frac{g}{4}$

B. $\frac{g}{2}$

C. $\frac{3g}{2}$

D. g

Answer: A



Watch Video Solution

78. A body of mass M is resting on a rough horizontal plane surface the coefficient of friction being equal to μ At $t = 0$ a horizontal force $F = F_0 t$ starts acting on it , where F_0 is a constant find the time T at which the motion starts?

A. $\frac{\mu M g}{F_0}$

B. $\frac{Mg}{\mu F_0}$

C. $\frac{\mu F}{Mg}$

D. $\frac{F}{F_0}$


Answer: A



Watch Video Solution

79. A plank with a box on it at one end is gradually raised about the other end. As the angle of inclination with the horizontal reaches 30° , the box starts to slip and slide $4.0m$

down the plank in $4.0s$. The coefficients of static and kinetic friction between the box and the plank will be, respectively.



- A. 0.4 and 0.3
- B. 0.6 and 0.6
- C. 0.6 and 0.5

D. 0.5 and 0.6

Answer: C



Watch Video Solution

80. There is a small hole at the bottom of tank filled with water. If total pressure at the bottom is $3atm$ ($1atm = 10^5 Nm^{-2}$), then find the velocity of water flowing from hole.

A. $\sqrt{400}m / s$

B. $\sqrt{600}m / s$

C. $\sqrt{60}m / s$

D. None of the above

Answer: A



Watch Video Solution

81. The working of venturimeter is based on

A. Torricelli's law

B. Pascal's law.

C. Bernoulli's theorem.

D. Stokes's law.

Answer: C



Watch Video Solution

82. In dimension of circular velocity v_0 liquid flowing through a tube are expressed as $(\eta^x \rho^y r^z)$ where η , ρ and r are the coefficient of viscosity of liquid density of liquid and

radius of the tube respectively then the value of x , y and z are given by

A. $1, 1, 1$

B. $1, -1, -1$

C. $-1, -1, 1$

D. $-1, -1, -1$

Answer: B



Watch Video Solution

83. The upper half of an inclined plane with inclination ϕ is perfectly smooth while the lower half is rough. A body starting from rest at the top will again come to rest at the bottom if the coefficient of friction for the lower half is given by

A. $\mu = 2 \tan \theta$

B. $\mu = \tan \theta$

C. $\mu = \frac{2}{\tan \theta}$

D. $\mu = \frac{1}{\tan \theta}$

Answer: A



Watch Video Solution

84. A bullet of mass 50 g moving horizontally with a velocity 210ms^{-1} gets embedded in a block of mass 1 kg kept on a rough horizontal surface. If the coefficient of kinetic friction between the block and surface is 0.5. The block- bullet system will move a distance of _____ before coming to rest (Acceleration due to gravity $= 10\text{ms}^{-2}$).

A. 40 m

B. 30 m

C. 20 m

D. 10 m

Answer: D



Watch Video Solution

85. A body of weight 64 N is pushed with just enough force to start it moving across a horizontal floor and the same force continues

to act afterwards. If the coefficients of static and dynamic friction are 0.6 and 0.4 respectively, the acceleration of the body will be (Acceleration due to gravity = g)

A. $\frac{g}{6.4}$

B. $0.64g$

C. $\frac{g}{32}$

D. $0.2g$

Answer: D



Watch Video Solution

86. The three vessels shown in figure have same base area. Equal volumes of a liquid are poured in the three vessels. The force on the base will be



- A. maximum at vessel A.
- B. maximum at vessel B.
- C. maximum at vessel C.

D. equal in all vessels.


Answer: C



Watch Video Solution

87. A fireman of mass 60 kg slides down a pole. As shown in figure. He is pressing the pole with a force of 600 N. The coefficient of friction between the hands and the pole is 0.5. With what acceleration will the fireman

slide down ($g = 10m / s^2$)



A. $1m / s^2$

B. $2.5m / s^2$

C. $5m / s^2$

D. $10m / s^2$


Answer: C



Watch Video Solution

88. A tank is filled with water up to a height H . Water is allowed to come out of a hole P in one of the walls at a depth D below the surface of water. Express the horizontal

distance x in terms of H and D



A. $x = \sqrt{D(H - D)}$

B. $x = \sqrt{\frac{D(H - D)}{2}}$

C. $x = 2\sqrt{D(H - D)}$


D. $x = 4\sqrt{D(H - D)}$

Answer: C



89. A point particle of mass m , moves along the uniformly rough track PQR as shown in figure. The coefficient of friction, between the particle and the rough track equals μ . The particle is released, from rest from the point P and it comes to rest at a point R. The energies, lost by the ball, over the parts, PQ and QR, of the track, are equal to each other, and no energy is lost when particle changes direction from PQ to QR.

The value of the coefficient of friction μ and the distance x ($= QR$), are, respectively close to:



A. 0.2 and 3.5m

B. 0.29 and 3.5m

C. 0.29 and 6.5m

D. 0.2 and 6.5m

Answer: B



Watch Video Solution

90. A 60 kg weight is dragged on a horizontal surface by a rope upto 2 metres. If coefficient of friction is $\mu = 0.5$ the angle of rope with the surface is 60° and $g = 9.8m/sec^2$, then work done is

A. 294 joule

B. 315 joule

C. 588 joule

D. 197 joule

Answer: B



Watch Video Solution

91. When a body is moving on a surface, the force of friction is called

- A. Static friction.
- B. Dynamic friction.
- C. Limiting friction.
- D. Rolling friction.

Answer: B



Watch Video Solution

92. The speed of the wind passing over the wings of a small aeroplane is 70 m/s and below the wing is 60 m/s . If the mass of the

plane is 1000 kg and the area of wing is $14m^2$,
then what will be the net vertical force on the
aeroplane ?

? (Density of air
 $= 1.2kg/m^3$ and $g = 10m/s^2$)

- A. 620 N upward
- B. 920 N upward
- C. 620 N downward
- D. 920 N downward

Answer: B



93. A rectangular vessel when full of water takes 10 minutes to be emptied through an orifice in its bottom. How much time will it take to be emptied when half filled with water

- A. 9 minutes
- B. 7 minutes
- C. 5 minutes
- D. 3 minutes

Answer: B



Watch Video Solution

94. A vessel completely filled with water has holes 'A' and 'B' at depths 'h' and '3h' from the top respectively. Hole 'A' is a square of side 'L' and 'B' is circle of radius 'r'. The water flowing out per second from both the holes is same. Then 'L' is equal to

A. $r^{\frac{1}{2}} (\pi)^{\frac{1}{2}} (3)^{\frac{1}{2}}$

B. $r(\pi)^{\frac{1}{4}}(3)^{\frac{1}{4}}$

C. $r(\pi)^{\frac{1}{2}}(3)^{\frac{1}{4}}$

D. $r^{\frac{1}{2}}(\pi)^{\frac{1}{3}}(3)^{\frac{1}{2}}$

Answer: C



Watch Video Solution

95. A large open tank has two holes in the wall. One is a square hole of side L at a depth y from the top and the other is a circular hole of radius R at a depth $4y$ from the top. When the

tank is completely filled with water, the quantities of water flowing out per second from both holes are the same. Then, R is equal to

A. $2\pi L$

B. $\frac{L}{\sqrt{2\pi}}$

C. L

D. $\frac{L}{2\pi}$

Answer: B



Watch Video Solution

96. A body of mass 2 kg is being dragged with uniform velocity of 2 m / s on a rough horizontal plane. The coefficient of friction between the body and the surface is 0.20. The amount of heat generated in 5 sec is . (4.2joule / cal and $g = 9.8m / s^2$)



A. 9.33 cal

B. 10.21 cal

C. 12.67 cal

D. 13.34 cal

Answer: A



Watch Video Solution

97. A spherical solid of volume V is made of a material of density ρ_1 . It is falling through a liquid of density ρ_2 ($\rho_2 < \rho_1$). Assume that the liquid applies a viscous force on the ball that is proportional to its speed v , i.e.,

$F_{viscous} = -kv^2$ ($k > 0$). The terminal speed of the ball is

A. $\frac{Vg\rho_1}{k}$

B. $\sqrt{\frac{Vg\rho_1}{k}}$

C. $\frac{Vg(\rho_1 - \rho_2)}{k}$

D. $\sqrt{\frac{Vg(\rho_1 - \rho_2)}{k}}$

Answer: D



Watch Video Solution

98. The heart of a man pumps 5 liters of blood through the arteries per minute at a pressure of 150 mm of mercury. If the density of mercury be $13.6 \times 10^3 \text{ kg/m}^3$ and $g = 10 \text{ m/s}^2$ then the power of heart in watt is :

A. 1.50

B. 1.70

C. 2.35

D. 3.0

Answer: B



Watch Video Solution

Mcq Evaluation Test

1. A body while sliding went straight down an incline of 45° to the horizontal and was subjected to a coefficient of kinetic friction of 0.20. Starting from rest, how long did it take him to reach a speed of 50km/hr ? (Ignore air resistance and take $h = 10\text{m/s}^2$)

A. $2.45s$

B. $9.2s$

C. $1.5 \times 10^2 s$

D. $2.0s$

Answer: A



Watch Video Solution

2. As an air bubble comes from the bottom of a lake to the top, its radius _____

A. increases

B. decreases

C. does not change


D. becomes zero

Answer: A



Watch Video Solution

3. Four container of honey are shown in the figure. The pressure at depth h is



A. greatest in A.

B. greatest in D.

C. least in B and C both.

D. equal in all the containers.


Answer: D



Watch Video Solution

4. What is the maximum value of the force F such that the block shown in the arrangement does not move ?

(Given $\mu = \frac{1}{2\sqrt{3}}$)



A. 20 N

B. 40 N

C. 12 N

D. 45 N

Answer: B



Watch Video Solution

5. You want to apply a force on a box so that it moves with constant speed across a horizontal floor. The coefficient of kinetic friction between the box and the floor is μ_k .

Of the following cases, the force you apply on the box will be smallest when you

A. push it with a force applied horizontally
at an angle $0 < \theta < 90^\circ$ in downward
direction.

B. pull it with a force applied horizontally
at the same angle as in (A), in upward
direction.

C. do either (A) or (B) since the applied
force is the same.

D. push or pull with a force applied
horizontally.

Answer: B



Watch Video Solution

6. A block of mass M is pulled along a horizontal surface by applying a force at angle θ with the horizontal. The friction coefficient between the block and the surface is μ . If the block travels at a uniform velocity, find the work done by this applied force during a displacement d of the block.

A. $\frac{\mu g s}{\cos \theta + \sin \theta}$

B. $\frac{\mu m g s \cos \theta}{\cos \theta + \mu \sin \theta}$

C. $\frac{\mu g s \sin \theta}{\cos \theta + \mu \sin \theta}$

D. $\frac{\mu g s \cos \theta}{\cos \theta - \mu \sin \theta}$

Answer: B



Watch Video Solution

7. Consider a car moving along a straight horizontal road with a speed of 36 km/h. If the coefficient of static friction between road and

tyers is 0.4, the shortest distance in which the car can be stopped is (Take $g = 10m / s^2$)

A. $33.8m$

B. $12.5m$

C. $58.6m$


D. $20m$

Answer: B



Watch Video Solution

8. Water and gasoline surfaces are open to the atmosphere at the same elevation. Find the height h of the third liquid in the right leg.
 (Density of gasoline $600\text{kg}/\text{m}^3$ and of liquid is $1600\text{kg}/\text{m}^3$)



A. 3.8m

B. 1.9m

C. 2.5m

D. $0.45m$

Answer: C



Watch Video Solution

9. A U-tube containing a liquid moves with a horizontal acceleration a along a direction joining the two vertical limbs. The separation between these limbs is d . The difference in their liquid levels is

A. xyg

B. $\frac{xg}{y}$

C. $\frac{yg}{x}$

D. $\frac{xy}{g}$

Answer: D



Watch Video Solution

10. Equal volume of two immiscible liquids of densities ρ and 2ρ are filled in a vessel as shown in the figure. Two small holes are punched at depths $h/2$ and $3h/2$ from the

surface of lighter liquid. If v_1 and v_2 are the velocities of efflux at these two holes, then

v_1 / v_2 is




Fig. 4.

A. $\frac{1}{2\sqrt{2}}$

B. $\sqrt{2}$

C. $1/4$

D. $\frac{1}{\sqrt{2}}$

Answer: B



Watch Video Solution

11. Assertion : A horse has to pull a cart harder during the first few steps of his motion.

Reason: The first few steps are always difficult.

A. Assertion is True, Reason is True, Reason

is a correct explanation for Assertion

B. Assertion is True, Reason is True, Reason

is not a correct explanation for Assertion

C. Assertion is True, Reason is False

D. Assertion is False, Reason is False.

Answer: C



Watch Video Solution

12. Digboi mine in Assam is the deepest in India. In this mine the conditions as compared to those at the surface are,

A. lower air pressure, higher acceleration
due to gravity.

B. higher air pressure, lower acceleration
due to gravity.

C. higher air pressure, higher acceleration
due to gravity.

D. lower air pressure, lower acceleration
due to gravity.

Answer: B



Watch Video Solution

13. Consider a liquid at rest in a container which opens into atmosphere. At any point A at a depth h below the surface of the liquid,

(i) the gauge pressure at point A is ρgh .

(ii) the gauge pressure at point a is $P_a + \rho gh$.

(iii) The pressure at point A is ρgh .

(iv) The pressure at point A is $P_a + \rho gh$.

The correct alternative is

A. (i) and (iii)

B. (ii) and (iv)

C. (iii) and (iv)

D. (i) and (iv)

Answer: D



Watch Video Solution

14. At two points on a horizontal tube of varying cross section carrying water, the radii are 1 cm and 0.5cm . The pressure difference between these points is 6 cm of water. How

much liquid flows through the tube per second ?

A. 100c. c. per s

B. 98c. c. per s

C. 88c. c. per s

D. 70c. c. per s

Answer: C



Watch Video Solution

15. An aquarium is filled with water. The lateral wall of the aquarium is 50 cm long and 40 cm high. Using $10g/cm^2$ for the acceleration due to gravity, and $1 g/cm^2$ for the density of water, the force on the lateral wall of the aquarium is

A. 36 N

B. 400 N

C. 180 N

D. 1500 N

Answer: B



Watch Video Solution

16. The fire fighters have a jet attached to the held of their water pipes

A. to increase the velocity of water flowing out of the pipe.

B. to decrease the velocity of water flowing out of the pipe.

C. to have a streamline flow of water.

D. to have a turbulent flow of water.

Answer: A



Watch Video Solution

17. The nature of graph between terminal velocity of spherical body vs the square of its radius is

A. exponentially increasing curve

B. straight line

C. parabola

D. exponentially decreasing curve

Answer: B



Watch Video Solution

18. Assertion: The accumulation of snow on the wings of an aeroplane reduces the lift.

Reason: Pressure difference depends upon the curvature.


- A. Assertion is True, Reason is True, Reason is a correct explanation for Assertion
- B. Assertion is True, Reason is True, Reason is not a correct explanation for Assertion
- C. Assertion is True, Reason is False
- D. Assertion is False, Reason is False.

Answer: A



Watch Video Solution

19. The figure below indicates flow of water through a tube of uniform cross-section with a constant speed in the direction as shown by the arrows. On the tube, water exerts



- A. a net force to the left.
- B. a net force to the right.
- C. an anticlockwise torque.

D. a clockwise torque.

Answer: D



Watch Video Solution

20. A water barrel having water up to a depth d is placed on a table of height H . A small hole is made on the wall of the barrel at its bottom. If the stream of water coming out of the hole falls on the ground at a horizontal distance r from the barrel, then the value of d is

A. $\frac{4H}{r^2}$

B. $4Hr^2$

C. $\frac{r^2}{4H}$

D. $\frac{H}{4r^2}$


Answer: C



Watch Video Solution

21. An incompressible, non-viscous liquid of density ρ flows through a horizontal tube as shown below. The flow condition is steady.

Area of cross section at A and B are a_1 and a_2 respectively. The height difference of liquid in two tubes inserted at A and B is h .



A. Pressure at A is more than at B by ρgh .

B. The diagram has an error as the height of liquid in tube at B will be higher than in tube at A.

C. For the given data, if flow is streamlined

Bernoulli's principle can be applied.

D. Both (A) and (C)

Answer: D



Watch Video Solution