



MATHS

NCERT - FULL MARKS MATHS(TAMIL)

VECTOR ALGEBRA

Example

1. Represent graphically a displacement of 40 km, 30° west of south.



Watch Video Solution

2. Classify the following measures as scalars and vectors.

(i) 5seconds (ii) 1000cm^3 (iii) 10 Newton

(iv) $30\frac{\text{km}}{\text{hr}}$ (v) $10\text{g}/\text{cm}^2$ (vi) 20 m/s towards North




[Watch Video Solution](#)

3. In the figure, which of the vectors are

(i) Collinear vectors

(ii) Equal vectors

(iii) Coplanar vectors



Watch Video Solution

4. Find the values of x , y and z so that the vectors

$$\vec{a} = x\hat{i} + 2\hat{j} + z\hat{k} \text{ and } \vec{b} = 2\hat{i} + y\hat{j} + \hat{k} \text{ are equal.}$$



Watch Video Solution

5. Let $\vec{a} = \hat{i} + 2\hat{j}$ and $\vec{b} = 2\hat{i} + \hat{j}$. Is $|\vec{a}| = |\vec{b}|$? Are the vectors \vec{a} and \vec{b} equal?

 [Watch Video Solution](#)

6. Find unit vector in the direction of vector $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$

 [Watch Video Solution](#)

7. Find a vector in the direction of $\vec{a} = \hat{i} - 2\hat{j}$ that has a magnitude 7 units.

 [Watch Video Solution](#)

8. Find the unit vector in the direction of the sum of vectors ,

$$\vec{a} = 2\hat{i} + 2\hat{j} - 5\hat{k} \text{ and } \vec{b} = 2\hat{i} + \hat{j} + 3\hat{k}.$$



[Watch Video Solution](#)

9. Write the direction ratio's of the vector $\vec{a} = \hat{i} + \hat{j} - 2\hat{k}$ and hence calculate its direction cosines.



[Watch Video Solution](#)

10. Find the vector joining the points P(2, 3, 0) and Q(-1, -2, -4) directed from P to Q



[Watch Video Solution](#)

11. Consider two points P and Q with position vectors $\vec{OP} = 3\vec{a} - 2\vec{b}$ and $\vec{OQ} = \vec{a} + \vec{b}$. Find the position vector of a point R which divides the line joining P and Q in the ratio 2 : 1
- (i) internally (ii) externally.

 [Watch Video Solution](#)

12. Show that the points whose position vectors $4\hat{i} - 3\hat{j} + \hat{k}$, $2\hat{i} - 4\hat{j} + 5\hat{k}$, $\hat{i} - \hat{j}$ form a right angled triangle.

 [Watch Video Solution](#)

13. Find the angle between two vectors \vec{a} and \vec{b} with magnitudes 1 and 2 respectively and when $\vec{a} \cdot \vec{b} = 1$.

 [Watch Video Solution](#)

14. Find angle θ between the vectors

$$\vec{a} = \hat{i} + \hat{j} - \hat{k} \text{ and } \vec{b} = \hat{i} + \hat{j} + \hat{k}.$$

 [Watch Video Solution](#)

15. If $\vec{a} = 5\hat{i} - \hat{j} - 3\hat{k}$ and $\vec{b} = \hat{i} + 3\hat{j} - 5\hat{k}$, then show that the vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ are perpendicular.

 [Watch Video Solution](#)

16. Find the projection of the vector $\vec{a} = 2\hat{i} + 3\hat{j} + 2\hat{k}$ on the vector $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$.

 [Watch Video Solution](#)

17. Find $\left| \vec{a} - \vec{b} \right|$, if two vectors \vec{a} and \vec{b} are such that $\left| \vec{a} \right| = 2$, $\left| \vec{b} \right| = 3$ and $\vec{a} \cdot \vec{b} = 4$.

 [Watch Video Solution](#)

18. If \vec{a} is a unit vector and $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 8$, then find $\left| \vec{x} \right|$.

 [Watch Video Solution](#)

19. For any two vectors \vec{a} and \vec{b} , we always have $\left| \vec{a} \cdot \vec{b} \right| \leq \left| \vec{a} \right| \left| \vec{b} \right|$ (Cauchy-Schwartz inequality).

 [Watch Video Solution](#)

20. For any two vectors \vec{a} and \vec{b} , we always have $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$ (triangle inequality).

 [Watch Video Solution](#)

21. Show that the points $A(-2\hat{i} + 3\hat{j} + 5\hat{k})$, $B(\hat{i} + 2\hat{j} + 3\hat{k})$ and $C(7\hat{i} - \hat{k})$ are collinear.

 [Watch Video Solution](#)

22. Find the magnitude of $\vec{a} \times \vec{b}$ if $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}$ and $\vec{b} = 3\hat{i} + 5\hat{j} - 2\hat{k}$.

 [Watch Video Solution](#)

23. Find the unit vectors perpendicular to each of the vectors

$$\vec{a} + \vec{b} \quad \text{and} \quad \vec{a} - \vec{b}, \text{ where } \vec{a} = \hat{i} + \hat{j} + \hat{k} \text{ and}$$

$$\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}.$$

 [Watch Video Solution](#)

24. Find the area of a triangle having the points A(1,1,1), B (1,2,3) and C(2,3,1) as its vertices.

 [Watch Video Solution](#)

25. Find the area of the parallelogram whose adjacent sides are given by the vectors $\vec{a} = 3\hat{i} + \hat{j} + 4\hat{k}$ and $\vec{b} = \hat{i} - \hat{j} + \hat{k}$.

 [Watch Video Solution](#)

26. Write all the unit vectors in XY - plane.

 [Watch Video Solution](#)

27. If $\hat{i} + \hat{j} + \hat{k}$, $2\hat{i} + 5\hat{j}$, $3\hat{i} + 2\hat{j} - 3\hat{k}$ and $\hat{i} - 6\hat{j} - \hat{k}$ are position vectors of points A,B,C and D respectively, then find the angle between \overrightarrow{AB} and \overrightarrow{CD} . Deduce that \overrightarrow{AB} and \overrightarrow{CD} are collinear.

 [Watch Video Solution](#)

28. Let \vec{a} , \vec{b} and \vec{c} be the three vectors such that $|\vec{a}| = 3$, $|\vec{b}| = 4$, $|\vec{c}| = 5$, and each one of them being perpendicular to the sum of the other two. Find $|\vec{a} + \vec{b} + \vec{c}|$.

 [Watch Video Solution](#)

29. Three vectors \vec{a} , \vec{b} and \vec{c} satisfy the condition

$\vec{a} + \vec{b} + \vec{c} = \vec{0}$. Evaluate the quantity

$\mu = \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$, if

$|\vec{a}| = 3$, $|\vec{b}| = 4$ and $|\vec{c}| = 2$.

 [Watch Video Solution](#)

30. If $\vec{\alpha} = 3\hat{i} - \hat{j}$ and $\vec{\beta} = 2\hat{i} + \hat{j} - 3\hat{k}$, then express, $\vec{\beta}$ in the form $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$, where $\vec{\beta}_1$ is parallel to $\vec{\alpha}$ and $\vec{\beta}_2$ is perpendicular to $\vec{\alpha}$.

 [Watch Video Solution](#)

1. Represent graphically a displacement of 40 km, 30° east of north.



[Watch Video Solution](#)

2. Classify the following measures as scalars and vectors .

(i) 10 kg

(ii) 2 meters north

(iii) 40°

(iv) 40 watt

(v) 10^{19} coulomb

(vi) $20m / s^2$



[Watch Video Solution](#)

3. Classify the following as scalar and vector quantities.

(i) Time period (ii) Distance

(iii) Force (iv) Velocity

(v) Work done



[Watch Video Solution](#)

4. Answer the followings true or false.

(i) \vec{a} and $-\vec{a}$ are collinear.

(ii) Two collinear vectors are always equal in magnitude.

(iii) Two vectors having same magnitude are collinear.

(iv) Two collinear vectors having the same magnitude are equal.



[Watch Video Solution](#)

1. Compute the magnitude of the following vectors :

$$\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 2\hat{i} - 7\hat{j} - 3\hat{k}, \vec{c} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} - \frac{1}{\sqrt{3}}\hat{k}$$



[Watch Video Solution](#)

2. Write two different vectors having same magnitude.



[Watch Video Solution](#)

3. Write two different vectors having the same direction.



[Watch Video Solution](#)

4. Find the values of x and y so that the vectors $2\hat{i} + 3\hat{j}$ and $x\hat{i} + y\hat{j}$ are equal.

 [Watch Video Solution](#)

5. Find the scalar and vector components of the vector with initial point (2,1) and terminal point (-5,7)

 [Watch Video Solution](#)

6. Find the sum of the vectors

$$\vec{a} = \hat{i} - 2\hat{j} + \hat{k}, \vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k} \text{ and } \vec{c} = \hat{i} - 6\hat{j} - -7\hat{k}$$

.

 [Watch Video Solution](#)

7. Find the unit vector in the direction of the vector

$$\vec{a} = \hat{i} + \hat{j} + 2\hat{k}.$$



[Watch Video Solution](#)

 [Watch Video Solution](#)

8. Find the unit vector in the direction of vector \overrightarrow{PQ} , where P and Q are the points (1, 2, 3) and (4, 5, 6) respectively.

 [Watch Video Solution](#)

9. For given vectors $\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + \hat{j} - \hat{k}$, find the unit vector in the direction of the vector $\vec{a} + \vec{b}$.

 [Watch Video Solution](#)

10. Find a vector in the direction of vector $5\hat{i} - \hat{j} + 2\hat{k}$ which has magnitude 8 units.

 [Watch Video Solution](#)

11. Show that the vectors $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $-4\hat{i} + 6\hat{j} - 8\hat{k}$ are collinear.



Watch Video Solution

12. Find the direction cosines of the vector $\hat{i} + 2\hat{j} + 3\hat{k}$.



Watch Video Solution

13. Find the direction cosines of the vector joining the points A (1,2,-3) and B(-1,-2,1) directed from A to B.



Watch Video Solution

14. Show that the vector $\hat{i} + \hat{j} + \hat{k}$ is equally inclined with the coordinate axes.

 [Watch Video Solution](#)

15. Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are $\hat{i} + 2\hat{j} - \hat{k}$ and $-\hat{i} + \hat{j} + \hat{k}$ respectively, in the ratio 2 : 1

(i) internally (ii) externally

 [Watch Video Solution](#)


16. Find the position vector of the mid point of the vector joining the points P(2,3,4) and Q(4,1,-2).

 [Watch Video Solution](#)

17. Show that the points A, B and C with position vectors ,
 $\vec{a} = 3\hat{i} - 4\hat{j} - 4\hat{k}$, $\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - 3\hat{j} + 5\hat{k}$
 , respectively form the vertices of a right angled triangle.

 Watch Video Solution

18. In triangle ABC, which of the following is not true?



A. $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$

B. $\vec{AB} + \vec{BC} - \vec{AC} = \vec{0}$

C. $\vec{AB} + \vec{BC} - \vec{AC} = \vec{0}$

$$D. \vec{AB} - \vec{CB} + \vec{CA} = \vec{0}$$

Answer: C

 [Watch Video Solution](#)

19. If \vec{a} and \vec{b} are collinear vectors, then which of the following are incorrect?

A. $\vec{b} = \lambda \vec{a}$, for some scalar λ

B. $\vec{a} = \pm \vec{b}$

C. the respective components of \vec{a} and \vec{b} are not proportional

D. both the vectors \vec{a} and \vec{b} have same direction, but different magnitudes.

Answer: B::C::D

 [Watch Video Solution](#)

Exercise 10 3

1. Find the angle between two vectors \vec{a} and \vec{b} with magnitudes $\sqrt{3}$ and 2, respectively having $\vec{a} \cdot \vec{b} = \sqrt{6}$.

 [Watch Video Solution](#)

2. Find the angle between the vectors $\hat{i} - 2\hat{j} + 3\hat{k}$ and $3\hat{i} - 2\hat{j} + \hat{k}$

 [Watch Video Solution](#)

3. Find the projection of the vector $\hat{i} - \hat{j}$ on the vector $7\hat{i} + \hat{j}$.



[Watch Video Solution](#)

4. Find the projection of the vector $\hat{i} + 3\hat{j} + 7\hat{k}$ on the vector $7\hat{i} - \hat{j} + 8\hat{k}$.



[Watch Video Solution](#)

5. Show that each of the given three vectors is a unit vector: $\frac{1}{7}(2\hat{i} + 3\hat{j} + 6\hat{k})$, $\frac{1}{7}(3\hat{i} - 6\hat{j} + 2\hat{k})$, $\frac{1}{7}(6\hat{i} + 2\hat{j} - 3\hat{k})$. Also, show they are mutually perpendicular to each other.



[Watch Video Solution](#)

6. Find $|\vec{a}|$ and $|\vec{b}|$, if $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 8$ and $|\vec{a}| = 8|\vec{b}|$.

 [Watch Video Solution](#)

7. Evaluate the product $(3\vec{a} - 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$.

 [Watch Video Solution](#)

8. Find the magnitude of two vectors \vec{a} and \vec{b} , having the same magnitude and such that the angle between them is 60° and their scalar product is $\frac{1}{2}$.

 [Watch Video Solution](#)

9. Find $|\vec{x}|$ if for a unit vector \vec{a} , $(\vec{x} - \vec{a})(\vec{x} + \vec{a}) = 12$

 [Watch Video Solution](#)

10. Given

$\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 3\hat{i} + \hat{j}$ such that $\vec{a} + \lambda\vec{b}$ is perpendicular to \vec{c} , then find the value of λ .

 [Watch Video Solution](#)

11. Show that $|\vec{a}|\vec{b} + |\vec{b}|\vec{a}$ is perpendicular to $|\vec{a}|\vec{b} - |\vec{b}|\vec{a}$, for any two nonzero vectors \vec{a} and \vec{b} .

 [Watch Video Solution](#)

12. If $\vec{a} \cdot \vec{a} = 0$ and $\vec{a} \cdot \vec{b} = 0$ then what can be concluded about the vector \vec{b} ?

 [Watch Video Solution](#)

13. If $\vec{a}, \vec{b}, \vec{c}$ are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, find the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$.

 [Watch Video Solution](#)

14. If either vector $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then $\vec{a} \cdot \vec{b} = 0$. But the converse need not be true. Justify your answer with an example.

 [Watch Video Solution](#)

15. If either vector A,B,C of a triangle ABC are $(1,2,3),(-1,0,0),(0,1,2)$, respectively , then find $\angle ABC$. [$\angle ABC$ is the angle between the vectors \overrightarrow{BA} and \overrightarrow{BC}].

 [Watch Video Solution](#)

16. Show that the points $A(1,2,7)$, $B(2,6,3)$ and $C(3, 10, -1)$ are collinear.

 [Watch Video Solution](#)

17. Show that the vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $3\hat{i} - 4\hat{k}$ form the vertices of a right angled triangle.

 [Watch Video Solution](#)

18. If \vec{a} is a non zero vector of magnitude 'a' and λ a nonzero scalar, then $\lambda \vec{a}$ is unit vector if

A. $\lambda = 1$

B. $\lambda = -1$

C. $a = |\lambda|$

D. $a = 1/|\lambda|$

Answer: D



[Watch Video Solution](#)

Exercise 10 4

1. Find $\left| \vec{a} \times \vec{b} \right|$, if $\vec{a} = \hat{i} - 7\hat{j} + 7\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 2\hat{k}$.



[Watch Video Solution](#)

 Watch Video Solution

2. Find a unit vector perpendicular to each of the vector $\vec{a} + \vec{b}$ and $\vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$.

 Watch Video Solution

3. If a unit vector \vec{a} makes angles $\frac{\pi}{3}$ with \hat{i} , $\frac{\pi}{4}$ with \hat{j} and an acute angle θ with \hat{k} , then find θ and hence, the components of \vec{a} .

 Watch Video Solution

4. Show that $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2(\vec{a} \times \vec{b})$

 Watch Video Solution

5. Find λ and μ if $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + \lambda\hat{j} + \mu\hat{k}) = \vec{0}$.

 [Watch Video Solution](#)

6. Given that $\vec{a} \cdot \vec{b} = 0$ and $\vec{a} \times \vec{b} = \vec{0}$. What can you conclude about the vectors \vec{a} and \vec{b} ?

 [Watch Video Solution](#)

7. Let the vectors $\vec{a}, \vec{b}, \vec{c}$ be given as $a_1\hat{i} + a_2\hat{j} + a_3\hat{k}, b_1\hat{i} + b_2\hat{j} + b_3\hat{k}, c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$. Then show that $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$

 [Watch Video Solution](#)

8. If either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then $\vec{a} \times \vec{b} = \vec{0}$. Is the converse true? Justify your answer with an example.



Watch Video Solution

9. Find the area of the triangle with vertices A(1,1,2) B(2,3,5) and C(1,5,5).



Watch Video Solution

10. Find the area of the parallelogram whose adjacent sides are determined by the vectors $\vec{a} = \hat{i} - \hat{j} + 3\hat{k}$ and $\vec{b} = 2\hat{i} - 7\hat{j} + \hat{k}$.



Watch Video Solution

11. Let the vectors \vec{a} and \vec{b} be such that $|\vec{a}| = 3$ and $|\vec{b}| = \frac{\sqrt{2}}{3}$, then $\vec{a} \times \vec{b}$ is a unit vector, if the angle between \vec{a} and \vec{b} is

A. $\pi/6$

B. $\pi/4$

C. $\pi/3$

D. $\pi/2$

Answer: B



Watch Video Solution

12. Area of a rectangle having vertices A,B,C and D with positions vectors $-\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k}$, $\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k}$, $\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}$ and $-\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}$, respectively is

A. $\frac{1}{2}$

B. 1

C. 2

D. 4

Answer: C



[Watch Video Solution](#)

Miscellaneous Exercise On Chapter 10

1. Write down a unit vector in XY - plane making an angle of 30° with the positive direction of x- axis.



[Watch Video Solution](#)

2. Find the scalar components and magnitude of the vector joining the points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$.

 [Watch Video Solution](#)

3. A girl walks 4 km towards west, then she walks 3 km in a direction 30° east of north and stops. Determine the girl's displacement from her initial point of departure.

 [Watch Video Solution](#)

4. If $\vec{a} = \vec{b} + \vec{c}$, then is it true that $|\vec{a}| = |\vec{b}| + |\vec{c}|$?

Justify your answer.

 [Watch Video Solution](#)

5. Find the value of x for which $x(\hat{i} + \hat{j} + \hat{k})$ is a unit vector.

 [Watch Video Solution](#)

6. Find a vector of magnitude 5 units and parallel to the resultant of the vectors $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$.

 [Watch Video Solution](#)

7. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - \hat{j} + 3\hat{k}$ and $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$, find a unit vector parallel to the vector $2\vec{a} - \vec{b} + 3\vec{c}$.

 [Watch Video Solution](#)

8. Show that the points A(1, -2, -8) B (5, 0, -2) and C(11, 3, 7) are collinear and find the ratio in which B divides AC.

 [Watch Video Solution](#)

9. Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are $(2\vec{a} + \vec{b})$ and $(\vec{a} - 3\vec{b})$ externally in the ratio 1:2 Also , show that P is the mid point of the line segment RQ.

 [Watch Video Solution](#)

10. The two adjacent sides of a parallelogram are $2\hat{i} - 4\hat{j} + 5\hat{k}$ and $\hat{i} - 2\hat{j} - 3\hat{k}$. Find the unit vector parallel to its diagonal. Also, find its area.

 [Watch Video Solution](#)

 Watch Video Solution

11. Show that the direction cosines of a vector equally inclined to the axes OX,OY and OZ are $\pm \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right)$.

 Watch Video Solution

12. Let
 $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$
Find a vector \vec{d} which is perpendicular to both \vec{a} and \vec{b} , and $\vec{c} \cdot \vec{d} = 15$.

 Watch Video Solution

13. The scalar product of the vector $\hat{i} + \hat{j} + \hat{k}$ with a unit vector along the sum of vectors $2\hat{i} + 4\hat{j} - 5\hat{k}$ and $\lambda\hat{i} + 2\hat{j} + 3\hat{k}$ is equal to one. Find the value of λ .

 [Watch Video Solution](#)

14. If $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular vectors of equal magnitudes, show that the vector $\vec{c} \cdot \vec{d} = 15$ is equally inclined to \vec{a}, \vec{b} and \vec{c} .

 [Watch Video Solution](#)

15. Prove that $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a}|^2 + |\vec{b}|^2$, if and only if \vec{a}, \vec{b} are perpendicular, given $\vec{a} \neq \vec{0}, \vec{b} \neq \vec{0}$.

 [Watch Video Solution](#)

16. If θ is the angle between two vectors \vec{a} and \vec{b} then $\vec{a} \cdot \vec{b} \geq 0$ only when

A. $0 < \theta < \frac{\pi}{2}$

B. $0 \leq \theta \leq \frac{\pi}{2}$

C. $0 < \theta < \pi$

D. $0 \leq \theta \leq \pi$

Answer: B



[Watch Video Solution](#)

17. Let \vec{a} and \vec{b} be two unit vectors and θ is the angle between them Then $\vec{a} + \vec{b}$ is a unit vector if

A. $\theta = \frac{\pi}{4}$

B. $\theta = \frac{\pi}{3}$

C. $\theta = \frac{\pi}{2}$

D. $\theta = \frac{2\pi}{3}$

Answer: D



Watch Video Solution

18. The value of $\hat{i} \cdot (\hat{j} \times \hat{k}) + \hat{j} \cdot (\hat{k} \times \hat{i}) + \hat{k} \cdot (\hat{j} \times \hat{i}) = \dots\dots\dots$

A. 0

B. -1

C. 1

D. 3

Answer: C



Watch Video Solution

19. If θ is the angle between any two vectors \vec{a} and \vec{b} , then

$$\left| \vec{a} \cdot \vec{b} \right| = \left| \vec{a} \times \vec{b} \right|, \text{ when } \theta \text{ is equal to}$$

A. 0

B. $\frac{\pi}{4}$

C. $\frac{\pi}{2}$

D. π

Answer: B



Watch Video Solution