



PHYSICS

BOOKS - PSEB

GRAVITATION

Exercise

1. Answer the following :- You can shield a charge from electrical forces by putting it inside a hollow conductor. Can you shield a

body from the gravitational influence of nearby matter by putting it inside a hollow sphere or by some other means ?



[Watch Video Solution](#)

2. Answer the following :- An astronaut inside a small space ship orbiting around the earth cannot detect gravity. If the space station orbiting around the earth has a large size, can he hope to detect gravity ?



[Watch Video Solution](#)

3. Answer the following :- If you compare the gravitational force on the earth due to the sun to that due to the moon, you would find that the Sun's pull is greater than the moon's pull. (you can check this yourself using the data available in the succeeding exercises). However, the tidal effect of the moon's pull is greater than the tidal effect of sun. Why ?



[Watch Video Solution](#)

4. Choose the correct alternative :-
Acceleration due to gravity
 \in *creases / decreases* with increasing
altitude.



Watch Video Solution

5. Choose the correct alternative :-
Acceleration due to gravity
 \in *creases / decreases* with increasing depth

(assume the earth to be a sphere of uniform density).



[Watch Video Solution](#)

6. Choose the correct alternative :-

Acceleration due to gravity is independent of mass of the *earth* / *mass* of the body.



[Watch Video Solution](#)

7. Choose the correct alternative :- The formula $-GMm(1/r_2 - 1/r_1)$ is more accurate than the formula $mg(r_2 - r_1)$ for the difference of potential energy between two points r_2 and r_1 , distance away from the centre of the earth.



[Watch Video Solution](#)

8. Suppose there existed a planet that went around the sun twice as fast as the earth.

What would be its orbital size as compared to that of the earth ?



[Watch Video Solution](#)

9. Io, one of the satellites of Jupiter, has an orbital period of 1.769 days and the radius of the orbit is $4.22 \times 10^8 m$. Show that the mass of Jupiter is about one-thousandth that of the sun.



[Watch Video Solution](#)

10. Let us assume that our galaxy consists of 2.5×10^{11} stars each of one solar mass. How long will a star at a distance of 50,000 ly from the galactic centre take to complete one revolution ? Take the diameter of the Milky Way to be 10^5 ly.



Watch Video Solution

11. Choose the correct alternative:- If the zero of potential energy is at infinity, the total

energy of an orbiting satellite is negative of its

$k \in \text{etic} / \text{potential}$ energy.



[Watch Video Solution](#)

12. Choose the correct alternative:- The energy required to launch an orbiting satellite out of earth's gravitational influence is m or $e / \le ss$ than the energy required to project a stationary object at the same height (as the satellite) out of earth's influence.



[Watch Video Solution](#)

13. Does the escape speed of a body from the earth depend on:- the mass of the body



Watch Video Solution

14. Does the escape speed of a body from the earth depend on:- the location from where it is projected,



Watch Video Solution

15. Does the escape speed of a body from the earth depend on:- the direction of projection,



Watch Video Solution

16. Does the escape speed of a body from the earth depend on:- the height of the location from where the body is launched?



Watch Video Solution

17. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- linear speed,



Watch Video Solution

18. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- angular speed,



Watch Video Solution

19. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- angular momentum,



Watch Video Solution

20. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- kinetic energy,



Watch Video Solution

21. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- potential energy,



Watch Video Solution

22. A comet orbits the sun in a highly elliptical orbit. Does the comet have a constant:- total energy throughout its orbit? Neglect any mass loss of the comet when it comes very close to the Sun.



Watch Video Solution

23. Which of the following symptoms is likely to afflict an astronaut in space:- swollen feet,



Watch Video Solution

24. Which of the following symptoms is likely to afflict an astronaut in space:- swollen face



Watch Video Solution

25. Which of the following symptoms is likely to afflict an astronaut in space:- headache,



Watch Video Solution

26. Which of the following symptoms is likely to afflict an astronaut in space:- orientational problem.



Watch Video Solution

27. A rocket is fired from the earth towards the sun. At what distance from the earth's centre is the gravitational force on the rocket zero ?

Mass of the sun = $2 \times 10^{30} \text{ kg}$, mass of the earth = $6 \times 10^{24} \text{ kg}$. Neglect the effect of other planets etc. (orbital radius = $1.5 \times 10^{11} \text{ m}$).



Watch Video Solution

28. How will you 'weigh the sun', that is estimate its mass? The mean orbital radius of

the earth around the sun is $1.5 \times 10^8 km$.



[Watch Video Solution](#)

29. A saturn year is 29.5 times the earth year.
How far is the saturn from the sun if the earth
is $1.50 \times 10^8 km$ away from the sun ?



[Watch Video Solution](#)

30. A body weighs 63 N on the surface of the
earth. What is the gravitational force on it due

to the earth at a height equal to half the radius of the earth ?



[Watch Video Solution](#)

31. Assuming the earth to be a sphere of uniform mass density, how much would a body weigh half way down to the centre of the earth if it weighed 250 N on the surface ?



[Watch Video Solution](#)

32. A rocket is fired vertically with a speed of 5km s^{-1} from the earth's surface. How far from the earth does the rocket go before returning to the earth ? Mass of the earth = $6.0 \times 10^{24}\text{kg}$, mean radius of the earth = $6.4 \times 10^6\text{m}$, $G = 6.67 \times 10^{-11}\text{Nm}^2\text{kg}^{-2}$.



Watch Video Solution

33. The escape speed of a projectile on the earth's surface is 11.2km s^{-1} . A body is

projected out with thrice this speed. What is the speed of the body far away from the earth? Ignore the presence of the sun and other planets.



[Watch Video Solution](#)

34. A satellite orbits the earth at a height of 400 km above the surface. How much energy must be expended to rocket the satellite out of the earth's gravitational influence? Mass of the satellite = 200 kg, mass of the earth

$= 6.0 \times 10^{24} \text{ kg}$, radius of the earth = $6.4 \times 10^6 \text{ m}$, $G = 6.67 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$.



Watch Video Solution

35. Two stars each of one solar mass ($= 2 \times 10^{30} \text{ kg}$) are approaching each other for a head on collision. When they are a distance 10^9 km , their speeds are negligible. What is the speed with which they collide ? The radius of each star is 10^4 km . Assume the

stars to remain undistorted until they collide.

(Use the known value of G).



Watch Video Solution

36. Two heavy spheres each of mass 100 kg and radius 0.10 m are placed 1.0 m apart on a horizontal table. What is the gravitational force and potential at the mid point of the line joining the centres of the spheres ? Is an object placed at that point in equilibrium? If so, is the equilibrium stable or unstable ?



[Watch Video Solution](#)

37. As you have learnt in the text, a geostationary satellite orbits the earth at a height of nearly 36,000 km from the surface of the earth. What is the potential due to earth's gravity at the site of this satellite ? (Take the potential energy at infinity to be zero). Mass of the earth = $6.0 \times 10^{24} \text{ kg}$, radius = 6400 km.



[Watch Video Solution](#)

38. A star 2.5 times the mass of the sun and collapsed to a size of 12 km rotates with a speed of 1.2 rev. per second. (Extremely compact stars of this kind are known as neutron stars. Certain stellar objects called pulsars belong to this category). Will an object placed on its equator remain stuck to its surface due to gravity ? (mass of the sun $= 2 \times 10^{30} \text{ kg}$).



Watch Video Solution

39. A spaceship is stationed on Mars. How much energy must be expended on the spaceship to launch it out of the solar system ? Mass of the space ship = 1000 kg, mass of the sun = $2 \times 10^{30} \text{ kg}$, mass of mars = $6.4 \times 10^{23} \text{ kg}$, radius of mars = 3395 km, radius of the orbit of mars = $2.28 \times 10^8 \text{ km}$, $G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$.



Watch Video Solution

40. A rocket is fired 'vertically' from the surface of mars with a speed of 2 km s^{-1} . If 20% of its initial energy is lost due to martian atmospheric resistance, how far will the rocket go from the surface of mars before returning to it ? Mass of mars = $6.4 \times 10^{23} \text{ kg}$, radius of mars = 3395 km, $G = 6.67 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$.



[Watch Video Solution](#)