



PHYSICS

BOOKS - PSEB

OSCILLATIONS

Exercise

1. Which of the following examples represent periodic motion?

A. A swimmer completing one (return) trip from one bank of a river to the other and back

B. A freely suspended bar magnet displaced from its N-S direction and released.

C. A hydrogen molecule rotating about its centre of mass

D. An arrow released from a bow.

Answer:



[Watch Video Solution](#)

2. Which of the following examples represent (nearly) simple harmonic motion and which represent periodic but not simple harmonic motion? The rotation of earth about its axis.



[Watch Video Solution](#)

3. Which of the following examples represent (nearly) simple harmonic motion and which

represent periodic but not simple harmonic motion? motion of an oscillating mercury column in a U-tube.



[Watch Video Solution](#)

4. Which of the following examples represent (nearly) simple harmonic motion and which represent periodic but not simple harmonic motion? motion of a ball bearing inside a smooth curved bowl, when released from a point slightly above the lower most point



[Watch Video Solution](#)


5. Which of the following examples represent (nearly) simple harmonic motion and which represent periodic but not simple harmonic motion? General vibrations of a polyatomic molecule about its equilibrium position.



[Watch Video Solution](#)

6. Fig. 14.23 depicts four $x-t$ plots for linear motion of a particle. Which of the plots

represent periodic motion? What is the period of motion (in case of periodic motion)



[Watch Video Solution](#)

7. Which of the following functions of time represent (a) simple harmonic, (b) periodic but not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (co is any positive constant): $\sin \omega t - \cos \omega t$



[Watch Video Solution](#)

8. Which of the following functions of time represent (a) simple harmonic, (b) periodic but

not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (co is any positive constant): $\sin^3 \omega t$



[Watch Video Solution](#)

9. Which of the following functions of time represent (a) simple harmonic, (b) periodic but not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (co is any positive constant): $3 \cos \left(\left(\frac{\pi}{4} \right) - (2\omega t) \right)$



[Watch Video Solution](#)

10. Which of the following functions of time represent (a) simple harmonic, (b) periodic but not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (ω is any positive constant):

$$\cos \omega t + \cos 3\omega t = \cos 5\omega t$$



[Watch Video Solution](#)

11. Which of the following functions of time represent (a) simple harmonic, (b) periodic but not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (co is any positive constant): $\exp(-\omega^2 t^2)$



Watch Video Solution

12. Which of the following functions of time represent (a) simple harmonic, (b) periodic but

not simple harmonic, and (c) non-periodic motion? Give period for each case of periodic motion (ω is any positive constant):

$$1 + \omega t + \omega^2 t^2$$



[Watch Video Solution](#)

13. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of

velocity, acceleration and force on the particle when it is at the end A,



[Watch Video Solution](#)

14. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of velocity, acceleration and force on the particle when it is at the end B,



[Watch Video Solution](#)

15. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of velocity, acceleration and force on the particle when it is at the mid-point of AB going towards A,



Watch Video Solution

16. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of velocity, acceleration and force on the particle when it is at 2 cm away from B going towards A,



Watch Video Solution

17. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of velocity, acceleration and force on the particle when it is at 3 cm away from A going towards B, and



[Watch Video Solution](#)

18. A particle is in linear simple harmonic motion between two points, A and B, 10 cm apart. Take the direction from A to B as the positive direction and give the signs of velocity, acceleration and force on the particle when it is at 4 cm away from B going towards A



Watch Video Solution

19. Which of the following relationships between the acceleration and the displacement x of a particle involve simple harmonic motion?

A. $a=0.7x$

B. $a=-200x^2$

C. $a = -10x$

D. $a = 100x^3$

Answer:



Watch Video Solution

20. The motion of a particle executing simple harmonic motion is described by the displacement function, $x(t) = A \cos (\omega t + \phi)$. If the initial ($t = 0$) position of the particle is 1 cm and its initial velocity is $\omega \text{ cm} / \text{s}$, what are its amplitude and initial phase angle ? The angular frequency of the particle is $\pi \text{ s}^{-1}$. If instead of the cosine function, we choose the sine function to describe the SHM : $x = B \sin (\omega t + \alpha)$, what are the amplitude and initial

phase of the particle with the above initial conditions.




[Watch Video Solution](#)

21. A spring balance has a scale that reads from 0 to 50 kg. The length of the scale is 20 cm. A body suspended from this balance, when displaced and released, oscillates with a period of 0.6 s. What is the weight of the body ?




[Watch Video Solution](#)

22. A spring having with a spring constant $1200Nm^{-1}$ is mounted on a horizontal table as shown in Fig. 14.24. A mass of 3 kg is attached to the free end of the spring. The mass is then pulled sideways to a distance of 2.0 cm and released. Determine the frequency of oscillations,




Watch Video Solution

23. A spring having with a spring constant $1200Nm^{-1}$ is mounted on a horizontal table as shown in Fig. 14.24. A mass of 3 kg is attached to the free end of the spring. The mass is then pulled sideways to a distance of 2.0 cm and released. Determine maximum acceleration of the mass,



Watch Video Solution

24. A spring having with a spring constant $1200Nm^{-1}$ is mounted on a horizontal table as shown in Fig. 14.24. A mass of 3 kg is attached to the free end of the spring. The mass is then pulled sideways to a distance of 2.0 cm and released. Determine the maximum speed of the mass



Watch Video Solution

25. Plot the corresponding reference circle for each of the following simple harmonic motions. Indicate the initial ($t = 0$) position of the particle, the radius of the circle, and the angular speed of the rotating particle. For simplicity, the sense of rotation may be fixed to be anticlockwise in every case: (x is in cm and t is in s). $x = -2 \sin (3t + (\pi / 3))$



Watch Video Solution

26. Plot the corresponding reference circle for each of the following simple harmonic motions. Indicate the initial ($t = 0$) position of the particle, the radius of the circle, and the angular speed of the rotating particle. For simplicity, the sense of rotation may be fixed to be anticlockwise in every case: (x is in cm and t is in s). $x = \cos(\pi / (6 - t))$



Watch Video Solution

27. Plot the corresponding reference circle for each of the following simple harmonic motions. Indicate the initial ($t = 0$) position of the particle, the radius of the circle, and the angular speed of the rotating particle. For simplicity, the sense of rotation may be fixed to be anticlockwise in every case: (x is in cm and t is in s). $x = 3 \sin(2\pi t + (\pi / 4))$




[Watch Video Solution](#)

28. Plot the corresponding reference circle for each of the following simple harmonic motions. Indicate the initial ($t = 0$) position of the particle, the radius of the circle, and the angular speed of the rotating particle. For simplicity, the sense of rotation may be fixed to be anticlockwise in every case: (x is in cm and t is in s). $x = 2 \cos \pi t$



Watch Video Solution

29. Figure 14.26 (a) shows a spring of force constant k clamped rigidly at one end and a mass m attached to its free end. A force F applied at the free end stretches the spring. Figure 14.26 (b) shows the same spring with both ends free and attached to a mass m at either end. Each end of the spring in Fig. 14.26(b) is stretched by the same force F . What is the maximum extension of the spring in the two cases ?



(a)





(b)



Watch Video Solution

30. Figure 14.26 (a) shows a spring of force constant k clamped rigidly at one end and a

mass m attached to its free end. A force F applied at the free end stretches the spring. Figure 14.26 (b) shows the same spring with both ends free and attached to a mass m at either end. Each end of the spring in Fig. 14.26(b) is stretched by the same force F . If the mass in Fig. (a) and the two masses in Fig. (b) are released, what is the period of oscillation in each case ?



 [Watch Video Solution](#)

31. The piston in the cylinder head of a locomotive has a stroke (twice the amplitude)

of 1.0 m. If the piston moves with simple harmonic motion with an angular frequency of 200 rad/min , what is its maximum speed ?



[Watch Video Solution](#)

32. The acceleration due to gravity on the surface of moon is 1.7 ms^{-2} . What is the time period of a simple pendulum on the surface of moon if its time period on the surface of earth is 3.5 s? (g on the surface of earth is 9.8 ms^{-2})



Watch Video Solution

33. Answer the following questions : Time period of a particle in SHM depends on the force constant k and mass m of the particle: $T = 2\pi \sqrt{\frac{m}{k}}$. A simple pendulum executes SHM approximately. Why then is the time period of a pendulum independent of the mass of the pendulum?



Watch Video Solution

34. Answer the following questions : The motion of a simple pendulum is approximately simple harmonic for small angle oscillations. For larger angles of oscillation, a more involved analysis shows that T is greater than $2\pi\sqrt{\frac{l}{g}}$. Think of a qualitative argument to appreciate this result



Watch Video Solution

35. Answer the following questions : A man with a wristwatch on his hand falls from the top of a tower. Does the watch give correct time during the free fall ?



Watch Video Solution

36. Answer the following questions : What is the frequency of oscillation of a simple pendulum mounted in a cabin that is freely falling under gravity ?





[Watch Video Solution](#)

37. A simple pendulum of length l and having a bob of mass M is suspended in a car. The car is moving on a circular track of radius R with a uniform speed v . If the pendulum makes small oscillations in a radial direction about its equilibrium position, what will be its time period ?



[Watch Video Solution](#)

38. A cylindrical piece of cork of density ρ and base area A and height h floats in a liquid of density ρ_1 . The cork is depressed slightly and then released. Show that the cork oscillates up and down simple harmonically with a period $T =$

$$2\pi \sqrt{\frac{h\rho}{\rho_1 g}}$$

where ρ is the density of cork.

(Ignore damping due to viscosity of the liquid)



Watch Video Solution


39. One end of a U-tube containing mercury is connected to a suction pump and the other end to atmosphere. A small pressure difference is maintained between the two columns. Show that, when the suction pump is removed, the column of mercury in the U-tube executes simple harmonic motion.



Watch Video Solution

40. An air chamber of volume V has a neck area of cross section a into which a ball of mass m just fits and can move up and down without any friction (Fig.14.27). Show that when the ball is pressed down a little and released, it executes SHM. Obtain an expression for the time period of oscillations assuming pressure-volume variations of air to

be isothermal [see Fig. 14.27



[Watch Video Solution](#)

41. You are riding in an automobile of mass 3000 kg. Assuming that you are examining the oscillation characteristics of its suspension

system. The suspension sags 15 cm when the entire automobile is placed on it. Also, the amplitude of oscillation decreases by 50% during one complete oscillation. Estimate the values of : the spring constant k .



[Watch Video Solution](#)

42. You are riding in an automobile of mass 3000 kg. Assuming that you are examining the oscillation characteristics of its suspension system. The suspension sags 15 cm when the

entire automobile is placed on it. Also, the amplitude of oscillation decreases by 50% during one complete oscillation. Estimate the values of : the damping constant b for the spring and shock absorber system of one wheel, assuming that each wheel supports 750 kg.



[Watch Video Solution](#)

43. Show that for a particle in linear SHM the average kinetic energy over a period of

oscillation equals the average potential energy over the same period.



[Watch Video Solution](#)

44. A circular disc of mass 10 kg is suspended by a wire attached to its centre. The wire is twisted by rotating the disc and released. The period of torsional oscillations is found to be 1.5 s. The radius of the disc is 15 cm. Determine the torsional spring constant of the wire. (Torsional spring constant α is defined by the

relation $J = -\alpha\theta$, where J is the restoring couple and θ the angle of twist).



[Watch Video Solution](#)

45. A body describes simple harmonic motion with an amplitude of 5 cm and a period of 0.2 s. Find the acceleration and velocity of the body when the displacement is 5 cm.



[Watch Video Solution](#)

46. A body describes simple harmonic motion with an amplitude of 5 cm and a period of 0.2 s. Find the acceleration and velocity of the body when the displacement is 3 cm.



Watch Video Solution

47. A body describes simple harmonic motion with an amplitude of 5 cm and a period of 0.2 s. Find the acceleration and velocity of the body when the displacement is 0 cm.





Watch Video Solution

48. A mass attached to a spring is free to oscillate, with angular velocity ω , in a horizontal plane without friction or damping. It is pulled to a distance x_0 and pushed towards the centre with a velocity v_0 at time $t = 0$. Determine the amplitude of the resulting oscillations in terms of the parameters ω , x_0 and v_0 .



Watch Video Solution

