



PHYSICS

BOOKS - PSEB

UNITS AND MEASUREMENT

Exercise

1. Fill in the blanks:- The volume of a cube side 1 cm is equal to m^3



Watch Video Solution

2. Fill in the blanks:- The surface area of a solid cylinder of radius 2.0 cm and height 10.0 cm is equal to.....(mm)²



[Watch Video Solution](#)

3. Fill in the blanks:- A vehicle moving with a speed of 18 km 'h⁻¹' covers.....m in 1 s



[Watch Video Solution](#)

4. Fill in the blanks:- The relative density of lead is

11.3. its density is gcm^{-3} or kgm^{-3}



Watch Video Solution

5. Fill in the blanks by suitable conversion of

units:- $1kgm^2s^{-2} = \dots\dots gcm^2s^{-2}$



Watch Video Solution

6. Fill in the blanks by suitable conversion of

units:- $1m = \dots .1y$

 [Watch Video Solution](#)

7. Fill in the blanks by suitable conversion of units:-

$$3.0ms^{-2} = \dots kmh^{-2}$$

 [Watch Video Solution](#)

8. Fill in the blanks by suitable conversion of units:-

$$G = 6.67 \times 10^{-11} Nm^2(kg)^{-2} = \dots (cm)^3 s^{-2} g^{-1}$$

 [Watch Video Solution](#)

9. A calorie is a unit of heat or energy and it equal about 4.2 J where $1J = 1kgm^2s^{-2}$. Suppose we employ a system of units in which the unit of mass equals α kg, the unit of length equals β m, the unit of time is γ s. Show that a calorie has a magnitude $4.2 \alpha^{-1} \beta^{-2} \gamma^2$ in terms of the new units.



[Watch Video Solution](#)

10. Explain this statement clearly :- “To call a dimensional quantity ‘large’ or ‘small’ is

meaningless without specifying a standard for comparison”. In view of this, reframe the following statements wherever necessary :- atoms are very small objects



[Watch Video Solution](#)

11. Explain this statement clearly :- “To call a dimensional quantity ‘large’ or ‘small’ is meaningless without specifying a standard for comparison”. In view of this, reframe the following statements wherever necessary :- a jet plane moves with great speed



[Watch Video Solution](#)

12. Explain this statement clearly :- “To call a dimensional quantity ‘large’ or ‘small’ is meaningless without specifying a standard for comparison”. In view of this, reframe the following statements wherever necessary :- atoms are very small objects



[Watch Video Solution](#)

13. Explain this statement clearly :- “To call a dimensional quantity ‘large’ or ‘small’ is

meaningless without specifying a standard for comparison". In view of this, reframe the following statements wherever necessary :- a jet plane moves with great speed



[Watch Video Solution](#)

14. Explain this statement clearly :- " To call a dimensional quantity 'large' or 'small' is meaningless without specifying a standard for comparison". In view of this, reframe the following statements wherever necessary :- a proton is much more massive than an electron



[Watch Video Solution](#)

15. Explain this statement clearly :- " To call a dimensional quantity 'large' or 'small' is meaningless without specifying a standard for comparison". In view of this, reframe the following statements wherever necessary :- the speed of sound is much smaller than the speed of light.



[Watch Video Solution](#)

16. A new unit of length is chosen such that the speed of light in vacuum is unity. What is the

distance between the Sun and the Earth in terms of the new unit if light takes 8 min and 20 s to cover this distance ?



[Watch Video Solution](#)

17. Which of the following is the most precise device for measuring length :-

A. a vernier callipers with 20 divisions on the sliding scale

B. a screw gauge of pitch 1 mm and 100 divisions on the circular scale

C. an optical instrument that can measure length to within a wavelength of light ?

D.

Answer:



Watch Video Solution

18. A student measures the thickness of a human hair by looking at it through a microscope of magnification 100. He makes 20 observations and finds that the average width of the hair in the field

of view of the microscope is 3.5 mm. What is the estimate on the thickness of hair ?



[Watch Video Solution](#)

19. Answer the following :- You are given a thread and a metre scale. How will you estimate the diameter of the thread ?



[Watch Video Solution](#)

20. Answer the following :- screw gauge has a pitch of 1.0 mm and 200 divisions on the circular

scale. Do you think it is possible to increase the accuracy of the screw gauge arbitrarily by increasing the number of divisions on the circular scale ?



[Watch Video Solution](#)

21. Answer the following :- The mean diameter of a thin brass rod is to be measured by vernier callipers. Why is a set of 100 measurements of the diameter expected to yield a more reliable estimate than a set of 5 measurements only ?



[Watch Video Solution](#)

22. The photograph of a house occupies an area of 1.75cm^2 on a 35 mm slide. The slide is projected on to a screen, and the area of the house on the screen is 1.55m^2 . What is the linear magnification of the projector-screen arrangement.



Watch Video Solution

23. State the number of significant figures in the following :- 0.007m^2



Watch Video Solution

24. State the number of significant figures in the following :- $2.64 \times 10^{24} \text{ kg}$

 [Watch Video Solution](#)

25. State the number of significant figures in the following :- 0.2370 g cm^{-3}

 [Watch Video Solution](#)

26. State the number of significant figures in the following :- 6.320 J

 [Watch Video Solution](#)

27. State the number of significant figures in the following :- $6.032Nm^{-2}$

 [Watch Video Solution](#)

28. State the number of significant figures in the following :- $0.0006032m^2$

 [Watch Video Solution](#)

29. The length, breadth and thickness of a rectangular sheet of metal are 4.234 m, 1.005 m, and 2.01 cm respectively. Give the area and volume of the sheet to correct significant figures.



Watch Video Solution

30. The mass of a box measured by a grocer's balance is 2.300 kg. Two gold pieces of masses 20.15 g and 20.17 g are added to the box. What is:- the difference in the masses of the pieces to correct significant figures ?



Watch Video Solution



[Watch Video Solution](#)

31. The mass of a box measured by a grocer's balance is 2.300 kg. Two gold pieces of masses 20.15 g and 20.17 g are added to the box. What is:- the difference in the masses of the pieces to correct significant figures ?



[Watch Video Solution](#)

32. A physical quantity P is related to four observables a , b , c and d as follows :-

$P = a^3 b^2 / ((\sqrt{c})d)$ The percentage errors of

measurement in a, b, c and d are 1%, 3%, 4% and 2%, respectively. What is the percentage error in the quantity P? If the value of P calculated using the above relation turns out to be 3.763, to what value should you round off the result ?



[Watch Video Solution](#)

33. A book with many printing errors contains four different formulas for the displacement y of a particle undergoing a certain periodic motion :-
 $y = a \sin 2\pi t / T$ (a = maximum displacement of the particle, v = speed of the particle. T = time-

period of motion). Rule out the wrong formulas on dimensional grounds.



[Watch Video Solution](#)

34. A book with many printing errors contains four different formulas for the displacement y of a particle undergoing a certain periodic motion :-
 $y = a \sin vt$ (a = maximum displacement of the particle, v = speed of the particle. T = time-period of motion). Rule out the wrong formulas on dimensional grounds.



[Watch Video Solution](#)

35. A book with many printing errors contains four different formulas for the displacement y of a particle undergoing a certain periodic motion :-
 $y = (a/T)\sin t/a$ (a = maximum displacement of the particle, v = speed of the particle. T = time-period of motion). Rule out the wrong formulas on dimensional grounds.



Watch Video Solution

36. A book with many printing errors contains four different formulas for the displacement y of a

particle undergoing a certain periodic motion :-

$$y = (a\sqrt{2})\left(\sin \frac{2\pi t}{T} + \cos \frac{2\pi t}{T}\right) \quad (a = \text{maximum displacement of the particle, } v = \text{speed of the particle. } T = \text{time-period of motion}).$$

Rule out the wrong formulas on dimensional grounds.



[Watch Video Solution](#)

37. A famous relation in physics relates 'moving mass' m to the 'rest mass' m_0 of a particle in terms of its speed v and the speed of light, c . (This relation first arose as a consequence of special relativity due to Albert Einstein). A boy recalls the

relation almost correctly but forgets where to put

the constant c . He writes :- $m = \frac{m_0}{(1 - v^2)^{\frac{1}{2}}}$ Guess

where to put the missing c .



[Watch Video Solution](#)

38. The unit of length convenient on the atomic scale is known as an angstrom and is denoted by

$\overset{\circ}{\text{A}}$: $1\overset{\circ}{\text{A}} = 10^{-10}m$. The size of a hydrogen atom is

about $0.5\overset{\circ}{\text{A}}$. What is the total atomic volume in

m^3 of a mole of hydrogen atoms ?



[Watch Video Solution](#)

39. One mole of an ideal gas at standard temperature and pressure occupies 22.4 L (molar volume). What is the ratio of molar volume to the atomic volume of a mole of hydrogen ? (Take the size of hydrogen molecule to be about $1\overset{\circ}{\text{A}}$). Why is this ratio so large ?



Watch Video Solution

40. Explain this common observation clearly : If you look out of the window of a fast moving train, the nearby trees, houses etc. seem to move rapidly

in a direction opposite to the train's motion, but the distant objects (hill tops, the Moon, the stars etc.) seem to be stationary. (In fact, since you are aware that you are moving, these distant objects seem to move with you).



[Watch Video Solution](#)

41. The nearest star to our solar system is 4.29 light years away. How much is this distance in terms of parsecs? How much parallax would this star (named Alpha Centauri) show when viewed

from two locations of the Earth six months apart in its orbit around the Sun ?



Watch Video Solution

42. Precise measurements of physical quantities are a need of science. For example, to ascertain the speed of an aircraft, one must have an accurate method to find its positions at closely separated instants of time. This was the actual motivation behind the discovery of radar in World War II. Think of different examples in modern science where precise measurements of length,

time, mass etc. are needed. Also, wherever you can, give a quantitative idea of the precision needed.



Watch Video Solution

43. Just as precise measurements are necessary in science, it is equally important to be able to make rough estimates of quantities using rudimentary ideas and common observations. Think of ways by which you can estimate the following (where an estimate is difficult to obtain, try to get an upper bound on the quantity) :- the total mass of rain-bearing clouds over India during the Monsoon



[Watch Video Solution](#)

44. Just as precise measurements are necessary in science, it is equally important to be able to make rough estimates of quantities using rudimentary ideas and common observations. Think of ways by which you can estimate the following (where an estimate is difficult to obtain, try to get an upper bound on the quantity) :- the wind speed during a storm



[Watch Video Solution](#)

45. Just as precise measurements are necessary in science, it is equally important to be able to make rough estimates of quantities using rudimentary ideas and common observations. Think of ways by which you can estimate the following (where an estimate is difficult to obtain, try to get an upper bound on the quantity) :- the number of strands of hair on your head



Watch Video Solution

46. Just as precise measurements are necessary in science, it is equally important to be able to make rough estimates of quantities using rudimentary ideas and common observations. Think of ways by which you can estimate the following (where an estimate is difficult to obtain, try to get an upper bound on the quantity) :- the number of air molecules in your classroom.



Watch Video Solution

47. The Sun is a hot plasma (ionized matter) with its inner core at a temperature exceeding 10^7 K, and its outer surface at a temperature of about 6000 K. At these high temperatures, no substance remains in a solid or liquid phase. In what range do you expect the mass density of the Sun to be, in the range of densities of solids and liquids or gases ? Check if your guess is correct from the following data : mass of the Sun = $2.0 \times 10^{30} \text{ kg}$, radius of the Sun = $7.0 \times 10^8 \text{ m}$.



[Watch Video Solution](#)

48. When the planet Jupiter is at a distance of 824.7 million kilometers from the Earth, its angular diameter is measured to be $35.72''$ of arc. Calculate the diameter of Jupiter.



Watch Video Solution

49. A man walking briskly in rain with speed v must slant his umbrella forward making an angle θ with the vertical. A student derives the following relation between θ and v : $\tan \theta = v$ and checks that the relation has a correct limit: as $v \rightarrow 0$, $\theta \rightarrow 0$, as expected. (We are assuming there is no

strong wind and that the rain falls vertically for a stationary man). Do you think this relation can be correct ? If not, guess the correct relation.



[Watch Video Solution](#)

50. It is claimed that two cesium clocks, if allowed to run for 100 years, free from any disturbance, may differ by only about 0.02 s. What does this imply for the accuracy of the standard cesium clock in measuring a time-interval of 1 s ?



[Watch Video Solution](#)

51. Estimate the average mass density of a sodium atom assuming its size to be about 2.5\AA . (Use the known values of Avogadro's number and the atomic mass of sodium). Compare it with the density of sodium in its crystalline phase : 970kgm^{-3} . Are the two densities of the same order of magnitude ? If so, why ?



Watch Video Solution

52. The unit of length convenient on the nuclear scale is a fermi : $1\text{ f} = 10^{-15}\text{m}$. Nuclear sizes obey roughly the following empirical relation :-

$r = (r_0 A)^{\frac{1}{3}}$ where r_0 is the radius of the nucleus, A is mass number, and r_0 is a constant equal to about, 1.2 f. Show that the rule implies that nuclear mass density is nearly constant for different nuclei. Estimate the mass density of sodium nucleus. Compare it with the average mass density of a sodium atom obtained in Exercise. 2.27.



[Watch Video Solution](#)

53. A LASER is a source of very intense, monochromatic, and unidirectional beam of light.

These properties of a laser light can be exploited to measure long distances. The distance of the Moon from the Earth has been already determined very precisely using a laser as a source of light. A laser light beamed at the Moon takes 2.56 s to return after reflection at the Moon's surface. How much is the radius of the lunar orbit around the Earth ?



[Watch Video Solution](#)

54. A SONAR (sound navigation and ranging) uses ultrasonic waves to detect and locate objects

under water. In a submarine equipped with a SONAR the time delay between generation of a probe wave and the reception of its echo after reflection from an enemy submarine is found to be 77.0 s. What is the distance of the enemy submarine? (Speed of sound in water = 1450m s^{-1}).



[Watch Video Solution](#)

55. The farthest objects in our Universe discovered by modern astronomers are so distant that light emitted by them takes billions of years to reach

the Earth. These objects (known as quasars) have many puzzling features, which have not yet been satisfactorily explained. What is the distance in km of a quasar from which light takes 3.0 billion years to reach us ?



[Watch Video Solution](#)

56. A great physicist of this century (P.A.M. Dirac) loved playing with numerical values of fundamental constants of nature. This led him to an interesting observation. Dirac found that from the basic constants of atomic physics (c , e , mass of

electron, mass of proton) and the gravitational constant G , he could arrive at a number with the dimension of time. Further, it was a very large number, its magnitude being close to the present estimate on the age of the universe (*~15 billion years*). From the table of fundamental constants in this book, try to see if you too can construct this number (or any other interesting number you can think of). If its coincidence with the age of the universe were significant, what would this imply for the constancy of fundamental constants ?



[Watch Video Solution](#)

