



PHYSICS

BOOKS - PSEB

WORK, ENERGY AND POWER

Exercise

1. The sign of work done by a force on a body is important to understand. State carefully if the following quantities are positive or negative:-

work done by a man in lifting a bucket out of a well by means of a rope tied to the bucket.



[Watch Video Solution](#)

2. The sign of work done by a force on a body is important to understand. State carefully if the following quantities are positive or negative:-
work done by friction on a body sliding down an inclined plane.



[Watch Video Solution](#)

3. The sign of work done by a force on a body is important to understand. State carefully if the following quantities are positive or negative:-
work done by an applied force on a body moving on a rough horizontal plane with uniform velocity.



[Watch Video Solution](#)

4. The sign of work done by a force on a body is important to understand. State carefully if the following quantities are positive or negative:-

work done by the resistive force of air on a vibrating pendulum in bringing it to rest.



[Watch Video Solution](#)

5. A body of mass 2 kg initially at rest moves under the action of an applied horizontal force of 7 N on a table with coefficient of kinetic friction = 0.1. Compute the:- work done by the applied force in 10 s.



[Watch Video Solution](#)

6. A body of mass 2 kg initially at rest moves under the action of an applied horizontal force of 7 N on a table with coefficient of kinetic friction = 0.1. Compute the:- work done by friction in 10 s.



[Watch Video Solution](#)

7. A body of mass 2 kg initially at rest moves under the action of an applied horizontal force of 7 N on a table with coefficient of kinetic friction = 0.1. Compute the:- work done by the net force on the body in 10 s.




 [Watch Video Solution](#)

8. A body of mass 2 kg initially at rest moves under the action of an applied horizontal force of 7 N on a table with coefficient of kinetic friction = 0.1. Compute the:- change in kinetic energy of the body in 10 s.

 [Watch Video Solution](#)

9. The potential energy function for a particle executing linear simple harmonic motion is given by $V(x) = kx^2 / 2$, where k is the force constant of

the oscillator. For $k = 0.5 \text{ Nm}^{-1}$, the graph of $V(x)$ versus x is shown in Fig. 6.12. Show that a particle of total energy 1 J moving under this potential must 'turn back' when it reaches $x = \pm 2 \text{ m}$.



Watch Video Solution


10. Answer the following :- The casing of a rocket in flight burns up due to friction. At whose expense is the heat energy required for burning obtained? The rocket or the atmosphere?



Watch Video Solution

11. Answer the following :- Comets move around the sun in highly elliptical orbits. The gravitational force on the comet due to the sun is not normal to the comet's velocity in general. Yet the work done by the gravitational force over every


complete orbit of the comet is zero. Why ?



[Watch Video Solution](#)

12. Answer the following :- An artificial satellite orbiting the earth in very thin atmosphere loses its energy gradually due to dissipation against atmospheric resistance, however small. Why then does its speed increase progressively as it comes


closer and closer to the earth ?



Watch Video Solution

13. Answer the following :- In Fig. 6.13(i) the man walks 2 m carrying a mass of 15 kg on his hands. In Fig. 6.13(ii), he walks the same distance pulling the rope behind him. The rope goes over a pulley, and a mass of 15 kg hangs at its other end. In which

case is the work done greater ?



Watch Video Solution

14. Underline the correct alternative :- When a conservative force does positive work on a body, the potential energy of the body \in creases / decreases / remains the same.



Watch Video Solution

15. Underline the correct alternative :- Work done by a body against friction always results in a loss of its kinetic / potential energy.



Watch Video Solution

16. Underline the correct alternative :- The rate of change of total momentum of a many-particle system is proportional to the external force / sum of the internal forces on the system.



Watch Video Solution

17. Underline the correct alternative :- In an inelastic collision of two bodies, the quantities which do not change after the collision are the total _____ kinetic _____

$e \neq rgy / \rightarrow tall \in earmomentum / \rightarrow tale \neq rgy$
of the system of two bodies.



[Watch Video Solution](#)

18. State if each of the following statements is true or false. Give reasons for your answer:- In an

elastic collision of two bodies, the momentum and energy of each body is conserved.



[Watch Video Solution](#)

19. State if each of the following statements is true or false. Give reasons for your answer:- Total energy of a system is always conserved, no matter what internal and external forces on the body are present.



[Watch Video Solution](#)

20. State if each of the following statements is true or false. Give reasons for your answer:- Work done in the motion of a body over a closed loop is zero for every force in nature.



Watch Video Solution

21. State if each of the following statements is true or false. Give reasons for your answer:- In an inelastic collision, the final kinetic energy is always less than the initial kinetic energy of the system.



Watch Video Solution

22. Answer carefully, with reasons :-In an elastic collision of two billiard balls, is the total kinetic energy conserved during the short time of collision of the balls (i.e. when they are in contact) ?



Watch Video Solution

23. State if each of the following statements is true or false. Give reasons for your answer:- In an elastic collision of two bodies, the momentum and energy of each body is conserved.



[Watch Video Solution](#)

24. State if each of the following statements is true or false. Give reasons for your answer:- In an elastic collision of two bodies, the momentum and energy of each body is conserved.



[Watch Video Solution](#)

25. A body is initially at rest. It undergoes one-dimensional motion with constant acceleration.

The power delivered to it at time t is proportional to

A. $t^{\frac{1}{2}}$

B. t

C. $t^{\frac{3}{2}}$

D. t^2

Answer:



Watch Video Solution

26. A body is moving unidirectionally under the influence of a source of constant power. Its displacement in time t is proportional to

A. $t^{\frac{1}{2}}$

B. t

C. $t^{\frac{3}{2}}$

D. t^2

Answer:



Watch Video Solution

27. A body constrained to move along the z-axis of a coordinate system is subject to a constant force F given by $F = -\hat{i} + 2\hat{j} + 3\hat{k}N$ where $\hat{i}, \hat{j}, \hat{k}$ are unit vectors along the x-, y- and z-axis of the system respectively. What is the work done by this force in moving the body a distance of 4 m along the z-axis ?



[Watch Video Solution](#)

28. An electron and a proton are detected in a cosmic ray experiment, the first with kinetic energy 10 keV, and the second with 100 keV. Which

is faster, the electron or the proton ? Obtain the ratio of their speeds, (electron mass = $9.11 \times 10^{-31} \text{ kg}$, proton mass = $1.67 \times 10^{-27} \text{ kg}$, $1\text{eV} = 1.60 \times 10^{-19} \text{ J}$)



[Watch Video Solution](#)

29. A rain drop of radius 2 mm falls from a height of 500 m above the ground. It falls with decreasing acceleration (due to viscous resistance of the air) until at half its original height, it attains its maximum (terminal) speed, and moves with uniform speed thereafter. What is the work

done by the gravitational force on the drop in the first and second half of its journey ? What is the work done by the resistive force in the entire journey if its speed on reaching the ground is 10ms^{-1} ?



[Watch Video Solution](#)

30. A molecule in a gas container hits a horizontal wall with speed 200 m s^{-1} and angle 30° with the normal, and rebounds with the same speed. Is momentum conserved in the collision ? Is the collision elastic or inelastic ?



[Watch Video Solution](#)


31. A pump on the ground floor of a building can pump up water to fill a tank of volume $30m^3$ in 15 min. If the tank is 40 m above the ground, and the efficiency of the pump is 30%, how much electric power is consumed by the pump ?



[Watch Video Solution](#)

32. Two identical ball bearings in contact with each other and resting on a frictionless table are


hit head-on by another ball bearing of the same mass moving initially with a speed V . If the collision is elastic, which of the following (Fig. 6.14) is a possible result after collision ?



 [Watch Video Solution](#)

33. The bob A of a pendulum released from 30° to the vertical hits another bob B of the same mass

at rest on a table as shown in Fig. 6.15. How high does the bob A rise after the collision ? Neglect the size of the bobs and assume the collision to be elastic.



[Watch Video Solution](#)

34. The bob of a pendulum is released from a horizontal position. If the length of the pendulum is 1.5 m, what is the speed with which the bob arrives at the lowermost point, given that it dissipated 5% of its initial energy against air resistance ?



Watch Video Solution

35. A trolley of mass 300 kg carrying a sandbag of 25 kg is moving uniformly with a speed of $27\text{km} / \text{h}$ on a frictionless track. After a while, sand

starts leaking out of a hole on the floor of the trolley at the rate of 0.05kg s^{-1} . What is the speed of the trolley after the entire sand bag is empty ?



[Watch Video Solution](#)

36. A body of mass 0.5 kg travels in a straight line with velocity $v = ax \left(\frac{3}{2}\right)$ where $a = 5\text{m} \left(\frac{1}{2}\right) \text{s}^{-1}$. What is the work done by the net force during its displacement from $x = 0$ to $x = 2\text{ m}$?



[Watch Video Solution](#)

37. The blades of a windmill sweep out a circle of area:- If the wind flows at a velocity v perpendicular to the circle, what is the mass of the air passing through it in time t ?

 [Watch Video Solution](#)

38. The blades of a windmill sweep out a circle of area :-What is the kinetic energy of the air ?

 [Watch Video Solution](#)

39. The blades of a windmill sweep out a circle of area :- Assume that the windmill converts 25% of the wind's energy into electrical energy, and that $A = 30m^2$, $v = 36km/h$ and the density of air is $1.2kg, m^{-3}$. What is the electrical power produced ?



Watch Video Solution

40. A person trying to lose weight (dieter) lifts a 10 kg mass, one thousand times, to a height of 0.5 m each time. Assume that the potential energy lost each time she lowers the mass is dissipated:-

How much work does she do against the gravitational force ?



[Watch Video Solution](#)

41. A person trying to lose weight (dieter) lifts a 10 kg mass, one thousand times, to a height of 0.5 m each time. Assume that the potential energy lost each time she lowers the mass is dissipated:- Fat supplies $3.8 \times 10^7 J$ of energy per kilogram which is converted to mechanical energy with a 20% efficiency rate. How much fat will the dieter use up?



[Watch Video Solution](#)

42. A family uses 8 kW of power:- Direct solar energy is incident on the horizontal surface at an average rate of 200 W per square meter. If 20% of this energy can be converted to useful electrical energy, how large an area is needed to supply 8 kW?



[Watch Video Solution](#)


43. A bullet of mass 0.012 kg and horizontal speed 70ms^{-1} strikes a block of wood of mass 0.4 kg and instantly comes to rest with respect to the block. The block is suspended from the ceiling by means of thin wires. Calculate the height to which the block rises. Also, estimate the amount of heat produced in the block.



Watch Video Solution

44. Two inclined frictionless tracks, one gradual and the other steep meet at A from where two


stones are allowed to slide down from rest, one on each track (Fig. 6.16). Will the stones reach the bottom at the same time? Will they reach there with the same speed? Explain. Given $\theta_1 = 30^\circ$, $\theta_2 = 60^\circ$, and $h = 10$ m, what are the speeds and times taken by the two stones ?



[Watch Video Solution](#)

45. A 1 kg block situated on a rough incline is connected to a spring of spring constant $100Nm^{-1}$ as shown in Fig. 6.17. The block is released from rest with the spring in the unstretched position. The block moves 10 cm down the incline before coming to rest. Find the coefficient of friction between the block and the incline. Assume that the spring has a negligible

mass and the pulley is frictionless.



[Watch Video Solution](#)

46. A bolt of mass 0.3 kg falls from the ceiling of an elevator moving down with a uniform speed of 7 m s^{-1} . It hits the floor of the elevator (length of the elevator = 3 m) and does not rebound. What is the heat produced by the impact? Would

your answer be different if the elevator were stationary ?



Watch Video Solution

47. A trolley of mass 200 kg moves with a uniform speed of 36 km/h on a frictionless track. A child of mass 20 kg runs on the trolley from one end to the other (10 m away) with a speed of 4 m s^{-1} relative to the trolley in a direction opposite to the its motion, and jumps out of the trolley. What is the final speed of the trolley ? How much has



the trolley moved from the time the child begins to run ?



[Watch Video Solution](#)

48. Which of the following potential energy curves in Fig. 6.18 cannot possibly describe the elastic collision of two billiard balls ? Here r is the

distance between centres of the balls.



[Watch Video Solution](#)