

## CHEMISTRY

### BOOKS - VIKRAM PUBLICATION ( ANDHRA PUBLICATION)

## STOICHIOMETRY

### Solved Problems

1. Calculate molecular mass of glucose ( $C_6H_{12}O_6$ ) molecule.

 [Watch Video Solution](#)

2. A compound contains 4.07% hydrogen, 24.27% carbon and 71.65% chlorine. Its molar mass is 98.96 g. What are its empirical and molecular formulas ?

 [Watch Video Solution](#)

3. Calculate the amount of water (g) produced by the combustion of 16 g of methane.

 [Watch Video Solution](#)

4. How many moles of methane are required to produce 22 g  $CO_2(g)$  after combustion?

 [Watch Video Solution](#)

5. 50.0 kg of  $N_2(g)$  and 10.0 kg of  $H_2(g)$  are mixed to produce  $NH_2(g)$ . Calculate the  $NH_2(g)$  formed. Identify the limiting reagent in the production of  $NH_3$  in this situation.

 [Watch Video Solution](#)

6. A solution is prepared by adding 2 g of a substance A to 18 g of water. Calculate the mass per cent of the solute.

 [Watch Video Solution](#)

7. Calculate the molarity of NaOH in the solution prepared by dissolving its 4 g in enough water to form 250 mL of the solution.

 [Watch Video Solution](#)

8. The density of 3M solution of NaCl is  $1.25 \text{ gmL}^{-1}$ . Calculate molality of the solution.

 [Watch Video Solution](#)

9. Calculate the normality of oxalic acid solutions containing 6.3g of  $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  in 500 ml of solutions.






[Watch Video Solution](#)

10. Calculate the mass of  $Na_2CO_3$  required to prepare 250 ml of 0.5 N solution.

[Watch Video Solution](#)

11. In the reactions given below, identify the species undergoing oxidation and reduction :

[Watch Video Solution](#)

12. Justify that the reaction :  $2Na(s) + H_2(g) \rightarrow 2NaH(s)$  is a redox change.

[Watch Video Solution](#)

13. Using Stock notation, represent the following compounds :

$H AuCl_4$ ,  $Tl_2O$ ,  $FeO$ ,  $Fe_2O_3$ ,  $CuI$ ,  $CuO$ ,  $MnO$  and  $MnO_2$ .

 [Watch Video Solution](#)

14. Justify that the reaction :

$2Cu_2O(s) + Cu_2S(s) \rightarrow 6Cu(s) + SO_2(g)$  is a redox reaction. Identify

the species oxidised/reduced, which acts as an oxidant and which acts as a reductant.

 [Watch Video Solution](#)




15. Which of the following species, do not show disproportionation reaction and why ?

$ClO^-$ ,  $ClO_2^-$ ,  $ClO_3^-$  and  $ClO_4^-$

Also write reaction for each of the species that disproportionates.



 [Watch Video Solution](#)

16. Suggest a scheme of classification of the following redox reactions



Watch Video Solution

17. Why do the following reactions proceed differently?



Watch Video Solution

18. Write the net ionic equation for the reaction of potassium dichromate (VI),  $K_2Cr_2O_7$  with sodium sulphite,  $Na_2SO_3$ , in an acid solution to give

chromium (III) ion and the sulphate ion.

 [Watch Video Solution](#)

19. Permanganate ion reacts with bromide ion in basic medium to give manganese dioxide and bromate ion. Write the balanced ionic equation for the reaction.

 [Watch Video Solution](#)

20. Permanganate (VII) ion,  $MnO_4^-$  in basic solution oxidises iodide ion,  $I^-$  to produce molecular iodine ( $I_2$ ) and manganese (IV) oxide ( $MnO_2$ ). Write a balanced ionic equation to represent this redox reaction.

 [Watch Video Solution](#)

Very Short Answer Questions

1. How many number of moles of glucose are present in 540 gms of glucose ?

 [Watch Video Solution](#)

2. Calculate the weight of 0.1 mole of sodium carbonate.

 [Watch Video Solution](#)

3. How many molecules of glucose are present in 5.23g of glucose (Molecular weight of glucose 180 u).

 [Watch Video Solution](#)

4. Calculate the number of molecules present in  $1.12 \times 10^{-7}$  c.c. of a gas at STP (c.c.- cubic centimeters =  $cm^3$ ).


 [Watch Video Solution](#)



5. The empirical formula of a compound is  $CH_2O$ . Its molecular weight is 90. Calculate the molecular formula of the compound.

 [Watch Video Solution](#)

6. Balance the following equation by the oxidation number method.



 [Watch Video Solution](#)

7. What volume of  $H_2$  at STP is required to reduce 0.795 g of  $CuO$  to give  $Cu$  and  $H_2O$ .

 [Watch Video Solution](#)

8. Calculate the volume of  $O_2$  at STP required to completely burn 100 ml. of acetylene.

 [Watch Video Solution](#)

9. Now a days it is thought that oxidation is simply decrease in electron density and reduction is increase in electron density.

How would you justify this?

 [Watch Video Solution](#)

10. What is a redox concept? Give an example.

 [Watch Video Solution](#)

11. Calculate the mass percent of the different elements present in sodium sulphate ( $Na_2SO_4$ ).

 [Watch Video Solution](#)

12. What do you mean by significant figures?

 [Watch Video Solution](#)

13. If the speed of light is  $3.0 \times 10^8 \text{ms}^{-1}$ . Calculate the distance covered by light in  $2.00 \text{ns}$ .

 [Watch Video Solution](#)

## Short Answer Questions

1. The approximate production of sodium carbonate per month is  $424 \times 10^6 \text{g}$ . While that of methyl alcohol is  $320 \times 10^6 \text{ gm}$ . Which is produced more in terms of moles ?

 [Watch Video Solution](#)

2. How much minimum volume of CO at STP is needed to react completely with 0.112 L of  $O_2$  at 1.5 atm. Pressure and  $127^\circ C$  to give  $CO_2$ .

 [Watch Video Solution](#)

3. Chemical analysis of a carbon compound gave the following percentage composition by weight of the elements present, carbon = 10.06 %, hydrogen = 0.84 %, chlorine = 89.10 %. Calculate the empirical formula of the compound.

 [Watch Video Solution](#)

4. A carbon compound on analysis gave the following percentage composition, carbon 14.5%, hydrogen 1.8%, chlorine 64.46%, oxygen 19.24%. Calculate the empirical formula of the compound.

 [Watch Video Solution](#)

5. Calculate the empirical formula of a compound having percentage composition:

Potassium (K) = 26.57, Chromium (Cr) = 35.36, Oxygen (O) = 38.07.

(Given the Atomic weights of K, Cr and O are 39, 52 and 16 respectively)

 [Watch Video Solution](#)


6. A carbon compound contains 12.8% Carbon, 2.1% Hydrogen, 85.1% Bromine. The molecular weight of the compound is 187.9. Calculate the molecular formula.

 [Watch Video Solution](#)

7. 0.188 g of an organic compound having an empirical formula  $CH_2Br$  displaced 24.2 cc of air at  $14^\circ C$  and 752 mm pressure. Calculate the molecular formula of the compound. (Aqueous tension at  $14^\circ C$  is 12 mm)

 [Watch Video Solution](#)

8. Calculate the amount of 90%  $H_2SO_4$  required for the preparation of 420 kg HCl.



 [Watch Video Solution](#)

9. An astronaut receives the energy required in his body by the combustion of 34gm of sucrose per hour. How much oxygen he has to carry along with him for his energy requirement in a day?

 [Watch Video Solution](#)

10. What volume of  $CO_2$  is obtained at STP by heating 4 g of  $CaCO_3$  ?

 [Watch Video Solution](#)

11. When 50 gm of a sample of sulphur was burnt in air 4% of the sample was left over. Calculate the volume of air required at STP containing 21% oxygen by volume.

 [Watch Video Solution](#)

12. Calculate the volume of oxygen gas required at STP conditions for the complete combustion of 10 cc of methane gas at  $20^{\circ}C$  and 770mm pressure.

 [Watch Video Solution](#)

13. Calculate the volume of  $H_2$  liberated at  $27^{\circ}C$  and 760 mm of Hg pressure by action by 0.6 g magnesium with excess of dil HCl.

 [Watch Video Solution](#)

14. Explain the role of redox reactions in titrimetre processes and galvanic cells.

 [Watch Video Solution](#)

15. Define and explain molar mass.

 [Watch Video Solution](#)

16. What are disproportionation reactions? Give example.

 [Watch Video Solution](#)

17. What is comproportionation reactions? Give example.

 [Watch Video Solution](#)



18. Determine the empirical formula of an oxide of iron which has 69.9% iron and 30.1% dioxygen by mass.

 [Watch Video Solution](#)

19. Calculate the mass of sodium acetate ( $CH_3COONa$ ) required to make 500 ml. of 0.375 molar aqueous solution. Molar mass of sodium acetate is  $82.0245g\ mol^{-1}$ .

 [Watch Video Solution](#)

20. What is the concentration of sugar ( $C_{12}H_{22}O_{11}$ ) in  $mol\ L^{-1}$  if 20 g are dissolved in enough water to make a final volume upto 2L?

 [Watch Video Solution](#)

**21.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034



**Watch Video Solution**

**22.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034



**Watch Video Solution**

**23.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034



**Watch Video Solution**

**24.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034





[Watch Video Solution](#)

**25.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034



[Watch Video Solution](#)

**26.** How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034



[Watch Video Solution](#)

**27.** Round up the following upto three significant figures :

i) 34.216, ii) 10.4107, iii) 0.04597, iv) 2808



[Watch Video Solution](#)

**28.** Round up the following upto three significant figures :

i) 34.216, ii) 10.4107, iii) 0.04597, iv) 2808



[Watch Video Solution](#)

**29.** Round up the following upto three significant figures :

i) 34.216, ii) 10.4107, iii) 0.04597, iv) 2808



[Watch Video Solution](#)

**30.** Round up the following upto three significant figures :

i) 34.216, ii) 10.4107, iii) 0.04597, iv) 2808



[Watch Video Solution](#)

**31.** Calculate the molarity of a solution of ethanol in water in which the mole fraction of ethanol is 0.040 (assume the density of water to be one).

Use the data given in the following table to calculate the molar mass of naturally occurring argon isotopes :


Isotope	Isotopic molar mass	Abundance
$^{36}\text{Ar}$	$35.96755 \text{ g mol}^{-1}$	0.337%
$^{38}\text{Ar}$	$37.96272 \text{ g mol}^{-1}$	0.063%
$^{40}\text{Ar}$	$39.9624 \text{ g mol}^{-1}$	99.600 %

 [Watch Video Solution](#)

32. A welding fuel gas contains carbon and hydrogen only. Burning a small sample of it in oxygen gives 3.38 g carbon dioxide, 0.690 g of water and no other products. A volume of 10.0 L (measured at STP) of this welding gas is found to weigh 11.6 g. Calculate (i) empirical formula, (ii) molar mass of the gas, and (iii) molecular formula.

 [Watch Video Solution](#)


33. Calcium Carbonate reacts with aqueous HCl to give  $\text{CaCl}_2$  and  $\text{CO}_2$  according to the reaction,



What mass of  $\text{CaCO}_3$  is required to react completely with 25 ml of 0.75 M HCl?

 [Watch Video Solution](#)

**34.** Chlorine is prepared in the laboratory by treating manganese dioxide ( $\text{MnO}_2$ ) with aqueous hydrochloric acid according to the reaction



How many grams of HCl react with 5.0g of manganese dioxide ?

 [Watch Video Solution](#)

**35.** To 50 ml. of 0.1 N  $\text{Na}_2\text{CO}_3$  solution 150 ml. of  $\text{H}_2\text{O}$  is added. Then calculate the normality of resultant solution.

 [Watch Video Solution](#)

36. Calculate the volume of 0.1 N  $H_2SO_4$  required to neutralise 200 ml. of 0.2 N NaOH solution.

It is an acid base neutralisation reaction.

Hence, at the neutralisation point.

Number of equivalents of acid = Number of equivalents of base.

 [Watch Video Solution](#)








37. Calculate normality of  $H_2SO_4$  solutions if 50 ml of it completely neutralise 250 ml. of 0.1 N  $Ba(OH)_2$  solutions.

 [Watch Video Solution](#)

38. Calculate the volume of 0.1M  $KMnO_4$  required to react with 100 ml. of 0.1 M  $H_2C_2O_4$ .  
 $2H_2O$  solution in the presence of  $H_2SO_4$ .







 [Watch Video Solution](#)

39. Assign oxidation number to the underlined elements in each of the following species.



Watch Video Solution




40. Assign oxidation number to the underlined elements in each of the following species.



Watch Video Solution




41. Assign oxidation number to the underlined elements in each of the following species.







 [Watch Video Solution](#)

**42.** Assign oxidation number to the underlined elements in each of the following species.



 [Watch Video Solution](#)









**43.** Assign oxidation number to the underlined elements in each of the following species.












Watch Video Solution

44. Assign oxidation number to the underlined elements in each of the following species.



Watch Video Solution

45. Assign oxidation number to the underlined elements in each of the following species.



Watch Video Solution

46. Assign oxidation number to the underlined elements in each of the following species.

- a)  $NaH_2\underline{P}O_4$       b)  $NaH\underline{S}O_4$       c)  $H_4\underline{P}_2O_7$   
d)  $K_2\underline{Mn}O_4$       e)  $Ca\underline{O}_2$       f)  $Naun \partial \in e(B)\underline{H}_4$   
g)  $H_2\underline{S}_2O_7$       h)  $K\underline{A}l\underline{S}O_{4_2} \cdot 12H_2O$

 [Watch Video Solution](#)

47. What are the oxidation number to the underlined elements in each of the following and how do you rationalise your results?

- a)  $K\underline{I}_3$     b)  $H_2\underline{S}_4O_6$     c)  $\underline{F}e_3O_4$

 [Watch Video Solution](#)

48. What are the oxidation numbers of the underlined elements in each of the following and how do you rationalise your results?

- a)  $H_2\underline{S}_4O_6$       b)  $\underline{F}e_3O_4$   
c)  $\underline{C}H_3\underline{C}H_2OH$       d)  $\underline{C}H_3\underline{C}OOH$






 [Watch Video Solution](#)

49. What are the oxidation numbers of the underlined elements in each of the following and how do you rationalise your results?








 Watch Video Solution

50. Justify that the following reactions are redox reactions.



 Watch Video Solution

51. Justify that the following reactions are redox reactions.



 [Watch Video Solution](#)


52. Justify that the following reactions are redox reactions :

 [Watch Video Solution](#)

53. Justify that the following reactions are redox reactions :


 [Watch Video Solution](#)

54. Justify that the following reactions are redox reactions :



 [Watch Video Solution](#)

55. Fluorine reacts with ice and results in the change.



Justify that this reaction is a redox reaction.

 [Watch Video Solution](#)

56. Calculate the oxidation number of sulphur, chromium and nitrogen ion  $\text{H}_2\text{SO}_5$ ,  $\text{Cr}_2\text{O}_7^{2-}$  and  $\text{NO}_3^-$ . Suggest structure of those compounds.

 [Watch Video Solution](#)

**57.** Write the formulae for the following compounds.

a) Mercury (II) chloride

b) Nickel (II) sulphate

c) Tin (IV) oxide

d) Thallium (I) sulphate

e) Iron (III) sulphate

f) Chromium (III) oxide.

 [Watch Video Solution](#)

**58.** Suggest a list of the substances where carbon exhibit oxidation states from -4 to +4 and nitrogen from -3 to +5.



 [Watch Video Solution](#)

**59.** While sulphur dioxide and hydrogen peroxide act as oxidising as well as reducing agents in their reactions, ozone and nitric acid act only as oxidants. Why?




Watch Video Solution

**60.** Consider the reactions



Why it is more appropriate to write these reaction as



Also suggest a technique to investigate the path of the above (a) and (b) redox reactions.



Watch Video Solution

**61.** The compound  $AgF_2$  is unstable compound. However, if formed, the compound acts as a very strong oxidising agent. Why ?



Watch Video Solution



**62.** Whenever a reaction between an oxidising agent and a reducing agent is carried out, a compound of lower oxidation state is formed if the reducing agent is in excess and a compound of higher oxidation state is formed if the oxidising agent is in excess. Justify this statement giving three illustrations.

 [Watch Video Solution](#)


**63.** How do you count the following observations?

a) Though alkaline potassium permanganate and acidic potassium permanganate both are used as oxidants, yet in the manufacture of benzoic acid from toluene we use alcoholic potassium permanganate as an oxidant. Why? Write balanced redox equation for the reaction.


b) When concentrated sulphuric acid is added to inorganic mixture containing chloride, we get colourless pungent smelling gas HCl, but if the mixture contains bromide then we get red vapour of bromine. Why?

 [Watch Video Solution](#)




64. Identify the substance oxidised, reduced, oxidising agent and reducing agent for each of the following reactions :



b)




c)




Watch Video Solution


65. Identify the substance oxidised, reduced, oxidising agent and reducing agent for each of the following reactions :





b)



c)






 [Watch Video Solution](#)




**66.** Identify the substance oxidised, reduced, oxidising agent and reducing agent for each of the following reactions :



b)




c)




 [Watch Video Solution](#)




**67.** Identify the substance oxidised, reduced, oxidising agent and reducing agent for each of the following reactions :



b)




c)




Watch Video Solution




**68.** Identify the substance oxidised, reduced, oxidising agent and reducing agent for each of the following reactions :



b)





c)





Watch Video Solution

69. Consider the reactions



Why does the same reductant, thiosulphate react differently with iodine and bromine ?




Watch Video Solution

70. Justify giving reactions that among halogens, fluorine is the best oxidant and among hydrohalic compounds, hydroiodic acid is the best reductant.



Watch Video Solution

71. Why does the following reaction occur ?




What conclusion about the compound  $Na_4XeO_6$  (of which  $XeO_6^{4-}$  is a part) can be drawn from the reaction.


 [Watch Video Solution](#)

72. Consider the reactions :



a)



b)




c)



What inference do you draw about the behaviour of  $Ag^+$  and  $Cu^{2+}$  from these reactions?


 [Watch Video Solution](#)

**73.** Balance the following redox reaction in basic medium by ion-electron method :




[Watch Video Solution](#)

**74.** Balance the following redox reactions by ion-electron method :




[Watch Video Solution](#)

**75.** Balance the following redox reactions by ion-electron method :






[Watch Video Solution](#)

76. Balance the following Redox reaction by ion-electron method in acidic medium.




 [Watch Video Solution](#)

77. Balance the following equations in basic medium by ion-electron method and oxidation number methods and identify the oxidising agent and the reducing agent.



 [Watch Video Solution](#)

78. What sorts of information can you draw from the following reaction ?



 [Watch Video Solution](#)





Watch Video Solution

79. The  $Mn^{3+}$  ion is unstable solution and undergoes disproportionation to give  $Mn^{2+}$ ,  $MnO_2$  and  $H^+$  ion. Write balanced ionic equation for the reaction.



Watch Video Solution

80. Consider the elements Cs, Ne, I and F.

- Identify the element that exhibits only negative oxidation state.
- Identify the element that exhibits only positive oxidation state.
- Identify the element that exhibit both positive and negative oxidation states
- Identify the element which neither exhibit the negative nor does the positive oxidation state.



Watch Video Solution

**81.** Consider the elements Cs, Ne, I and F.

- a) Identify the element that exhibits only negative oxidation state.
- b) Identify the element that exhibits only positive oxidation state.
- c) Identify the element that exhibit both positive and negative oxidation states
- d) Identify the element which neither exhibit the negative nor does the positive oxidation state.

 [Watch Video Solution](#)

**82.** Consider the elements Cs, Ne, I and F.

- a) Identify the element that exhibits only negative oxidation state.
- b) Identify the element that exhibits only positive oxidation state.
- c) Identify the element that exhibit both positive and negative oxidation states
- d) Identify the element which neither exhibit the negative nor does the positive oxidation state.

 [Watch Video Solution](#)

**83.** Consider the elements Cs, Ne, I and F.

- a) Identify the element that exhibits only negative oxidation state.
- b) Identify the element that exhibits only positive oxidation state.
- c) Identify the element that exhibit both positive and negative oxidation states
- d) Identify the element which neither exhibit the negative nor does the positive oxidation state.

 [Watch Video Solution](#)

**84.** Chlorine is used to purify drinking water. Excess of Chlorine is harmful.

The excess of Chlorine is removed by treating with sulphur dioxide.

Present a balanced equation for this redox change taking place in water.

 [Watch Video Solution](#)

**85.** Refer to the periodic table given in your book and now answer the following questions.

- a) Select the possible non metals that can show disproportionation reaction
- b) Select the metals that can show disproportionation

 [Watch Video Solution](#)

**86.** Refer to the periodic table given in your book and now answer the following questions.

- a) Select the possible non metals that can show disproportionation reaction
- b) Select the metals that can show disproportionation

 [Watch Video Solution](#)

**87.** In Ostwal's process for the manufacture of nitric acid the first step involves the oxidation of ammonia gas by oxygen gas to give nitric oxide

gas and steam. What is the maximum weight of nitric oxide that can be obtained starting only with 10.00 g of ammonia and 20.00 g of oxygen.

 [Watch Video Solution](#)

88. i) Arrange the following metals in the order in which they displace each other from the solution of their salts.

Al, Cu, Fe, Mg and Zn

ii) Calculate the molarity of sodium carbonate in a solution prepared by dissolving 5.3 g in enough water to form 250 ml of the solution.

 [Watch Video Solution](#)

## Long Answer Questions

1. Write the balanced ionic equation which represents the oxidation of iodine ( $I^-$ ) ion by per-manganate ion in basic medium to give iodine (I) and manganese dioxide ( $MnO_2$ ).

 [Watch Video Solution](#)

2. Write the balanced ionic equation for the oxidation of sulphite ions to sulphate ions in acid medium by permanganate ion.

 [Watch Video Solution](#)


3. Oxalic acid is oxidised by permanganate ion in acid medium of  $Mn^{2+}$  balance the reaction by ion-electron method.

 [Watch Video Solution](#)

4. Phosphorus when heated with NaOH solution gives Phosphine ( $PH_3$ ) and  $H_2PO_2^-$ . Give balanced equation.


 [Watch Video Solution](#)

5. Balance the following equation.



 [Watch Video Solution](#)

6. Balance the following equation by the oxidation number method.



 [Watch Video Solution](#)

### Important Question

1. How many number of moles of glucose are present in 540 gms of glucose ?

 [Watch Video Solution](#)

2. Calculate the weight of 0.1 mole of sodium carbonate.

 [Watch Video Solution](#)

3. How many molecules of glucose are present in 5.23g of glucose (Molecular weight of glucose 180 u).

 [Watch Video Solution](#)

4. The empirical formula of a compound is  $CH_2O$ . Its molecular weight is 90. Calculate the molecular formula of the compound.

 [Watch Video Solution](#)

5. Balance the following equation by the Oxidation number method

 [Watch Video Solution](#)



6. What volume of  $H_2$  at STP is required to reduce 0.795 g of CuO to give Cu and  $H_2O$ .

 [Watch Video Solution](#)

7. What is a redox concept? Give an example.

 [Watch Video Solution](#)

8. What do you mean by significant figures?

 [Watch Video Solution](#)

9. Chemical analysis of a carbon compound gave the following percentage composition by weight of the elements present, carbon = 10.06 %, hydrogen = 0.84 %, chlorine = 89.10 %. Calculate the empirical formula of the compound.

 [Watch Video Solution](#)



[Watch Video Solution](#)

10. A carbon compound on analysis gave the following percentage composition, carbon 14.5%, hydrogen 1.8%, chlorine 64.46%, oxygen 19.24%. Calculate the empirical formula of the compound.



[Watch Video Solution](#)

11. Calculate the empirical formula of a compound having percentage composition:

Potassium (K) = 26.57, Chromium (Cr) = 35.36, Oxygen (O) = 38.07.

(Given the Atomic weights of K, Cr and O are 39, 52 and 16 respectively)



[Watch Video Solution](#)

12. A carbon compound contains 12.8% Carbon, 2.1% Hydrogen, 85.1% Bromine. The molecular weight of the compound is 187.9. Calculate the molecular formula.

 [Watch Video Solution](#)

13. 0.188 g of an organic compound having an empirical formula  $CH_2Br$  displaced 24.2 cc of air at  $14^\circ C$  and 752 mm pressure. Calculate the molecular formula of the compound. (Aqueous tension at  $14^\circ C$  is 12 mm)

 [Watch Video Solution](#)

14. What volume of  $CO_2$  is obtained at STP by heating 4 g of  $CaCO_3$  ?

 [Watch Video Solution](#)

15. Calculate the volume of  $H_2$  liberated at  $27^\circ C$  and 760 mm of Hg pressure by action by 0.6 g magnesium with excess of dil HCl.

 [Watch Video Solution](#)

16. Define and explain molar mass.

 [Watch Video Solution](#)

17. What are disproportionate reactions? Give example.

 [Watch Video Solution](#)

18. What is comproportionation reactions? Give example.

 [Watch Video Solution](#)

19. How many significant figures are present in the following ?

i) 0.0025, ii) 208, iii) 5005, iv) 126,000 v) 500.0, vi) 2.0034

 [Watch Video Solution](#)


20. Round up the following upto three significant figures :

i) 34.216, ii) 10.4107, iii) 0.04597, iv) 2808

 [Watch Video Solution](#)

21. Chlorine is prepared in the laboratory by treating manganese dioxide

( $MnO_2$ ) with aqueous hydrochloric acid according to the reaction



How many grams of HCl react with 5.0g of manganese dioxide ?

 [Watch Video Solution](#)

22. To 50 ml. of 0.1 N  $Na_2CO_3$  solution 150 ml. of  $H_2O$  is added. Then calculate the normality of resultant solution.

 [Watch Video Solution](#)

23. Calculate the volume of 0.1 N  $H_2SO_4$  required to neutralise 200 ml. of 0.2 N NaOH solution.

It is an acid base neutralisation reaction.

Hence, at the neutralisation point.

Number of equivalents of acid = Number of equivalents of base.

 [Watch Video Solution](#)

24. Calculate normality of  $H_2SO_4$  solutions if 50 ml of it completely neutralise 250 ml. of 0.1 N  $Ba(OH)_2$  solutions.








 [Watch Video Solution](#)

25. Calculate the volume of 0.1M  $KMnO_4$  required to react with 100 ml. of 0.1 M  $H_2C_2O_4$ .

$2H_2O$  solution in the presence of  $H_2SO_4$ .






 [Watch Video Solution](#)

26. Assign oxidation number to the underlined elements in each of the following species.



Watch Video Solution

27. Justify that the following reactions are redox reactions.



Watch Video Solution

**28.** Explain thermal conductivity and coefficient of thermal conductivity. A copper bar of thermal conductivity  $401 \text{ W (mK)}$  has one end at  $104^\circ \text{C}$  and the other end at  $24^\circ \text{C}$ . The length of the bar is  $0.10 \text{ m}$  and the cross-sectional area is  $1.0 \times 10^{-6} \text{ m}^{-2}$ . What is the rate of heat conduction, along the bar?

 [Watch Video Solution](#)

**29.** How do you count the following observations?


a) Though alkaline potassium permanganate and acidic potassium permanganate both are used as oxidants, yet in the manufacture of benzoic acid from toluene we use alcoholic potassium permanganate as an oxidant. Why? Write balanced redox equation for the reaction.

b) When concentrated sulphuric acid is added to inorganic mixture containing chloride, we get colourless pungent smelling gas  $\text{HCl}$ , but if the mixture contains bromide then we get red vapour of bromine. Why?

 [Watch Video Solution](#)




**30.** Balance the following redox reaction in basic medium by ion-electron method :



 [Watch Video Solution](#)

**31.** What sorts of information can you draw from the following reaction ?



 [Watch Video Solution](#)

**32.** The  $\text{Mn}^{3+}$  ion is unstable in solution and undergoes disproportionation to give  $\text{Mn}^{2+}$ ,  $\text{MnO}_2$  and  $\text{H}^+$  ion. Write balanced ionic equation for the reaction.

 [Watch Video Solution](#)

33. In Ostwal's process for the manufacture of nitric acid the first step involves the oxidation of ammonia gas by oxygen gas to give nitric oxide gas and steam. What is the maximum weight of nitric oxide that can be obtained starting only with 10.00 g of ammonia and 20.00 g of oxygen.

 [Watch Video Solution](#)

34. Write the balanced ionic equation which represents the oxidation of iodine ( $I^-$ ) ion by per-manganate ion in basic medium to give iodine (I) and manganese dioxide ( $MnO_2$ ).

 [Watch Video Solution](#)

35. Write the balanced ionic equation for the oxidation of sulphite ions to sulphate ions in acid medium by permanganate ion.

 [Watch Video Solution](#)

36. Oxalic acid is oxidised by permanganate ion in acid medium of  $Mn^{2+}$

balance the reaction by ion-electron method.


 [Watch Video Solution](#)

37. Phosphorus when heated with NaOH solution gives Phosphine

( $PH_3$ ) and  $H_2PO_2^-$ . Give balanced equation.


 [Watch Video Solution](#)

38. Balance the following equation.



 [Watch Video Solution](#)

39. What are the oxidation numbers of the underlined elements in each of the following and how do you rationalise your results?



**Watch Video Solution**